



UNIVERSITY OF LEEDS

This is a repository copy of *KAI2 promotes Arabidopsis root hair elongation at low external phosphate by controlling local accumulation of AUX1 and PIN2*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/180315/>

Version: Supplemental Material

Article:

Villaécija-Aguilar, JA, Körösy, C, Maisch, L et al. (6 more authors) (2022) KAI2 promotes Arabidopsis root hair elongation at low external phosphate by controlling local accumulation of AUX1 and PIN2. *Current Biology*, 32 (1). pp. 228-236. ISSN 0960-9822

<https://doi.org/10.1016/j.cub.2021.10.044>

© 2021, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

