

This is a repository copy of *Mapping the protonation states of the histidine brace in an AA10 lytic polysaccharide monooxygenase using CW-EPR spectroscopy and DFT calculations*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/181159/>

Version: Accepted Version

Article:

Lindley, Peter J., Parkin, Alison orcid.org/0000-0003-4715-7200, Davies, Gideon J. orcid.org/0000-0002-7343-776X et al. (1 more author) (2022) Mapping the protonation states of the histidine brace in an AA10 lytic polysaccharide monooxygenase using CW-EPR spectroscopy and DFT calculations. FARADAY DISCUSSIONS. 336–348. ISSN 1364-5498

<https://doi.org/10.1039/D1FD00068C>

Reuse


Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Faraday Discussions

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

This article can be cited before page numbers have been issued, to do this please use: P. J. Lindley, A. Parkin, G. Davies and P. H. Walton, *Faraday Discuss.*, 2021, DOI: 10.1039/D1FD00068C.

ARTICLE

Mapping the protonation states of the histidine brace in an AA10 lytic polysaccharide monoxygenase using CW-EPR spectroscopy and DFT calculations

Peter J. Lindley^a, Alison Parkin^a, Gideon J. Davies^a and Paul H. Walton^{*a}

The active site of the polysaccharide-degrading lytic polysaccharide monoxygenase (LPMO) enzymes features a single copper ion coordinated by a histidine brace. The primary coordination sphere of the copper contains several ligating atoms which are bonded to ionisable protons (e.g. OH₂, NH₂), the pK_as of which are unknown. Using a combination of CW-EPR X-band spectroscopy over a range of pH and DFT calculations, we show that the active site of a chitin-active AA10 LPMO can exist in three different protonation states (pK_{a1} = 8.7, pK_{a2} ~ 11.5), representing the ionisation of the coordinating groups. The middle pH species (fully formed at pH ~10.5) is proposed to be Cu(II)(His)₂(OH)₂ (N₂O₂ coordination) with a decoordinated R-NH₃⁺ group at the amino terminus. This species also sees a rotation of the SOMO equatorial plane from the canonical histidine brace plane, whereby the nominal Cu d(x²-y²)-orbital has rotated some 45 degrees along the His-Cu(II)-His axis, driven by the elongation and decoordination of the amino group. The highest pH species (> 12) is proposed to exist as a Cu(II)-azanide, in which the NH₂ of the amino terminus has been deprotonated. The high pH means that this species is unlikely to be biologically relevant in the catalytic cycle of AA10 LPMOs.


Introduction

Lytic polysaccharide monoxygenases (LPMOs) are copper-containing enzymes which catalyse the oxidative cleavage of glycosidic bonds within polysaccharides.¹⁻³ The active site of LPMOs has been shown by crystallography to contain a single copper ion coordinated in a T-shaped configuration by a 'histidine brace', which consists of an N-terminal histidine chelating to the copper via its NH₂ and π-N imidazole atoms, along with the τ-N atom of a further histidine.⁴ In the copper(II) resting state, structures further show that one or two exogenous ligands, usually water or hydroxide, occupy the remaining copper coordination sites.

^aDepartment of Chemistry, University of York, Heslington, YORK, UK, YO10 5DD

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x





View Article Online
DOI: 10.1039/D1FD00068C

Figure 1. Active site of a chitin-active AA10 LPMO, depicting the histidine brace exogenous ligands (L) ligating to the Cu(II) ion, and conserved residues.

LPMOs are divided into Auxiliary Activity (AA) families in the Carbohydrate Active enZyme (CAZy) database (www.cazy.org) based on sequence similarity where so far eight distinct classes have been characterised (AA9-11, AA13-17).^{1,2,5-10} The number of exogenous ligands is partly dependent on the genomic class of the LPMO. For instance, structures of AA9-classified enzymes exhibit a single water/hydroxide to give a planar N₃O coordination geometry around the copper, whereas chitin-active AA10-classified enzymes appear to show the copper(II) in a distorted square-pyramidal N₃O₂ geometry, where two exogenous water/hydroxide ligands complete the coordination sphere as depicted in Figure 1. In both AA9 and AA10 cases, EPR spectroscopy establishes that the SOMO is dominated by the d(x²-y²) orbital, requiring the ligand field of the copper to have a large sigma-bonding component in the equatorial plane of the metal ion. In the case of AA9 LPMOs, the coordination geometry of the Cu is best described as axially elongated, whereas in the case of AA10 LPMOs EPR spectroscopy shows *ca* 3% of d(z²) mixing into the largely d(x²-y²) SOMO, commensurate with a distorted square-pyramidal geometry.^{11,12}

In chitin-active AA10 LPMOs the large number of ionisable protons of the ligands which coordinate to the copper ion in the histidine brace (NH₂, 2 × H₂O), opens up the possibility that the copper(II) centre exhibits a range of different protonation states, some of which may be relevant to the enzyme's catalytic cycle. Indeed, this reasoning has extended as far as suggesting that the amino terminus can deprotonate itself to form a copper(II)-azanide bond, the σ-donating power of which is expected to be high, possibly capable of stabilising a high valent copper species.¹³ Such a species may be a key intermediate in the catalytic cycle which is capable of abstracting a hydrogen atom from the polysaccharide substrate.^{2,14} As such, determining the different degrees of protonation of the active site is germane to any discussion about the catalytic mechanisms of LPMOs.

It is in this context that we aimed to establish the nature of the different protonation states of the active site of an AA10 LPMO from the bacterium *Bacillus licheniformis*, BI(AA10).¹⁵ To this end, we employed CW-EPR spectroscopy in conjunction with DFT calculations to show that the active site exists in three distinct protonation states across the pH range of 6.5 to 12.5. We further show that the highest pH species is most likely to be the copper(II)-azanide species proposed earlier,^{2,13} but that it only exists at pHs outside the usual biological range, and is not a biologically relevant species for the resting state of the enzyme.



Results and Discussion

Preparation and purity of *Bacillus licheniformis* AA10 LPMO.

Samples of this enzyme were prepared in *Escherichia coli* (BL21 competent cells) *via* periplasmic expression, previously reported by Courtade *et al.*¹⁶ Crude protein was obtained from the periplasm by osmotic shock and further purified using a 5 mL HiTrap® DEAE FF anion exchanger, eluting with a NaCl gradient from 0–500 mM over 90 column volumes. LPMO-containing fractions were collected and further purified on a Superdex S75 gel filtration column (GE Life Sciences).

The purity of *Bl*(AA10) was established by gel electrophoresis, which displayed a clear and single band at *ca.* 20 kDa (Supplementary Information). For the subsequent EPR experiments, it was important to determine that all copper in the sample was bound stoichiometrically to the active site of the enzyme. On this basis, a single equivalent of Cu was added to the enzyme from a solution of 10 mM CuSO₄ prior to size exclusion chromatography, from which any uncoordinated metal ions were separated from the purified enzyme.

CW-EPR X-band spectroscopy

A sample of *Bl*(AA10) (0.4 mM) was prepared in a mixed buffer system of MES, CHES, HEPES and CAPS, each at 5 mM. These buffers were chosen for their known poor-coordination properties toward metal ions and, when used synchronously, buffer a wide range of pHs. The pH of the solution was adjusted up and down by the addition of dilute solutions of NaOH and H₂SO₄, respectively. The pH of the resulting solution at room temperature was determined using a calibrated Fisherbrand™ accumet™ AE150 Benchtop pH Meter, from which 12 samples from pH 6.5 to 12.5 were prepared in 0.5 pH unit steps. To verify that the pH extremes did not denature the protein, EPR spectra were collected on samples starting at pH 6.5 and then raising to pH 12.5, and then recollected on the same sample at a lower pH, indeed restoring the previous species.

At each pH, the sample was rapidly (< 2 s) cooled¹⁷ to 150 K and its CW-EPR X band spectrum collected[‡]. The trend of EPR spectra shows a qualitative change across the pH range, in which it is evident that all species have a spectral envelope which is consistent with a predominantly $d(x^2-y^2)$ SOMO, showing that the Cu(II) ion at all pH has a ligand field dominated by sigma donation in its equatorial plane (Figure 2). Spin quantification by double integration of the spectra also shows that there are no redox processes that occur at the active site across the pH range (Supplementary Information).

Analysis of EPR spectra and distribution of pH dependent species

Between pHs 6.5 and 8.0, a single species is seen to dominate the spectrum, as evidenced by a single set of Cu-hyperfine split peaks at low field ($g_3 = 2.260$, $A_3 = 445$ MHz). Increasing the pH leads to the formation of a new set of peaks in the low field region ($g_3 = 2.233$, $A_3 = 554$ MHz) along with distinctive superhyperfine coupling around $g = 2.07$, commensurate with the formation of a second distinct species, the concentration of which is maximised at pH 10.5. A further increase in pH to 12.5 sees the formation of a further set of peaks in the low field region ($g_3 = 2.180$, $A_3 = 614$ MHz), consistent with the formation of a third distinct species. This final species is not fully formed at pH 12.5 but is dominant at that pH. The total number of species across the pH range was assessed using Singular Value Decomposition analysis of the collected spectra (Supplementary Information). From this analysis, three principal components emerged, corroborating the qualitative assessment of the pH dependence of the EPR spectra.



View Article Online
DOI: 10.1039/D1FD00068C





Figure 2. Frozen solution X-band EPR spectra (150 K) of *B/(AA10)* (0.4 mM) in a mixed buffer of MES, CHES, HEPES and CAPS (5 mM each) between pH 6.5 and 12.5.

On the basis of three components (species 1, 2 and 3), we sought to deconvolute the series of spectra. Given the clear separation between the different species across the pH range, it was assumed that the spectrum at pH 6.5 represented 100% of a single complex. Indeed, it was possible to simulate (with EasySpin) the lowest pH species (pH 6.5) with a single set of spin-Hamiltonian parameters (Figure 3a, Table 1). The same was also true of the spectra collected at pH 10.5 and 12.5 (Figure 3b and 3c, respectively, Table 1). In performing the simulation of species 3, we noted the appearance of a significant 'overshoot' feature which appears at high field in the species at pH 12.5. Such a feature has in the past been confused with an anomalously low g_1 value.¹⁸





View Article Online
DOI: 10.1039/D1FD00068C

Figure 3. X-band EPR a) low pH species 1, with simulation (red), b) intermediate pH species 2, with simulation (red), c) high pH species 3, with simulation (red).

Once spin Hamiltonian parameters of each species were obtained, it was possible to simulate the mixed species spectra (pHs 7 to 9.5 and >10.5) to determine species distributions at each pH, from which pK_a s of species 1 (low pH) and species 2 (middle pH), and the stoichiometry of protons for each ionisation were determined using standard methods (Supplementary Information). The deprotonation of species 1 to 2 occurs with a pK_a of 8.7 and is a single proton step. The deprotonation of species 2 to 3 occurs with a pK_a of ~ 11.5 and is also a single proton step.

Ligand field analyses of spin Hamiltonian parameters

The collected spin Hamiltonian parameters of the three species are shown in Table 1. In all cases the g values follow the pattern of $g_1 < g_2 < g_3$, confirming a SOMO with high $d(x^2-y^2)$ character. Some $d(z^2)$ mixing is evident for all species from the differences in the values of g_1 and g_2 . The degree of mixing, b^2 , can be calculated from the standard perturbative EPR formulae as described by Hitchman et al,¹⁹ which give the molecular orbital coefficients, a and b , for the SOMO defined in equation 1 (Table 1).

$$\psi(\text{SOMO}) = a\psi(x^2-y^2) + b\psi(z^2) \quad (1)$$

This mixing is low for species 2 and 3, revealing that the ligand field around the Cu in each of these is close to axial four-fold symmetry (note that this does not necessarily translate into four-fold axial symmetry of the identity of the coordinating atoms, as different coordinating atoms may have similar ligand field parameters). For species 1, more mixing is evident ($b^2 = 3.6\%$), which is in-line with previous EPR studies on AA10 LPMOs¹¹ and the subsequent structures of these enzymes which exhibit a distorted square pyramidal geometry around the copper ion (Figure 1), where the exogenous ligands are two water/hydroxide molecules, one of which forms a Cu-O bond of ca 1.9 Å, and the other a more distant bond of 2.2 Å.²⁰



Table 1. Spin Hamiltonian parameters of pH dependent species. It is assumed that the A_3 value is negative in sign, although this is not determined experimentally.

DOI: 10.1039/D1FD00068C

Species	g values			Hyperfine coupling constants /MHz			$b^2/\%$
	g_1	g_2	g_3	$ A_1 $	$ A_2 $	A_3	
1	2.030	2.125	2.260	60	90	-445	3.6
2	2.044	2.069	2.233	60	76	-554	0.8
3	2.038	2.062	2.180	70	90	-614	0.9

Taking the value of g_3 as a measure of the relative ligand field strengths in the equatorial plane of the copper, it is evident that this ligand field increases significantly moving from species 1 to 2 to 3, and—moreover, from the values of $d(z^2)$ mixing—that the ligand field in species 2 and 3 is relatively evenly distributed amongst the ligating atoms of the equatorial plane, suggestive of a near four-fold symmetric planar arrangement of the equatorial ligand field. Indeed, in corroboration of such a ligand field, the increase in A_3 value from species 1 to 2 is also known to be associated with a change in the Fermi coupling as the coordination number of the copper drops from five to four.¹⁵ Without access to d-d electronic transition energies, it is not possible to be more precise about the nature of the ligand field; nevertheless the trend in increasing ligand field around the equatorial plane is evident from the relative spin-Hamiltonian parameters.

Further information about the ligand field in species 2 can be obtained from the clearly delineated Cu-N superhyperfine coupling (Figure 3b). This coupling could be accurately simulated with the inclusion of two nitrogen atoms (^{14}N , $I = 1$) in the spin Hamiltonian, in which each N atom couples to the copper with a principal coupling value of 40 MHz. This value is similar to those already known for the Cu-His coupling constants seen in other LPMOs.^{17, 21} The simulation of species 2 could be adequately performed with the inclusion of two N nuclei, incommensurate with the expected N_3 coordination of the histidine brace. Additionally, the g_3 and A_3 values of species 2, place it squarely within those expected for a CuN_2O_2 coordination geometry on a canonical Peisach-Blumberg plot.²²

In assigning a putative structure to species 2, it is informative to consider previous EPR studies on the substrate-bound state of chitin-active AA10 enzymes. These species have been well characterised as having a planar CuN_3O coordination geometry with g_z value of 2.205 and an A_z value of ~ 620 MHz.^{15, 23} On this basis, a similar axial CuN_3O (sq pl) ligand field is unlikely for species 2, given the discrepancies in g_z and A_z values. This difference in spin Hamiltonian parameters necessitates an alternative ligand field assignment for species 2. Thus, taking together the Cu spin Hamiltonian values and superhyperfine couplings a more likely scenario is one in which the NH_2 group of the histidine brace is decoordinated from the copper (see DFT calculations below) and the coordination sphere around the Cu is made up of two histidines and two water/hydroxide ligands (see below for further discussion).

The spin Hamiltonian parameters for species 3 are distinct from other known LPMOs, particularly evident in the value of g_3 (2.18) which is lower than all known values, including the situation in which a chloride ion has entered the coordination sphere of the copper ($g_3 = 2.23$). Given the low value, we are directed towards the proposal that the ligand field around the copper in



species 3 is one that is highly σ -donating causing a large separation in $d(x^2-y^2)$ and $d(xy)$ orbital energies. Such a strong σ donation might be expected of a Cu-azanide interaction, which would arise from the recoordination of the NH_3^+ amino terminus in species 2 which had been doubly deprotonated to give -NH^- ligating group. Commensurate with this proposal is the simulation of the superhyperfine coupling in species 3 which could only be satisfactorily achieved with the inclusion of three nitrogen atoms in the spin Hamiltonian with principal coupling constants of 35, 35 and 33 MHz (see DFT calculations below).

From the overall ligand field analysis, a hypothesis emerges for the nature of species 1, 2 and 3 and the pH-dependent ionisation processes between them. The hypothesis is depicted in Figure 4, in which it is shown that species 2 forms via deprotonation of a coordinating water molecule followed by the decoordination and subsequent protonation of the -NH_2 group, driven by an internal proton transfer process from one of two coordinating water molecules in species 1. The formation of species 3 from species 2, on the other hand, sees the reverse proton transfers from the decoordinated -NH_3^+ to a coordinated hydroxide with the loss of the resulting water molecule, and the subsequent deprotonation of an N-H proton from the -NH_2 and recoordination to the metal as an azanide. In support of such an assignment, the formation of species 3 was accompanied by an unusual colour change from colourless to pale pink (Supplementary Information) due to the formation of a semi-intense absorption band ($\epsilon \approx 2500 \text{ M}^{-1} \text{ cm}^{-1}$) centred at 520 nm. The intensity of this band is in keeping with that of a charge transfer transition – likely arising from a ligand to metal charge transfer (LMCT) from the proposed azanide (-NH^-) group to the copper (see TD-DFT calculations, Supplementary Information). Thus, the proposed ligand field changes depicted in Figure 4 are revealed by the differences in spin Hamiltonian parameters across the species, in which g_z decreases and A_z increases (going from species 1 to 3) commensurate with an increasing equatorial ligand field.




Figure 4. Hypothetical protonation states and coordination geometries of species 1, 2 and 3.

DFT calculations

Building on the hypothesis depicted in Figure 4, Density Functional Theory geometry optimisations (uBP86) and EPR calculations (uB3LYP with 38% HF exchange) were performed on the proposed species 1, 2 and 3.

In the absence of a high-resolution crystal structure for the *Bacillus licheniformis* AA10, crystallographic coordinates of the spectroscopically similar enzyme, *Bacillus amyloliquefaciens* AA10 (PDB 5IJU)²⁰ were used as the basis for these calculations, in keeping with the experiments of Courtade *et al.*¹⁵ Importantly in this regard, the structure of BaAA10 is one of the few LPMO structures in which the copper ion has not been photoreduced from Cu(II) to Cu(I) during the X-ray experiment.²⁴ Accordingly, key residues from the crystallographic coordinates were used for the basis of calculations for species 1; retaining the primary



coordination sphere and supporting residues (full information and coordinates in Supplementary Information). Following standard practice, the spatial coordinates of certain atoms at the periphery of the model were kept fixed throughout the calculation to prevent unfettered movement of residues. These atoms are depicted in Figure S7 (Supplementary Information), and were selected on the basis of earlier DFT studies on a closely related LPMO.¹⁵

For species 1, the geometry was initially modelled with the inclusion of two ligating water molecules with bond distances based on the positioning of the oxygen atoms in the crystal structure. Notably in this regard, the water situated beneath the histidine brace plane is expected to participate in strong hydrogen bonding with a nearby aspartate residue (glutamate in *BaAA10*), the potential of which can affect the protonation state and ligating properties of the water molecule. Indeed, upon optimisation, despite the fact that Glu and Asp residues are expected to be deprotonated across the studied pH range owing to the low pK_a s of their carboxylate groups ($pK_a = 2-4.5$),²⁵ one of the protons bound to the water molecule migrated to the neighbouring carboxylate group (figure 5) to give a mixed hydroxide/water coordination at the Cu. Confidence in this structural assignment comes from the resulting geometry of the optimised structure which is a close match to the crystal structure coordinates. Thus, for species 1, the coordination sphere of the Cu is best described as a distorted square-pyramid, where water and hydroxide act as exogenous ligands.

While a good match was found between the optimised and experimental structures, the match between calculated and experimental EPR parameters (Table 2) was only fair (e.g. calculated $g_3 = 2.216$, experimental $g_3 = 2.26$). Moreover, the calculated Cu hyperfine values, A_1 and A_2 have relative magnitudes opposite to that observed experimentally. Notwithstanding the known difficulties in DFT-calculated EPR parameters matching with those from experiment, the relatively large differences seen between experimental and calculated EPR parameters for species 1 are a recurring theme in DFT calculations on AA10 LPMO active sites in their Cu(II) resting states, even with careful selection of basis sets and functionals.²⁶ As such, while the optimised structure of 1 matches well that of experiment and is likely a good representation of the actual structure, caution must be applied in interpreting the calculated EPR data for species 1, where the reasons for the discrepancy between theory and experiment remain unclear. This situation is in contrast to the calculations of EPR parameters for species 2 and 3 which give a somewhat closer match to experimental data (see below).




Figure 5. DFT model of species 1 (truncated for clarity) showing distorted square-pyramidal geometry and hydroxide/water coordination of exogenous ligands.



Table 2. Calculated spin Hamiltonian parameters of pH dependent species. The x axis of the calculated g tensor lies along the His-Cu-His direction.

View Article Online
DOI: 10.1039/D1FD00068C

Species	g values			Hyperfine coupling constants /MHz		
	g_x	g_y	g_z	$ A_x $	$ A_y $	A_z
1	2.049	2.081	2.216	110	29	-598
2	2.070	2.079	2.244	94	3	-582
3	2.037	2.070	2.188	15	140	-496

Following the ligand field analysis from above, species 2 was initially modelled by removing the proton shared between the water and glutamate, and by transferring the proton attached to the second water ligand to the $-NH_2$ group (representing an overall removal of a single proton from species 1, as per the experimental findings). Upon optimisation, the angle between the hydroxide ligands $\angle O-Cu-O$ increased from *ca.* 90° to 157° . The divergence of the hydroxide ligands was also accompanied by the expected decoordination of the now protonated $-NH_3^+$ group from the metal and a rotation of the $d(x^2-y^2)$ SOMO by *ca.* 45° (Figure 6), such that the equatorial plane is now defined by the two hydroxides and the nitrogen atoms of two histidine ligands, thus completing the CuN_2O_2 coordination geometry expected from the ligand field analysis. Calculated EPR parameters (Table 2) for species 2 reflect the experimental data in that g_x and g_y are closer in values, and that the value of g_z is *ca.* 2.24.




Figure 6. Hypothetical $d(x^2-y^2)$ dominated SOMO of species 2 showing the orbital plane has rotated. (Colours denote orbital phase).

Species 3 was modelled as a Cu(II)-azanide species formed by expulsion of water from species 2, followed by the further removal of an $-NH_2$ proton resulting in a short Cu-NH bond of 1.94 Å (full computational details can be found in the Supplementary Information). The optimised geometry is best described as CuN_3O planar, albeit distorted ($\angle HN-Cu-O \sim 160^\circ$). This distortion away from planar has been previously reported for four-coordinate LPMO structures when modelled by DFT.¹⁵ Calculated EPR parameters for species 3 match well with those of experiment, most notably in the $g_{3,z}$ value of 2.18. Also, from a spectroscopic perspective, Species 3 exhibits a semi-intense UV/visible absorption band centred at 520 nm ($\epsilon \approx 2500 \text{ M}^{-1} \text{ cm}^{-1}$). This band is proposed to arise from a LMCT transition from the proposed azanide group to the Cu, corroborated with TD-DFT calculations (Supplementary Information). Both d-d and LMCT transition energies



appear to be significantly blue-shifted with respect to the experimental transitions. Exact transition energies are known to carry significant error when calculated in this way for open shell systems. Nevertheless, these calculations reveal that an intense absorption band is expected for the Cu(II)-azanide complex proposed as species 3 and its energy is predicted to be positioned between the π - π^* and the d-d transitions. The three optimised structures for species 1, 2 and 3 are shown in Figure 7.




Figure 7. DFT optimised geometries of species 1, 2 and 3. Models are truncated, and certain atoms are hidden for clarity.

Conclusions

Herein, we establish three discrete protonation states for an AA10 LPMO within the pH range of 6.5 and 12.5, two of which are biologically relevant and may be important in the context of the LPMO mechanism. The first species exists at acidic-neutral pH and is believed to be consistent with a 5-coordinate (N_3O_2); in keeping with the current understanding of the chitin-active AA10 coordination sphere. A second species is formed by deprotonating the ligating water molecule (pK_a 8.7); yielding a near-axial EPR spectral envelope ($g_1=2.044$, $g_2=2.069$ and $g_3=2.233$). We hypothesise that this species exists as a N_2O_2 , 4-coordinate ligand field with two hydroxide ligands and an elongated/decoordinated $-R-NH_3^+$ following proton transfer from the coordinating hydroxide to the amino terminus. This gives rise to a rotated SOMO no longer defined by the chelating histidine brace. The third species formed at high pH ($pK_a \sim 11.5$) is proposed to exist as a 4-coordinate, N_3O complex, whereby species 2 has lost a water molecule (from the coordination of an OH^- and NH_3^+) and the $-NH_2$ group has been deprotonated to form a Cu(II)-NH azanide complex. The high pH at which this species becomes accessible is outside of that which is biologically relevant and as such is unlikely to be pertinent in catalysis.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements



We acknowledge the University of York for support of PJL's PhD studentship. We thank Z. Forsberg, G. Courtade, V. G. H. Eijsink and F. L. Aachmann for providing the clone/cell colonies and purification protocol for the *BIAA10* enzyme. GJD thanks the Royal Society for the Ken Murray Research Professorship.

DOI: 10.1039/D1FD00068C

Notes and references

‡Continuous wave (cw) X-band EPR spectra were obtained using a 100 μ L frozen solution of *BIAA10* at a concentration of 0.4 mM. Data collection was carried out on a Bruker micro EMX spectrometer using a microwave frequency of *ca.* 9.3 GHz, modulation amplitude of 4 G, modulation frequency of 100 kHz and microwave power of 10.02 mW. Spectra were intensity averaged over three scans and simulated using the EasySpin 5.2.28²⁷ open-source toolbox implemented by MATLAB R2020a software on a desktop PC. Raw EPR data are available at 10.15124/7a485d50-ea4e-4cd1-8fc2-650057c076af

1. G. Vaaje-Kolstad, B. Westereng, S. J. Horn, Z. Liu, H. Zhai, M. Sørli and V. G. H. Eijsink, *Science*, 2010, **330**, 219-222.
2. R. J. Quinlan, M. D. Sweeney, L. Lo Leggio, H. Otten, J.-C. N. Poulsen, K. S. Johansen, K. B. R. M. Krogh, C. I. Jørgensen, M. Tovborg, A. Anthonen, T. Tryfona, C. P. Walter, P. Dupree, F. Xu, G. J. Davies and P. H. Walton, *Proc. Nat. Acad. Sci. USA*, 2011, **108**, 15079-15084.
3. C. M. Phillips, W. T. Beeson, J. H. Cate and M. A. Marletta, *ACS Chem. Biol.*, 2011, **6**, 1399-1406.
4. L. Ciano, G. J. Davies, W. B. Tolman and P. H. Walton, *Nat. Catal.*, 2018, **1**, 571-577.
5. G. R. Hemsworth, B. Henrissat, G. J. Davies and P. H. Walton, *Nat. Chem. Biol.*, 2014, **10**, 122-126.
6. V. V. Vu, W. T. Beeson, E. A. Span, E. R. Farquhar and M. A. Marletta, *Proc. Nat. Acad. Sci. USA*, 2014, **111**, 13822-13827.
7. M. Couturier, S. Ladevèze, G. Sulzenbacher, L. Ciano, M. Fanuel, C. Moreau, A. Villares, B. Cathala, F. Chaspoul, K. E. Frandsen, A. Labourel, I. Herpoël-Gimbert, S. Grisel, M. Haon, N. Lenfant, H. Rogniaux, D. Ropartz, G. J. Davies, M.-N. Rosso, P. H. Walton, B. Henrissat and J.-G. Berrin, *Nat. Chem. Biol.*, 2018, **14**, 306.
8. F. Sabbadin, G. R. Hemsworth, L. Ciano, B. Henrissat, P. Dupree, T. Tryfona, R. D. S. Marques, S. T. Sweeney, K. Besser, L. Elias, G. Pesante, Y. Li, A. A. Dowle, R. Bates, L. D. Gomez, R. Simister, G. J. Davies, P. H. Walton, N. C. Bruce and S. J. McQueen-Mason, *Nat. Commun.*, 2018, **9**, 756.
9. C. Filiatrault-Chastel, D. Navarro, M. Haon, S. Grisel, I. Herpoël-Gimbert, D. Chevret, M. Fanuel, B. Henrissat, S. Heiss-Blanquet, A. Margeot and J.-G. Berrin, *Biotechnol. Biofuels*, 2019, **12**, 55.
10. F. Sabbadin, S. Urresti, B. Henrissat, A. O. Avrova, L. R. J. Welsh, P. Lindley, M. Csukai, J. N. Squires, P. H. Walton, G. J. Davies, N. C. Bruce, S. C. Whisson and S. J. McQueen-Mason, *Science*, 2021, **373**, 774-779.
11. G. R. Hemsworth, E. J. Taylor, R. Q. Kim, R. C. Gregory, S. J. Lewis, J. P. Turkenburg, A. J. Parkin, G. J. Davies and P. H. Walton, *J. Am. Chem. Soc.*, 2013, **135**, 6069-6077.
12. F. L. Aachmann, M. Sørli, G. Skjåk-Bræk, V. G. H. Eijsink and G. Vaaje-Kolstad, *Proc. Nat. Acad. Sci. USA*, 2012, **106**, 18779-18784.
13. N. Gagnon and W. B. Tolman, *Acc. Chem. Res.*, 2015, **48**, 2126-2131.
14. J.-P. Bacik, S. Mekasha, Z. Forsberg, A. Y. Kovalevsky, G. Vaaje-Kolstad, V. G. H. Eijsink, J. C. Nix, L. Coates, M. J. Cuneo, C. J. Unkefer and J. C. H. Chen, *Biochemistry*, 2017, **56**, 2529-2532.
15. G. Courtade, L. Ciano, A. Paradisi, P. J. Lindley, Z. Forsberg, M. Sørli, R. Wimmer, G. J. Davies, V. G. H. Eijsink, P. H. Walton and F. L. Aachmann, *Proc. Nat. Acad. Sci. USA*, 2020, **117**, 19178-19189.
16. G. Courtade, S. B. Le, G. I. Sætrum, T. Brautaset and F. L. Aachmann, *Carbohydr. Res.*, 2017, **448**, 212-219.



17. G. R. Hemsworth, L. Ciano, G. J. Davies and P. H. Walton, in *Methods Enzymol.*, ed. F. Armstrong, Academic Press, 2018, vol. 613, pp. 63-90.
18. B. Bissaro, I. Isaksen, G. Vaaje-Kolstad, V. G. H. Eijsink and Å. K. Røhr, *Biochemistry*, 2018, **57**, 1893-1906. View Article Online
DOI: 10.1039/D1FD00068C
19. M. A. Hitchman and R. L. Belford, *Inorg. Chem.*, 1969, **8**, 958-965.
20. R. C. Gregory, G. R. Hemsworth, J. P. Turkenburg, S. J. Hart, P. H. Walton and G. J. Davies, *Dalton Transactions*, 2016, **45**, 16904-16912.
21. K. E. H. Frandsen, T. J. Simmons, P. Dupree, J.-C. N. Poulsen, G. R. Hemsworth, L. Ciano, E. M. Johnston, M. Tovborg, K. S. Johansen, P. von Freiesleben, L. Marmuse, S. Fort, S. Cottaz, H. Driguez, B. Henrissat, N. Lenfant, F. Tuna, A. Baldansuren, G. J. Davies, L. Lo Leggio and P. H. Walton, *Nat. Chem. Biol.*, 2016, **12**, 298.
22. J. Peisach and W. E. Blumberg, *Arch. Biochem. Biophys.*, 1974, **165**, 691-708.
23. B. Bissaro, B. Streit, I. Isaksen, V. G. H. Eijsink, G. T. Beckham, J. L. DuBois and Å. K. Røhr, *Proc. Nat. Acad. Sci. USA*, 2020, **117**, 1504.
24. A. Paradisi, M. J. Steward, P. Lindley, G. J. Davies and P. H. Walton, in *Reference Module in Chemistry, Molecular Sciences and Chemical Engineering*, Elsevier, 2020, DOI: <https://doi.org/10.1016/B978-0-12-409547-2.14928-5>.
25. M. J. Harms, C. A. Castañeda, J. L. Schlessman, G. R. Sue, D. G. Isom, B. R. Cannon and B. García-Moreno E, *J. Mol. Biol.*, 2009, **389**, 34-47.
26. Y. A. Theibich, S. P. A. Sauer, L. L. Leggio and E. D. Hedegård, *Comput. Struct. Biotechnol. J.*, 2021, **19**, 555-567.
27. S. Stoll and A. Schweiger, *Journal of Magnetic Resonance*, 2006, **178**, 42-55.

