

This is a repository copy of Should we use weight-based vitamin D treatment in children?.

White Rose Research Online URL for this paper: https://eprints.whiterose.ac.uk/182717/

Version: Supplemental Material

Article:

Arshad, F., Arundel, P., Bishop, N. orcid.org/0000-0001-7263-8546 et al. (1 more author) (2022) Should we use weight-based vitamin D treatment in children? Archives of Disease in Childhood, 107 (6). pp. 620-621. ISSN 0003-9888

https://doi.org/10.1136/archdischild-2021-322852

This article has been accepted for publication in Archives of Disease in Childhood, 2021 following peer review, and the Version of Record can be accessed online at http://dx.doi.org/10.1136/archdischild-2021-322852. © Authors (or their employer(s)) 2021. Reuse of this manuscript version (excluding any databases, tables, diagrams, photographs and other images or illustrative material included where a another copyright owner is identified) is permitted strictly pursuant to the terms of the Creative Commons Attribution-Non Commercial 4.0 International (CC-BY-NC 4.0) https://creativecommons.org/licenses/by-nc/4.0/

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.






Figure 1. Plot of decimal age in years against increase in 25-OH vitamin D concentration following treatment of vitamin D deficiency on primary y axis (left) and total treatment dose on secondary y axis (right).