



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/182723/>

Version: Accepted Version

---

**Article:**

Casanova, N, Beaulieu, K, Oustric, P et al. (2023) Increases in physical activity are associated with a faster rate of weight loss during dietary energy restriction in women with overweight and obesity. *British Journal of Nutrition*, 129 (8). pp. 1451-1461. ISSN: 0007-1145

<https://doi.org/10.1017/S000711452200023X>

---

© The Authors 2022. This article has been published in a revised form in *British Journal of Nutrition* [<https://doi.org/10.1017/S000711452200023X>]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

1 **Increases in physical activity are associated with a faster rate of weight loss during dietary**  
2 **energy restriction in women with overweight and obesity**

3

4 Nuno Casanova<sup>1,2</sup>, Kristine Beaulieu<sup>3</sup>, Pauline Oustric<sup>3</sup>, Dominic O'Connor<sup>3</sup>, Catherine  
5 Gibbons<sup>3</sup>, John Blundell<sup>3</sup>, Graham Finlayson<sup>3</sup> and Mark Hopkins<sup>1</sup>

6

7 <sup>1</sup> School of Food Science and Nutrition, Faculty of Environment, University of Leeds, Leeds,  
8 LS2 9JT, UK

9 <sup>2</sup> KinesioLab, Research Unit in Human Movement Analysis, Piaget Institute, Av. Jorge Peixinho  
10 30 Quinta da Arreinel, 2805-059 Almada, Portugal

11 <sup>3</sup> Appetite Control and Energy Balance Research Group, School of Psychology, Faculty of  
12 Medicine and Health, University of Leeds, Leeds, LS2 9JT, UK

13

14

15 **Corresponding author:**

16 Mark Hopkins

17 School of Food Science and Nutrition,

18 Faculty of Environment,

19 University of Leeds,

20 Leeds,

21 LS2 9JT

22 United Kingdom.

23 Email: m.hopkins@leeds.ac.uk

24

25 **Short title:** Increases in physical activity accelerate weight loss

26

27 **Abstract (250 words)**

28 This secondary analysis examined the influence of changes in physical activity (PA), sedentary  
29 time and energy expenditure (EE) during dietary energy restriction on the rate of weight loss (WL)  
30 and 1-year follow-up weight change in women with overweight / obesity.

31 Measurements of body weight and composition (air-displacement plethysmography), resting  
32 metabolic rate (indirect calorimetry), total daily (TDEE) and activity EE (AEE), minutes of PA  
33 and sedentary time (PA monitor) were taken at baseline, after 2 weeks, after  $\geq 5\%$  WL or 12 weeks  
34 of continuous (25% daily energy deficit) or intermittent (75% daily energy deficit alternated with  
35 *ad libitum* day) energy restriction, and at 1-year post-WL. The rate of WL was calculated as total  
36 %WL / number of dieting weeks. Data from both groups were combined for analyses.

37 Thirty-seven participants (age=35 $\pm$ 10y; BMI=29.1 $\pm$ 2.3kg/m<sup>2</sup>) completed the intervention (WL=-  
38 5.9 $\pm$ 1.6%) and 18 returned at 1-year post-WL (weight change=+4.5 $\pm$ 5.2%). Changes in sedentary  
39 time at 2 weeks were associated with the rate of WL during energy restriction (r=-0.38; p=0.03).  
40 Changes in total (r=0.54; p<0.01), light (r=0.43; p=0.01) and moderate-to-vigorous PA (r=0.55;  
41 p<0.01), sedentary time (r=-0.52; p<0.01), steps per day (r=0.39; p=0.02), TDEE (r=0.46; p<0.01)  
42 and AEE (r=0.51; p<0.01) during energy restriction were associated with the rate of WL. Changes  
43 in total (r=-0.50; p=0.04) and moderate-to-vigorous PA (r=-0.61; p=0.01) between post-WL and  
44 follow-up were associated with 1-year weight change (r=-0.51; p=0.04).

45 These findings highlight that PA and sedentary time could act as modifiable behavioural targets  
46 to promote better weight outcomes during dietary energy restriction and/or weight maintenance.

47

48 **Keywords:** Physical activity; sedentary time; energy expenditure; weight loss; weight regain

## 49 **1. Introduction**

50 It has been reported that up to 80% of individuals who achieve clinically significant weight loss  
51 (WL) fail to sustain this WL after 1 year or more [1]. While researchers have attempted to identify  
52 predictors of WL and WL maintenance, inconsistent findings are reported and potential predictors  
53 of WL often have limited explanatory value [2, 3]. Identification of predictive factors is important  
54 as it would allow proactive changes to be made during a WL intervention, potentially improving  
55 longer-term weight management success. Two factors that have been previously highlighted as  
56 predictors of WL are early changes in body weight (2 to 6 weeks) [4, 5] and the amount of physical  
57 activity (PA) performed during periods of WL [6].

58 Previous research has reported that PA may decline during dietary-induced WL [7-9], with a  
59 systematic review by Silva et al. reporting decreases in PA and/or non-exercise activity  
60 thermogenesis in 50% (7 out of 14 studies) of diet-only interventions [7]. However, several  
61 studies have reported no changes in PA during WL [10, 11]. For instance, after 12 weeks of  
62 continuous or intermittent energy restriction to ~12.5% WL, Coutinho et al. did not observe any  
63 within or between group differences in the number of steps per day [10]. Inter-individual  
64 variability in WL and body composition outcomes is commonly observed in studies of dietary  
65 energy restriction [12, 13], but whether individual differences in changes in PA and sedentary  
66 behaviours influence WL and WL maintenance success remains unclear.

67 While the role of PA and exercise in weight management has been questioned [14], interventions  
68 combining both dietary-energy restriction and changes in PA usually promote a greater WL which  
69 is better sustained over time [15]. For instance, a systematic-review observed that combining  
70 dietary-energy restriction and exercise lead to a 20% greater total WL in comparison to dietary  
71 modifications alone [16]. Furthermore, during 6 months of a lifestyle WL intervention,  
72 participants on the higher PA group had an increase of 47min/day (and a reduction in sedentary  
73 time of 52min/day), achieving a greater total WL [17]. However, findings regarding the role of  
74 PA or exercise in weight management are not always consistent, with a recent systematic review  
75 reported no significant effects of exercise on WL maintenance [18].

76 Of note, few studies have objectively measured PA during dietary-induced WL, and in particular,  
77 during the early stages of WL, to examine whether changes in free-living PA influences the  
78 dynamics of WL e.g., rate, extent or composition of WL. Examining the early- and longer-term  
79 changes in PA at the individual level during dietary-induced energy restriction would allow for a  
80 better understanding of the role of PA in facilitating or resisting early and/or sustained WL, and  
81 would provide a framework in which effective behaviour change interventions could be designed  
82 to improve weight management success rates [19].

83 Therefore, the aim of this secondary analysis was to examine the influence of early (baseline to  
84 week 2) and post-intervention changes in objectively measured PA and sedentary time during  
85 dietary energy restriction on 1) the rate of WL and 2) 1-year follow-up weight change in women  
86 with overweight and obesity.

87

## 88 **2. Material and Methods**

89 Healthy women with overweight and obesity were recruited from the University of Leeds and the  
90 surrounding area via posters and email lists to take part in a study examining ‘the effects of a  
91 personalised weight loss meal plan on body composition and metabolism’ (NCT03447600). In  
92 this study, participants were randomised to either continuous (CER; daily 25% energy restriction  
93 – all foods were provided) or intermittent (IER; 75% energy restriction days alternated with *ad*  
94 *libitum* eating days – food was only provided on ‘fast’ days) energy restriction until  $\geq 5\%$  WL or  
95 12 weeks (even if WL target was not achieved). The present analyses represent exploratory  
96 analysis of secondary outcomes from this study, and previous findings from the main dietary  
97 energy restriction study have been reported elsewhere [20, 21]. Specific details of the dietary  
98 intervention during the WL phase are provided elsewhere [22], and for the purposes of this paper,  
99 findings from both dietary groups were combined as no group differences existed in the main  
100 outcomes reported here (see section 2.4). No instructions were given to nor contact kept with  
101 participants after the WL phase and thus they were not required to maintain the same dietary  
102 pattern. Participants that completed the WL phase ( $\geq 5\%$  WL or within 12 weeks) were invited for  
103 a 1-year follow-up 4 weeks before the measurements to avoid influencing their behaviours

104 throughout the 12 months. Therefore, while this was not a weight maintenance intervention, the  
105 aim of the follow-up measurement was to attempt to highlight factors (during and after dietary-  
106 induced energy restriction) associated with post-WL weight change as these could have important  
107 implications regarding weight management interventions.

108 Participants were excluded if they had health problems that could affect study outcomes; history  
109 of eating disorders; taking medication, supplements or treatment known to affect appetite/weight  
110 within the past month and/or during the study; pregnant, planning to become pregnant or  
111 breastfeeding; known food allergies/intolerances; smokers or had ceased smoking in the past 6  
112 months; lost significant amount of weight in the previous 6 months ( $\pm 4$ kg); exercised  $>3$  days per  
113 week, significantly changed their PA patterns in the past 6 months or intended to change them  
114 during the study; worked in appetite/feeding related areas; or were shift workers. Participants  
115 provided written informed consent before taking part and were remunerated £100 upon  
116 completion of the WL protocol, and £30 after the 1-year follow-up measurements. The study  
117 received approval from the School of Psychology Research Ethics Committee at the University  
118 of Leeds (ref: PSC-238, date: 10/01/2018; amendment to include 1-year follow-up - ref: PSC-  
119 669, date: 11/04/2019).

120

## 121 **2.1 Study design**

122 Participants completed a free-living week of measurements where a PA monitor was worn  
123 continuously to assess minutes of PA and to estimate total daily (TDEE) and activity energy  
124 expenditure (AEE). Upon completion of the free-living week of measurements, participants  
125 attended the laboratory for a testing day which took place after a 10-12 hour overnight fast. This  
126 day included assessments of body composition, resting metabolic rate (RMR), as well other  
127 variables (e.g., appetite ratings and eating behaviour traits) reported elsewhere as these were not  
128 the main aim of the current secondary analysis [20-22]. Upon completing both free-living and  
129 laboratory measurements, participants were randomised to either CER or IER until they reached  
130  $\geq 5\%$  WL or 12 weeks, as previously described [22]. Participants had weekly meetings with a

131 dietitian to monitor body weight and adjust the meal plan if needed. Upon reaching  $\geq 5\%$  WL on  
132 a weekly meeting, participants completed a final free-living week of measurements while still on  
133 CER or IER, emailing their fasted body weight each day to the research dietitian. Measurements  
134 were collected at baseline (before diet allocation), after 2 weeks of energy restriction (to examine  
135 the associations between early changes and longer-term outcomes), at  $\geq 5\%$  WL (or 12 weeks)  
136 and at 1-year post-WL. To assess the impact of early changes in physiological and psychological  
137 outcomes, measurements were collected after 2 weeks of the diet so as to avoid the first phase of  
138 WL in which rapid changes in body water and glycogen stores can occur, and because it is not  
139 uncommon for a 5% WL (the target WL in this study) to occur within 4-6 weeks [23].

140

## 141 **2.2 Free-living measurements**

### 142 **2.2.1 Physical activity**

143 Participants wore a PA monitor (SenseWear Armband; BodyMedia, Inc., Pittsburgh, USA) to  
144 assess PA and estimate TDEE and AEE over 7 days at baseline (before the diet intervention),  
145 after 2 weeks of dietary energy restriction, post-WL and at 1-year follow-up. The SenseWear  
146 Armband is a device which been shown to provide valid estimates of PA and EE [24]. The  
147 SenseWear Armband uses body weight, height and age, as well galvanic skin response, skin  
148 temperature, heat flux and complex pattern-recognition algorithms to determine activity type, to  
149 estimate TDEE. Minutes spent in sedentary ( $< 1.5$  METs), light ( $1.5$ - $2.0$  METs), moderate ( $3.0$ -  
150  $5.9$  METs) and vigorous ( $\geq 6.0$  METs) activities, as well daily steps and sleep duration were  
151 calculated using proprietary algorithms presented in the device's accompanying software (version  
152 8.0 professional), previously validated [24]. AEE was calculated using the following equation:

153

$$154 \quad \textit{Activity Energy Expenditure} = \textit{TDEE} \times 0.9 - \textit{RMR}$$

155

156 Participants were instructed to wear the monitor halfway between their elbow and shoulder for at  
157 least 23 hours per day (including overnight, although daily and nightly activities have not been  
158 discriminated), only removing during activities that involved contact with water (e.g., shower and

159 swimming). Compliance with utilising the monitor was defined as having a minimum of 22 hours  
160 of verifiable time per day for at least 5 days (including one weekend day). All participants wore  
161 the PA monitor for at least 5 days, with a mean wear time per day of 23 hours and 40 minutes  
162 (from 23 hours and 7 minutes to 23 hours and 54 minutes). Participants were instructed not to  
163 change their structured exercise habits for the duration of the WL phase e.g., start an exercise  
164 programme if this was not already part of their routine. However, no specific instructions were  
165 given regarding habitual daily PA behaviours, and these behaviours were not restricted or  
166 controlled throughout the intervention to allow quantification of the degree of spontaneous non-  
167 exercise PA changes. As changes in PA behaviours may naturally occur in response to periods of  
168 negative energy balance despite the absence of specific recommendations [7], the aim of this  
169 analysis was to examine how these spontaneous changes could influence body weight outcomes.  
170 An important factor to consider is that AEE and TDEE are influenced by changes in body weight.  
171 Therefore, when exporting the data from the SenseWear Armband, the value for body weight was  
172 updated to control for the reduction in EE induced by losses of body mass. Furthermore, steps per  
173 day and minutes of total, light and moderate-to-vigorous PA, and sedentary time, were examined  
174 as these are commonly used measurements of PA independent of body weight and body  
175 composition. No instructions were given to participants between the post-WL phase and the 1-  
176 year follow-up in terms of PA (or dietary) patterns and therefore, participants could have started  
177 or stopped any type of formal exercise routines during these 12 months.

178

## 179 **2.3 Laboratory measurements**

### 180 **2.3.1 Body weight and composition**

181 Body weight and composition were measured whilst participants were wearing tight fitting  
182 clothing and a swimming cap using air-displacement plethysmography (BodPod, COSMED Inc.,  
183 Concord, USA). Fat mass (FM) and FFM were estimated to the nearest 0.01kg, and  
184 manufacturer's instructions were followed and the Siri equation [25] was used to estimate body  
185 fat percentage.

186

187 **2.3.2 Rate of weight loss**

188 In the present study, total percentage of WL and the time to complete the intervention (i.e., final  
189 day of measurements) ranged from 3.2% to 8.3% and 35 to 93 days, respectively. As individuals  
190 with different starting body masses were being compared, which could alter the absolute amount  
191 of WL [26], relative changes in body weight were reported as a percentage. To control for the  
192 variability in intervention duration and total WL between participants, mean rate of WL  
193 throughout the intervention was calculated. In the scientific literature [27-31], rate of WL has  
194 been calculated using the following equation:

195

196 
$$\text{Rate of Weight Loss (\% per week)} = \frac{\text{Total Weight Loss (\%)}}{\text{Time (weeks)}}$$

197

198 The mean rate of WL was calculated at weeks 2 and post-WL. As the timing for the follow-up  
199 measurements was matched between participants (approximately 1 year), percentage of body  
200 weight change from post-WL to 1-year follow-up was calculated.

201

202 **2.3.3 Resting metabolic rate**

203 RMR was measured with an indirect calorimeter fitted with a ventilated hood (GEM, Nutren  
204 Technology Ltd). Participants were asked to remain in a supine position for 40 minutes without  
205 moving, talking or falling asleep. Before each measurement, an individual calibration process was  
206 performed. RMR was calculated using the 5-minute steady state method [32], and data was  
207 entered into the Weir equation [33].

208

209 **2.4 Statistical analyses**

210 Data are presented as mean  $\pm$  standard deviation. Data were analysed using SPSS software version  
211 25 (IBM Corp., Armonk, New York). The Shapiro-Wilk test was used to examine for normality  
212 of distribution and all data were normally distributed. Analyses were conducted with data from  
213 participants that completed the intervention ( $\geq 5\%$  WL or 12 weeks). Differences between  
214 intervention groups (CER and IER) at baseline were examined using Welch's t-tests. Changes

215 over time were analysed with repeated measures maximum-likelihood linear mixed models to  
216 account for missing data, using SPSS (version 26, IBM, USA). Measures day (baseline, week 2,  
217 post-WL and 1-year post-WL), intervention group (CER and IER) and their interaction were  
218 analysed as fixed factors and subject as random factor. Bonferroni adjustments were applied to  
219 post-hoc analyses. Data are presented as estimated marginal means and 95% confidence intervals.  
220 For the analyses pertaining to the rate of WL, data from both groups were combined as no  
221 statistical differences existed between groups [22]. Partial correlations (adjusted for WL group  
222 and baseline values) were conducted to examine the associations between baseline characteristics,  
223 changes from baseline to week 2 and from baseline to post-WL with the mean rate of WL, as the  
224 rate of WL was different between dietary groups (CER:  $0.8 \pm 0.3\%$ /week; IER:  $0.6 \pm 0.3\%$ /week;  
225  $p=0.01$ ). Pearson correlations were also conducted to examine the associations between changes  
226 from post-WL to follow-up and 1-year weight change. However, as 1-year weight change was  
227 similar between groups and these were not following a particular dietary pattern, these  
228 associations were not adjusted for group. The main study from which these secondary analyses  
229 have been conducted was originally powered to detect an interaction in self-selected meal size  
230 (*ad libitum* energy intake) between 2 groups and 2 repeated measurements [22], but power  
231 calculations (G\*Power v3.1) indicated that a sample size of 23 would be sufficient to see a  
232 correlation coefficient of 0.50 between PA and weight change with  $\alpha=0.05$  and  $1-\beta=0.8$  (based  
233 on a previous study that observed a correlation coefficient of  $r=-0.69$  [34]). Statistical significance  
234 was defined as  $p<0.05$ .

235

### 236 **3. Results**

#### 237 **3.1 Participant flow**


238 A total of 54 participants were enrolled in the trial, 46 completed baseline measurements, with no  
239 differences between groups (all  $p>0.18$ ) and were randomly allocated to a diet group (CER – 22;  
240 IER – 24), and 37 reached  $\geq 5\%$  WL or 12 weeks (CER – 19; IER – 18). Eighteen participants  
241 returned for the 1-year follow-up (CER – 11; IER – 7). Characteristics of the participants that

242 completed the WL intervention (n=37) and that returned after 1-year (n=18) can be found in table  
243 1 and a participant flow chart can be found in figure 1.

244

245 TABLE 1

246



247

248 **Figure 1** – Participant flow chart.

249

250 **3.2 Changes during the intervention**

251 Mean values for each group at each time point during the intervention can be seen in Table 2. No  
 252 baseline differences were observed between dietary groups (all  $p > 0.12$ ). Both groups achieved a  
 253 similar total WL (CER:  $6.2 \pm 0.8\%$ ; IER:  $5.5 \pm 2.1\%$ ;  $p = 0.17$ ). The mean rate of WL was similar

254 between groups at week 2 (CER:  $0.2\pm 0.1\%$ /week; IER:  $0.2\pm 0.1\%$ /week;  $p=0.79$ ), but different  
255 throughout the entire intervention (CER:  $0.8\pm 0.3\%$ /week; IER:  $0.6\pm 0.3\%$ /week;  $p=0.01$ ). Both  
256 groups presented a similar weight change from post-WL to 1-year follow-up (CER:  $5.0\pm 6.0\%$ ;  
257 IER:  $3.7\pm 4.0\%$ ;  $p=0.62$ ). One participant (CER) displayed weight regain of 19.7%, and when  
258 removed, weight regain was near identical (CER:  $3.6\pm 3.7\%$ ; IER:  $3.7\pm 4.0\%$ ;  $p=0.93$ ). Weight  
259 change from post-WL to 1-year follow-up in the whole group ranged from -2.1% to +19.7% (-1.4  
260 to +14.0kg), or from -2.1% to 9.7% (-1.4 to +8.2kg) when the outlier was removed.

261 There was a main effect of time ( $p<0.001$ ) but no effect of group or interaction ( $p\geq 0.15$ ) for body  
262 weight, fat mass, fat-free mass, body fat percentage, RMR, TDEE and AEE. Post-hoc analyses  
263 are shown in Table 2. There were no time, group, or interaction effects for daily steps, sleep  
264 duration, total PA, light PA, moderate-to-vigorous PA or sedentary time ( $p\geq 0.07$ ).

265

266

267


## TABLE 2

268

269 **3.4 Associations between changes at week 2 and mean rate of weight loss**

270 No associations were seen between baseline PA, sedentary time, sleep duration, TDEE or AEE  
271 with the mean rate of WL throughout the intervention ( $p>0.05$ ).


272 Changes in total PA ( $r=0.29$ ;  $p=0.10$ ), light ( $r=0.03$ ;  $p=0.86$ ) and moderate-to-vigorous PA  
273 ( $r=0.25$ ;  $p=0.16$ ), steps per day ( $r=0.19$ ;  $p=0.26$ ), sleep duration ( $r=0.18$ ;  $p=0.32$ ), TDEE ( $r=0.07$ ;  
274  $p=0.72$ ) and AEE ( $r=0.07$ ;  $p=0.70$ ) from baseline to week 2 were not associated with the mean  
275 rate of WL throughout the intervention. As shown in figure 2, mean rate of WL ( $r=0.42$ ;  $p=0.01$ )  
276 and changes in sedentary time ( $r=-0.37$ ;  $p=0.03$ ) from baseline to week 2 were associated with  
277 the mean rate of WL throughout the energy restriction phase.



278 **Figure 2** – Associations between mean rate of weight loss in the participants that completed the  
279 intervention with A) mean rate of weight loss at week 2 and B) changes in sedentary time at week  
280 2. Grey bands represent the 95% confidence intervals.  
281

282 **3.5 Associations between changes throughout the intervention and mean rate of weight loss**

283 Changes in sleep duration ( $r=0.06$ ;  $p=0.73$ ) were not associated with the mean rate of WL during  
284 the energy restriction phase. Changes in total PA ( $r=0.55$ ;  $p<0.01$ ), light PA ( $r=0.43$ ;  $p=0.01$ ),  
285 moderate-to-vigorous PA ( $r=0.51$ ;  $p<0.01$ ), sedentary time ( $r=-0.56$ ;  $p<0.01$ ), steps per day  
286 ( $r=0.39$ ;  $p=0.02$ ), TDEE ( $r=0.41$ ;  $p=0.02$ ) and AEE ( $r=0.47$ ;  $p<0.01$ ) were associated with the  
287 mean rate of WL (Figure 3). Associations were also found between the days to reach 5% WL  
288 (which ranged from 35 to 93 days) and changes throughout the energy restriction phase in total  
289 PA ( $r=-0.49$ ;  $p=0.004$ ), light PA ( $r=-0.43$ ;  $p=0.01$ ), moderate-to-vigorous PA ( $r=-0.47$ ;  $p=0.007$ ),  
290 sedentary time ( $r=0.55$ ;  $p=0.001$ ) and steps per day ( $r=0.36$ ;  $p=0.04$ ).




291  
 292  
 293  
 294  
 295  
 296  
 297  
 298

**Figure 3** – Associations between mean rate of weight loss in the participants that completed the intervention with changes throughout the intervention in A) total physical activity, B) moderate-to-vigorous physical activity, C) sedentary time, D) steps per day, E) total daily energy expenditure, and F) activity energy expenditure. Grey bands represent the 95% confidence intervals.

299 **3.6 Factors associated with post-WL 1-year weight change**

300 Changes in light PA ( $r=-0.32$ ;  $p=0.24$ ), sedentary time ( $r=0.39$ ;  $p=0.13$ ), steps per day ( $r=-0.39$ ;  
301  $p=0.12$ ), sleep duration ( $r=-0.08$ ;  $p=0.77$ ), TDEE ( $r=-0.07$ ;  $p=0.80$ ) and AEE ( $r=-0.06$ ;  $p=0.81$ )  
302 from post-WL to 1-year follow-up were not associated with 1-year weight change. However,  
303 changes in total PA ( $r=-0.50$ ;  $p=0.04$ ) and moderate-to-vigorous PA ( $r=-0.61$ ;  $p=0.01$ ), were  
304 associated with 1-year weight change (Figure 4).



305 **Figure 4** – Associations between 1-year weight change and changes between post-WL and 1-year  
306 follow-up in A) total physical activity and B) moderate-to-vigorous physical activity. Grey bands  
307 represent the 95% confidence intervals.

308

309 The mean rate of WL during the WL phase was not associated with 1-year weight change ( $r=-$   
310  $0.01$ ;  $p=0.97$ ). However, changes in total PA ( $r=-0.50$ ;  $p=0.04$ ), moderate-to-vigorous PA ( $r=-$   
311  $0.64$ ;  $p<0.01$ ), sedentary time ( $r=-0.71$ ;  $p<0.01$ ) and TDEE ( $r=-0.48$ ;  $p=0.04$ ) from baseline to  
312 post-WL were negatively associated with the changes from post-WL to 1-year follow-up, with  
313 greater increases in PA or TDEE during the WL phase being associated with greater decreases  
314 during the 1-year post-WL phase.

315

316 **4. Discussion**

317 The aim of this secondary analysis was to explore whether changes in objectively measured PA,  
318 sedentary time and EE were associated with the rate of WL during dietary energy restriction and  
319 1-year weight change post-WL. In these data, baseline characteristics were not associated with  
320 longer-term WL outcomes, but the rate of WL and changes in sedentary time after 2 weeks were

321 associated with the mean rate of WL during the dietary intervention period. Changes in total PA,  
322 light PA, moderate-to-vigorous PA, sedentary time, steps per day, TDEE and AEE from baseline  
323 to post-WL were associated with the mean rate of WL during the energy restriction phase, while  
324 changes in total PA and moderate-to-vigorous PA from post-WL to 1-year follow-up were  
325 associated with the change in body weight during the non-contact follow-up period. Changes in  
326 sleep duration were not associated with body weight outcomes at any timepoints. Data from this  
327 secondary analysis suggests that increases (or smaller reductions) in PA behaviours during dietary  
328 energy restriction may help facilitate WL and attenuate weight regain. As such, these data  
329 highlight the potential importance of considering PA and sedentary time in dietary weight  
330 management interventions.

#### 331 **4.1 Changes in physical activity during diet-induced weight loss**

332 It has been previously suggested that diet-induced WL may lead to reductions in PA, with a recent  
333 systematic review reporting that 7 out of 14 diet-only interventions observed decreases in non-  
334 exercise PA [7]. In the current study, no changes were observed in mean PA or sedentary time  
335 over time, and no differences in PA or sedentary time were seen between dietary groups. This  
336 corroborates the findings from a previous study comparing intermittent to continuous energy  
337 restriction [10] and other diet-only interventions that did not observe reductions in the amount of  
338 PA [11, 35]. However, despite the absence of mean changes in PA in the present study, a large  
339 inter-individual variability was observed. For instance, changes in total PA from baseline to post-  
340 WL ranged from -130 to +209min/day, while the mean change was only +5min/day ( $p=1.00$ ). As  
341 such, focusing on the changes in PA at the group level may mask important information regarding  
342 how individual differences in PA influence the rate of WL at the individual level.

#### 343 **4.2 Associations between early changes in physical activity and the mean rate of weight loss**

344 Several studies have reported that WL in the first weeks of an intervention (2-6 weeks) is a  
345 predictor of longer-term total WL [36-38]. For instance, Tronieri et al. observed that participants  
346 that lost more weight in the first 4 weeks, lost more weight at week 14 ( $r^2=0.61$ ;  $p<0.001$ ) and  
347 presented a faster rate of WL [38]. In the current study, a faster rate of WL during the first two

348 weeks of energy restriction and a decrease in sedentary time during the first 2 weeks were  
349 associated with a faster mean rate of WL during the total energy restriction period. Furthermore,  
350 early changes in PA (i.e., baseline to week 2) were strongly correlated with the baseline to post-  
351 WL changes ( $r=0.60-0.70$ ;  $p<0.001$ ), suggesting that the early changes in PA were maintained  
352 across the full dietary energy restriction period. While few studies have looked into the influence  
353 of early changes in PA or EE on WL outcomes, these findings are in agreement with a study by  
354 Reinhardt et al. in which changes in TDEE in response to a 24-hour fast were associated with WL  
355 after 6 weeks [39]. In this study, individuals that presented a greater decrease in TDEE (which is  
356 influenced by PA) during 24 hours of fasting, presented a slower rate of WL. However, as this  
357 was measured in a respiratory chamber (in which PA could be artificially limited), it remains  
358 unknown whether this association between changes in 24-hour TDEE and 6-week WL was due  
359 to changes in PAEE or some other TDEE component (although the authors reported that changes  
360 in sleeping metabolic rate were not associated with WL). These findings suggest that early  
361 changes in body weight and PA (2 weeks in the case of the current study) during diet-induced  
362 energy restriction may reflect how well someone will respond in terms of longer-term WL. If this  
363 is the case, this could improve weight management success as practitioners would be able to be  
364 proactive and adjust an intervention early based on shorter-term responses. However, future  
365 studies should aim to replicate these findings to confirm whether early changes in PA allow to  
366 predict how individuals will lose weight in the longer-term.

#### 367 **4.3 Associations between changes throughout the intervention and the mean rate of weight** 368 **loss**

369 An important finding from the current study was that changes in PA and sedentary time  
370 throughout the diet intervention were associated with the mean rate of WL, with participants that  
371 had greater increases in PA and decreases in sedentary time presenting faster mean rates of WL.  
372 These findings are in agreement with a previous WL study (meal plan and instructions to increase  
373 PA) in which the group of individuals that had greater increases in moderate-to-vigorous PA lost  
374 more weight after 6 months [6], suggesting that maintaining or increasing PA during periods of

375 dietary-induced energy restriction may be an important behavioural strategy to facilitate WL.  
376 Overall, these findings corroborate previous literature reporting that the combination of diet and  
377 PA leads to better WL outcomes [16, 18].

378 Although the amount of PA performed (e.g., minutes per day) and AEE are related, PA is a  
379 behaviour while AEE represents the EE associated with movement and is therefore also  
380 influenced by the mass and composition of an individual [40]. In this data, PA levels were strongly  
381 associated with AEE (total PA –  $r=0.70$ ;  $p<0.001$ ; moderate-to-vigorous PA –  $r=0.74$ ;  $p<0.001$ ;  
382 sedentary time –  $r=-0.64$ ;  $p<0.001$ ), suggesting that a potential mechanism to explain the current  
383 findings is that the changes in PA and sedentary time helped to better maintain or increase the  
384 energy deficit created via energy restriction. It is important to report these objectively measured  
385 effects since some recent pronouncements have claimed a limited relationship between PA and  
386 AEE which could undermine a rationale for promoting the beneficial effects of PA on body weight  
387 (or body fat) [41].

388 In the present study, participants exercised  $\leq 3$  days per week at baseline and were instructed not  
389 to change their exercise habits during the dietary intervention (e.g., start a structured exercise  
390 regime alongside the dietary intervention), but no strict restrictions were placed on other PA  
391 behaviours during the intervention. However, it is important to highlight that time spent  
392 performing moderate-to-vigorous PA in the current study was on average  $>60$ min/day, suggesting  
393 the participants included were relatively active. Whether the changes in PA and sedentary time  
394 during the WL intervention were intentional is unknown, and an important question that should  
395 be addressed in future research is whether individuals who demonstrate better WL outcomes  
396 during energy restriction actively increase their PA to augment WL. While PA levels are readily  
397 modifiable, it cannot be ruled out that individuals became more active as a result of their greater  
398 WL. Therefore, the hypothesis that PA may increase as a consequence of WL, rather than  
399 increases in PA leading to a faster rate of WL, cannot be ruled out and should be explored in  
400 future studies.

401

#### 402 **4.4 Factors associated with 1-year weight change**

403 Changes in moderate-to-vigorous PA from post-WL to 1-year follow-up were associated with 1-  
404 year weight change. These findings are in agreement with previous studies highlighting PA as a  
405 robust predictor of WL maintenance [3, 42, 43], but not all [44]. An interesting observation in the  
406 present study was that participants that increased PA during the WL phase had lower baseline  
407 values, but these individuals also demonstrated greater reductions in PA between the end of the  
408 WL phase and the 1-year follow-up point. This perhaps suggests that participants with a greater  
409 rate of WL consciously increased their PA, but after the WL phase terminated, the absence of a  
410 specific WL goal may have led to a return to baseline PA levels. However, it is important to  
411 consider that since the sample size was limited to 18 individuals (from 37 participants that finished  
412 the WL phase), these findings should be interpreted cautiously. Nonetheless, the observed  
413 associations in this secondary analysis should be viewed as an initial proof of concept highlighting  
414 the relevance of PA for sustained WL, and this enquiry should be replicated in future studies with  
415 larger sample sizes.

#### 416 **4.5 Limitations**

417 The equation used to calculate the rate of WL, as well assessing changes in PA at baseline, week  
418 2, post-WL and after 1-year assumes that these changes are linear over time. This may be  
419 inaccurate due to the daily fluctuations in both EI and EE that may occur during periods of  
420 negative energy balance [45, 46]. However, the main aim of this study was to identify the factors  
421 associated with WL variability and not with the intra-individual variability in weekly changes in  
422 body weight. Therefore, this calculation allowed for an examination of the factors that explain  
423 why some individuals lose weight faster (on average). Furthermore, changes in PA at week 2 and  
424 post-intervention were strongly associated (all  $r=0.60-0.70$ ;  $p<0.001$ ), as well with changes  
425 between post-WL and 1-year follow-up, suggesting that the individual changes in PA were  
426 consistent across the study period. It is also important to acknowledge that the sample size,  
427 especially at 1-year follow-up, was small and consisted only of women, potentially limiting the  
428 generalisability of the findings. Lastly, as there was no contact between post-WL and 1-year

429 follow-up, it is not known whether changes in PA and sedentary time were conscious and  
430 voluntary remains unknown.

431

## 432 **5. Conclusion**

433 The results from this secondary analysis corroborate previous findings demonstrating that  
434 baseline characteristics may not be good indicators of longer-term WL. However, increases in PA  
435 behaviours after 2 weeks and throughout the intervention were associated with a faster mean rate  
436 of WL. Furthermore, decreases in PA behaviours were associated with a greater 1-year weight  
437 regain. Conversely, an increase in sedentary time was associated with a slower rate of WL and  
438 greater weight regain. These findings highlight the potential contribution of PA during dietary  
439 weight management interventions, and as a potentially modifiable component of TDEE, may be  
440 an important behavioural target during dietary energy restriction that promotes better weight  
441 outcomes during dietary energy restriction.

442

### 443 **Financial support:**

444 This work was funded in part by a Research Fellowship awarded to Nuno Casanova by the  
445 Wellcome Trust's International Strategic Support Fund (204825/Z/16/Z) and by a Research  
446 Fellowship awarded to Kristine Beaulieu by the European Society for Clinical Nutrition and  
447 Metabolism (ESPEN).

448

### 449 **Conflict of interest:**

450 None.

451

### 452 **Authorship:**

453 The authors' responsibilities were as follows – NC, KB, PO, CG, JB, GF and MH: designed  
454 research; NC, KB, PO and DOC: conducted research; NC: analysed data; NC: wrote of the first

455 draft of the manuscript; NC: had primary responsibility for final content; and all authors: read and  
456 approved the final manuscript.  
457

458 **References**

459

460 [1] Wing, R. R., Phelan, S. Long-term weight loss maintenance. *The American*  
461 *Journal of Clinical Nutrition*. 2005,82:222S-5S.

462 [2] Greaves, C. J., Sheppard, K. E., Abraham, C., Hardeman, W., Roden, M.,  
463 Evans, P. H., et al. Systematic review of reviews of intervention components  
464 associated with increased effectiveness in dietary and physical activity  
465 interventions. *BMC Public Health*. 2011,11:119.

466 [3] Paixão, C., Dias, C. M., Jorge, R., Carraça, E. V., Yannakoulia, M., de Zwaan,  
467 M., et al. Successful weight loss maintenance: A systematic review of weight  
468 control registries. *Obes Rev*. 2020,21:e13003.

469 [4] Yank, V., Xiao, L., Wilson, S. R., Stafford, R. S., Rosas, L. G., Ma, J. Short-  
470 term weight loss patterns, baseline predictors, and longer-term follow-up within a  
471 randomized controlled trial. *Obesity (Silver Spring)*. 2014,22:45-51.

472 [5] Unick, J. L., Pellegrini, C. A., Demos, K. E., Dorfman, L. Initial Weight Loss  
473 Response as an Indicator for Providing Early Rescue Efforts to Improve Long-  
474 term Treatment Outcomes. *Current diabetes reports*. 2017,17:69.

475 [6] Fazzino, T. L., Fabian, C., Befort, C. A. Change in Physical Activity During a  
476 Weight Management Intervention for Breast Cancer Survivors: Association with  
477 Weight Outcomes. *Obesity (Silver Spring)*. 2017,25 Suppl 2:S109-S15.

478 [7] Silva, A. M., Judice, P. B., Carraca, E. V., King, N., Teixeira, P. J., Sardinha,  
479 L. B. What is the effect of diet and/or exercise interventions on behavioural  
480 compensation in non-exercise physical activity and related energy expenditure of  
481 free-living adults? A systematic review. *Br J Nutr*. 2018,119:1327-45.

482 [8] de Groot, L. C., van Es, A. J., van Raaij, J. M., Vogt, J. E., Hautvast, J. G.  
483 Energy metabolism of overweight women 1 mo and 1 y after an 8-wk slimming  
484 period. *Am J Clin Nutr*. 1990,51:578-83.

485 [9] Weigle, D. S. Contribution of decreased body mass to diminished thermic  
486 effect of exercise in reduced-obese men. *Int J Obes*. 1988,12:567-78.

487 [10] Coutinho, S. R., Halset, E. H., Gasbakk, S., Rehfeld, J. F., Kulseng, B.,  
488 Truby, H., et al. Compensatory mechanisms activated with intermittent energy  
489 restriction: A randomized control trial. *Clin Nutr*. 2018,37:815-23.

490 [11] van Dale, D., Schoffelen, P. F., ten Hoor, F., Saris, W. H. Effects of addition  
491 of exercise to energy restriction on 24-hour energy expenditure, sleeping  
492 metabolic rate and daily physical activity. *Eur J Clin Nutr*. 1989,43:441-51.

493 [12] Yancy, W. S., Jr., Olsen, M. K., Guyton, J. R., Bakst, R. P., Westman, E. C.  
494 A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and  
495 hyperlipidemia: a randomized, controlled trial. *Ann Intern Med*. 2004,140:769-77.

496 [13] Gardner, C. D., Trepanowski, J. F., Del Gobbo, L. C., Hauser, M. E., Rigdon,  
497 J., Ioannidis, J. P. A., et al. Effect of Low-Fat vs Low-Carbohydrate Diet on 12-  
498 Month Weight Loss in Overweight Adults and the Association With Genotype  
499 Pattern or Insulin Secretion: The DIETFITS Randomized Clinical Trial. *Jama*.  
500 2018,319:667-79.

501 [14] Malhotra, A., Noakes, T., Phinney, S. It is time to bust the myth of physical  
502 inactivity and obesity: you cannot outrun a bad diet. *British journal of sports*  
503 *medicine*. 2015:bjspports-2015-094911.

504 [15] Jakicic, J. M., Rogers, R. J., Davis, K. K., Collins, K. A. Role of Physical  
505 Activity and Exercise in Treating Patients with Overweight and Obesity. *Clin*  
506 *Chem*. 2018,64:99-107.

- 507 [16] Curioni, C. C., Lourenço, P. M. Long-term weight loss after diet and exercise:  
508 a systematic review. *Int J Obes (Lond)*. 2005,29:1168-74.
- 509 [17] DeLany, J. P., Kelley, D. E., Hames, K. C., Jakicic, J. M., Goodpaster, B. H.  
510 Effect of physical activity on weight loss, energy expenditure, and energy intake  
511 during diet induced weight loss. *Obesity (Silver Spring)*. 2014,22:363-70.
- 512 [18] Bellicha, A., van Baak, M. A., Battista, F., Beaulieu, K., Blundell, J. E.,  
513 Busetto, L., et al. Effect of exercise training on weight loss, body composition  
514 changes, and weight maintenance in adults with overweight or obesity: An  
515 overview of 12 systematic reviews and 149 studies. *n/a:e13256*.
- 516 [19] Stubbs, R. J., Duarte, C., O'Driscoll, R., Turicchi, J., Michalowska, J.  
517 Developing evidence-based behavioural strategies to overcome physiological  
518 resistance to weight loss in the general population. *Proc Nutr Soc*. 2019,78:576-  
519 89.
- 520 [20] Beaulieu, K., Casanova, N., Oustric, P., Hopkins, M., Varady, K., Finlayson,  
521 G., et al. An exploratory investigation of the impact of 'fast' and 'feed' days during  
522 intermittent energy restriction on free-living energy balance behaviours and  
523 subjective states in women with overweight/obesity. *Eur J Clin Nutr*. 2020.
- 524 [21] Oustric, P., Beaulieu, K., Casanova, N., O'Connor, D., Gibbons, C., Hopkins,  
525 M., et al. Food Liking but Not Wanting Decreases after Controlled Intermittent or  
526 Continuous Energy Restriction to  $\geq 5\%$  Weight Loss in Women with  
527 Overweight/Obesity. *Nutrients*. 2021,13.
- 528 [22] Beaulieu, K., Casanova, N., Oustric, P., Turicchi, J., Gibbons, C., Hopkins,  
529 M., et al. Matched Weight Loss Through Intermittent or Continuous Energy  
530 Restriction Does Not Lead To Compensatory Increases in Appetite and Eating  
531 Behavior in a Randomized Controlled Trial in Women with Overweight and  
532 Obesity. *The Journal of Nutrition*. 2019.
- 533 [23] Heymsfield, S. B., Thomas, D., Nguyen, A. M., Peng, J. Z., Martin, C., Shen,  
534 W., et al. Voluntary weight loss: systematic review of early phase body  
535 composition changes. *Obes Rev*. 2011,12:e348-61.
- 536 [24] O'Driscoll, R., Turicchi, J., Beaulieu, K., Scott, S., Matu, J., Deighton, K., et  
537 al. How well do activity monitors estimate energy expenditure? A systematic  
538 review and meta-analysis of the validity of current technologies. 2018:bjsports-  
539 2018-099643.
- 540 [25] Siri, W. Body composition from fluid space and density. J. Brozek & A.  
541 Hanschel (Eds.), In: *Techniques for measuring body composition*. Washington,  
542 DC National Academy of Science 1961 p. 223-2441961.
- 543 [26] Hatoum, I. J., Kaplan, L. M. Advantages of percent weight loss as a method  
544 of reporting weight loss after Roux-en-Y gastric bypass. *Obesity (Silver Spring)*.  
545 2013,21:1519-25.
- 546 [27] Finkler, E., Heymsfield, S. B., St-Onge, M.-P. Rate of weight loss can be  
547 predicted by patient characteristics and intervention strategies. *Journal of the*  
548 *Academy of Nutrition and Dietetics*. 2012,112:75-80.
- 549 [28] Nackers, L. M., Ross, K. M., Perri, M. G. The association between rate of  
550 initial weight loss and long-term success in obesity treatment: does slow and  
551 steady win the race? *Int J Behav Med*. 2010,17:161-7.
- 552 [29] Vink, R. G., Roumans, N. J., Arkenbosch, L. A., Mariman, E. C., van Baak,  
553 M. A. The effect of rate of weight loss on long-term weight regain in adults with  
554 overweight and obesity. *Obesity (Silver Spring)*. 2016,24:321-7.
- 555 [30] Coutinho, S. R., With, E., Rehfeld, J. F., Kulseng, B., Truby, H., Martins, C.  
556 The impact of rate of weight loss on body composition and compensatory

557 mechanisms during weight reduction: A randomized control trial. *Clin Nutr.*  
558 2018,37:1154-62.

559 [31] Turicchi, J., O'Driscoll, R., Finlayson, G., Beaulieu, K., Deighton, K., Stubbs,  
560 R. J. Associations between the rate, amount, and composition of weight loss as  
561 predictors of spontaneous weight regain in adults achieving clinically significant  
562 weight loss: A systematic review and meta-regression. *Obes Rev.* 2019,20:935-  
563 46.

564 [32] Sanchez-Delgado, G., Alcantara, J. M. A., Ortiz-Alvarez, L., Xu, H., Martinez-  
565 Tellez, B., Labayen, I., et al. Reliability of resting metabolic rate measurements  
566 in young adults: Impact of methods for data analysis. *Clin Nutr.* 2018,37:1618-  
567 24.

568 [33] Weir, J. B. New methods for calculating metabolic rate with special reference  
569 to protein metabolism. *J Physiol.* 1949,109:1-9.

570 [34] Aadland, E., Robertson, L. Physical activity is associated with weight loss  
571 and increased cardiorespiratory fitness in severely obese men and women  
572 undergoing lifestyle treatment. *Journal of obesity.* 2012,2012:810594-.

573 [35] Weinsier, R. L., Hunter, G. R., Zuckerman, P. A., Redden, D. T., Darnell, B.  
574 E., Larson, D. E., et al. Energy expenditure and free-living physical activity in  
575 black and white women: comparison before and after weight loss. *Am J Clin Nutr.*  
576 2000,71:1138-46.

577 [36] Barnes, R. D., Ivezaj, V., Pittman, B. P., Grilo, C. M. Early weight loss predicts  
578 weight loss treatment response regardless of binge-eating disorder status and  
579 pretreatment weight change. *Int J Eat Disord.* 2018,51:558-64.

580 [37] James, B. L., Roe, L. S., Loken, E., Rolls, B. J. Early predictors of weight  
581 loss in a 1-year behavioural weight-loss programme. *Obes Sci Pract.* 2018,4:20-  
582 8.

583 [38] Tronieri, J. S., Wadden, T. A., Chao, A. M., Pearl, R. L., Alamuddin, N.,  
584 Berkowitz, R. I. Early Weight Loss in Behavioral Treatment Predicts Later Rate  
585 of Weight Loss and Response to Pharmacotherapy. *Ann Behav Med.*  
586 2019,53:290-5.

587 [39] Reinhardt, M., Thearle, M. S., Ibrahim, M., Hohenadel, M. G., Bogardus, C.,  
588 Krakoff, J., et al. A Human Thrifty Phenotype Associated With Less Weight Loss  
589 During Caloric Restriction. *Diabetes.* 2015,64:2859-67.

590 [40] Westerterp, K. Physical activity and physical activity induced energy  
591 expenditure in humans: measurement, determinants, and effects. 2013,4.

592 [41] Pontzer, H., Durazo-Arvizu, R., Dugas, L. R., Plange-Rhule, J., Bovet, P.,  
593 Forrester, T. E., et al. Constrained Total Energy Expenditure and Metabolic  
594 Adaptation to Physical Activity in Adult Humans. *Current Biology.* 2016,26:410-  
595 7.

596 [42] Ostendorf, D. M., Lyden, K., Pan, Z., Wyatt, H. R., Hill, J. O., Melanson, E.  
597 L., et al. Objectively Measured Physical Activity and Sedentary Behavior in  
598 Successful Weight Loss Maintainers. *Obesity (Silver Spring).* 2018,26:53-60.

599 [43] Ostendorf, D. M., Blankenship, J. M., Grau, L., Arbet, J., Mitchell, N. S.,  
600 Creasy, S. A., et al. Predictors of long-term weight loss trajectories during a  
601 behavioral weight loss intervention: An exploratory analysis.n/a.

602 [44] Washburn, R. A., Szabo-Reed, A. N., Gorczyca, A. M., Sullivan, D. K.,  
603 Honas, J. J., Mayo, M. S., et al. A Randomized Trial Evaluating Exercise for the  
604 Prevention of Weight Regain. 2021,29:62-70.

605 [45] Melby, C. L., Paris, H. L., Foright, R. M., Peth, J. Attenuating the Biologic  
606 Drive for Weight Regain Following Weight Loss: Must What Goes Down Always  
607 Go Back Up? *Nutrients*. 2017,9.

608 [46] Casanova, N., Beaulieu, K., Finlayson, G., Hopkins, M. Metabolic  
609 adaptations during negative energy balance and their potential impact on appetite  
610 and food intake. *Proc Nutr Soc*. 2019,78:279-89.

611

612 **Table 1** – Participant characteristics of the completers at baseline and 1-year follow-up.

	Baseline			1-Year Follow-Up		
	CER (n=19)	IER (n=18)	Total (n=37)	CER (n=11)	IER (n=7)	Total (n=18)
Age (y)	34 ± 9	36 ± 11	35 ± 10	38 ± 9	37 ± 12	38 ± 10
Body weight (kg)	79.6 ± 10.3	80.1 ± 11.1	79.9 ± 10.6	73.7 ± 6.8	77.1 ± 13.3	75.0 ± 9.6
Height (cm)	165.1 ± 7.8	165.5 ± 8.7	165.3 ± 8.1	161.5 ± 4.6	161.5 ± 6.6	161.5 ± 5.2
BMI (kg/m <sup>2</sup> )	29.1 ± 2.4	29.1 ± 2.2	29.1 ± 2.3	28.2 ± 2.2	29.5 ± 4.0	28.7 ± 3.0
Fat mass (kg)	32.8 ± 8.1	33.5 ± 6.7	33.1 ± 7.4	28.8 ± 5.7	34.2 ± 10.0	30.9 ± 7.9
Fat mass (%)	40.7 ± 6.1	41.6 ± 4.1	41.2 ± 5.2	38.8 ± 5.2	43.6 ± 5.9	40.7 ± 5.8
Fat-free mass (kg)	46.9 ± 5.4	46.6 ± 6.1	46.7 ± 5.7	44.9 ± 3.9	42.9 ± 4.6	44.1 ± 4.2

613

614

615 **Table 2** – Mean values for participants of both CER and IER that completed the intervention at baseline, week 2, post-WL and at 1-year follow-up.

		<b>Baseline</b>	<b>Week 2</b>	<b>Post-WL</b>	<b>1-year follow-up</b>	<b>Baseline vs week 2</b>	<b>Baseline vs post-WL</b>	<b>Post-WL vs follow-up</b>
<b>Body weight (kg)</b>	<b>CER</b>	79.63 [74.82, 84.45]	77.72 [72.91, 82.53]	74.71 [69.89, 79.52]	78.21 [73.35, 83.07]	<0.001	<0.001	<0.001
	<b>IER</b>	80.09 [75.15, 85.03]	78.18 [73.24, 83.12]	75.66 [70.72, 80.61]	78.58 [73.52, 83.64]			
<b>Fat mass (kg)</b>	<b>CER</b>	32.75 [29.42, 36.08]	31.22 [27.89, 34.55]	29.08 [25.75, 32.41]	31.93 [28.53, 35.32]	0.008	<0.001	<0.001
	<b>IER</b>	33.52 [30.10, 36.94]	32.65 [29.23, 36.07]	30.48 [27.06, 33.90]	32.94 [29.38, 36.51]			
<b>Fat-free mass (kg)</b>	<b>CER</b>	46.88 [44.25, 49.51]	46.50 [43.87, 49.13]	45.63 [43.00, 48.26]	46.28 [43.63, 48.92]	<0.001	<0.001	0.051
	<b>IER</b>	46.57 [43.87, 49.27]	45.52 [42.82, 48.22]	45.19 [42.49, 47.89]	45.66 [42.92, 48.39]			
<b>Body fat (%)</b>	<b>CER</b>	40.73 [38.30, 43.15]	39.78 [37.36, 42.21]	38.52 [36.09, 40.94]	40.46 [37.97, 42.94]	0.568	<0.001	0.001
	<b>IER</b>	41.64 [39.15, 44.13]	41.58 [39.09, 44.08]	40.07 [37.58, 42.56]	41.36 [38.74, 43.98]			
<b>RMR (kcal/day)</b>	<b>CER</b>	1456 [1370, 1542]	1433 [1347, 1519]	1435 [1349, 1521]	1657 [1564, 1750]	1.000	1.000	<0.001
	<b>IER</b>	1435 [1346, 1523]	1459 [1371, 1548]	1478 [1389, 1566]	1638 [1533, 1742]			
<b>TDEE (kcal/day)</b>	<b>CER</b>	2352 [2214, 2489]	2311 [2173, 2448]	2263 [2125, 2400]	2376 [2232, 2520]	0.005	<0.001	0.015
	<b>IER</b>	2455 [2309, 2600]	2333 [2187, 2478]	2296 [2150, 2442]	2385 [2223, 2548]			
<b>AEE (kcal/day)</b>	<b>CER</b>	661 [568, 753]	647 [554, 740]	601 [509, 694]	472 [365, 579]	0.084	0.001	0.060
	<b>IER</b>	773 [675, 871]	639 [541, 737]	595 [495, 695]	514 [380, 647]			
<b>Total PA (min/day)</b>	<b>CER</b>	235 [194, 277]	244 [202, 285]	255 [214, 297]	226 [181, 271]	1.000	1.000	1.000
	<b>IER</b>	257 [213, 301]	241 [198, 285]	247 [203, 292]	239 [185, 293]			
<b>Sed time (min/day)</b>	<b>CER</b>	760 [714, 805]	739 [693, 784]	723 [677, 768]	791 [741, 841]	1.000	0.670	0.166
	<b>IER</b>	744 [696, 792]	739 [691, 787]	743 [695, 792]	746 [686, 806]			
<b>Light PA (min/day)</b>	<b>CER</b>	168 [137, 198]	173 [143, 204]	182 [152, 212]	141 [107, 174]	1.000	1.000	0.054
	<b>IER</b>	180 [148, 212]	177 [145, 209]	176 [143, 208]	162 [123, 202]			
<b>MVPA (min/day)</b>	<b>CER</b>	68 [53, 84]	72 [56, 87]	74 [58, 89]	86 [69, 104]	1.000	1.000	0.821
	<b>IER</b>	77 [61, 94]	65 [48, 81]	72 [56, 89]	77 [56, 98]			
<b>Steps per day</b>	<b>CER</b>	8623 [7380, 9865]	8715 [7473, 9958]	8455 [7213, 9698]	9643 [8266, 11019]	1.000	1.000	0.833
	<b>IER</b>	9262 [7949, 10576]	8469 [7155, 9782]	8578 [7251, 9905]	8712 [7060, 10364]			

<b>Sleep (min/day)</b>	<b>CER</b>	424 [398, 450]	437 [411, 463]	443 [417, 469]	405 [376, 434]	0.297	0.597	0.832
	<b>IER</b>	417 [389, 444]	433 [406, 461]	423 [395, 451]	430 [395, 466]			

616 Data are estimated marginal means [95% confidence interval]. AEE, activity energy expenditure; CER, continuous energy restriction; IER, intermittent energy  
617 restriction; MVPA, moderate-to-vigorous physical activity; RMR, resting metabolic rate; Sed time, sedentary time.

618

619

620