

This is a repository copy of *Effect of relative income price on smoking initiation among adolescents in Ghana:evidence from pseudo-longitudinal data.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/183861/>

Version: Accepted Version

Article:

Boachie, Micheal Kofi, Immurana, Mutsapha, Ngeh Tingum, Ernest et al. (2 more authors) (2022) Effect of relative income price on smoking initiation among adolescents in Ghana:evidence from pseudo-longitudinal data. *BMJ Open*. e054367. ISSN 2044-6055

<https://doi.org/10.1136/bmjopen-2021-054367>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

BMJ Open

Effect of relative income price on smoking initiation among adolescents in Ghana: evidence from pseudo-longitudinal data

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-054367.R2
Article Type:	Original research
Date Submitted by the Author:	22-Jan-2022
Complete List of Authors:	Boachie, Micheal Kofi; University of the Witwatersrand, Centre Health Economics and Decision Science - PRICELESS SA Immurana, Mustapha; University of Health and Allied Sciences, Institute of Health Research Tingum, Ernest; University of Cape Town, Economics Mdege, Noreen; University of York, Health Sciences Ross, Hana; University of Cape Town, School of Economics
Primary Subject Heading:	Global health
Secondary Subject Heading:	Health economics, Global health, Public health, Smoking and tobacco
Keywords:	HEALTH ECONOMICS, PUBLIC HEALTH, SOCIAL MEDICINE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3 **Effect of relative income price on smoking initiation among adolescents in Ghana:**
4
5 **evidence from pseudo-longitudinal data**
6
7

8 **Micheal Kofi Boachie^{1,2**}, Mustapha Immurana³, Ernest N. Tingum^{2,4}, Noreen Dadirai**
9 **Mdege⁵, Hana Ross²**
10
11

12
13
14 ¹ SAMRC/Wits Centre for Health Economics and Decision Science – PRICELESS SA,
15 School of Public Health, University of the Witwatersrand, Johannesburg 2193, South Africa;
16 mkboachie@gmail.com
17

18 ² Research Unit on the Economics of Excisable Products (REEP), School of Economics,
19 University of Cape Town, Cape Town, South Africa; hana.ross@uct.ac.za
20

21 ³ Institute of Health Research, University of Health and Allied Sciences, Ho, Ghana;
22 mimmurna@uhas.edu.gh; mustaphaimmurana@gmail.com
23
24

25 ⁴ Department of Economics, University of Namibia, Windhoek, Namibia;
26 ngehernest@gmail.com
27

28 ⁵ Department of Health Sciences, University of York, Heslington, United Kingdom;
29 noreen.mdege@york.ac.uk
30

31 **** Correspondence:** mkboachie@gmail.com; Tel.: +27 845 644 650.
32
33
34
35

36 **Word count:**
37

38
39 Abstract: 290
40

41 Introduction: 714
42

43 Materials and Methods: 1507
44

45 Results: 1330
46

47 Discussion and Conclusion: 508
48

49 Total word count: 4349
50

51 Number of References: 54
52
53
54
55
56
57
58
59
60

Abstract:

Objective: Many smokers initiate smoking during adolescence. Making tobacco products less affordable is one of the best ways to control tobacco use. Studies on the effect of relative income price (RIP (i.e., affordability)) of cigarettes on smoking initiation are scarce in low- and middle-income countries, especially in Sub-Saharan Africa where data are limited. The goal of this study is to examine the effect of cigarette RIP on adolescent smoking initiation in Ghana.

Setting: The study uses a pseudo-longitudinal dataset constructed from the Global Youth Tobacco Surveys (GYTS (2000-2009 and 2017)) and RIP for the most sold cigarette brand in Ghana.

Participants: The GYTS is a national survey on adolescents.

Primary and secondary outcome: Effect of RIP on adolescent smoking initiation in Ghana.

Results:

Using the GYTS 2000-2009 data, we find that the probability of smoking initiation falls significantly in response to a higher RIP, with an elasticity of -0.372 (95% CI: -0.701 to -0.042) for the unmatched sample and -0.490 (95% CI: -0.818 to -0.161) for the matched sample. The RIP elasticity for females [(-0.888) (95% CI: -1.384 to -0.392) and (-0.928) (95% CI: -1.434 to -0.422)] is statistically significant at 1% in both the unmatched and the matched samples, respectively, while the RIP elasticity for males is statistically insignificant in the 2000 - 2009 surveys. Analysis of the 2017 GYTS shows a similar outcome: a negative relationship between RIP and smoking initiation, and the results are statistically significant for both male and female, and for both matched and unmatched samples.

1
2
3
4
5
6 **Conclusion:** The affordability (RIP) of cigarettes is negatively related to the probability of
7
8 smoking initiation among adolescents in Ghana. Raising tobacco taxes in line with income
9
10 growth would make cigarettes less affordable and dissuade adolescents from initiating
11
12 smoking.
13
14

15
16 **Keywords:** prices; taxes; smoking initiation; adolescents; relative income price
17
18
19
20
21
22
23
24

25 **Strengths and Limitations of the study**

- 26
27 • This is the first study to explore the impact of Relative Income Price (i.e.,
28 affordability index) of cigarettes on youth smoking initiation in Sub-Saharan
29 Africa.
30
31
- 32 • Our analysis controls for variables that are known, empirically or theoretically, to
33 be associated with smoking initiation, and the relationship is tested using a pseudo-
34 longitudinal dataset of 17 years.
35
36
- 37 • We also examine potential sex differences in the effect of affordability on cigarette
38 smoking initiation: this is key to the implementation of tobacco control policies that
39 confer adequate protection across both genders.
40
41
- 42 • Since GYTS data are available in many low- and middle-income countries (LMIC),
43 our study provide a template on how to do analyses elsewhere in order to enhance
44 our understating of the impact of cigarette affordability on smoking uptake in
45 LMIC.
46
47
48
49
50
51
52
53
54
55
56
57

- The results are subject to self-reporting and recall biases as well as omitted variable bias due to lack of data on other factors affecting smoking uptake.

For peer review only

1. INTRODUCTION

Tobacco use is one of the major risk factors for many non-communicable diseases such as lung cancer and ischemic heart disease,^{1 2} and it accounts for over eight million deaths annually worldwide.³ Tobacco use also imposes huge financial burdens on households and governments.^{4 5} The majority (80-90%) of the one billion adult smokers globally began the habit during their adolescence.⁶ The current smoking prevalence in Ghana is still low relative to other African countries (3.2% among adults aged 15 years and older,⁷ and 6.4% among students aged 13-15 years⁸). However, the number of smokers is predicted to increase from 1.3 million to 1.7 million (i.e., by about 30%) between 2020 and 2025.⁹ The expected increase in the number of smokers will be partly driven by initiation among adolescents, therefore lowering initiation is key to slowing down the tobacco epidemic in Ghana.

Empirical evidence shows that tobacco consumption (initiation and intensity) is significantly inversely related to price.¹⁰⁻¹² In addition to being the most cost-effective measure to reduce tobacco use, increasing the taxes of tobacco products generates revenue for governments.¹³⁻¹⁵ Nevertheless, economic factors such as income growth can negatively affect the response of tobacco consumption to tax/price changes.^{16 17} Increasing tobacco prices can be more effective in reducing tobacco consumption if it reduces affordability.¹⁸ Affordability (relative income price (RIP)) elasticity which measures the sensitivity of consumers to real changes in both price and income, can therefore be a useful parameter to explain and predict the sensitivity of consumers to tobacco tax and price policies even in the presence of income growth.¹⁷ This is particularly important for tobacco control measures aimed at adolescents because they are more price sensitive than adults,^{12 19} for instance in Ghana where 71.3% of current cigarette smokers aged 13-15 buy their own cigarettes.⁸

1
2
3 Some Sub-Saharan African (SSA) studies, albeit few, have demonstrated that increasing
4 higher cigarette prices reduces smoking prevalence and intensity of use.^{12 20-25} However, there
5 is a scarcity of studies on the relationship between cigarette prices and smoking initiation in
6 the SSA context. One study, using data from 48 countries, including four from SSA (Kenya,
7 Nigeria, Senegal, and South Africa), concluded that higher cigarette prices reduce smoking
8 initiation in early youth, with girls being more responsive than boys.²⁶ However, findings
9 from the SSA countries were not reported separately from the overall study findings. A study
10 in South Africa reported a significant reduction in regular smoking initiation among males
11 due to higher cigarette prices, but not among females¹⁰. Another study in Nigeria and Ghana
12 concluded that increasing cigarette prices resulted in a reduction in both 30-day cigarette
13 smoking and cigarette smoking onset in both countries.²⁷

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Ghana has implemented a number of tobacco tax changes over the last 20 years. For example,
it implemented a specific excise tax in 2008 and subsequently switched to an ad valorem tax
structure in 2010.^{28 29} At the same time, per capita income in Ghana has been growing at an
average rate of 4.4% annually in the last decade.³⁰ These changes have implications for the
retail prices of tobacco products (e.g., cigarettes), and the affordability of cigarettes or other
tobacco products. However, to our knowledge, no study has analyzed the impact of cigarette
affordability on smoking initiation in Ghana.

We address this critical evidence gap by examining the effect of cigarette affordability on
smoking initiation among adolescents in Ghana. We hypothesize that making cigarettes less
affordable reduces the likelihood of smoking initiation among young people, and we make
use of the Global Youth Tobacco Survey (GYTS) data and other datasets to test that
hypothesis. The control variables used are sex, age, parents', and friends' smoking status,
being offered a cigarette for free, family/class discussion about tobacco, exposure to
antismoking messages, and exposure to tobacco advertisements. These controls are based on

1
2
3 variables that are known, empirically or theoretically, to be associated with smoking
4
5 initiation.^{27 31}
6
7

8 Our analysis addresses the potential endogeneity of price, if any, as a driver of cigarette
9 demand through i) using aggregate level prices and not self-reported prices³² and ii) the use
10 of propensity score matching techniques.^{33 34} We also examine potential sex differences in the
11 effect of affordability on cigarette smoking initiation. An understanding of these dynamics is
12 key to the implementation of context-specific tobacco and non-communicable disease control
13 policies in Ghana.
14
15
16
17
18
19
20
21
22

23 **2. MATERIALS AND METHODS**

24 **2.1. Data and variables**

25
26 We make use of three waves (2000, 2006 and 2009) of the GYTS and RIP (affordability) data
27 (1991–2009) to analyze the effect of affordability on smoking initiation among adolescents in
28 Ghana. The World Health Organization (WHO) defines adolescents as young people between
29 the ages of 10 and 19 years. The GYTS questionnaire specifies ages from 11 years or
30 younger to 17 years or higher for current age (i.e., age at the time of survey). It also captures
31 age at first puff, which ranges from 7 years or younger to 16 years or older. For the purposes
32 of this study, we classify respondents as adolescents, youth or young people. The terms are
33 used interchangeably in the study. The GYTS is a school-based survey developed to enhance
34 the capacity of countries to monitor tobacco use among the youth, as well as implement and
35 evaluate tobacco control and prevention programs.³⁵ These data provide representative trends
36 of tobacco use among adolescents. The GYTS is a cross-sectional survey and does not follow
37 individuals over time, but provides data on smoking patterns among adolescents. In countries
38 where it is conducted at regular intervals, it allows the monitoring of trends over time. We are
39 aware of the 2017 GYTS for Ghana, but we do not include it in the analysis of the pooled
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 2000 – 2009 surveys due to inconsistencies in the questions asked, compared to those in
4
5 previous GYTS surveys. There is no other survey on adolescents in Ghana with comparable
6
7 measures except the ones outlined. We analyze the 2017 GYTS separately while linking it
8
9 with RIP data for 2008 – 2017 based on age-at-risk criteria^{27 32 36} (as done for the 2000 –
10
11 2009 surveys).
12
13

14
15 Although the GYTS data contain adolescents whose first puff was at age 7 or younger, we
16
17 assume that a student is at risk at age 8 because that is the age at which the child is relatively
18
19 developed and is able to start out-of-home interaction with peers.^{27 32 36} Students who started
20
21 smoking before reaching age 8 and those below age 8 are therefore excluded from the pooled
22
23 sample and not followed. In line with previous studies, a student exits the sample once
24
25 smoking is initiated.^{10 27 32 33}
26
27
28
29

30 In Ghana, there was no law prohibiting the sale of cigarettes to minors until 2012 when
31
32 restrictions on sale to persons below age 18 years were introduced.³⁷ The GYTS sample is
33
34 drawn using a two-stage cluster-sampling design.^{35 38 39} Schools are selected with probability
35
36 proportional to school enrolment size during the first stage, and then classes within
37
38 participating schools are selected as a systematic equal probability sample with a random
39
40 start during the second stage. All students in the selected classes are eligible to participate in
41
42 the survey.
43
44
45

46 The Ghana GYTS questionnaire captures information on the use of tobacco products such as
47
48 cigarettes and shisha. The data also include information on parental and peer smoking habits,
49
50 perception about tobacco use (e.g., weight gain, health effects, and ease of quitting), money
51
52 spent on tobacco in the last 30 days before the survey, and second-hand smoking (SHS).⁴⁰
53
54 Studies vary widely on the way they define or measure smoking initiation.⁴¹⁻⁴³ For the GYTS,
55
56 smoking initiation is measured using the definition of a lifetime smoker, i.e., a person who
57
58
59
60

has ever tried smoking, *even one or two puffs of a cigarette*.^{27 43 44} Thus, for our study, smoking initiation (dependent variable) is a dichotomous variable generated from the following GYTS question where students answer Yes/No: *Have you ever tried or experimented with cigarette smoking, even one or two puffs?*.²⁷

The main independent variable is the affordability index or the RIP, measured as the percentage of GDP per capita (per capita income) required to buy 100 packs of cigarettes (20 sticks per pack, in total 2,000 sticks).^{17 18 45 46} Affordability is a relative measure and is calculated using nominal prices and nominal GDP per capita, or real prices and real GDP per capita. Data on per capita income are obtained from the World Bank's World Development Indicators,³⁰ and those of average cigarette prices (for the most-sold brand) come from the WHO, relevant publications of the tobacco industry⁴⁷ and the Government of Ghana⁴⁸. Years with missing data on prices were interpolated using the formula:

$$P_{t-1} = \frac{P_t}{(1 + Tob.Inflation_t)} \dots\dots\dots (1)$$

where P_{t-1} is the previous year's price of cigarette, P_t is current price of cigarette and $Tob.Inflation_t$ is the current tobacco inflation.⁴⁹ We then calculate RIP following methods used by preceding studies, with a lower affordability index (RIP) indicating that cigarettes have become more affordable and a higher value indicating that cigarettes have become less affordable relative to the reference year.^{17 18 45 46} The WHO uses the same approach to obtain its affordability index. Other independent variables used are sex, age, parents', and friends' smoking status, whether offered cigarettes for free, family/class discussion about tobacco, exposure to antismoking messages, and exposure to tobacco advertisements. These variables are selected as they have been shown, theoretically or empirically, to be associated with

1
2
3 smoking initiation.^{27 31} Except age and RIP, which are continuously measured, all variables
4
5 are dichotomous.
6
7

8 **2.2. Data analysis**

10 We construct a pseudo-longitudinal dataset based on current age and age at first puff. In
11
12 doing this, we create a historical dataset starting from age 8 (age-at-risk criteria) and follow
13
14 the person until s/he initiates smoking. This is done by inferring the year of first puff using
15
16 the GYTS question: “How old were you when you first tried a cigarette?” and the age at the
17
18 time of the survey.²⁷
19
20
21
22

23 STATA routine command, *expand*, is used to expand the person’s age at the time of the
24
25 survey after which an event variable indicating smoking status is created. We then link the
26
27 RIP (affordability index) data to this pseudo-longitudinal dataset.
28
29

30
31 Our statistical technique is duration or event history analysis where the timing of transition
32
33 into initiation is a function of the probability of initiating in period t conditional on not having
34
35 experienced a transition until period t , known as the hazard rate.¹⁰ Following previous
36
37 empirical studies,^{10 32 36} we employ the discrete time-hazard model, with logit specification
38
39 (see equation 2), to study the association between RIP (affordability) and smoking initiation
40
41 among adolescents.
42
43
44

$$45 \Pr(\textit{Initiation} = 1 | X') = \beta_0 + \beta_1 \textit{RIP} + \beta_i X' \dots (2)$$

46
47
48 where Initiation is defined as first cigarette puff, RIP is the affordability index, X' is a vector
49
50 of other independent variables affecting smoking initiation among adolescents, and β is a
51
52 vector of the regression coefficients. The predictors, X' , represent age, sex, whether offered
53
54 free cigarette, parental and peer smoking status, family/class discussion on the dangers of
55
56 tobacco, exposure to tobacco advertisements, and hearing of antismoking messages, and
57
58
59
60

1
2
3 awareness of tobacco control policies introduced in 2012 (for the 2017 GYTS). We report
4 odds ratios (ORs), and the statistical level of significance is set at $p \leq 0.1$. OR < 1 implies that
5 when a higher share of income is required to buy 2000 cigarettes (cigarettes are less
6 affordable), the risk of smoking initiation declines, and vice versa. The partial derivative of
7 equation 2 with respect to RIP gives the affordability elasticity.
8
9

10
11
12
13
14
15 To check the robustness of the logistic regression estimates, we employ a propensity score
16 matching (PSM) technique to match ever-smokers to never-smokers based on the propensity
17 scores. Our approach to matching follows previous studies.^{33 50 51} The propensity scores are
18 obtained by running a logit regression to estimate the probability of being a smoker based on
19 the variables in equation 2, except RIP, and the predicted probabilities are then used to match
20 ever-smokers to never-smokers. Using the neighborhood matching, ever-smokers are
21 matched to their two nearest neighbors. After matching the sample, we re-estimate the logit
22 model to assess the effect of affordability on the probability of initiating smoking, using
23 GYTS weights on the matched sample.^{50 51} With the matching approach, we are able to obtain
24 the effect of affordability on the probability of initiating smoking among adolescent smokers
25 and non-smokers who possess similar characteristics based on the propensity scores. This
26 technique addresses issues of endogeneity and concerns relating to the fact that some never-
27 smokers will never choose to smoke or use any form of tobacco irrespective of market
28 conditions.^{33 34} Further, we minimize the problem of endogeneity by not using self-reported
29 prices.³² Data analysis is conducted using STATA version 15. The study benefited immensely
30 from discrete time modelling guidelines and STATA code produced by Professor Stephen
31 Jenkins.⁵²
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 **2.3 Interpretation of RIP Elasticity**

55
56
57
58
59
60

1
2
3 Although the RIP is measured in percentages, interpretation of the affordability index follows
4 the same procedure for elasticity interpretation. The elasticity measures the percentage
5 change in probability of initiating smoking following a percentage change in RIP, *ceteris*
6 *paribus*. Assuming the current RIP is 6% or 0.06, then a 1% increase in RIP corresponds to
7 the current RIP increasing from 6% to 6.06%. When using a unit change interpretation, a unit
8 change will be RIP moving from 6% to 7% and therefore probabilities will change in
9 absolute units and not percentages. Such distinction is important in understanding the impact
10 of affordability on smoking behavior.
11
12
13
14
15
16
17
18
19
20
21

22 **Patient and Public Involvement**

23
24
25 No patient involved.
26
27
28
29
30

31 **3. RESULTS**

32 **3.1. Descriptive statistics**

33
34 A total of 20,202 adolescents were interviewed across the three GYTS waves (2000 – 2009).
35
36 54% of the respondents were males, while 76.47% were aged 15 years or less. In the 2017
37 GYTS, 5,664 people were interviewed, with about 48% being males. Overall, 12.46% and
38 8.9% of the respondents in the pooled (2000 – 2009) and 2017 surveys, respectively, had ever
39 smoked.
40
41
42
43
44
45
46
47
48

49 Given our age-at-risk criteria, 15,861 (2000 – 2009 GYTS) and 5,389 (2017 GYTS) people
50 were eligible for inclusion in our pseudo-longitudinal analysis. For surveys prior to 2017,
51 4.2% initiated smoking at some point between 1991 and 2009, and about 77% of smoking
52 initiators did so before reaching age 16. Further, 67% of initiators were males. Overall, males
53 represented 53.62% of the eligible respondents. In the 2017 survey, 4.72% of the respondents
54
55
56
57
58
59
60

initiated smoking at some point between 2008 and 2017. The characteristics of the samples are presented in Table 1. Due to incomplete information on some of the variables, the number of people used in the regression varies.

Table 1: Descriptive statistics

Variable	2000 – 2009 GYTS	2017 GYTS
	Students, n = 15 861	n=5389
Initiated smoking during the period	4.20%	4.72%
RIP (Affordability)	19.87 (SD=6.53)	7.63 (SD =0.86)
Offered free cigarettes	12.44%	8.13%
Sex (male =1)	53.62%	48.73%
At least one parent smoke	11.78%	-
Family/class discuss about tobacco	72.50%	51.47%
At least a friend smoke	15.94%	-
Exposed to tobacco adverts	40.46%	56.03%
Age (Years)	14.15 (SD=1.7)	14.10 (SD =1.03)
Heard anti-smoking campaigns	74.64%	57.26%
Age at initiation (Years)	11.95 (SD = 2.9)	11.26 (SD=2.41)
Percentage of initiators before age 16	77%	94%
Percentage of initiators who are males	67%	59%
Awareness of smoke free policies	-	78.24%

3.2. Regression Results

Results from the logit regressions for the unmatched and matched samples are reported in Tables 2 (GYTS 2000 – 2009) and 3 (GYTS 2017). The results show a statistically significant and negative relationship between RIP and smoking initiation. For instance, the results for the full unmatched sample (Table 2) show that OR on RIP is 0.98. This implies that a unit increase in affordability is associated with 0.98 odds of initiating smoking (OR = 0.98, $p < 0.05$). Thus, an adolescent who is subjected to a unit increase in RIP has 2% (i.e., $1 -$

0.98) $\times 100 = 2\%$) lower probability of initiating smoking than his/her counterpart who is not subjected to the same increase. Note that these results are not elasticities. Females have 40.1% (i.e., $1 - 0.599 \times 100 = 40.1\%$) lower probability of initiating smoking (OR = 0.599, $p < 0.01$) compared to their male counterparts in the unmatched sample.

Table 2: Effect of RIP on smoking initiation among adolescents (GYTS 2000 -2009)

VARIABLES	Unmatched Odds ratio	Matched Odds ratio
Affordability (RIP)	0.981** (0.009)	0.974*** (0.009)
Offered free cigarette (Ref=no)	1.491*** (0.216)	0.517*** (0.071)
Sex (Ref = male)	0.599*** (0.072)	0.615*** (0.076)
At least one parent smokes (Ref=no)	2.131*** (0.280)	0.862 (0.104)
Family/class discussion (Ref = no)	1.001 (0.133)	1.711*** (0.230)
At least one friend smokes (Ref=no)	4.109*** (0.520)	1.094 (0.126)
Exposure to adverts (Ref=no)	1.155 (0.140)	1.027 (0.121)
Age	1.150*** (0.042)	0.991 (0.031)
Heard of anti-smoking message/campaign (Ref=no)	1.342* (0.217)	2.048*** (0.321)
Survey cycle (ref=2000)		
2006	0.958 (0.146)	0.880 (0.138)
2009	1.108 (0.171)	1.003 (0.159)
Log (time)	1.110 (0.106)	1.393*** (0.146)
Constant	0.000*** (0.000)	0.048*** (0.024)
Observations	106,673	10,078
Number of people	15,201	1,611

Ever-smokers	611	611
Pseudo R-squared	0.0815	0.0448
Chi2	439.2***	91.84***

Robust standard errors in parentheses

*** p<0.01, ** p<0.05, * p<0.1

Table 3: **Effect of RIP on smoking initiation among adolescents (GYTS 2017)**

VARIABLES	Unmatched Odds ratio	Matched Odds ratio
Affordability (RIP)	0.821** (0.066)	0.804*** (0.065)
Sex (Ref=male)	0.659** (0.120)	0.902 (0.177)
Offered free cigarettes (Ref= no)	3.403*** (0.726)	0.978 (0.221)
Heard of anti-smoking message (Ref=no)	1.165 (0.213)	1.009 (0.192)
Exposed to tobacco adverts (Ref= no)	3.030*** (0.622)	1.893*** (0.421)
Smoke free policies awareness (Ref=no)	1.250 (0.329)	1.160 (0.332)
Age	1.847*** (0.294)	1.793*** (0.279)
Class discussion on tobacco harms (Ref=no)	0.795 (0.138)	1.278 (0.241)
Log(time)	0.104*** (0.067)	0.119*** (0.076)
Constant	0.000*** (0.000)	0.002*** (0.002)
Observations	37,654	4,850
Number of people	5,301	747
Ever-smokers	231	206
Pseudo R-squared	0.0599	0.0292
Chi2	158***	38.72***

Robust standard errors in parentheses

*** p<0.01, ** p<0.05, * p<0.1

1
2
3 Similarly, in Table 3, in the unmatched sample, an adolescent faced with a unit increase in
4 RIP has about 18% (i.e., $1 - 0.821 \times 100 = 17.9\%$) lower probability of starting smoking
5
6 than his/her counterpart who is not subjected to the same increase.
7
8
9

10 Other significant factors that influence smoking initiation in our two samples include whether
11 the adolescent's parents (OR = 2.131, $p < 0.01$) or friends (OR = 4.109, $p < 0.01$) smoke. In
12 addition, adolescents who are offered free cigarettes have a high probability of initiating
13 smoking (OR = 1.491, $p < 0.01$) compared to those who receive no such offer for the 2000 -
14 2009 wave (Table 2). In the 2017 wave, the odds of adolescents starting smoking when given
15 cigarettes freely is 3.403 ($p < 0.01$) (Table 3).
16
17
18
19
20
21
22
23
24

25 In the matched sample (Table 2), 611 ever-smokers were matched to their two nearest
26 neighbors (1000 never-smokers) which produced a sample of 1,611 adolescents with similar
27 characteristics. An adolescent subjected to a unit increase in RIP has a 0.97 times lower
28 chance of initiating smoking (OR = 0.974, $p < 0.01$) compared to those not exposed to the
29 same increase in RIP. Similarly in the matched sample of the GYTS 2017 (Table 3), a unit
30 increase in RIP is associated with 0.80 times lower odds of smoking initiation (OR = 0.804,
31 $p < 0.01$).
32
33
34
35
36
37
38
39
40
41

42 Getting free cigarettes (OR = 0.517, $p < 0.01$), family/class discussion on tobacco (OR =
43 1.711, $p < 0.01$), and hearing antismoking messages (OR = 2.048, $p < 0.01$) are all found to be
44 statistically significant in influencing smoking initiation in the matched sample (Table 2).
45
46
47
48

49 However, the odds for these variables are contrary to *a priori* expectations. Similarly, in the
50 matched sample (Table 2), the likelihood of initiating smoking among females is lower. The
51 results imply that females have about 39% lower probability of initiating smoking (OR =
52 0.615, $p < 0.01$) than males.
53
54
55
56
57
58

59 3.3. Affordability Elasticities 60

In the unmatched sample, the estimated affordability elasticity is -0.372 [CI: -0.701 to -0.042] for the 2000-2009 sample and -1.247 [-2.248 to -0.246] for the 2017 sample. These elasticities are statistically significant at the 5% level. By sex, the affordability elasticity is -0.137 for males and -0.888 for females for the 2000-2009 sample, but only that of females is statistically significant ($p < 0.01$). The elasticities are higher for 2017 GYTS (-0.938 for males and -1.610 for females). The results are presented in Table 4.

Table 4: Affordability Elasticity estimates

	Both sexes		Male		Female	
Panel A: 2000 -2009						
	Percentage changes (d(lny)/d(lnx))					
VARIABLES	Unmatched	Matched	Unmatched	Matched	Unmatched	Matched
Affordability	-0.372** (0.168)	-0.490*** (0.168)	-0.137 (0.219)	-0.326 (0.216)	-0.888*** (0.253)	-0.928*** (0.258)
95% CI	-0.701 to -0.042	-0.818 to -0.161	-0.567 to 0.292	-0.749 to 0.097	-1.384 to -0.392	-1.434 to -0.422
Observations	106,673	10,078	55,396	5,648	51,277	4,430
Panel B: 2017 GYTS						
Affordability	-1.247** (0.511)	-1.349*** (0.500)	-0.938** (0.474)	-1.045** (0.484)	-1.610* (0.866)	1.518** (0.778)
95% CI	-2.248 to -0.246	-2.328 to -0.369	-1.867 to -0.008	-1.993 to -0.096	-3.307 to -0.087	-3.043 to -0.007
Observations	37,654	4,850	18,084	2,807	19,570	2,043

Standard errors in parentheses

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

In the matched sample for the 2000-2009 GYTS, the overall elasticity is -0.490 [CI: -0.818 to -0.161] for both sexes (Table 4), which is similar to that of the unmatched sample. For males, the effect of changes in RIP is statistically insignificant. Among females, a percentage increase in RIP is associated with a 0.928% lower probability of smoking initiation. The elasticities for both males and females in the 2017 GYTS were negative, statistically significant, and more than unity.

1
2
3 The mean and median standardized difference for the covariates used in matching show that
4 the matching satisfies the balancing test (results not reported). The mean and median
5 standardized difference is 2.3%, which is below the normal 10% threshold. Therefore, the
6 balancing property is satisfied.
7
8
9
10
11

12 **5. DISCUSSION AND CONCLUSION**

13
14
15
16 In this study, increasing the RIP of cigarettes is significantly associated with a lower
17 probability of initiating smoking. This finding is consistent with international literature,
18 including the few existing studies in SSA that have reported that making cigarettes less
19 affordable lowers the likelihood of smoking initiation among young people.^{17 18 26} In addition,
20 the results from the unmatched 2000-2009 sample suggest that males are not responsive to
21 changes in RIP whilst females are. Nevertheless, in the matched sample analysis, especially
22 using the 2017 GYTS, both males and females are responsive to changes in RIP. Indeed, the
23 issue of affordability becomes more important given that Ghana's per capita income has been
24 growing at an average of 4.4% annually in the last decade.³⁰
25
26
27
28
29
30
31
32
33
34
35

36
37 Parental smoking increased the odds of smoking initiating. This points to the parental
38 influence on the lifestyle of adolescents. Adolescents whose parents smoke may perceive
39 smoking as acceptable behavior. Previous studies have reported similar findings.^{10 27 31 53} The
40 odds of smoking initiation are higher for those whose friends' smoke. This points to the
41 influence of peers, and is consistent, for example, with Mak, et al.³¹. Those who were offered
42 free cigarettes by sales representatives were more likely to initiate smoking. All forms of
43 tobacco promotion and advertising were banned in Ghana in 1982. However, the tobacco
44 industry seems to be breaking these laws, because 12.44% of youth reported being offered a
45 cigarette for free. This observation, together with our results, suggest the need to strengthen
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the enforcement of the existing ban on all forms of tobacco advertising and promotion in
4
5 Ghana.
6
7

8 This study has several limitations. The GYTS is a self-reporting survey, which means the
9
10 responses are prone to recall bias even in cases where the adolescents are required to answer
11
12 questions about events that occurred in the past 30 days. For instance, students may not recall
13
14 the exact age at which they tried their first puff. There is also a social desirability bias when
15
16 self-reporting behaviors such as smoking, especially among females. The weakness of our
17
18 measure of smoking initiation is that it may not predict regular smoking behavior well.^{42 43} In
19
20 addition, there are other important factors affecting smoking uptake among the youth that are
21
22 not included in this study. For instance, changing community norms regarding smoking, the
23
24 enforcement of laws regarding the sale of cigarettes to minors, and changes in the social
25
26 image of smoking are key factors that may influence smoking participation,^{54 55} but are not
27
28 included in the models estimated.
29
30
31
32
33

34 In conclusion, making cigarettes less affordable is associated with a lower probability of
35
36 smoking initiation among adolescents in Ghana. This supports the use of price measures,
37
38 through higher excise taxes, as effective strategies to decrease smoking initiation among
39
40 adolescents in Ghana. Since incomes are rising at the average of 4.4% annually³⁰, tobacco
41
42 taxes need to be adjusted regularly to ensure that cigarettes or other tobacco products become
43
44 less affordable over time in order to discourage young people from initiating smoking and to
45
46 encourage smokers to quit.
47
48
49
50

51 **Author Contributions:** Conceptualization, M.K.B.; methodology, M.K.B.; software, M.K.B.;
52 validation, M.K.B., M.I., E.N.T., N.D.M., and H.R.; formal analysis, M.K.B.; investigation,
53 M.K.B., M.I., E.N.T., N.D.M., and H.R.; data curation, M.K.B.; writing—original draft
54 preparation, M.K.B., M.I., E.N.T., N.D.M., and H.R.; writing—review and editing, M.K.B.,
55 M.I., E.N.T., N.D.M., H.R.; project administration, M.K.B.; supervision, H.R.; funding
56 acquisition, H.R. All authors have read and approved the manuscript.
57
58
59
60

Funding: Funding for this study was provided by Cancer Research UK (Grant number: C62640/A24723) and International Development Research Centre (Grant Number: 108820-001) with additional support from the SAMRC/Wits Centre for Health Economics and Decision Science – PRICELESS SA (grant number 23108). Funding for ENT and NDM is acknowledged from the Tobacco Control Capacity Programme (Grant Number: MR/P027946/2) supported by UK Research and Innovation (UKRI) with funding from the Global Challenges Research Fund (GCRF). The funders had no role in study design, data collection and analysis, or preparation of the manuscript.

Conflicts of Interest: The authors declare that they have no conflict of interest.

Research Ethics Approval: Human Participants

This study does not involve human participants.

Research Ethics Approval: Animals

This study does not involve animal subjects.

Patient consent for publication: No patient involved.

Data Availability statement: The publicly available data can be accessed:

<https://nccd.cdc.gov/GTSSDataSurveyResources/Ancillary/DataReports.aspx?CAID=2>. The relative income price data are available from the authors on reasonable request.

REFERENCES

1. Doll R, Peto R, Boreham J, et al. Mortality in relation to smoking: 50 years' observations on male British doctors. *Bmj* 2004;328(7455):1519.
2. Winkler V, Mangolo NJ, Becher H. Lung cancer in South Africa: a forecast to 2025 based on smoking prevalence data. *BMJ open* 2015;5(3):e006993.

- 1
2
3
4 3. World Health Organization. Tobacco: Key Facts. 2020 [updated 27 May 2020. Available
5
6 from: <https://www.who.int/news-room/fact-sheets/detail/tobacco> accessed 22
7
8 November 2020.
- 9
10 4. Goodchild M, Nargis N, d'Espaignet ET. Global economic cost of smoking-attributable
11
12 diseases. *Tobacco Control* 2018;27(1):58-64.
- 13
14 5. Boachie MK, Rossouw L, Ross H. The Economic Cost of Smoking in South Africa, 2016.
15
16 *Nicotine Tob Res* 2021;23(2):286 - 93. doi: 10.1093/ntr/ntaa162
- 17
18 6. USDHHS. Preventing tobacco use among youth and
19
20 young adults: A report of the surgeon general 2012 [Available from:
21
22 [http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-](http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf)
23
24 [report.pdf](http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf) (accessed: Jan 3, 2019).
- 25
26 7. World Health Organization. WHO global report on trends in prevalence of tobacco use 2000-
27
28 2025. 3rd ed. Geneva: World Health Organization, 2019.
- 29
30 8. CDC, WHO, MOH. GLOBAL YOUTH TOBACCO SURVEY: Factsheet, Ghana 2017,
31
32 2018.
- 33
34 9. World Health Organization. WHO global report on trends in prevalence of tobacco smoking
35
36 2015. Geneva: World Health Organization, 2015.
- 37
38 10. Vellios N, Van Walbeek C. Determinants of regular smoking onset in South Africa using
39
40 duration analysis. *BMJ Open* 2016;6(7):e011076.
- 41
42 11. IARC. Effectiveness of tax and price policies for tobacco control. In: Cancer. IAFRo, ed.
43
44 USA: International Agency for Research on Cancer, 2011.
- 45
46 12. Boachie MK, Ross H. Determinants of smoking intensity in South Africa: evidence from
47
48 township communities. *Preventive Medicine Reports* 2020;19:101099.
- 49
50 13. Van Walbeek C. A simulation model to predict the fiscal and public health impact of a
51
52 change in cigarette excise taxes. *Tobacco control* 2010;19(1):31-36.
- 53
54 14. Yurekli A, de Beye J. Design and administer tobacco taxes. *World Bank economics of*
55
56 *tobacco toolkit* 2001(4)
- 57
58 15. Van Walbeek C. Raising Additional Government Revenues in Ghana by Raising the Excise
59
60 Tax on Tobacco and Alcohol Washington, D.C.: WBG Global Tobacco Control

- 1
2
3
4 Program, World Bank Group; 2014 [Available from:
5 [http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-](http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-and-Alcohol.pdf)
6 [Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-](http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-and-Alcohol.pdf)
7 [and-Alcohol.pdf](http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-and-Alcohol.pdf) accessed 15 July 2019.
8
9
10
11
12 16. He Y, Shang C, Chaloupka FJ. The association between cigarette affordability and
13 consumption: An update. *PLOS ONE* 2018;13(12):e0200665. doi:
14 10.1371/journal.pone.0200665
15
16
17 17. Nargis N, Stoklosa M, Shang C, et al. Price, Income, and Affordability as the Determinants
18 of Tobacco Consumption: A Practitioner's Guide to Tobacco Taxation. *Nicotine and*
19 *Tobacco Research* 2021;23(1):40-47.
20
21
22 18. Blecher E, Van Walbeek C. An international analysis of cigarette affordability. *Tobacco*
23 *Control* 2004;13(4):339-46.
24
25
26 19. Ding A. Youth are more sensitive to price changes in cigarettes than adults. *The Yale*
27 *journal of biology and medicine* 2003;76(3):115.
28
29
30 20. Ho L-M, Schafferer C, Lee J-M, et al. The effect of cigarette price increases on cigarette
31 consumption, tax revenue, and smoking-related death in Africa from 1999 to 2013.
32 *International journal of public health* 2017;62(8):899-909.
33
34
35 21. Chelwa G, van Walbeek C. Does cigarette demand respond to price increases in Uganda?
36 Price elasticity estimates using the Uganda National Panel Survey and Deaton's method.
37 *BMJ open* 2019;9(3):e026150. doi: 10.1136/bmjopen-2018-026150 [published Online
38 First: 2019/03/23]
39
40
41 22. Stoklosa M, Goma F, Nargis N, et al. Price, tax and tobacco product substitution in Zambia:
42 findings from the ITC Zambia Surveys. *Tob Control* 2019;28(Suppl 1):s45-s52. doi:
43 10.1136/tobaccocontrol-2017-054037 [published Online First: 2018/03/27]
44
45
46 23. Mukong AK, Tingum EN. The demand for cigarettes: New evidence from South Africa.
47 *Development Southern Africa* 2020;37(1):40-54.
48
49
50 24. Tingum EN, Mukong AK, Mdege N. The effects of price and non-price policies on cigarette
51 consumption in South Africa. *Tob Induc Dis* 2020;18(July) doi: 10.18332/tid/123424
52
53
54
55
56
57
58
59
60

- 1
2
3
4 25. Kidane A, Mduma J, Naho A, et al. The Demand for Cigarettes in Tanzania and
5 Implications for Tobacco Taxation Policy. *Advances in economics and business*
6 2015;3(10):428.
7
8
9
10 26. Kostova D. A (nearly) global look at the dynamics of youth smoking initiation and cessation:
11 the role of cigarette prices. *Applied Economics* 2013;45(28):3943-51.
12
13
14 27. Asare S, Stoklosa M, Drope J, et al. Effects of Prices on Youth Cigarette Smoking and
15 Tobacco Use Initiation in Ghana and Nigeria. *International Journal of Environmental*
16 *Research and Public Health* 2019;16(17):3114.
17
18
19
20 28. Ministry of Health. Needs assessment for the Implementation of WHO FCTC in Ghana.
21 Accra: Ministry of Health,, 2010.
22
23
24 29. Gov't. of Ghana. Customs and Excise (Duties and Other Taxes) Amendment Act 739. ,
25 2007.
26
27
28 30. World Bank. World Development Indicators.: World Bank; 2019 [Available from:
29 <https://databank.worldbank.org/source/world-development-indicators>. accessed 01 Jan
30 2020.
31
32
33
34 31. Mak K-K, Ho S-Y, Day JR. Smoking of parents and best friend—-independent and
35 combined effects on adolescent smoking and intention to initiate and quit smoking.
36 *Nicotine Tob Res* 2012;14(9):1057-64.
37
38
39
40 32. Guindon GE, Paraje GR, Chávez R. Prices, inflation, and smoking onset: the case of
41 Argentina. *Economic Inquiry* 2018;56(1):424-45.
42
43
44 33. Dauchy E, Ross H. The Effect of Price and Tax Policies on the Decision to Smoke among
45 Men in Kenya. *Addiction* 2019;114(7):1249-63.
46
47
48 34. Li M. Using the propensity score method to estimate causal effects: A review and practical
49 guide. *Organizational Research Methods* 2013;16(2):188-226.
50
51
52 35. Warren CW. Tobacco use among youth: a cross country comparison. *Tobacco Control*
53 2002;11(3):252-70.
54
55
56 36. Boachie MK, Immurana M, Agyemang JK, et al. Cigarette prices and smoking
57 experimentation in Sierra Leone: an exploratory study. *Tobacco Use Insights*
58 2022;forthcoming
59
60

- 1
2
3
4 37. Gov't. of Ghana. Public Health Act, 2012 Act 851. Ghana 2012.
5
6 38. Wellington E. Ghana Global Youth Tobacco Survey
7
8 (GYTS), Factsheet. 2006 [Available from:
9
10 <http://www.afro.who.int/en/downloads/docdownload/1887-ghana-gyts-2006->
11
12 [factsheet.html](http://www.afro.who.int/en/downloads/docdownload/1887-ghana-gyts-2006-factsheet.html) (Retrieved Jan 28, 2019).
13
14 39. Cadmus EO, Ayo-Yusuf OA. The effect of smokeless tobacco use and exposure to cigarette
15 promotions on smoking intention among youths in Ghana. *Cogent Medicine*
16 2018;5(1):1531459.
17
18 40. CDC and WHO. Ghana global youth tobacco survey. Atlanta, United States: Centers for
19 Disease Control and Prevention (CDC). 2006.
20
21 41. Freedman KS, Nelson NM, Feldman LL. Smoking initiation among young adults in the
22 United States and Canada, 1998-2010: a systematic review. *Preventing chronic disease*
23 2012;9
24
25 42. Azagba S, Baskerville NB, Minaker L. A comparison of adolescent smoking initiation
26 measures on predicting future smoking behavior. *Preventive medicine reports*
27 2015;2:174-77.
28
29 43. Reidpath DD, Davey TM, Kadirvelu A, et al. Does one cigarette make an adolescent smoker,
30 and is it influenced by age and age of smoking initiation? Evidence of association from
31 the US Youth Risk Behavior Surveillance System (2011). *Preventive medicine*
32 2014;59:37-41.
33
34 44. Centers for Disease Control Prevention. Selected cigarette smoking initiation and quitting
35 behaviors among high school students--United States, 1997. *MMWR Morbidity and*
36 *mortality weekly report* 1998;47(19):386.
37
38 45. Blecher E. Targeting the affordability of cigarettes: a new benchmark for taxation policy
39 in low-income and-middle-income countries. *Tobacco control* 2010;19(4):325-30.
40
41 46. Nargis N, Stoklosa M, Drope J, et al. Trend in the affordability of tobacco products in
42 Bangladesh: findings from the ITC Bangladesh Surveys. *Tobacco control*
43 2019;28(Suppl 1):s20-s30.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 47. Philip Morris International. World summary of cigarette retail prices, tax burdens and tax
5 incidence.: Philips Moris Records 1996 [Available from:
6 <https://www.industrydocuments.ucsf.edu/docs/#id=glhf0066>.
7
8
9
- 10 48. Ghana Gto. Report card on the WHO Framework Convention on Tobacco Control., 2009.
11
- 12 49. Ghana Statistical Service. Consumer Price Index (CPI): Time Series In: Service GS, ed.
13 Accra, 2012.
14
- 15 50. Austin PC, Jembere N, Chiu M. Propensity score matching and complex surveys. *Statistical*
16 *methods in medical research* 2018;27(4):1240-57.
17
- 18 51. DuGoff EH, Schuler M, Stuart EA. Generalizing observational study results: applying
19 propensity score methods to complex surveys. *Health services research*
20 2014;49(1):284-303.
21
- 22 52. Jenkins SP. Survival Analysis with Stata. [Available from:
23 <https://www.iser.essex.ac.uk/resources/survival-analysis-with-stata> accessed 01
24 January 2020.
25
- 26 53. Gilman SE, Rende R, Boergers J, et al. Parental smoking and adolescent smoking initiation:
27 an intergenerational perspective on tobacco control. *Pediatrics* 2009;123(2):e274-e81.
28
- 29 54. Gilpin EA, Pierce JP. Trends in adolescent smoking initiation in the United States: is
30 tobacco marketing an influence? *Tobacco Control* 1997;6(2):122-27.
31
- 32 55. Cawley J, Markowitz S, Tauras J. Obesity, cigarette prices, youth access laws and
33 adolescent smoking initiation. *Eastern Economic Journal* 2006;32(1):149-70.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5-7
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	7
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7-8
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	7-9
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	8-9
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	8-9
Bias	9	Describe any efforts to address potential sources of bias	10
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	9
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	9-10
		(b) Describe any methods used to examine subgroups and interactions	9-10
		(c) Explain how missing data were addressed	9-10
		(d) If applicable, describe analytical methods taking account of sampling strategy	9-10
		(e) Describe any sensitivity analyses	11
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	12
		(b) Give reasons for non-participation at each stage	
		(c) Consider use of a flow diagram	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	13
		(b) Indicate number of participants with missing data for each variable of interest	
Outcome data	15*	Report numbers of outcome events or summary measures	
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	13-17

		(b) Report category boundaries when continuous variables were categorized	
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	
Discussion			
Key results	18	Summarise key results with reference to study objectives	18-19
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	19
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	11-12
Generalisability	21	Discuss the generalisability (external validity) of the study results	19
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	20

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

1
2
3 **Effect of relative income price on smoking initiation among adolescents in Ghana:**
4
5 **evidence from pseudo-longitudinal data**
6
7

8 **Micheal Kofi Boachie^{1,2**}, Mustapha Immurana³, Ernest N. Tingum^{2,4}, Noreen Dadirai**
9 **Mdege⁵, Hana Ross²**
10
11

12
13
14 ¹ SAMRC/Wits Centre for Health Economics and Decision Science – PRICELESS SA,
15 School of Public Health, University of the Witwatersrand, Johannesburg 2193, South Africa;
16 mkboachie@gmail.com
17

18
19 ² Research Unit on the Economics of Excisable Products ([REEP](#)), School of Economics,
20 University of Cape Town, Cape Town, South Africa; hana.ross@uct.ac.za
21

22 ³ Institute of Health Research, University of Health and Allied Sciences, Ho, Ghana;
23 mimmurna@uhas.edu.gh; mustaphaimmurana@gmail.com
24

25 ⁴ Department of Economics, University of Namibia, Windhoek, Namibia;
26 ngehernest@gmail.com
27

28 ⁵ Department of Health Sciences, University of York, Heslington, United Kingdom;
29 noreen.mdege@york.ac.uk
30

31 **** Correspondence:** mkboachie@gmail.com; Tel.: +27 845 644 650.
32
33
34
35

36 **Word count:**
37

38
39 Abstract: 290
40

41 Introduction: ~~71441~~
42

43 Materials and Methods: ~~15047~~
44

45 Results: 1330
46

47 Discussion and Conclusion: 508
48

49 Total word count: 43~~4986~~
50

51 Number of References: 54
52
53
54
55
56
57
58
59
60

Abstract:

Objective: Many smokers initiate smoking during adolescence. Making tobacco products less affordable is one of the best ways to control tobacco use. Studies on the effect of relative income price (RIP (i.e., affordability)) of cigarettes on smoking initiation are scarce in low- and middle-income countries, especially in Sub-Saharan Africa where data are limited. The goal of this study is to examine the effect of cigarette RIP on adolescent smoking initiation in Ghana.

Setting: The study uses a pseudo-longitudinal dataset constructed from the Global Youth Tobacco Surveys (GYTS (2000-2009 and 2017)) and RIP for the most sold cigarette brand in Ghana.

Participants: The GYTS is a national survey on adolescents.

Primary and secondary outcome: Effect of RIP on adolescent smoking initiation in Ghana.

Results:

Using the GYTS 2000-2009 data, we find that the probability of smoking initiation falls significantly in response to a higher RIP, with an elasticity of -0.372 (95% CI: -0.701 to -0.042) for the unmatched sample and -0.490 (95% CI: -0.818 to -0.161) for the matched sample. The RIP elasticity for females [(-0.888) (95% CI: -1.384 to -0.392) and (-0.928) (95% CI: -1.434 to -0.422)] is statistically significant at 1% in both the unmatched and the matched samples, respectively, while the RIP elasticity for males is statistically insignificant in the 2000 - 2009 surveys. Analysis of the 2017 GYTS shows a similar outcome: a negative relationship between RIP and smoking initiation, and the results are statistically significant for both male and female, and for both matched and unmatched samples.

1
2
3
4
5
6 **Conclusion:** The affordability (RIP) of cigarettes is negatively related to the probability of
7
8 smoking initiation among adolescents in Ghana. Raising tobacco taxes in line with income
9
10 growth would make cigarettes less affordable and dissuade adolescents from initiating
11
12 smoking.
13
14

15
16 **Keywords:** prices; taxes; smoking initiation; adolescents; relative income price
17
18
19
20
21
22
23
24

25 **Strengths and Limitations of the study**

- 26
27 • This is the first study to explore the impact of Relative Income Price (i.e.,
28 affordability index) of cigarettes on youth smoking initiation in Sub-Saharan
29
30 Africa.
31
32
- 33
34 • Our analysis controls for variables that are known, empirically or theoretically, to
35
36 be associated with smoking initiation, and the relationship is tested using a pseudo-
37
38 longitudinal dataset of 17 years.
39
40
- 41
42 • We also examine potential sex differences in the effect of affordability on cigarette
43
44 smoking initiation: this is key to the implementation of tobacco control policies that
45
46 confer adequate protection across both genders.
47
48
- 49
50 • Since GYTS data are available in many low- and middle-income countries (LMIC),
51
52 our study provide a template on how to do analyses elsewhere in order to enhance
53
54 our understating of the impact of cigarette affordability on smoking uptake in
55
56 LMIC.
57
58
59
60

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
- The results are subject to self-reporting and recall biases as well as omitted variable bias due to lack of data on other factors affecting smoking uptake.

For peer review only

1. INTRODUCTION

Tobacco use is one of the major risk factors for many non-communicable diseases such as lung cancer and ischemic heart disease,^{1 2} and it accounts for over eight million deaths annually worldwide.³ Tobacco use also imposes huge financial burdens on households and governments.^{4 5} The majority (80-90%) of the one billion adult smokers globally began the habit during their adolescence.⁶ The current smoking prevalence in Ghana is still low relative to other African countries (3.2% among adults aged 15 years and older,⁷ and 6.4% among students aged 13-15 years⁸). However, the number of smokers is predicted to increase from 1.3 million to 1.7 million (i.e., by about 30%) between 2020 and 2025.⁹ The expected increase in the number of smokers will be partly driven by initiation among adolescents, therefore lowering initiation is key to slowing down the tobacco epidemic in Ghana.

Empirical evidence shows that tobacco consumption (initiation and intensity) is significantly inversely related to price.¹⁰⁻¹² In addition to being the most cost-effective measure to reduce tobacco use, increasing the taxes of tobacco products generates revenue for governments.¹³⁻¹⁵ Nevertheless, economic factors such as income growth can negatively affect the response of tobacco consumption to tax/price changes.^{16 17} Increasing tobacco prices can be more effective in reducing tobacco consumption if it reduces affordability.¹⁸ Affordability (relative income price (RIP)) elasticity which measures the sensitivity of consumers to real changes in both price and income, can therefore be a useful parameter to explain and predict the sensitivity of consumers to tobacco tax and price policies even in the presence of income growth.¹⁷ This is particularly important for tobacco control measures aimed at adolescents because they are more price sensitive than adults,^{12 19} for instance in Ghana where 71.3% of current cigarette smokers aged 13-15 buy their own cigarettes.⁸

1
2
3 Some Sub-Saharan African (SSA) studies, albeit few, have demonstrated that increasing
4
5 higher cigarette prices reduces smoking prevalence and intensity of use.^{12 20-25} However, there
6
7 is a scarcity of studies on the relationship between cigarette prices and smoking initiation in
8
9 the SSA context. One study, using data from 48 countries, including four from SSA (Kenya,
10
11 Nigeria, Senegal, and South Africa), concluded that higher cigarette prices reduce smoking
12
13 initiation in early youth, with girls being more responsive than boys.²⁶ However, findings
14
15 from the SSA countries were not reported separately from the overall study findings. A study
16
17 in South Africa reported a significant reduction in regular smoking initiation among males
18
19 due to higher cigarette prices, but not among females¹⁰. Another study in Nigeria and Ghana
20
21 concluded that increasing cigarette prices resulted in a reduction in both 30-day cigarette
22
23 smoking and cigarette smoking onset in both countries.²⁷

24
25
26
27
28
29 Ghana has implemented a number of tobacco tax changes over the last 20 years. For example,
30
31 it ~~implemented~~introduced a specific excise tax in 2008⁷ and subsequently switched to an ad
32
33 valorem tax structure in 2010.^{28 29} At the same time, per capita income in Ghana has been
34
35 growing at an average rate of 4.4% annually in the last decade.³⁰ These changes have
36
37 implications for the retail prices of tobacco products (e.g., cigarettes), and the affordability of
38
39 cigarettes or other tobacco products. However, to our knowledge, no study has analyzed the
40
41 impact of cigarette affordability on smoking initiation in Ghana.

42
43
44
45
46 We address this critical evidence gap by examining the effect of cigarette affordability on
47
48 smoking initiation among adolescents in Ghana ~~using~~. We hypothesize that making cigarettes
49
50 less affordable reduces the likelihood of smoking initiation among young people, and ~~we~~
51
52 make use of the Global Youth Tobacco Survey (GYTS) data and other datasets to test that
53
54 hypothesis. The control variables used are sex, age, parents', and friends' smoking status,
55
56 being offered a cigarette for free, family/class discussion about tobacco, exposure to
57
58 antismoking messages, and exposure to tobacco advertisements. These controls are based on
59
60

1
2
3 variables that are known, empirically or theoretically, to be associated with smoking
4
5 initiation.^{27 31}

6
7
8 -Our analysis addresses the potential endogeneity of price, if any, as a driver of cigarette
9
10 demand through i) using aggregate level prices and not self-reported prices³²; and ii) the use
11
12 of propensity score matching techniques. ~~and iii) the fact that some people will never choose~~
13
14 ~~to smoke or use any form of tobacco, for example, for reasons of health or religious belief~~^{33 34}

15
16
17 We also examine potential sex differences in the effect of affordability on cigarette smoking
18
19 initiation. An understanding of these dynamics is key to the implementation of context-
20
21 specific tobacco and non-communicable disease control policies in Ghana.
22
23

24 25 **2. MATERIALS AND METHODS**

26 27 28 **2.1. Data and variables**

29
30
31 We make use of three waves (2000, 2006 and 2009) of the GYTS and RIP (affordability) data
32
33 (1991–2009) to analyze the effect of affordability on smoking initiation among adolescents in
34
35 Ghana. The World Health Organization (WHO) defines adolescents as young people between
36
37 the ages of 10 and 19 years. The GYTS questionnaire specifies ages from 11 years or
38
39 younger to 17 years or higher for current age (i.e., age at the time of survey). It also captures
40
41 age at first puff, which ranges from 7 years or younger to 16 years or older. For the purposes
42
43 of this study, we classify respondents as adolescents, youth or young people. The terms are
44
45 used interchangeably in the study. The GYTS is a school-based survey developed to enhance
46
47 the capacity of countries to monitor tobacco use among the youth, as well as implement and
48
49 evaluate tobacco control and prevention programs.³⁵ These data provide representative trends
50
51 of tobacco use among adolescents. The GYTS is a cross-sectional survey and does not follow
52
53 individuals over time, but provides data on smoking patterns among adolescents. In countries
54
55 where it is conducted at regular intervals, it allows the monitoring of trends over time. We are
56
57
58
59
60

1
2
3 aware of the 2017 GYTS for Ghana, but we do not include it in the analysis of the pooled
4
5 2000 – 2009 surveys due to inconsistencies in the questions asked, compared to those in
6
7 previous GYTS surveys. There is no other survey on adolescents in Ghana with comparable
8
9 measures except the ones outlined. We analyze the 2017 GYTS separately while linking it
10
11 with RIP data for 2008 – 2017 based on age-at-risk criteria^{27 32 36} (as done for the 2000 –
12
13 2009 surveys).
14
15

16
17 Although the GYTS data contain adolescents whose first puff was at age 7 or younger, we
18
19 assume that a student is at risk at age 8 because that is the age at which the child is relatively
20
21 developed and is able to start out-of-home interaction with peers.^{27 32 36} Students who started
22
23 smoking before reaching age 8 and those below age 8 are therefore excluded from the pooled
24
25 sample and not followed. In line with previous studies, a student exits the sample once
26
27 smoking is initiated.^{10 27 32 33}
28
29
30
31

32
33 In Ghana, there was no law prohibiting the sale of cigarettes to minors until 2012 when
34
35 restrictions on sale to persons below age 18 years were introduced.³⁷ The GYTS sample is
36
37 drawn using a two-stage cluster-sampling design.^{35 38 39} Schools are selected with probability
38
39 proportional to school enrolment size during the first stage, and then classes within
40
41 participating schools are selected as a systematic equal probability sample with a random
42
43 start during the second stage. All students in the selected classes are eligible to participate in
44
45 the survey.
46
47

48
49 The Ghana GYTS questionnaire captures information on the use of tobacco products such as
50
51 cigarettes and shisha. The data also include information on parental and peer smoking habits,
52
53 perception about tobacco use (e.g., weight gain, health effects, and ease of quitting),
54
55 amount money spent on tobacco in the last 30 days before the survey, and second-hand
56
57 smoking (SHS).⁴⁰ Studies vary widely on the way they define or measure smoking
58
59
60

1
2
3 initiation.⁴¹⁻⁴³ For the GYTS, smoking initiation is measured using the definition of a lifetime
4 smoker, i.e., a person who has ever tried smoking, *even one or two puffs of a cigarette*.^{27 43 44}
5
6 Thus, for our study, smoking initiation (dependent variable) is a dichotomous variable
7
8 generated from the following GYTS question where students answer Yes/No: *Have you ever*
9
10 *tried or experimented with cigarette smoking, even one or two puffs?*.²⁷

11
12
13
14
15 The main independent variable is the affordability index or the RIP, measured as the
16
17 percentage of GDP per capita (per capita income) required to buy 100 packs of cigarettes (20
18
19 sticks per pack, in total 2,000 sticks).^{17 18 45 46} Affordability is a relative measure and is
20
21 calculated using nominal prices and nominal GDP per capita, or real prices and real GDP per
22
23 capita. Data on per capita income are obtained from the World Bank's World Development
24
25 Indicators,³⁰ and those of average cigarette prices (for the most-sold brand) come from the
26
27 WHO, relevant publications of the tobacco industry⁴⁷ and the Government of Ghana⁴⁸. Years
28
29 with missing data on prices were interpolated using the formula:
30
31
32

$$P_{t-1} = \frac{P_t}{(1 + Tob.Inflation_t)} \dots\dots\dots (1)$$

33
34
35 where P_{t-1} is the previous year's price of cigarette, P_t is current price of cigarette and
36
37
38
39 $Tob.Inflation_t$ is the current tobacco inflation.⁴⁹ We then calculate RIP following methods
40
41
42 used by preceding studies, with a lower affordability index (RIP) indicating that cigarettes
43
44
45 have become more affordable and a higher value indicating that cigarettes have become less
46
47
48 affordable relative to the reference year.^{17 18 45 46} The WHO uses the same approach to obtain
49
50
51 its affordability index. Other independent variables used are sex, age, parents', and friends'
52
53
54 smoking status, whether offered cigarettes for free, family/class discussion about tobacco,
55
56
57 exposure to antismoking messages, and exposure to tobacco advertisements. These variables
58
59
60 are selected as they have been shown, theoretically or empirically, to be associated with

1
2
3 smoking initiation.^{27 31} Except age and RIP, which are continuously measured, all variables
4 are dichotomous.
5
6

7 8 **2.2. Data analysis** 9

10 We construct a pseudo-longitudinal dataset based on current age and age at first puff. In
11 doing this, we create a historical dataset starting from age 8 (age-at-risk criteria) and follow
12 the person until s/he initiates smoking. This ~~was~~ done by inferring the year of first puff using
13 the GYTS question: “How old were you when you first tried a cigarette?” and the age at the
14 time of the survey.²⁷
15

16 STATA routine command, *expand*, is used to expand the person’s age at the time of the
17 survey after which an event variable indicating smoking status is created. ~~Students who are
18 below age 8 and who started smoking before age 8 are removed from the dataset.~~ We then
19 link the RIP (affordability index) data to this pseudo-longitudinal dataset. ~~Although the
20 GYTS data contain adolescents whose first puff was at age 7 or younger, we assume that a
21 student is at risk at age 8 and exits the sample once smoking is initiated.^{10 27 32 33} We choose
22 age 8 as the age-at-risk because that is the age at which the child is relatively developed and
23 able to start out-of-home interaction with peers, according to a previous study.³² Students
24 who started smoking before reaching age 8 were therefore excluded from the pooled sample
25 and not followed.~~
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 Our statistical technique is duration or event history analysis where the timing of transition
47 into initiation is a function of the probability of initiating in period t conditional on not having
48 experienced a transition until period t , known as the hazard rate.¹⁰ Following previous
49 empirical studies,^{10 32 36} ~~Vellios and Van Walbeek¹⁰ and Guindon, et al.³²~~, we employ the
50 discrete time-hazard model, with logit specification (see equation 2), to study the association
51 between RIP (affordability) and smoking initiation among adolescents.
52
53
54
55
56
57
58
59
60

$$\Pr(\textit{Initiation} = 1 | X') = \beta_0 + \beta_1 \textit{RIP} + \beta_i X' \dots (2)$$

where *Initiation* is defined as first cigarette puff, *RIP* is the affordability index, X' is a vector of other independent variables affecting smoking initiation among adolescents, and β is a vector of the regression coefficients. The predictors, X' , represent age, sex, whether offered free cigarette, parental and peer smoking status, family/class discussion on the dangers of tobacco, exposure to tobacco advertisements, and hearing of antismoking messages, and awareness of tobacco control policies introduced in 2012 (for the 2017 GYTS). We report odds ratios (ORs), and the statistical level of significance is set at $p \leq 0.1$. $OR < 1$ implies that when a higher share of income is required to buy 2000 cigarettes (cigarettes are less affordable), the risk of smoking initiation declines, and vice versa. The partial derivative of equation 2 with respect to *RIP* gives the affordability elasticity.

To check the robustness of the logistic regression estimates, we employ a propensity score matching (PSM) technique to match ever-smokers to never-smokers based on the propensity scores. Our approach to matching follows previous studies.^{33 50 51} The propensity scores are obtained by running a logit regression to estimate the probability of being a smoker based on the variables in equation 2, except *RIP*, and the predicted probabilities are then used to match ever-smokers to never-smokers. Using the neighborhood matching, ever-smokers are matched to their two nearest neighbors. After matching the sample, we re-estimate the logit model to assess the effect of affordability on the probability of initiating smoking, using GYTS weights on the matched sample.^{50 51} With the matching approach, we are able to obtain the effect of affordability on the probability of initiating smoking among adolescent smokers and non-smokers who possess similar characteristics based on the propensity scores. This technique addresses issues of endogeneity and concerns relating to the fact that some never-smokers will never choose to smoke or use any form of tobacco irrespective of market conditions.^{33 34} Further, we minimize the problem of endogeneity by not using self-reported

1
2
3 prices.³² Data analysis is conducted using STATA version 15. The study benefited immensely
4
5 from discrete time modelling guidelines and STATA code produced by Professor Stephen
6
7 Jenkins.⁵²
8
9

10 **2.3 Interpretation of RIP Elasticity**

11
12 Although the RIP is measured in percentages, interpretation of the affordability index follows
13
14 the same procedure for elasticity interpretation. The elasticity measures the percentage
15
16 change in probability of initiating smoking following a percentage change in RIP, ceteris
17
18 paribus. Assuming the current RIP is 6% or 0.06, then a 1% increase in RIP corresponds to
19
20 the current RIP increasing from 6% to 6.06%. When using a unit change interpretation, a unit
21
22 change will be RIP moving from 6% to 7% and therefore probabilities will change in
23
24 absolute units and not percentages. Such distinction is important in understanding the impact
25
26 of affordability on smoking behavior.
27
28
29
30
31

32 **Patient and Public Involvement:**

33
34 No patient involved.
35
36
37
38
39
40
41

42 **3. RESULTS**

43 **3.1. Descriptive statistics**

44
45 A total of 20,202 adolescents were interviewed across the three GYTS waves (2000 – 2009).
46
47 54% of the respondents were males, while 76.47% were aged 15 years or less. In the 2017
48
49 GYTS, 5,664 people were interviewed, with about 48% being males. Overall, 12.46% and
50
51 8.9% of the respondents in the pooled (2000 – 2009) and 2017 surveys, respectively, had ever
52
53 smoked.
54
55
56
57
58
59
60

Given our age-at-risk criteria, 15,861 (2000 – 2009 GYTS) and 5,389 (2017 GYTS) people were eligible for inclusion in our pseudo-longitudinal analysis. For surveys prior to 2017, 4.2% initiated smoking at some point between 1991 and 2009, and about 77% of smoking initiators did so before reaching age 16. Further, 67% of initiators were males. Overall, males represented 53.62% of the eligible respondents. In the 2017 survey, 4.72% of the respondents initiated smoking at some point between 2008 and 2017. The characteristics of the samples are presented in Table 1. Due to incomplete information on some of the variables, the number of people used in the regression varies.

Table 1: Descriptive statistics

Variable	2000 – 2009 GYTS	2017 GYTS
	Students, n = 15 861	n=5389
Initiated smoking during the period	4.20%	4.72%
RIP (Affordability)	19.87 (SD=6.53)	7.63 (SD =0.86)
Offered free cigarettes	12.44%	8.13%
Sex (male =1)	53.62%	48.73%
At least one parent smoke	11.78%	-
Family/class discuss about tobacco	72.50%	51.47%
At least a friend smoke	15.94%	-
Exposed to tobacco adverts	40.46%	56.03%
Age (Years)	14.15 (SD=1.7)	14.10 (SD =1.03)
Heard anti-smoking campaigns	74.64%	57.26%
Age at initiation (Years)	11.95 (SD = 2.9)	11.26 (SD=2.41)
Percentage of initiators before age 16	77%	94%
Percentage of initiators who are males	67%	59%
Awareness of smoke free policies	-	78.24%

3.2. Regression Results

Results from the logit regressions for the unmatched and matched samples are reported in Tables 2 (GYTS 2000 – 2009) and 3 (GYTS 2017). The results show a statistically significant and negative relationship between RIP and smoking initiation. For instance, the results for the full unmatched sample (model 1, Table 2) show that OR on RIP is 0.98. This implies that a unit increase in affordability is associated with 0.98 odds of initiating smoking (OR = 0.98, $p < 0.05$). Thus, an adolescent who is subjected to a unit increase in RIP has 2% (i.e., $1 - 0.98 \times 100 = 2\%$) lower probability of initiating smoking than his/her counterpart who is not subjected to the same increase. Note that these results are not elasticities. Females have 40.1% (i.e., $1 - 0.599 \times 100 = 40.1\%$) lower probability of initiating smoking (OR = 0.599, $p < 0.01$) compared to their male counterparts in the unmatched sample.

Table 2: Effect of RIP on smoking initiation among adolescents (GYTS 2000 -2009)

VARIABLES	Unmatched Odds ratio	Matched Odds ratio
Affordability (RIP)	0.981** (0.009)	0.974*** (0.009)
Offered free cigarette (Ref=no)	1.491*** (0.216)	0.517*** (0.071)
Sex (Ref = male)	0.599*** (0.072)	0.615*** (0.076)
At least one parent smokes (Ref =no)	2.131*** (0.280)	0.862 (0.104)
Family/class discussion (Ref = no)	1.001 (0.133)	1.711*** (0.230)
At least one friend smokes (Ref=no)	4.109*** (0.520)	1.094 (0.126)
Exposure to adverts (Ref=no)	1.155 (0.140)	1.027 (0.121)
Age	1.150*** (0.042)	0.991 (0.031)
Heard of anti-smoking message/campaign (Ref=no)	1.342* (0.217)	2.048*** (0.321)
Survey cycle (ref=2000)		

2006	0.958	0.880
	(0.146)	(0.138)
2009	1.108	1.003
	(0.171)	(0.159)
Log (time)	1.110	1.393***
	(0.106)	(0.146)
Constant	0.000***	0.048***
	(0.000)	(0.024)
Observations	106,673	10,078
Number of people	15,201	1,611
Ever-smokers	611	611
Pseudo R-squared	0.0815	0.0448
Chi2	439.2***	91.84***

Robust standard errors in parentheses

*** p<0.01, ** p<0.05, * p<0.1

Table 3: Effect of RIP on smoking initiation among adolescents (GYTS 2017)

	Unmatched	Matched
VARIABLES	Odds ratio	Odds ratio
Affordability (RIP)	0.821**	0.804***
	(0.066)	(0.065)
Sex (Ref=male)	0.659**	0.902
	(0.120)	(0.177)
Offered free cigarettes (Ref= no)	3.403***	0.978
	(0.726)	(0.221)
Heard of anti-smoking message (Ref=no)	1.165	1.009
	(0.213)	(0.192)
Exposed to tobacco adverts (Ref= no)	3.030***	1.893***
	(0.622)	(0.421)
Smoke free policies awareness (Ref=no)	1.250	1.160
	(0.329)	(0.332)
Age	1.847***	1.793***
	(0.294)	(0.279)
Class discussion on tobacco harms (Ref=no)	0.795	1.278
	(0.138)	(0.241)
Log(time)	0.104***	0.119***
	(0.067)	(0.076)
Constant	0.000***	0.002***
	(0.000)	(0.002)
Observations	37,654	4,850

Number of people	5,301	747
Ever-smokers	231	206
Pseudo R-squared	0.0599	0.0292
Chi2	158***	38.72***

Robust standard errors in parentheses

*** p<0.01, ** p<0.05, * p<0.1

Similarly, in Table 3, in the unmatched sample, an adolescent faced with a unit increase in RIP has about 18% (i.e., $1 - 0.821 \times 100 = 17.9\%$) lower probability of starting smoking than his/her counterpart who is not subjected to the same increase.

Other significant factors that influence smoking initiation in our two samples include whether the adolescent's parents (OR = 2.131, p<0.01) or friends (OR = 4.109, p<0.01) smoke. In addition, adolescents who are offered free cigarettes have a high probability of initiating smoking (OR = 1.491, p<0.01) compared to those who receive no such offer for the 2000 - 2009 wave (Table 2). In the 2017 wave, the odds of adolescents starting smoking when given cigarettes freely is 3.403 (p<0.01) (Table 3).

In the matched sample (~~model 2~~, Table 2), 611 ever-smokers were matched to their two nearest neighbors (1000 never-smokers) which produced a sample of 1,611 adolescents with similar characteristics. An adolescent subjected to a unit increase in RIP has a 0.97 times lower chance of initiating smoking (OR = 0.974, p<0.01) compared to those not exposed to the same increase in RIP. Similarly in the matched sample of the GYTS 2017 (Table 3), a unit increase in RIP is associated with 0.80 times lower odds of smoking initiation (OR = 0.804, p<0.01).

Getting free cigarettes (OR = 0.517, p<0.01), family/class discussion on tobacco (OR = 1.711, p<0.01), and hearing antismoking messages (OR = 2.048, p<0.01) are all found to be statistically significant in influencing smoking initiation in the matched sample (Table 2).

However, the odds for these variables are contrary to *a priori* expectations. Similarly, in the

matched sample (Table 2), the likelihood of initiating smoking among females is lower. The results imply that females have about 39% lower probability of initiating smoking (OR = 0.615, $p < 0.01$) than males.

3.3. Affordability Elasticities

In the unmatched sample, the estimated affordability elasticity is -0.372 [CI: -0.701 to -0.042] for the 2000-2009 sample and -1.247 [-2.248 to -0.246] for the 2017 sample. These elasticities are statistically significant at the 5% level. By sex, the affordability elasticity is -0.137 for males and -0.888 for females for the 2000-2009 sample, but only that of females is statistically significant ($p < 0.01$). The elasticities are higher for 2017 GYTS (-0.938 for males and -1.610 for females). The results are presented in Table 4.

Table 4: Affordability Elasticity estimates

	Both sexes		Male		Female	
Panel A: 2000 -2009						
	Percentage changes (d(lny)/d(lnx))					
VARIABLES	Unmatched	Matched	Unmatched	Matched	Unmatched	Matched
Affordability	-0.372**	-0.490***	-0.137	-0.326	-0.888***	-0.928***
	(0.168)	(0.168)	(0.219)	(0.216)	(0.253)	(0.258)
95% CI	-0.701 to -0.042	-0.818 to -0.161	-0.567 to 0.292	-0.749 to 0.097	-1.384 to -0.392	-1.434 to -0.422
Observations	106,673	10,078	55,396	5,648	51,277	4,430
Panel B: 2017 GYTS						
Affordability	-1.247**	-1.349***	-0.938**	-1.045**	-1.610*	1.518**
	(0.511)	(0.500)	(0.474)	(0.484)	(0.866)	(0.778)
95% CI	-2.248 to -0.246	-2.328 to -0.369	-1.867 to -0.008	-1.993 to -0.096	-3.307 to -0.087	-3.043 to -0.007
Observations	37,654	4,850	18,084	2,807	19,570	2,043

Standard errors in parentheses
 *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

In the matched sample for the 2000-2009 GYTS, the overall elasticity is -0.490 [CI: -0.818 to -0.161] for both sexes (Table 4), which is similar to that of the unmatched sample. For males, the effect of changes in RIP is statistically insignificant. Among females, a percentage

1
2
3 increase in RIP is associated with a 0.928% lower probability of smoking initiation. The
4
5 elasticities for both males and females in the 2017 GYTS were negative, statistically
6
7 significant, and more than unity.
8
9

10 The mean and median standardized difference for the covariates used in matching show that
11
12 the matching satisfies the balancing test (results not reported). The mean and median
13
14 standardized difference is 2.3%, which is below the normal 10% threshold. Therefore, the
15
16 balancing property is satisfied.
17
18

19 20 21 **5. DISCUSSION AND CONCLUSION**

22
23 In this study, increasing the RIP of cigarettes is significantly associated with a lower
24
25 probability of initiating smoking. This finding is consistent with international literature,
26
27 including the few existing studies in SSA that have reported that making cigarettes less
28
29 affordable lowers the likelihood of smoking initiation among young people.^{17 18 26} In addition,
30
31 the results from the unmatched 2000-2009 sample suggest that males are not responsive to
32
33 changes in RIP whilst females are. Nevertheless, in the matched sample analysis, especially
34
35 using the 2017 GYTS, both males and females are responsive to changes in RIP. Indeed, the
36
37 issue of affordability becomes more important given that Ghana's per capita income has been
38
39 growing at an average of 4.4% annually in the last decade.³⁰
40
41
42
43
44

45 Parental smoking increased the odds of smoking initiating. This points to the parental
46
47 influence on the lifestyle of adolescents. Adolescents whose parents smoke may perceive
48
49 smoking as acceptable behavior. Previous studies have reported similar findings.^{10 27 31 53} The
50
51 odds of smoking initiation are higher for those whose friends' smoke. This points to the
52
53 influence of peers, and is consistent, for example, with Mak, et al.³¹. Those who were offered
54
55 free cigarettes by sales representatives were more likely to initiate smoking. All forms of
56
57 tobacco promotion and advertising were banned in Ghana in 1982. However, the tobacco
58
59
60

1
2
3 industry seems to be breaking these laws, because 12.44% of youth reported being offered a
4 cigarette for free.- This observation, together with our results, suggest the need to strengthen
5
6 the enforcement of the existing ban on all forms of tobacco advertising and promotion in
7
8
9
10 Ghana.

11
12
13 This study has several limitations. The GYTS is a self-reporting survey, which means the
14 responses are prone to recall bias even in cases where the adolescents are required to answer
15 questions about events that occurred in the past 30 days. For instance, students may not recall
16 the exact age at which they tried their first puff. There is also a social desirability bias when
17 self-reporting behaviors such as smoking, especially among females. The weakness of our
18 measure of smoking initiation is that it may not predict regular smoking behavior well.^{42 43} In
19 addition, there are other important factors affecting smoking uptake among the youth that are
20 not included in this study. For instance, changing community norms regarding smoking, the
21 enforcement of laws regarding the sale of cigarettes to minors, and changes in the social
22 image of smoking are key factors that may influence smoking participation,^{54 55} but are not
23 included in the models estimated.

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39 In conclusion, making cigarettes less affordable is associated with a lower probability of
40 smoking initiation among adolescents in Ghana. This supports the use of price measures,
41 through higher excise taxes, as effective strategies to decrease smoking initiation among
42 adolescents in Ghana. Since incomes are rising at the average of 4.4% annually³⁰, tobacco
43 taxes need to be adjusted regularly to ensure that cigarettes or other tobacco products become
44 less affordable over time in order to discourage young people from initiating smoking and to
45 encourage smokers to quit.

46
47
48
49
50
51
52
53
54
55
56 **Author Contributions:** Conceptualization, M.K.B.; methodology, M.K.B; software, M.K.B.;
57 validation, M.K.B., M.I., E.N.T., N.D.M., and H.R.; formal analysis, M.K.B.; investigation,
58 M.K.B., M.I., E.N.T., N.D.M., and H.R.; data curation, M.K.B; writing—original draft
59 preparation, M.K.B., M.I., E.N.T., N.D.M., and H.R.; writing—review and editing, M.K.B.,
60

1
2
3 M.I., E.N.T., N.D.M., H.R.; project administration, M.K.B.; supervision, H.R.; funding
4 acquisition, H.R. All authors have read and approved the manuscript.
5
6
7

8 **Funding:** Funding for this study was provided by Cancer Research UK (Grant number:
9 C62640/A24723) and International Development Research Centre (Grant Number: 108820-
10 001); with additional support from the SAMRC/Wits Centre for Health Economics and
11 Decision Science – PRICELESS SA (grant number 23108). Funding for ENT and NDM is
12 acknowledged from the Tobacco Control Capacity Programme (Grant Number:
13 MR/P027946/2) supported by UK Research and Innovation (UKRI) with funding from the
14 Global Challenges Research Fund (GCRF). The funders had no role in study design, data
15 collection and analysis, or preparation of the manuscript.
16
17
18
19
20
21
22
23
24
25
26

27 **Conflicts of Interest:** The authors declare that they have no conflict of interest.
28
29

30 **Research Ethics Approval: Human Participants**

31
32
33 This study does not involve human participants.
34
35

36 **Research Ethics Approval: Animals**

37
38
39 This study does not involve animal subjects.
40
41

42 **Patient consent for publication:** No patient involved.
43
44

45 **Data Availability statement:** The publicly available data can be accessed:

46
47 <https://nccd.cdc.gov/GTSSDataSurveyResources/Ancillary/DataReports.aspx?CAID=2>. The
48 relative income price data are available from the authors on reasonable request.
49
50
51
52
53
54

55 **REFERENCES**

56
57
58
59
60

1. Doll R, Peto R, Boreham J, et al. Mortality in relation to smoking: 50 years' observations on male British doctors. *Bmj* 2004;328(7455):1519.
2. Winkler V, Mangolo NJ, Becher H. Lung cancer in South Africa: a forecast to 2025 based on smoking prevalence data. *BMJ open* 2015;5(3):e006993.
3. World Health Organization. Tobacco: Key Facts. 2020 [updated 27 May 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/tobacco> accessed 22 November 2020.
4. Goodchild M, Nargis N, d'Espaignet ET. Global economic cost of smoking-attributable diseases. *Tobacco Control* 2018;27(1):58-64.
5. Boachie MK, Rossouw L, Ross H. The Economic Cost of Smoking in South Africa, 2016. *Nicotine Tob Res* 2021;23(2):286 - 93. doi: 10.1093/ntr/ntaa162
6. USDHHS. Preventing tobacco use among youth and young adults: A report of the surgeon general 2012 [Available from: <http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf> (accessed: Jan 3, 2019).
7. World Health Organization. WHO global report on trends in prevalence of tobacco use 2000-2025. 3rd ed. Geneva: World Health Organization, 2019.
8. CDC, WHO, MOH. GLOBAL YOUTH TOBACCO SURVEY: Factsheet, Ghana 2017, 2018.
9. World Health Organization. WHO global report on trends in prevalence of tobacco smoking 2015. Geneva: World Health Organization, 2015.
10. Vellios N, Van Walbeek C. Determinants of regular smoking onset in South Africa using duration analysis. *BMJ Open* 2016;6(7):e011076.
11. IARC. Effectiveness of tax and price policies for tobacco control. In: Cancer. IAFRo, ed. USA: International Agency for Research on Cancer, 2011.
12. Boachie MK, Ross H. Determinants of smoking intensity in South Africa: evidence from township communities. *Preventive Medicine Reports* 2020;19:101099.
13. Van Walbeek C. A simulation model to predict the fiscal and public health impact of a change in cigarette excise taxes. *Tobacco control* 2010;19(1):31-36.

- 1
2
3
4 14. Yurekli A, de Beye J. Design and administer tobacco taxes. *World Bank economics of*
5
6 *tobacco toolkit* 2001(4)
7
- 8 15. Van Walbeek C. Raising Additional Government Revenues in Ghana by Raising the Excise
9
10 Tax on Tobacco and Alcohol Washington, D.C.: WBG Global Tobacco Control
11
12 Program, World Bank Group; 2014 [Available from:
13
14 [http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-](http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-and-Alcohol.pdf)
15
16 [Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-](http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-and-Alcohol.pdf)
17
18 [and-Alcohol.pdf](http://documents.worldbank.org/curated/en/133561561707742558/pdf/Raising-Additional-Government-Revenues-in-Ghana-by-Raising-the-Excise-Tax-on-Tobacco-and-Alcohol.pdf) accessed 15 July 2019.
19
- 20 16. He Y, Shang C, Chaloupka FJ. The association between cigarette affordability and
21
22 consumption: An update. *PLOS ONE* 2018;13(12):e0200665. doi:
23
24 10.1371/journal.pone.0200665
25
- 26 17. Nargis N, Stoklosa M, Shang C, et al. Price, Income, and Affordability as the Determinants
27
28 of Tobacco Consumption: A Practitioner's Guide to Tobacco Taxation. *Nicotine and*
29
30 *Tobacco Research* 2021;23(1):40-47.
31
- 32 18. Blecher E, Van Walbeek C. An international analysis of cigarette affordability. *Tobacco*
33
34 *Control* 2004;13(4):339-46.
35
- 36 19. Ding A. Youth are more sensitive to price changes in cigarettes than adults. *The Yale*
37
38 *journal of biology and medicine* 2003;76(3):115.
39
- 40 20. Ho L-M, Schafferer C, Lee J-M, et al. The effect of cigarette price increases on cigarette
41
42 consumption, tax revenue, and smoking-related death in Africa from 1999 to 2013.
43
44 *International journal of public health* 2017;62(8):899-909.
45
- 46 21. Chelwa G, van Walbeek C. Does cigarette demand respond to price increases in Uganda?
47
48 Price elasticity estimates using the Uganda National Panel Survey and Deaton's method.
49
50 *BMJ open* 2019;9(3):e026150. doi: 10.1136/bmjopen-2018-026150 [published Online
51
52 First: 2019/03/23]
53
- 54 22. Stoklosa M, Goma F, Nargis N, et al. Price, tax and tobacco product substitution in Zambia:
55
56 findings from the ITC Zambia Surveys. *Tob Control* 2019;28(Suppl 1):s45-s52. doi:
57
58 10.1136/tobaccocontrol-2017-054037 [published Online First: 2018/03/27]
59
60

- 1
2
3
4 23. Mukong AK, Tingum EN. The demand for cigarettes: New evidence from South Africa.
5
6 *Development Southern Africa* 2020;37(1):40-54.
7
- 8 24. Tingum EN, Mukong AK, Mdege N. The effects of price and non-price policies on cigarette
9
10 consumption in South Africa. *Tob Induc Dis* 2020;18(July) doi: 10.18332/tid/123424
11
- 12 25. Kidane A, Mduma J, Naho A, et al. The Demand for Cigarettes in Tanzania and
13
14 Implications for Tobacco Taxation Policy. *Advances in economics and business*
15
16 2015;3(10):428.
17
- 18 26. Kostova D. A (nearly) global look at the dynamics of youth smoking initiation and cessation:
19
20 the role of cigarette prices. *Applied Economics* 2013;45(28):3943-51.
21
- 22 27. Asare S, Stoklosa M, Drope J, et al. Effects of Prices on Youth Cigarette Smoking and
23
24 Tobacco Use Initiation in Ghana and Nigeria. *International Journal of Environmental*
25
26 *Research and Public Health* 2019;16(17):3114.
27
- 28 28. Ministry of Health. Needs assessment for the Implementation of WHO FCTC in Ghana.
29
30 Accra: Ministry of Health,, 2010.
31
- 32 29. Gov't. of Ghana. Customs and Excise (Duties and Other Taxes) Amendment Act 739. ,
33
34 2007.
35
- 36 30. World Bank. World Development Indicators.: World Bank; 2019 [Available from:
37
38 <https://databank.worldbank.org/source/world-development-indicators>. accessed 01 Jan
39
40 2020.
41
- 42 31. Mak K-K, Ho S-Y, Day JR. Smoking of parents and best friend—-independent and
43
44 combined effects on adolescent smoking and intention to initiate and quit smoking.
45
46 *Nicotine Tob Res* 2012;14(9):1057-64.
47
- 48 32. Guindon GE, Paraje GR, Chávez R. Prices, inflation, and smoking onset: the case of
49
50 Argentina. *Economic Inquiry* 2018;56(1):424-45.
51
- 52 33. Dauchy E, Ross H. The Effect of Price and Tax Policies on the Decision to Smoke among
53
54 Men in Kenya. *Addiction* 2019;114(7):1249-63.
55
- 56 34. Li M. Using the propensity score method to estimate causal effects: A review and practical
57
58 guide. *Organizational Research Methods* 2013;16(2):188-226.
59
60

- 1
2
3
4 35. Warren CW. Tobacco use among youth: a cross country comparison. *Tobacco Control*
5 2002;11(3):252-70.
6
7
8 36. Boachie MK, Immurana M, Agyemang JK, et al. Cigarette prices and smoking
9 experimentation in Sierra Leone: an exploratory study. *Tobacco Use Insights*
10 2022;forthcoming
11
12
13
14 37. Gov't. of Ghana. Public Health Act, 2012 Act 851. Ghana 2012.
15
16 38. Wellington E. Ghana Global Youth Tobacco Survey
17 (GYTS), Factsheet. 2006 [Available from:
18 [http://www.afro.who.int/en/downloads/docdownload/1887-ghana-gyts-2006-](http://www.afro.who.int/en/downloads/docdownload/1887-ghana-gyts-2006-factsheet.html)
19 [factsheet.html](http://www.afro.who.int/en/downloads/docdownload/1887-ghana-gyts-2006-factsheet.html) (Retrieved Jan 28, 2019).
20
21
22
23
24 39. Cadmus EO, Ayo-Yusuf OA. The effect of smokeless tobacco use and exposure to cigarette
25 promotions on smoking intention among youths in Ghana. *Cogent Medicine*
26 2018;5(1):1531459.
27
28
29
30 40. CDC and WHO. Ghana global youth tobacco survey. Atlanta, United States: Centers for
31 Disease Control and Prevention (CDC). 2006.
32
33
34 41. Freedman KS, Nelson NM, Feldman LL. Smoking initiation among young adults in the
35 United States and Canada, 1998-2010: a systematic review. *Preventing chronic disease*
36 2012;9
37
38
39
40 42. Azagba S, Baskerville NB, Minaker L. A comparison of adolescent smoking initiation
41 measures on predicting future smoking behavior. *Preventive medicine reports*
42 2015;2:174-77.
43
44
45
46 43. Reidpath DD, Davey TM, Kadirvelu A, et al. Does one cigarette make an adolescent smoker,
47 and is it influenced by age and age of smoking initiation? Evidence of association from
48 the US Youth Risk Behavior Surveillance System (2011). *Preventive medicine*
49 2014;59:37-41.
50
51
52
53
54 44. Centers for Disease Control Prevention. Selected cigarette smoking initiation and quitting
55 behaviors among high school students--United States, 1997. *MMWR Morbidity and*
56 *mortality weekly report* 1998;47(19):386.
57
58
59
60

- 1
2
3
4 45. Blecher E. Targeting the affordability of cigarettes: a new benchmark for taxation policy
5 in low-income and-middle-income countries. *Tobacco control* 2010;19(4):325-30.
6
7
8 46. Nargis N, Stoklosa M, Drope J, et al. Trend in the affordability of tobacco products in
9 Bangladesh: findings from the ITC Bangladesh Surveys. *Tobacco control*
10 2019;28(Suppl 1):s20-s30.
11
12
13
14 47. Philip Morris International. World summary of cigarette retail prices, tax burdens and tax
15 incidence.: Philips Moris Records 1996 [Available from:
16 <https://www.industrydocuments.ucsf.edu/docs/#id=glhf0066>.
17
18
19
20 48. Ghana Gto. Report card on the WHO Framework Convention on Tobacco Control., 2009.
21
22 49. Ghana Statistical Service. Consumer Price Index (CPI): Time Series In: Service GS, ed.
23 Accra, 2012.
24
25
26 50. Austin PC, Jembere N, Chiu M. Propensity score matching and complex surveys. *Statistical*
27 *methods in medical research* 2018;27(4):1240-57.
28
29
30 51. DuGoff EH, Schuler M, Stuart EA. Generalizing observational study results: applying
31 propensity score methods to complex surveys. *Health services research*
32 2014;49(1):284-303.
33
34
35
36 52. Jenkins SP. Survival Analysis with Stata. [Available from:
37 <https://www.iser.essex.ac.uk/resources/survival-analysis-with-stata> accessed 01
38 January 2020.
39
40
41
42 53. Gilman SE, Rende R, Boergers J, et al. Parental smoking and adolescent smoking initiation:
43 an intergenerational perspective on tobacco control. *Pediatrics* 2009;123(2):e274-e81.
44
45
46 54. Gilpin EA, Pierce JP. Trends in adolescent smoking initiation in the United States: is
47 tobacco marketing an influence? *Tobacco Control* 1997;6(2):122-27.
48
49
50 55. Cawley J, Markowitz S, Tauras J. Obesity, cigarette prices, youth access laws and
51 adolescent smoking initiation. *Eastern Economic Journal* 2006;32(1):149-70.
52
53
54
55
56
57
58
59
60

1
2
3 Dear Editor
4

5 Thank you and the reviewers for the opportunity to revise our manuscript for consideration.
6 We have addressed the reviewers' comments and made changes in the main manuscript.
7 The responses to the reviewers on the changes made are provided below.
8
9

10
11
12 Kind regards

13
14 Micheal K Boachie
15
16
17
18
19

20 **Reviewer: 2**

21 Dr. Shaoying Ma, OSUMC
22
23

24 **Comments to the Author:**

25 This is a revised version of the paper that I reviewed in November, and the authors study
26 the impact of cigarette affordability (measured by relative income price, i.e. RIP) on youth
27 initiation in Ghana, using data from Global Youth Tobacco Surveys (GYTS).
28 The authors made some major changes to the original version of their manuscript, which
29 much improved the paper.
30
31

32
33
34 Author response

35 Thank you very much for the suggestions.
36
37
38
39

40 Please see below for specific comments.
41

42 **Comment**

43 1. Page 9 line 13, I appreciate that the authors added more details about how they
44 addressed the endogeneity issue, and I wonder how does "iii) the fact that some people will
45 never choose to smoke or use any form of tobacco, for example, for reasons of health or
46 religious belief" help to alleviate the endogeneity concern?
47
48

49 Author response

50
51 We have deleted the third item "iii) the fact that some people will never
52 choose to smoke or use any form of tobacco, for example, for reasons of
53 health or religious belief" Now the endogeneity is tackled using the aggregate
54 prices and propensity score matching.
55
56
57
58
59
60

Comment

2. Page 10 line 40, “amount spent on tobacco in the last 30 days before the survey”
Is it money amount, time amount, or something else?

Author response

Thank you very much for the observation. We were referring to money spent on tobacco. Thus, the expenditure on tobacco. We have revised it as “...
money spent on tobacco ...”

Comment

3. Page 10 line 12 and page 11 line 60, “age-at-risk criteria”

I see that “age-at-risk criteria” was explained in more details on page 12 lines 19-26. I would recommend moving the explanation in the text, and explain it when it is first mentioned, i.e. page 10 line 12; otherwise readers may not understand the basis of choosing to start “from age 8 (age-at-risk criteria)”. Also citations should be added to “age-at-risk criteria” on page 10 line 12 and page 11 line 60.

Author response

Thank you for the suggestion. We have revised the section and also included the citations at the first mention of the age-at-risk criteria. The text now reads as (page 8):

Although the GYTS data contain adolescents whose first puff was at age 7 or younger, we assume that a student is at risk at age 8 because that is the age at which the child is relatively developed and is able to start out-of-home interaction with peers.[1-3] Students who started smoking before reaching age 8 and those below age 8 are therefore excluded from the pooled sample and not followed. In line with previous studies, a student exits the sample once smoking is initiated.[1,2,4,5]

References

1. Guindon GE, Paraje GR, Chávez R. Prices, inflation, and smoking onset: the case of Argentina. *Economic Inquiry*. 2018;56(1):424-445.
2. Asare S, Stoklosa M, Drope J, Larsen A. Effects of Prices on Youth Cigarette Smoking and Tobacco Use Initiation in Ghana and Nigeria. *International Journal of Environmental Research and Public Health*. 2019;16(17):3114.
3. Boachie MK, Immurana M, Agyemang JK, Ross H. Cigarette prices and smoking experimentation in Sierra Leone: an exploratory study. *Tobacco Use Insights*. 2022;forthcoming.
4. Vellios N, Van Walbeek C. Determinants of regular smoking onset in South Africa using duration analysis. *BMJ Open*. 2016;6(7):e011076.

- 1
- 2
- 3
- 4 5. Dauchy E, Ross H. The Effect of Price and Tax Policies on the Decision to Smoke
- 5 among Men in Kenya. *Addiction*. 2019;114(7):1249-1263.
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

For peer review only