



UNIVERSITY OF LEEDS

This is a repository copy of *Kv7.4 channel is a key regulator of vascular inflammation and remodeling in neointimal hyperplasia and abdominal aortic aneurysms*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/184197/>

Version: Supplemental Material

---

**Article:**

Fan, XZ, Wang, YY, Cui, ZY et al. (5 more authors) (2022) Kv7.4 channel is a key regulator of vascular inflammation and remodeling in neointimal hyperplasia and abdominal aortic aneurysms. *Free Radical Biology and Medicine*, 178. pp. 111-124. ISSN 0891-5849

<https://doi.org/10.1016/j.freeradbiomed.2021.11.041>

---

© 2021, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

**Reuse**


This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



Schematic representation of a working model which impaired Kv7.4 channel inhibits vascular inflammation and remodeling.