



This is a repository copy of *Effective cosmology from one-body operators in group field theory*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/184317/>

Version: Accepted Version

Article:

Gielen, S. orcid.org/0000-0002-8653-5430, Marchetti, L., Oriti, D. et al. (1 more author) (2022) Effective cosmology from one-body operators in group field theory. *Classical and Quantum Gravity*, 39 (7). 075002. ISSN 0264-9381

<https://doi.org/10.1088/1361-6382/ac5052>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

ACCEPTED MANUSCRIPT • OPEN ACCESS

Effective cosmology from one-body operators in group field theory

To cite this article before publication: Steffen Gielen *et al* 2022 *Class. Quantum Grav.* in press <https://doi.org/10.1088/1361-6382/ac5052>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2022 The Author(s). Published by IOP Publishing Ltd..

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

Effective cosmology from one-body operators in group field theory

Steffen Gielen,^{1,*} Luca Marchetti,^{2,3,4,†} Daniele Oriti,^{3,‡} and Axel Polaczek^{1,§}

¹*School of Mathematics and Statistics, University of Sheffield,*

Hicks Building, Hounsfield Road, Sheffield S3 7RH, United Kingdom

²*Università di Pisa, Lungarno Antonio Pacinotti 43, 56126 Pisa, Italy, EU*

³*Arnold Sommerfeld Center for Theoretical Physics,*

Ludwig-Maximilians-Universität München,

Theresienstrasse 37, 80333 München, Germany, EU

⁴*Istituto Nazionale di Fisica Nucleare sez. Pisa,*

Largo Bruno Pontecorvo 3, 56127 Pisa, Italy, EU

(Dated: January 18, 2022)

We propose a new method for obtaining an effective Friedmann–Lemaître–Robertson–Walker (FLRW) cosmology from the quantum gravity dynamics of group field theory (GFT), based on the idea that an FLRW universe is characterised by a few macroscopic observables. Rather than relying on assuming a particular type of quantum state and computing expectation values in such a state, here we directly start from relations between macroscopic observables (defined as one-body operators) and formulate dynamics only for those observables. We apply the effective approach to constrained quantum systems (as developed by Bojowald and collaborators) to GFT, providing a systematic expansion in powers of \hbar . We obtain a kinematical phase space of expectation values and moments, which does not require an *a priori* choice of clock variable. We identify a gauge fixing of the system which corresponds to choosing one of the cosmological variables (with the role of extrinsic curvature) as a clock and which allows us to rewrite the effective dynamics in relational form. We show necessary and sufficient conditions for the resulting dynamics of expectation values to be compatible with those of classical FLRW cosmology and discuss the impact of quantum fluctuations.

* s.c.gielen@sheffield.ac.uk

† luca.marchetti@phd.unipi.it

‡ daniele.oriti@physik.lmu.de

§ apolaczek1@sheffield.ac.uk

I. INTRODUCTION

Many modern approaches to quantum gravity suggest that the classical continuum notions of space and time are emergent concepts, to be obtained from an underlying fundamental description in terms of different quantum degrees of freedom. Extracting continuum physics from such theories is generally a very complicated task, because of both conceptual and technical problems. Some of the most important outstanding issues, in relation to this task, are the problem of time, the problem of Hilbert space and the problem of the classical limit [1, 2]. There are of course many other issues, such as the implementation of diffeomorphism invariance in a discrete formalism [3] or the reliance on preferred foliations, but in the following we will focus only on these three. They are present in one form or another in any formulation of quantum gravity, but they are particularly visible in a canonical formalism.

The problem of time refers to the fact that, due to the diffeomorphism invariance of general relativity, there are no external structures with respect to which one can define (time) evolution. As a result, the Hamiltonian of general relativity (or, more in general, of any background independent theory) is forced to vanish in absence of boundaries [4–6]. At the quantum level, the vanishing of the Hamiltonian results in “frozen” dynamics, where states do not evolve in “time”. Change and evolution can however still be defined in terms of internal physical quantities. This is the relational strategy, which is based on the use of evolving constants of motion [7, 8] (see [9] for a review), i.e., Dirac observables encoding correlations between the physical variable chosen as a clock and the remaining ones. Though these observables can be formally defined in the classical theory [10, 11], their explicit construction, especially at the quantum level, is extremely complicated. One main reason is that one has in general very little control over the physical Hilbert space on which they should be represented. The construction of a Hilbert space by endowing the space of solutions to the quantum constraint with an appropriate inner product is a poorly understood issue; this is the Hilbert space problem mentioned above. Lastly, the problem of the classical limit refers to the extraction of (semi)classical physics from full quantum gravity, which has been often addressed through an explicit construction of specific classes of states. Semiclassicality is however not a sharp notion, and it is therefore important to clarify which results are general to semiclassical states and which ones are instead only valid for the states considered. Moreover, in approaches (like GFT or spin foam models or lattice quantum gravity) in which the basic structures of the theory are discrete, the semiclassical approximation leading to general relativity is necessarily accompanied by some form of continuum approximation, implemented via coarse graining or renormalisation group

1
2
3 techniques. The two kinds of approximations are conceptually and technically distinct [12], albeit
4 intertwined.
5

6 Effective techniques, describing quantum systems in terms of (a finite set of) expectation values
7 and moments of observables associated to a given state, may play a crucial role in order to practically
8 bypass the above difficulties, while still allowing to extract physical semiclassical predictions from the
9 theory. These techniques have been employed already in the context of simple quantum systems for
10 instance in [13–18]. In principle, in order to apply the framework developed in [13, 15] to quantum
11 gravity, it should be generalised to a field theoretic context, which will in general pose formidable
12 technical challenges. However, in quantum gravity one is often not interested in all degrees of freedom
13 in the theory, but rather in situations where only some degrees of freedom are relevant. In particular,
14 the relevant physical scales where quantum gravity effects might be seen are far away from the
15 microscopic Planck scale, and will thus involve a form of collective or coarse-grained characterisation
16 of the fundamental quantum gravity theory. One may then develop an effective framework only for
17 those observables which are most relevant at this coarse-grained (“hydrodynamic”) level: this extends
18 the effective framework in a more minimal way, essentially replacing the algebra of observables
19 of the quantum particle with a similarly finitely generated algebra of collective and macroscopic
20 observables. Expectation values and moments of these observables are then the quantities in terms
21 of which the effective hydrodynamic description is defined. This focus on such a coarse-grained
22 regime can also be seen as a way to implement a continuum approximation, as part of the extraction
23 of semiclassical physics via the chosen effective framework.
24
25
26
27
28
29
30
31
32
33
34
35
36

37 In quantum gravity, a system which can be characterised by a few collective observables is the
38 homogeneous and isotropic FLRW universe [19]. The hydrodynamic description of the fundamental
39 quantum gravity dynamics to capture homogeneous and isotropic cosmology has been employed
40 especially in the context of group field theories (GFTs) [20]. GFTs can be seen as quantum field
41 theories whose fundamental quanta are three-dimensional discrete chunks of space, i.e., quantum
42 tetrahedra, whose geometric properties are encoded in group variables (these variables are equivalent
43 to the data on open spin networks in loop quantum gravity [21]). Continuum spacetime notions are
44 expected to emerge from these theories only after an appropriate coarse-graining is performed. Their
45 Fock space structure is perfectly suited for the construction of non-perturbative collective states,
46 such as coherent states, which have an interpretation in terms of continuous and homogeneous
47 three-geometries [22, 23]. Using these states and mean field methods one can obtain (in the weak
48 interaction limit) an effectively hydrodynamic description characterised by Friedmann dynamics
49 with a quantum bounce for the averaged volume of the universe [24–27]. These results have sparked
50
51
52
53
54
55
56
57
58
59
60

1
2
3 further works addressing for instance the role of quantum fluctuations, especially for late times and
4 around the bounce [28, 29] and the definition of relational evolution in such emergent quantum
5 gravity theories [25, 30].
6
7

8 In this paper, we aim to provide a new perspective on the derivation of cosmological dynamics
9 from GFT by extending the effective techniques developed in [13–15] to the GFT setting. As in [24]
10 and many subsequent papers, we consider a GFT model minimally coupled with a massless scalar
11 field, and assume that the self-interactions of the GFT field are negligible in the regime of interest.
12 As mentioned above, the effective techniques were developed for a quantum algebra generated by
13 a finite set of operators. We propose to use a suitable finite set of one-body GFT operators¹. In
14 particular, the quantum constraint of [13] is given by the integrated (free) equation of motion in our
15 set-up. Truncating the system in this manner is motivated by the interest in only some collective
16 (hydrodynamic) variables, and is the most crucial approximation that we perform; we see it as a
17 requirement to be imposed on the kind of states employed in the effective description. One important
18 consequence of this assumption is that the number operator of GFT quanta² trivially commutes
19 with the constraint, so that its expectation value and all its moments will be constant. The quantum
20 algebra thus obtained allows to construct a quantum phase space coordinatised by expectation
21 values and moments of the relevant operators, according to standard techniques [13, 15, 31]. By
22 imposing that the states of interest are also semiclassical, one can truncate the hierarchy of moments
23 to a desired order in \hbar . The quantum constraint then produces a finite set of effective constraints,
24 generating gauge flows on the variables of the truncated quantum phase space. Similarly to [15–17],
25 all but one of these gauge flows are fixed so that one of the geometric operators in the algebra (here
26 chosen to classically match the extrinsic curvature) is effectively classical, and can be used as a
27 proper clock. Through this procedure, one ends up with a relational description of a reduced system
28 of variables which effectively represents the physical Hilbert space.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 We show how to recover the Friedmann dynamics of a flat FLRW universe at the level of
45 expectation values. Moreover, assuming that quantum fluctuations are small at small curvature,
46 they remain under control also large curvature, a feature shared with loop quantum cosmology
47 (LQC) models [32, 33]. We compare these results on fluctuations with previous works [28, 29] which
48
49
50
51

52
53 ¹ By “one-body operator” we mean an operator which can be written as an integral over a single combination of
54 creation and annihilation operators.
55

56 ² Note that this is the fundamental number operator and not the effective relational number operator, a function of a
57 relational clock value, typically used in GFT cosmology [24].
58
59
60

1
2
3 stressed the importance of a changing particle number in order to suppress or enhance quantum
4 fluctuations. This is in contrast with our approach here in which the particle number is constant
5 (keeping in mind that the notions of dynamics and time dependence differ in these different papers –
6 see next paragraph). Our results also provide additional insights into the role of relational dynamics
7 in emergent quantum gravity theories, emphasising the importance of coarse-graining (in our case
8 given by explicit truncation) before defining a notion of relationality, as suggested in [25].
9

10
11 Our method for implementing relational dynamics in GFT differs from what was done in previous
12 papers such as [24, 25, 27]. More precisely, while the approach proposed in this paper shares
13 with [25] the fact that relational evolution is defined only in an effective sense, in contrast to
14 [25] it does not require a specific form of states. In [24], instead, a relational notion of dynamics
15 was implemented by defining “relational operators” (which however need smearing in order to be
16 well-defined [25, 34]), while [27] employed a deparametrised framework from the start (the two
17 last approaches are connected in [30]). In our approach, we do not attempt to define microscopic
18 relational observables, nor we try to construct a reduced physical Hilbert space; these difficulties are
19 in fact bypassed by the effective approach and the notion of relationality is defined in the usual
20 sense of constrained dynamics on a phase space. A complete understanding of how these approaches
21 to relational dynamics are related to each other is still missing, and certainly deserves more study.
22
23

24
25 Finally, let us stress again that in order to obtain our results one does not need to specify a state
26 – only general semiclassicality and “hydrodynamic” conditions are assumed – thus suggesting that
27 the connection between GFTs and cosmological solutions may go beyond the choice of coherent
28 states (as in the simpler setting of [29]).
29
30
31

32 33 34 35 36 37 38 39 40 41 **II. REVIEW OF THE EFFECTIVE APPROACH**

42
43
44 In this section we review the effective approach to constrained systems [14, 15] and to relational
45 dynamics [16] (see also [17, 18]). While there are many fascinating aspects of this approach that
46 deserve to be discussed at length, we touch only the points necessary for the understanding of the
47 procedure developed in this paper, referring to [14–18] for more details.
48
49
50

51 52 53 **A. Effective constraints**

54
55
56 The effective approach to quantum constrained systems is formulated in the quantum phase
57 space, which can be constructed from first principles as follows [35, 36].
58
59
60

The starting point is a set of elements \hat{A}_i with $i = 1, \dots, n$ generating an associative unital algebra \mathcal{A} (which contains all finite polynomials in the \hat{A}_i) and satisfying the commutation relations

$$[\hat{A}_i, \hat{A}_j] = i\hbar f_{ij}{}^k \hat{A}_k \quad (2.1)$$

for some real constants $f_{ij}{}^k$. One usually assumes that this is an involutive algebra, i.e., there is a $*$ operation corresponding to the adjoint (where the generators are often thought of as self-adjoint).

Monomials of the form $\hat{A}_1^{m_1} \hat{A}_2^{m_2} \dots \hat{A}_n^{m_n}$ with $m_i \geq 0$ form a linear countable basis for \mathcal{A} . Now a quantum state ψ can be seen as a linear functional on \mathcal{A} , i.e., a map $\psi : \mathcal{A} \rightarrow \mathbb{C}$, with the conventional notation $\langle \hat{A}_i \rangle_\psi = \psi(\hat{A}_i)$ (and with $\langle \hat{1} \rangle_\psi = 1$). From an algebraic viewpoint this is the definition of the notion of state. One connection between this algebraic framework and the canonical description in terms of operators acting on Hilbert spaces is provided by the Gelfand, Naimark and Segal (GNS) theorem [37, 38] according to which, given any algebraic state α , it is always possible to construct an Hilbert space \mathcal{H}_α on which the algebra of observables act as an algebra of linear bounded operators $\pi_\alpha(\hat{A}_i)$, such that π_α is a $*$ -representation of \mathcal{A} . This Hilbert space \mathcal{H}_α contains a cyclic vector $|\alpha\rangle \in \mathcal{H}_\alpha$ such that $\alpha(\hat{A}_i) = \langle \alpha | \pi_\alpha(\hat{A}_i) | \alpha \rangle$. More generally, one may start from a Hilbert space \mathcal{H} on which the elements in \mathcal{A} act as operators; then one can again think of each $\psi \in \mathcal{H}$ as a linear functional by defining $\psi(\hat{A}_i) = \langle \psi | \hat{A}_i | \psi \rangle$. In either viewpoint, the statement relevant in the effective approach is that one is interested in elements of $\bar{\mathcal{A}}$, the dual vector space to \mathcal{A} . A subtle point here is that the existence of a Hilbert space representation would imply that $\langle \hat{A}_i^* \hat{A}_i \rangle_\psi \geq 0$ but it is mentioned in [35, 36] that this condition need not hold in the more general algebraic framework.

Because of linearity, any $\psi \in \bar{\mathcal{A}}$ is determined by the values it assigns to a linear basis of \mathcal{A} . So, one could coordinatise $\bar{\mathcal{A}}$ through the *expectation values* $\langle \hat{A}_i \rangle$ with $i = 1, \dots, n$ and *moments*

$$\Delta(A_1^{m_1} \dots A_n^{m_n}) = \left\langle \left(\hat{A}_1 - \langle \hat{A}_1 \rangle \right)^{m_1} \dots \left(\hat{A}_n - \langle \hat{A}_n \rangle \right)^{m_n} \right\rangle_{\text{Weyl}}, \quad (2.2)$$

where the subscript indicates a total symmetrisation of the product of operators inside the expectation value above, and where $\sum_{i=1}^n m_i \geq 2$. This choice of coordinatisation proves especially useful in effective approaches, where the states chosen are such that there is a clear hierarchy of moments, as we will discuss below.

The Poisson structure turning the state space $\bar{\mathcal{A}}$ coordinatised by expectation values and moments into the *quantum phase space* is directly inherited from the commutation relations of the elements

in \mathcal{A} : one defines³

$$\{\langle \hat{A} \rangle, \langle \hat{B} \rangle\} = (i\hbar)^{-1} \langle [\hat{A}, \hat{B}] \rangle \quad (2.3)$$

for any $\hat{A}, \hat{B} \in \mathcal{A}$ (it is enough to demand this for monomials in the generators \hat{A}_i). For constrained systems, the quantum phase space corresponds to a kinematical Hilbert space, since no constraint has been imposed yet. In this sense, variables in \mathcal{A} are kinematical quantum variables.

In the algebraic framework, we can now define a quantum constraint \hat{C} as an algebra element satisfying⁴ $\langle \hat{A}\hat{C} \rangle = 0$ for any $\hat{A} \in \mathcal{A}$ [15]. In other words, the constraint can be characterised by requiring that its expectation value and all of its fluctuations are zero:

$$C(\langle \hat{A}_i \rangle, \Delta(\dots)) = \langle \hat{C} \rangle = 0, \quad (2.4a)$$

$$C_{\text{pol}}(\langle \hat{A}_i \rangle, \Delta(\dots)) = \langle (\widehat{\text{pol}} - \langle \widehat{\text{pol}} \rangle) \hat{C} \rangle = 0, \quad (2.4b)$$

with $\widehat{\text{pol}}$ being a polynomial in the algebra elements. The notation here emphasises that the constraint functions should be expressed in terms of expectation values and moments⁵. Notice that, contrary to what is done for moments, the quantum constraints are not defined by a symmetric ordering. In turn, this implies that the quantum constraint functions can take complex values. This was however shown not to be an issue for deparametrisable systems, basically because the moments appearing in the constraints are *kinematical* moments and do not need to be restricted to real values. On the other hand, physical expectation values and moments must be real⁶ [15, 17].

The reason for this ordering choice is that a symmetric ordering would not in general lead to a first class system of constraints [15]. On the other hand, with the definition above it is easy to show that for any C_{pol} and $C_{\text{pol}'}$ defined as in (2.4b) we have

$$\{C_{\text{pol}}, C_{\text{pol}'}\} = (i\hbar)^{-1} \langle [\widehat{\text{pol}} \hat{C}, \widehat{\text{pol}'} \hat{C}] \rangle \approx 0$$

where \approx is Dirac's "weak equality" (equality up to functions that are multiplied by constraints), and so the system closes. As a consequence, these quantum constraint functions induce *quantum gauge*

³ This form of Poisson brackets can also be motivated by the geometrical formulation of quantum mechanics [31].

⁴ This can be seen as the algebraic counterpart of the Dirac condition $\hat{C}|\psi\rangle = 0$ for any state $|\psi\rangle$ in a Hilbert space, the connection between algebraic and Dirac formulation being provided by the GNS construction discussed above.

⁵ For non-polynomial constraints, this can be achieved via a Taylor expansion.

⁶ Notice that in the usual Hilbert space approach, $\hat{C}|\psi\rangle = 0$ likewise only implies that $\hat{A}\hat{C}|\psi\rangle = 0$ for any \hat{A} but not necessarily $(\hat{A}\hat{C} + \hat{C}\hat{A})|\psi\rangle = 0$, which could lead to an analogous discussion.

transformations on their solution space through their Hamiltonian flows. The presence of gauge flows even after solving the quantum constraint functions in the quantum phase space approach can be understood again as an effect of working with kinematical variables, which are indeed affected by gauge even at the classical level. Moments and expectation values of Dirac observables, instead, are gauge invariant also from this effective quantum phase space perspective. Indeed, if \hat{O} is a Dirac observable, the flow of its expectation value,

$$\begin{aligned} \langle \hat{O} \rangle'(\tau) &= \left\{ \langle \hat{O} \rangle, \left\langle \left(\widehat{\text{pol}} - \langle \widehat{\text{pol}} \rangle \right) \hat{C} \right\rangle \right\} \\ &= (i\hbar)^{-1} \left[\left\langle \left(\widehat{\text{pol}} - \langle \widehat{\text{pol}} \rangle \right) [\hat{O}, \hat{C}] \right\rangle + \left\langle [\hat{O}, \widehat{\text{pol}}] \left(\hat{C} - \langle \hat{C} \rangle \right) \right\rangle \right] \end{aligned} \quad (2.5)$$

where τ is parametrising the flow, vanishes weakly. The same of course is true for moments of Dirac observables. In this approach, the difficulty in solving the quantum constraint and explicitly constructing a physical Hilbert space has been traded for an infinite set of variables and constraint functions. These are in general intractable, unless one can identify some approximation that allows to reduce them to a finite set, neglecting subdominant contributions, and thus characterising the approach as an effective one. An example of such approximations is the semiclassical one, where it is assumed that moments of higher order are also higher order in \hbar . More precisely, for the semiclassical expansion one assumes that $\Delta(A_1^{m_1} \dots A_n^{m_n}) \sim \hbar^{(m_1 + \dots + m_n)/2}$. At a given order in \hbar , only a finite number of constraints and moments are relevant. By construction this hierarchy is also preserved by the Poisson structure. Indeed, one can show from (2.3) that the Poisson bracket of a moment of order $\hbar^{m/2}$ and a moment of order $\hbar^{n/2}$ is of order $\hbar^{(m+n-2)/2}$.

In the following, we will assume that a truncation at order \hbar captures the dynamics of suitable semiclassical states to a good approximation. This means that we will neglect all terms explicitly proportional to $\hbar^{3/2}$ or higher and all third or higher order moments. For an algebra generated by n elements \hat{A}_i , we then have $n + 1$ constraints $C = \langle \hat{C} \rangle = 0$ and $C_{A_i} = \langle (\hat{A}_i - \langle \hat{A}_i \rangle) \hat{C} \rangle = 0$ and $n + n(n + 1)/2 = n(n + 3)/2$ expectation values and moments.

The extension of the Poisson bracket (2.3) for an algebra with commutation relations (2.1) is, at the given truncation to order \hbar , given by the relations

$$\{ \langle \hat{A}_i \rangle, \Delta(A_j A_k) \} = f_{ij}{}^l \Delta(A_l A_k) + (j \leftrightarrow k), \quad (2.6a)$$

$$\{ \Delta(A_i A_j), \Delta(A_k A_l) \} = \left(f_{ik}{}^m \langle \hat{A}_m \rangle \Delta(A_j A_l) + (i \leftrightarrow j) \right) + (k \leftrightarrow l). \quad (2.6b)$$

B. Effective relational dynamics

Within the effective framework described above, the way physical variables are selected is through an effective local deparametrisation of the system [15, 17]. This is obtained via a partial gauge fixing of the aforementioned gauge freedom (*Zeitgeist*), leading to a projection into a classical “time”, and consequently to the definition of transient relational observables called *fashionables*. In [15, 17] most of the features of this effective approach to relational dynamics are explained through different (classical deparametrisable and non-deparametrisable) examples. Here we will try to provide a general summary of such features, most of which will also be true for the system we aim to study in the rest of this paper. An important feature of the effective approach explained above is that the quantum phase space generically has a degenerate Poisson structure, i.e., is not symplectic. Indeed, moments of any arbitrary order are characterised, in general, by a degenerate Poisson tensor⁷. A crucial consequence of this degeneracy of the Poisson structure is that even though the constraints C_{pol}, C are independent, the gauge flows generated by them are generally not. The Poisson geometry of constrained systems is discussed in [39].

In the examples studied in [17], at order \hbar there are $n + 1$ constraints, but only n gauge flows generated by them are independent. This means that of the $n(n + 3)/2$ variables relevant at that order, $2n$ variables can be eliminated using the constraints ($n + 1$ conditions) and $n - 1$ by fixing all the gauge flows but one. This leaves $n(n + 3)/2 - 2n = n(n - 1)/2$ remaining variables. The importance of this counting can be understood when one is interested, as we have mentioned above, in an effective deparametrisation of the system. Classically, such a deparametrisation would result in the complete elimination of a (possibly canonical) pair of variables. At the effective level, eliminating all variables related to a pair of operators having closed non-trivial commutation relations among themselves leads to a total of $(n - 2)(n + 1)/2 = n(n - 1)/2 - 1$ variables. So, by appropriately fixing $n - 1$ gauge flows and using the $n + 1$ constraints (see below), one could obtain a set of $(n - 2)(n + 1)/2$ physical variables and one classical “clock variable”, all of them subject to the remaining, unfixed gauge flow. In turn, this allows constructing local relational observables (*fashionables*) describing the evolution of the physical variables with respect to the clock.

Let us discuss in more detail how this procedure can be realised concretely. The theory is entirely clock neutral, since all physical variables are treated on equal footing. The partial gauge fixing mentioned above, or *Zeitgeist*, has exactly the purpose of choosing a specific clock, projecting the

⁷ As an example, consider a canonical pair \hat{q}, \hat{p} . At order \hbar , we have 3 moments, so the space is odd-dimensional and the Poisson structure of second order moments must be degenerate.

quantum clock we are interested in onto a classical parameter. Of course, such a clock, say \hat{T} , needs to have non-trivial commutation relations with \hat{C} , i.e., the clock may not be a Dirac observable since otherwise it is constant (cf. (2.5)).

We will assume this, and we will again further restrict to an order \hbar truncation. If the clock also forms a closed subalgebra with another operator, say \hat{P} (which is the situation studied in [15, 17] and considered below), the $n - 1$ conditions

$$\phi_\alpha = \Delta(TA_\alpha) = 0, \quad \alpha \in \mathcal{A} \setminus \{\hat{P}\}, \quad (2.7)$$

fix $n - 1$ gauge flows at order \hbar , and make the clock classical on the space of physical variables. Then, if the expectation value and the n moments of the variable \hat{P} can be eliminated using the $n + 1$ effective constraints⁸, one indeed arrives at a physical description involving the $(n - 2)(n + 1)/2$ expectation values and order \hbar moments of the remaining $n - 2$ variables, and $\langle \hat{T} \rangle$.

Once the gauge fixing conditions (2.7) are imposed, one is left with a mixture of first and second class constraints. However, since

$$\begin{aligned} \{\phi_\alpha, \phi_\beta\} &= \frac{1}{i\hbar} \left\langle \left[(\hat{T} - \langle \hat{T} \rangle)(\hat{A}_\alpha - \langle \hat{A}_\alpha \rangle), (\hat{T} - \langle \hat{T} \rangle)(\hat{A}_\beta - \langle \hat{A}_\beta \rangle) \right] \right\rangle \\ &= \frac{1}{i\hbar} \left\langle (\hat{T} - \langle \hat{T} \rangle)^2 \left[\hat{A}_\alpha - \langle \hat{A}_\alpha \rangle, \hat{A}_\beta - \langle \hat{A}_\beta \rangle \right] \right\rangle = f_{\alpha\beta}{}^\gamma \left\langle (\hat{T} - \langle \hat{T} \rangle)^2 \hat{A}_\gamma \right\rangle \\ &\simeq f_{\alpha\beta}{}^\gamma \langle \hat{A}_\gamma \rangle \Delta(T^2) \approx 0, \end{aligned}$$

where in the third line we have neglected higher order terms, we notice that the matrix of Poisson brackets $\Delta_{ij} = \{\chi_i, \chi_j\}$, where χ_i include first order constraints and the gauge fixing conditions (2.7), is actually off-block diagonal, with non-zero values only involving physical constraints and gauge fixing conditions. This, together with the fact that the $n(n - 1)/2$ remaining variables have weakly vanishing Poisson brackets with the gauge fixing constraints (which can be proven in an analogous way), shows that the Dirac brackets

$$\{f, g\}_{\text{Dirac}} = \{f, g\} - \{f, \chi_i\} \Delta^{ij} \{\chi_j, g\}, \quad \Delta^{ij} = (\Delta^{-1})_{ij} \quad (2.8)$$

for the $n(n - 1)/2$ remaining variables are identical to the Poisson brackets introduced above. Notice that the remaining moments must, at this point, be real, contrary to what may happen for moments of \hat{P} (see for instance [15, 17]) or for the expectation value of \hat{T} itself (see [17]).

⁸This for instance happens if the quantum constraint is linear in the variable \hat{P} , which will be a case of interest in the following.

Once a Zeitgeist has been chosen, for instance $T = \langle \hat{T} \rangle$ as specified above, one looks for the remaining gauge flow that preserves the gauge fixing conditions and, through that, one constructs relational quantities of the form $\langle \hat{A}_\alpha \rangle \circ T^{-1}$ and $\Delta(A_\alpha A_\beta) \circ T^{-1}$. These observables are transient in nature, since they can only be meaningful as long as the clock does not reach any turning point and maintains a monotonic evolution. When this is the case, one is forced to move from one Zeitgeist to another one; correspondingly, these observables may “fall out of fashion”. As such, these “fashionables” can only be state dependent. We refer again to [16, 17] for more details on this topic, and [18] for an explicit example of a model that does not have a globally defined clock. In the case we will be interested in, we will choose a clock variable with globally monotonic evolution, so that the Zeitgeist is valid globally.

C. Extending the effective approach to field theory

The effective approach to constrained systems is usually applied to quantum-mechanical systems of a finite number of degrees of freedom, often with a single constraint \hat{C} . The generators \hat{A}_i of \mathcal{A} then usually represent the quantum analogue of phase space coordinates for a classical system one proposes to quantise (e.g., positions and momenta of a system of particles). We saw above that the quantum phase space $\bar{\mathcal{A}}$ can be identified with an (algebraically constructed) set of quantum states on the algebra \mathcal{A} , or with a Hilbert space that the elements of \mathcal{A} could be seen as acting on.

In this paper, we want to extend the application of this approach to a field theory setting. A quantum field theory can be seen as the quantisation of a system with already classically infinitely many degrees of freedom, and would therefore be associated with a very large observable algebra \mathcal{A}_{QFT} (as is the starting point, e.g., in algebraic quantum field theory). We will however restrict to a much smaller algebra representing only a few coarse-grained observables we are interested in. That is, our algebra \mathcal{A} will only be a very small subalgebra of the full \mathcal{A}_{QFT} . It then follows that the quantum phase space $\bar{\mathcal{A}}$ of interest to us is much smaller than the field theory quantum phase space $\bar{\mathcal{A}}_{\text{QFT}}$. The smaller $\bar{\mathcal{A}}$ can then perhaps still be identified with some Hilbert space, but this will not be the Hilbert space of the field theory \mathcal{H}_{QFT} , which is much larger. Indeed there are in general different elements of \mathcal{H}_{QFT} that would be identified with the same element of $\bar{\mathcal{A}}$ in the effective approach; these are states that agree on expectation values and moments of macroscopic observables even though their microscopic definition may be different⁹.

⁹ This is consistent with a coarse-graining interpretation of the reduction to an observable sub-algebra.

1
2
3 Since the effective approach as reviewed here starts from the algebra \mathcal{A} and a set of constraints
4 only involving elements in \mathcal{A} , the question of how the quantum phase space $\bar{\mathcal{A}}$ might sit inside a
5 much larger Hilbert space is not of direct relevance for its application. In this sense, we will not
6 need to give an explicit construction of $\bar{\mathcal{A}}$ as a subspace of \mathcal{H}_{QFT} which in our case will be the GFT
7 Fock space. Our interpretation is that $\bar{\mathcal{A}}$ is a quantum phase space resulting from a coarse graining
8 of the field theory, which leads to an effective description characterised by a few operators (among
9 the infinitely many that can be constructed out of the field operators) whose expectation values and
10 moments coordinatise an effective quantum phase space.
11
12
13
14
15
16
17
18
19

20 III. TRUNCATING THE DYNAMICS TO ONE-BODY OPERATORS

21
22
23 In this paper we apply the effective approach outlined in the previous section to the dynamics of
24 group field theory, in order to obtain a systematic semiclassical expansion that does not require
25 specifying either a particular state or a particular relational clock. Instead, we will only assume
26 that the state is sufficiently semiclassical for an \hbar expansion to make sense, and we will construct a
27 clock using the strategy of deparametrisation and gauge fixing by a Zeitgeist explained above.
28
29
30
31

32 As we already outlined at the end of the previous section, out of the infinitely many independent
33 observables that can be constructed out of the GFT field operators we will only study a few “global”
34 one-body operators, and truncate the dynamics to a system of equations that only makes reference
35 to those operators. Clearly the choice of operators in our algebra is to some extent determined by the
36 dynamics of the theory. We implicitly assume that those few global quantities are sufficient to capture
37 the most relevant dynamical information of the field theory. Such an assumption is conceptually
38 similar to assuming a homogeneous cosmology in general relativity, which amounts to the assumption
39 that a few global quantities (the volume of space, a homogeneous energy density, etc.) capture the
40 dynamics of the Universe. Here we see this truncation as representing a coarse-grained description of
41 the field theory in which one is only interested in some macroscopic variables, which is justified for
42 states for which the coarse-grained description is “reasonably close” to the full microscopic dynamics
43 of the field theory. This viewpoint is then the quantum analogue of the classical perspective that
44 a homogeneous, isotropic universe in classical cosmology is not a very symmetric solution to the
45 Einstein equations but should rather be seen as obtained from averaging over inhomogeneities on
46 small scales [40, 41]. In the setting of quantum field theory, understanding the conditions for this
47 truncation to be valid is a more challenging problem.
48
49
50
51
52
53
54
55
56
57
58
59
60

A. Effective GFT quantum constraint

We now introduce the details of our effective approach to GFT dynamics. First, we introduce the type of field theories we are studying. We consider GFT models with domain $G^4 \times \mathbb{R}$, where G is a Lie group representing gravitational degrees of freedom and \mathbb{R} represents a scalar matter field. Writing the field as $\varphi(g_I, \chi)$ the dynamics are then schematically defined by an action

$$S[\varphi, \bar{\varphi}] = \int d^4g d\chi \bar{\varphi}(g_I, \chi) \mathcal{D}\varphi(g_I, \chi) + I[\varphi, \bar{\varphi}] \quad (3.1)$$

where \mathcal{D} is a differential operator on $G^4 \times \mathbb{R}$ and $I[\varphi, \bar{\varphi}]$ is the “interaction” part of the action which includes terms of higher than second order in the fields. We will focus on the free part and mostly neglect the contribution from interaction terms in the following. Such an action gives rise to equations of motion

$$\mathcal{D}\varphi(g_I, \chi) + \frac{\delta I[\varphi, \bar{\varphi}]}{\delta \bar{\varphi}(g_I, \chi)} = 0 \quad (3.2)$$

and the complex conjugate equation for $\bar{\varphi}$, but rather than solving this local equation (either at the full quantum level or in a mean-field approximation) we can multiply from the left with a conjugate field and integrate to obtain

$$\int d^4g d\chi \bar{\varphi}(g_I, \chi) \mathcal{D}\varphi(g_I, \chi) + \text{higher order} = 0, \quad (3.3)$$

i.e., the quadratic part of the GFT action has to vanish if we again neglect the effect of higher order contributions. (3.3), with the higher order parts assumed to be negligible, is our starting point.

We can now specify the form of \mathcal{D} further in order to make the form of the resulting dynamics more explicit. A minimal non-trivial \mathcal{D} would include a “mass term” and second derivatives with respect to the group arguments, i.e.,

$$\mathcal{D} = m^2 + \frac{\hbar^2}{4} \sum_{I=1}^4 \Delta_{g_I} + \lambda \hbar^2 \partial_\chi^2 \quad (3.4)$$

where Δ_{g_I} is a Laplace–Beltrami operator on G , acting on the argument I (the reason for the factor $\frac{1}{4}$ will be clear shortly). (3.4) is a standard form for the kinetic term of a GFT model, which has been studied in previous papers on GFT cosmology such as, e.g., [22, 42, 43]. It can be motivated by results from GFT renormalisation which show that Laplace–Beltrami operators with respect to group arguments are generated by radiative corrections [44, 45], even if one started with a pure mass term as is often done in the spin foam literature [46]. It could also be motivated by an effective field theory expansion in derivatives, with the further assumption that the theory should be invariant

with respect to arbitrary translations in $G^4 \times \mathbb{R}$ so that explicit dependence on g_I or χ is excluded. We will consider (3.4) for now but discuss possible extensions below. Notice the appearance of factors of \hbar associated to each derivative, which are standard in quantum mechanics but often dropped in quantum field theory by setting $\hbar = 1$. In our setting we will perform a semiclassical expansion in powers of \hbar so it is important to keep these powers manifest.

The discussion so far was classical. In the quantum theory, we would expect any physical state $|\psi\rangle$ to satisfy

$$\left(m^2 + \frac{\hbar^2}{4} \sum_{I=1}^4 \Delta_{g_I} + \lambda \hbar^2 \partial_\chi^2 \right) \hat{\varphi}(g_I, \chi) |\psi\rangle = 0 \quad (3.5)$$

but we will now replace this local condition by a much weaker, integrated version

$$\hat{C}|\psi\rangle = \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) \left(m^2 + \frac{\hbar^2}{4} \sum_{I=1}^4 \Delta_{g_I} + \lambda \hbar^2 \partial_\chi^2 \right) \hat{\varphi}(g_I, \chi) |\psi\rangle = 0 \quad (3.6)$$

which we can see as a constraint equation $\hat{C}|\psi\rangle = 0$ for a certain one-body operator \hat{C} .

It is clear that (3.6) is only one out of an in principle infinitely long list of similar conditions¹⁰, obtained by integrating the local equation of motion with different operators from the left¹¹. Having made this truncation, the next step is an expansion in powers of \hbar in which the still fully-quantum mechanical condition (3.6) is replaced by a series of statements about expectation values. This expansion is the essence of the effective approach which we are importing into the GFT setting.

B. Defining the observable algebra \mathcal{A}

In the discussion of Section II we presented the construction of a quantum phase space as starting from an observable algebra \mathcal{A} generated by a finite number of elements; quantum constraints were constructed from the elements of \mathcal{A} . In the field theory setting of GFT, it is advantageous to proceed in the opposite direction: having defined a constraint (3.6) we now introduce a number of one-body operators whose dynamics are controlled by (3.6). This set should be large enough to capture at least some physically interesting and non-trivially evolving quantities, but still small enough to keep calculations under control. We will find that this balance between making non-trivial statements about effective dynamics and keeping the complexity of the approach under control will already

¹⁰ It can be seen as the counterpart of the infinite set of Schwinger–Dyson equations.

¹¹ Here, due to the linearity of (3.5), the constraints obtained by this procedure only relate operators constructed out of the same number of field operators: each integrated constraint only involves n -body operators with the same n .

impose quite stringent limitations on how many quantities we can and should include. The basic operators we define are then the \hat{A}_i appearing in (2.1), i.e., the generators of \mathcal{A} .

The form of (3.6) suggests the definitions

$$\hat{N} = \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) \hat{\varphi}(g_I, \chi), \quad (3.7a)$$

$$\hat{\Lambda}_I = -\hbar^2 \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) \Delta_{g_I} \hat{\varphi}(g_I, \chi), \quad (3.7b)$$

$$\hat{\Pi}_2 = -\hbar^2 \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) \partial_\chi^2 \hat{\varphi}(g_I, \chi). \quad (3.7c)$$

\hat{N} corresponds to the GFT number operator. As a further simplification we will assume that the state $|\psi\rangle$ we are working with satisfies $\hat{\Lambda}_I|\psi\rangle = \hat{\Lambda}_J|\psi\rangle$ for any I and J (which one might interpret as a notion of isotropy). We will then simply write $\hat{\Lambda}$ instead of $\hat{\Lambda}_I$, given that all equations involving any of the operators $\hat{\Lambda}_I$ will be independent of the label I , and the sum over I will cancel the factor $\frac{1}{4}$ in (3.9). With these considerations the quantum constraint operator defined in (3.6) is given by

$$\hat{C} = m^2 \hat{N} - \hat{\Lambda} - \lambda \hat{\Pi}_2. \quad (3.8)$$

We will later replace the operator equation $\hat{C}|\psi\rangle = 0$ by a series of constraints for expectation values and higher moments, based on the effective approach introduced in Section II. This set of equations corresponds to a semiclassical expansion in powers of \hbar . At lowest order \hbar^0 , we have the non-trivial requirement that the sum of expectation values

$$C = \langle \hat{C} \rangle = m^2 \langle \hat{N} \rangle - \langle \hat{\Lambda} \rangle - \lambda \langle \hat{\Pi}_2 \rangle \quad (3.9)$$

should vanish.

The one-body operators we have defined so far all commute with each other, and thus are not governed by any non-trivial dynamics. We therefore also introduce a one-body operator

$$\hat{X} = \int d^4g d\chi \chi \hat{\varphi}^\dagger(g_I, \chi) \hat{\varphi}(g_I, \chi) \quad (3.10)$$

which we will interpret as corresponding to the value of a matter scalar field (summed over all quanta in a state, since this is an extensive quantity). If we assume the commonly used canonical commutation relations

$$[\hat{\varphi}(g_I, \chi), \hat{\varphi}^\dagger(g'_I, \chi')] = \delta(\chi - \chi') \int dh \prod_{I=1}^4 \delta(g_I h (g'_I)^{-1}) \quad (3.11)$$

for the field operators, where the slightly unusual integration over h ensures that this commutator is compatible with the gauge invariance property [22]

$$\hat{\varphi}(g_I, \chi) = \hat{\varphi}(g_I h, \chi) \quad \forall h \in \text{SU}(2), \quad (3.12)$$

we find

$$[\hat{X}, \hat{\Pi}_2] = 2i\hbar \left(-i\hbar \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) \partial_\chi \hat{\varphi}(g_I, \chi) \right) = 2i\hbar \hat{\Pi}_1 \quad (3.13)$$

so that $\hat{\Pi}_1$ defined by this equation also needs to be added to our algebra of one-body operators.

We then also find

$$[\hat{X}, \hat{\Pi}_1] = i\hbar \hat{N} \quad (3.14)$$

while all other commutators among quantities we have considered so far (including all commutators of \hat{N} or $\hat{\Lambda}$) are zero. We also add a one-body operator \hat{K} which has a non-trivial commutator with $\hat{\Lambda}$ but commutes with all other operators, such that the full set of operators forms a closed algebra. We will interpret the operator \hat{K} as corresponding to a notion of extrinsic curvature. The precise form of the commutator is unspecified for now; later on, we will constrain it by physical arguments and find explicit realisations of such an operator in particular examples.

The six one-body operators we have defined provide the basis for the extraction of effective cosmological dynamics. The operators are subject to a single constraint (3.8). In general, it is easy to find examples of systems of one-body operators subject to more than one constraint¹², but the specific form of the equations of motion and the algebra defined above guarantee that (3.8) is the only constraint that involves operators in the algebra alone. Indeed, consider an action on (3.5) from the left resulting in an integrated constraint which features an operator included in the above algebra but different from \hat{N} , $\hat{\Lambda}$, and $\hat{\Pi}_2$ (e.g., \hat{K}). This new constraint would inevitably involve also new operators whose matrix elements depend both on the group and the scalar field variables, which are however not included in the algebra of operators of interest. In this sense, we are considering a minimal self-consistent set of operators and dynamical relations.

Notice again that in this approach there is no dependence on any time parameter, nor a viewpoint that any of these operators define by themselves a notion of relational observable. Any notion of relational dynamics will only arise from comparing the dynamics of these operators to each other. Importantly, and in line with the philosophy of the effective approach, we make no explicit assumption about the quantum state, only that it is sufficiently semiclassical (with respect to these global observables). We do however see our approximation (of not solving local field equations but

¹² For instance, consider a hypothetical GFT model with quantum field equation $(f(g_I, \chi) + 1)\hat{\varphi}(g_I, \chi) = 0$, and consider the operators \hat{N} , $\hat{F}_+ = \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) f(g_I, \chi) \hat{\varphi}(g_I, \chi)$ and $\hat{F}_- = \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) f^{-1}(g_I, \chi) \hat{\varphi}(g_I, \chi)$. The equations of motion would then imply two separate constraints $\hat{F}_+ + \hat{N} = 0$ and $\hat{N} + \hat{F}_- = 0$.

only demanding that our one-body operators satisfy certain “averaged” relations) as an implicit condition on the quantum state.

C. Fixing the free commutator by analogy with classical Friedmann cosmology

The lowest-order truncation to the full quantum constraint would be to only require that the expectation value of the constraint vanishes. This leads to the constraint

$$C = m^2 \langle \hat{N} \rangle - \langle \hat{\Lambda} \rangle - \lambda \langle \hat{\Pi}_2 \rangle \approx 0 \quad (3.15)$$

that we already defined in (3.9). At the same level of truncation, the dynamics of our six quantum one-body operators can be reduced to the dynamics of the expectation values of these operators. The resulting dynamical system is then equivalent to a classical constrained system: the expectation values represent phase space variables subject to a single Hamiltonian constraint. The constraint generates gauge transformations, which we will assume can be interpreted as reparametrisation in time or as the time evolution flow generated by a Hamiltonian built from this constraint.

We can then demand as a minimal requirement that this system corresponds to the dynamics of a homogeneous isotropic Friedmann–Lemaître–Robertson–Walker (FLRW) universe in general relativity or a modified version of it which includes higher-curvature corrections. We will use this minimal requirement to fix the so far unspecified commutator between $\hat{\Lambda}$ and \hat{K} .

The flow generated by (3.15) is

$$\langle \hat{N} \rangle'(t) = \langle \hat{\Pi}_1 \rangle'(t) = \langle \hat{\Pi}_2 \rangle'(t) = \langle \hat{\Lambda} \rangle'(t) = 0 \quad (3.16)$$

so that we immediately have four constants of motion, and

$$\langle \hat{X} \rangle'(t) = -2\lambda \langle \hat{\Pi}_1 \rangle(t), \quad \langle \hat{K} \rangle'(t) = \left\{ \langle \hat{\Lambda} \rangle, \langle \hat{K} \rangle \right\}(t). \quad (3.17)$$

As we have stated when introducing it, the variable \hat{X} represents a matter scalar field whose value is summed over all quanta in a GFT state. Normally we would have to worry about the fact that this is an extensive variable and perhaps divide by the average particle number to obtain an intrinsic notion of matter scalar field. In our framework here, the variable $\langle \hat{N} \rangle$ is always a constant of motion, since the operator \hat{N} commutes with all other one-body operators we consider, hence we do (for now) not have to worry about the distinction between intensive and extensive quantities.

An immediate consequence from (3.16) and (3.17) is then that the evolution of $\langle \hat{X} \rangle$ is monotonic if we assume that the expectation value $\langle \hat{\Pi}_1 \rangle$ does not vanish, indeed we have

$$\langle \hat{X} \rangle(t) = \langle \hat{X} \rangle(0) - 2\lambda \langle \hat{\Pi}_1 \rangle(0) t. \quad (3.18)$$

$\langle \hat{X} \rangle$ can then be seen as a good clock, very similar to the way massless scalar fields often appear as clocks in quantum gravity and quantum cosmology.

Let us now compare this system explicitly with the dynamics of a flat FLRW universe filled with a massless scalar field in general relativity. For this system, the Hamiltonian is (see, e.g., [47])

$$\mathcal{H}_{\text{GR}} = N \left(-\frac{2\pi G}{3} \frac{p_a^2}{a} + \frac{p_\varphi^2}{2a^3} \right) \quad (3.19)$$

where a is the scale factor, p_a is its canonical momentum and p_φ is the conjugate momentum of the scalar field φ . A convenient lapse choice, $N = a^3$, transforms this to the Hamiltonian

$$\mathcal{H}_{\text{GR}}|_{N=V} = -6\pi G V^2 p_V^2 + \frac{p_\varphi^2}{2} \quad (3.20)$$

where we switch to a volume variable $V = a^3$ with conjugate momentum p_V . The equation of motion for the scalar field is now $\dot{\varphi} = p_\varphi$ and p_φ is a constant of motion, a similar form to the first equation in (3.17). This similarity would motivate further the identification of $\langle \hat{X} \rangle$ as corresponding to a matter scalar field. How about the equations for the spacetime geometry? We have

$$\dot{V} = \{V, \mathcal{H}_{\text{GR}}|_{N=V}\} = -12\pi G (V p_V) V \quad (3.21)$$

so that, taking into account that $V p_V$ is clearly a constant of motion, on-shell we have $\dot{V} = cV$ for some constant of motion c , leading to exponential solutions. Very similarly we have

$$\dot{p}_V = \{p_V, \mathcal{H}_{\text{GR}}|_{N=V}\} = 12\pi G (V p_V) p_V \quad (3.22)$$

and again exponential solutions (with inverse sign in the exponent, again ensuring that $V p_V$ is a constant of motion). Notice that from the vanishing of (3.20) we have $V p_V = \pm p_\varphi / \sqrt{12\pi G}$ where the sign is determined by initial conditions.

The explicit solutions for the scalar field φ and extrinsic curvature p_V are then given by

$$\varphi(t) = p_\varphi t + \varphi(0), \quad (3.23a)$$

$$p_V(t) = p_V(0) \exp(\pm \sqrt{12\pi G} p_\varphi t), \quad (3.23b)$$

where p_φ and the sign \pm are constants of motion. This explicit form shows in particular that φ and p_V are globally monotonic functions, and so either of them could be used as a clock for the other degrees of freedom. If we are only interested in comparing the variables φ and p_V to each other, we can give the relation between the two as

$$(\varphi \circ p_V^{-1})(p_V) = \pm \frac{1}{\sqrt{12\pi G}} \log \left(\frac{p_V}{p_V(0)} \right) + \varphi(0), \quad (3.24a)$$

$$(p_V \circ \varphi^{-1})(\varphi) = p_V(0) \exp \left(\pm \sqrt{12\pi G} (\varphi - \varphi(0)) \right). \quad (3.24b)$$

Notice that p_V cannot change sign, so the logarithm in the first relation is always well-defined.

We now want to reproduce these classical cosmological solutions, expressed in relational language. Comparing (3.22) with the second equation in (3.17), if we assume that the quantity $\langle \hat{K} \rangle$ is analogous to p_V in the classical cosmological model, and hence represents extrinsic curvature, dynamics analogous to those of general relativity will be obtained from assuming that

$$\{\langle \hat{\Lambda} \rangle, \langle \hat{K} \rangle\} \propto \langle \hat{K} \rangle. \quad (3.25)$$

In the example of general relativity, this proportionality is only given on-shell (identifying the phase space variable Vp_V with a conserved quantity) so *a priori* it would make sense to assume that also (3.25) only holds on-shell. However, the basic premise of our effective approach is to assume that the commutator of two elementary variables is a linear combination of elementary variables, as in (2.1). The proportionality factor in (3.25) then needs to be just a numerical constant, which we will denote by α ; we then add the commutator

$$[\hat{\Lambda}, \hat{K}] = i\hbar \alpha \hat{K} \quad (3.26)$$

to our algebra of basic operators. This commutator would also suggest that $\langle \hat{\Lambda} \rangle$ is analogous to Vp_V in the classical theory, consistent with the fact that $\langle \hat{\Lambda} \rangle$ is constant (cf. (3.16)).

For completeness, we also discuss the most general case in which the commutator takes the form

$$[\hat{\Lambda}, \hat{K}] = i\hbar \sum_{a \in \mathcal{V}} \alpha_a \hat{a} \quad (3.27)$$

where \mathcal{V} is a set of labels (“variables”), $\mathcal{V} = \{N, \Pi_1, \Pi_2, X, \Lambda, K\}$. Interestingly, even this most general form can be solved exactly, with the solution to (3.17) in the case of $\alpha_K \neq 0$ given by

$$\langle \hat{K} \rangle(t) = \left(\frac{x}{\alpha_K} - \frac{\alpha_X}{\alpha_K^2} 2\lambda \langle \hat{\Pi}_1 \rangle(0) \right) (e^{\alpha_K t} - 1) + \langle \hat{K} \rangle(0) e^{\alpha_K t} + \frac{\alpha_X}{\alpha_K} 2\lambda \langle \hat{\Pi}_1 \rangle(0) t \quad (3.28)$$

and in the case $\alpha_K = 0$ by the expression

$$\langle \hat{K} \rangle(t) = \langle \hat{K} \rangle(0) + xt - \alpha_X \lambda \langle \hat{\Pi}_1 \rangle(0) t^2, \quad (3.29)$$

where in both cases we defined

$$x = \alpha_N \langle \hat{N} \rangle(0) + \alpha_{\Pi_1} \langle \hat{\Pi}_1 \rangle(0) + \alpha_{\Pi_2} \langle \hat{\Pi}_2 \rangle(0) + \alpha_X \langle \hat{X} \rangle(0) + \alpha_\Lambda \langle \hat{\Lambda} \rangle(0). \quad (3.30)$$

This result would suggest that a more general form of the commutator between $\hat{\Lambda}$ and \hat{K} could lead to a modified effective cosmology, different from the classical case. In principle, this is what we would expect: the effective cosmology of a quantum gravity theory such as GFT does not need to

exactly reproduce classical Friedmann cosmology, and high-energy corrections such as those leading to a bounce [24, 29, 48] are expected and desirable. In the closely related setting of loop quantum cosmology, the expression for the extrinsic curvature also receives such corrections compared to the classical case, and is proportional to $\arctan(\exp(\pm\sqrt{12\pi G}\phi))$ rather than the classical exponential (3.24b) [49, 50], and these corrections are responsible for the resolution of the cosmological singularity.

However, if we at least require the resulting effective dynamics to reduce to classical cosmology when curvature is low, so that there is a regime for t in which $\langle\hat{K}\rangle(t)$ asymptotically approaches $e^{\alpha\kappa t} \ll 1$, we do obtain quite stringent constraints from (3.28): we must have

$$x = \alpha_X = 0. \quad (3.31)$$

The first condition $x = 0$ can be satisfied by fine-tuning initial conditions or, if one requires it to hold for generic initial states, by setting

$$\alpha_N = \alpha_{\Pi_1} = \alpha_{\Pi_2} = \alpha_\Lambda = 0 \quad (3.32)$$

reducing us to the case (3.26), which we will therefore assume in the following. It would be interesting to study in future work whether the more general solution (3.28) can be given a physical interpretation perhaps in terms of a modified theory of gravity.

We conclude this section with a few remarks about the operator \hat{K} . Firstly, the above discussion tentatively puts forward an identification of \hat{K} with the extrinsic curvature. However, it should also be possible to motivate the operator representing extrinsic curvature by considering the microscopic degrees of freedom. Note that such a motivation from microscopic degrees of freedom has been employed previously in GFT where the definition of the volume operator has been imported from loop quantum gravity (LQG) [24, 48]. In our framework this would mean that we are able to specify the integral kernel in the definition

$$\hat{K} = \int d^4g d^4g' d\chi d\chi' \hat{\varphi}^\dagger(g_I, \chi) K(g_I, g'_I, \chi, \chi') \hat{\varphi}(g'_I, \chi'). \quad (3.33)$$

Since such one-body operators describe local quantities from the perspective of an LQG spin network, the identification of curvature degrees of freedom is not straightforward (see, e.g., the discussion in [22]). A starting point could be the *twisted geometries* parametrisation of the LQG phase space, which identifies a discrete analogue of extrinsic curvature for each spin network link [51].

Secondly, it is not possible to find operators representing the algebra (3.26) on compact spaces.

To see that this is true, note that on a compact space the eigenbasis of the Laplacian will be countable. Assume that we have Hermitian matrices K_{ij} and $\Lambda_{ij} = \Lambda_i \delta_{ij}$ representing the algebra

on this countable eigenbasis of the Laplacian. Then the commutator gives

$$(\Lambda_i - \Lambda_j)K_{ij} = i\hbar\alpha K_{ij} \quad (3.34)$$

which implies that $K_{ij} = 0$. In our GFT setting, this argument would imply that a representation of (3.26) only exists if the Lie group G chosen to define the GFT model is non-compact.

Let us stress, however, how crucial the specific form of the microscopic dynamics is also for this interpretation of \hat{K} as an extrinsic curvature operator. Indeed, as an example consider, instead of (3.5), the constraint equation

$$\left(m(g_I)^2 + \frac{\hbar^2}{4} \sum_{I=1}^4 \Delta_{g_I} + \lambda(g_I) \hbar^2 \partial_\chi^2 \right) \hat{\varphi}(g_I, \chi) |\psi\rangle = 0, \quad (3.35)$$

where both $m(g_I)^2$ and $\lambda(g_I)$ are non-trivial¹³ and non-singular functions of the group variables, such that their ratio is a constant, $m(g_I)^2/\lambda(g_I) = \tilde{\lambda}$. We could then consider the one-body quantum constraint

$$\hat{C} = \hat{N} - \hat{\Lambda} - \tilde{\lambda} \hat{\Pi}_2 \quad (3.36)$$

with

$$\hat{\Lambda} = -\hbar^2 \int d^4g d\chi \hat{\varphi}^\dagger(g_I, \chi) m(g_I)^{-2} \Delta_{g_I} \hat{\varphi}(g_I, \chi). \quad (3.37)$$

The dynamics generated by the expectation value of the above constraint would be the same as before (putting $m^2 = 1$), thus implying that an operator \hat{K} such that

$$[\hat{\Lambda}, \hat{K}] = i\hbar\alpha \hat{K} \quad (3.38)$$

could be interpreted classically as extrinsic curvature. This operator, however, would be substantially different from \hat{K} introduced above: since $\hat{\Lambda}$ would not be diagonal on the Laplacian eigenbasis, one could not exclude a representation of \hat{K} on a compact Lie group G on the basis of the argument given above for \hat{K} .

This should make clear that the precise geometric interpretation of both fundamental and effective observables is a non-trivial matter and should be taken with some caution.

¹³ The dependence of m^2 and λ on the group variables is generally expected for GFTs constructed by path-integral identification with simplicial gravity models minimally coupled with a free massless scalar field [52]. This dependence actually encodes the non-trivial coupling between matter and geometry degrees of freedom.

IV. ONE-BODY RELATIONAL COSMOLOGY

We now apply the effective approach reviewed in Section II to the one-body group field theory model introduced in Section III¹⁴. The quantum constraint is given in (3.8),

$$\hat{C} = m^2 \hat{N} - \hat{\Lambda} - \lambda \hat{\Pi}_2, \quad (4.1)$$

and the quantum algebra is generated by $\mathcal{A} = \{\hat{a}\}_{a \in \mathcal{V}}$, where \mathcal{V} is the set of labels introduced in (3.27). In the following it sometimes proves useful to make an identification $\Pi_0 = N$ allowing for a very compact notation. This further necessitates the introduction of the symbol Π_{-1} which however will always be multiplied by zero.

Following the arguments from classical cosmology, we now assume that the non-trivial commutation relations of the algebra are given by

$$[\hat{X}, \hat{\Pi}_1] = i\hbar \hat{N}, \quad (4.2a)$$

$$[\hat{X}, \hat{\Pi}_2] = i\hbar 2\hat{\Pi}_1, \quad (4.2b)$$

$$[\hat{\Lambda}, \hat{K}] = i\hbar \alpha \hat{K}. \quad (4.2c)$$

We will assume that an expansion in orders of moments corresponds to an expansion in orders of \hbar , as anticipated, and truncate the quantum phase space at second order moments. The resulting quantum phase space is therefore 27-dimensional, consisting of 6 expectation values and 21 moments. The Poisson structure on the quantum phase space is given by the Poisson bracket induced by the commutator as in (2.3). We collect the explicit form of the resulting Poisson structure in Appendix A. The 7 effective constraints, as defined in (2.4), are given by

$$C = m^2 \langle \hat{N} \rangle - \langle \hat{\Lambda} \rangle - \lambda \langle \hat{\Pi}_2 \rangle, \quad (4.3a)$$

$$C_N = m^2 \Delta(N^2) - \Delta(N\Lambda) - \lambda \Delta(N\Pi_2), \quad (4.3b)$$

$$C_{\Pi_1} = m^2 \Delta(N\Pi_1) - \Delta(\Pi_1\Lambda) - \lambda \Delta(\Pi_1\Pi_2), \quad (4.3c)$$

$$C_{\Pi_2} = m^2 \Delta(N\Pi_2) - \Delta(\Pi_2\Lambda) - \lambda \Delta(\Pi_2^2), \quad (4.3d)$$

$$C_X = m^2 \Delta(NX) - \Delta(X\Lambda) - i\hbar \lambda \langle \hat{\Pi}_1 \rangle - \lambda \Delta(\Pi_2 X), \quad (4.3e)$$

$$C_\Lambda = m^2 \Delta(N\Lambda) - \Delta(\Lambda^2) - \lambda \Delta(\Pi_2 \Lambda), \quad (4.3f)$$

$$C_K = m^2 \Delta(NK) + \frac{i\hbar}{2} \alpha \langle \hat{K} \rangle - \Delta(\Lambda K) - \lambda \Delta(\Pi_2 K). \quad (4.3g)$$

¹⁴ We performed the calculation using Wolfram Mathematica. A Mathematica package and a program containing the calculations of this section are available at <https://gitlab.com/qggftc/effective-constraints>.

Although the algebra considered here does not have any canonically conjugate pair, we will nevertheless closely mimic the treatment of a system with a canonical pair as discussed in Section II. As a “clock” we choose the operator \hat{K} and as the “conjugate” we choose $\hat{\Lambda}$. From the effective constraints C and C_a , $a \in \mathcal{V}$, it is then possible to eliminate the expectation value $\langle \hat{\Lambda} \rangle$ and the 6 moments $\Delta(\Lambda a)$, $a \in \mathcal{V}$. Likewise, we choose a gauge in which all but one of the moments of the operator \hat{K} vanish. This can be achieved by the gauge fixing conditions

$$G_a = \Delta(Ka) = 0, \quad a \in \mathcal{V} \setminus \{\Lambda\}. \quad (4.4)$$

To see that this does fix the gauge sufficiently, we compute the following Poisson brackets for any $a \in \mathcal{V} \setminus \{\Lambda\}$ (with the identification $\Pi_0 = N$),

$$\{G_a, C\} \approx 0, \quad (4.5a)$$

$$\{G_a, C_{\Pi_n}\} \approx (1 - \delta_{aK})\alpha\langle \hat{K} \rangle \left(\Delta(\Pi_n a) - \delta_{aX} \frac{i\hbar}{2} n \langle \hat{\Pi}_{n-1} \rangle \right) \quad (4.5b)$$

$$\{G_a, C_X\} \approx (1 - \delta_{aK})\alpha\langle \hat{K} \rangle \left(\Delta(Xa) + \delta_{a\Pi_n} \frac{i\hbar}{2} n \langle \hat{\Pi}_{n-1} \rangle \right), \quad (4.5c)$$

$$\{G_{\Pi_n}, C_\Lambda\} \approx \alpha\langle \hat{K} \rangle (m^2 \Delta(N\Pi_n) - \lambda \Delta(\Pi_2 \Pi_n)), \quad (4.5d)$$

$$\{G_X, C_\Lambda\} \approx \alpha\langle \hat{K} \rangle (m^2 \Delta(NX) - \lambda \Delta(\Pi_2 X) - 3i\hbar\lambda \langle \hat{\Pi}_1 \rangle), \quad (4.5e)$$

$$\{G_K, C_\Lambda\} \approx 2i\hbar\alpha^2 \langle \hat{K} \rangle^2, \quad (4.5f)$$

$$\{G_a, C_K\} \approx 0. \quad (4.5g)$$

From this we see that both C and C_K remain unfixed by our choice of gauge. However, it turns out that only C has a non-trivial flow on the gauge-fixed constraint hypersurface. Therefore, we may study the flow generated by C on the gauge-fixed constraint hypersurface and interpret this as the dynamics of the system. After imposing the 7 effective constraints (cf. (4.3)) and the 5 gauge fixing conditions (cf. (4.4)), the reduced quantum phase space has coordinates given by the $27 - 7 - 5 = 15$ variables $\tilde{\mathcal{V}} = \{\langle \hat{N} \rangle, \langle \hat{\Pi}_1 \rangle, \langle \hat{\Pi}_2 \rangle, \langle \hat{X} \rangle, \langle \hat{K} \rangle, \Delta(N^2), \Delta(N\Pi_1), \Delta(\Pi_1^2), \Delta(N\Pi_2), \Delta(\Pi_1\Pi_2), \Delta(\Pi_2^2), \Delta(NX), \Delta(\Pi_1 X), \Delta(\Pi_2 X), \Delta(X^2)\}$.

We now study the flow generated by the constraint which has not been fixed by our choice of gauge. That is, we are interested in solving the set of differential equations

$$a'(t) = \{a, C\}(t), \quad a \in \tilde{\mathcal{V}}. \quad (4.6)$$

The equations with non-vanishing right-hand side are given by

$$\langle \hat{X} \rangle'(t) = -2\lambda \langle \hat{\Pi}_1 \rangle(t), \quad (4.7a)$$

$$\langle \hat{K} \rangle'(t) = \alpha \langle \hat{K} \rangle(t), \quad (4.7b)$$

$$\Delta(NX)'(t) = -2\lambda \Delta(N\Pi_1)(t), \quad (4.7c)$$

$$\Delta(\Pi_1 X)'(t) = -2\lambda \Delta(\Pi_1^2)(t), \quad (4.7d)$$

$$\Delta(\Pi_2 X)'(t) = -2\lambda \Delta(\Pi_1 \Pi_2)(t), \quad (4.7e)$$

$$\Delta(X^2)'(t) = -4\lambda \Delta(\Pi_1 X)(t). \quad (4.7f)$$

The variables which are non-constant under the gauge flow have the following dependence on the gauge flow parameter:

$$\langle \hat{X} \rangle(t) = \langle \hat{X} \rangle(0) - 2\lambda \langle \hat{\Pi}_1 \rangle(0)t, \quad (4.8a)$$

$$\langle \hat{K} \rangle(t) = \langle \hat{K} \rangle(0)e^{\alpha t}, \quad (4.8b)$$

$$\Delta(NX)(t) = \Delta(NX)(0) - 2\lambda \Delta(N\Pi_1)(0)t, \quad (4.8c)$$

$$\Delta(\Pi_1 X)(t) = \Delta(\Pi_1 X)(0) - 2\lambda \Delta(\Pi_1^2)(0)t, \quad (4.8d)$$

$$\Delta(\Pi_2 X)(t) = \Delta(\Pi_2 X)(0) - 2\lambda \Delta(\Pi_1 \Pi_2)(0)t, \quad (4.8e)$$

$$\Delta(X^2)(t) = \Delta(X^2)(0) - 4\lambda \Delta(\Pi_1 X)(0)t + 4\lambda^2 \Delta(\Pi_1^2)(0)t^2. \quad (4.8f)$$

All the remaining variables remain constant along the flow. It is interesting to note that both (4.8a) and (4.8b) can be inverted. Hence both $\langle \hat{X} \rangle$ and $\langle \hat{K} \rangle$ could be viewed as a ‘‘clock’’ from a relational point of view, in the sense that their values can be used to parametrise all other dynamical variables. Of course, there is still a difference between the two since we used a gauge fixing of moments following the perspective presented in Section II in which $\langle \hat{K} \rangle$ is seen as the clock, so that in our gauge the variable X has non-trivial quantum fluctuations. If we want to interpret (4.8a) and (4.8b) as representing effective cosmological dynamics constructed from our set of one-body operators, which would be analogous to the classical Friedmann cosmology discussed in Section III C, we could view either $\langle \hat{X} \rangle$ as a function of $\langle \hat{K} \rangle$ or vice versa; since the relation between the two is a globally invertible function both functional relations are equivalent.

Concretely, we can express $\langle \hat{X} \rangle$ as a function of $\langle \hat{K} \rangle$,

$$(\langle \hat{X} \rangle \circ \langle \hat{K} \rangle^{-1})(\langle \hat{K} \rangle) = \langle \hat{X} \rangle(0) - \frac{2\lambda \langle \hat{\Pi}_1 \rangle}{\alpha} \log \left(\frac{\langle \hat{K} \rangle}{\langle \hat{K} \rangle(0)} \right) \quad (4.9)$$

which can be compared with the classical solution we found in Section III C,

$$(\varphi \circ p_V^{-1})(p_V) = \varphi(0) \pm \frac{1}{\sqrt{12\pi G}} \log \left(\frac{p_V}{p_V(0)} \right). \quad (4.10)$$

1
2
3 The classical solution for a flat FLRW cosmology is hence reproduced by the solutions (4.8a)
4 and (4.8b) of our effective one-body dynamics if we identify $\langle \hat{X} \rangle$ with a constant multiple of φ and
5 $\langle \hat{K} \rangle$ with a constant multiple of p_V . The ratio λ/α , which is determined by the GFT dynamics
6 and by the operators appearing in the algebra (4.2), corresponds to Newton's constant in classical
7 cosmology, with the relation between the two determined by the conserved quantity $\langle \hat{\Pi}_1 \rangle$ and the
8 identification between $\langle \hat{X} \rangle$ and the classical φ . (Recall from the discussion below (3.17) that we
9 need to define an intensive quantity from the extensive $\langle \hat{X} \rangle$ in order to justify the interpretation as
10 a physical scalar field.) If one identifies the quotient $\langle \hat{X} \rangle / \langle \hat{\Pi}_1 \rangle$ of two extensive quantities with the
11 classical scalar field φ and the parameters λ and/or α are chosen appropriately, arbitrary initial
12 conditions of our effective equations lead to a solution of classical FLRW cosmology. This is the
13 analogue of the recovery of (modified) Friedmann cosmologies in previous work on GFT cosmology
14 such as [24, 29, 48], where a similar identification of parameters of the fundamental theory with an
15 emergent Newton's constant is made¹⁵.

16
17 An interesting property of the system of equations (4.8a)–(4.8e) is that expectation values and
18 moments are entirely decoupled: the inclusion of $O(\hbar)$ moments does not alter at all the dynamics of
19 the original variables in \mathcal{V} which correspond to $O(\hbar^0)$. Indeed, the solutions (4.8a)–(4.8b) are exactly
20 the ones already discussed in Section III C where we fixed the form of the commutator involving
21 \hat{K} . This behaviour is due to the linearity of the initial constraint in all variables, and one might
22 see it as representing an unphysical aspect of the truncation we have been using. Indeed, it would
23 seem to imply that semiclassicality properties of the fundamental state are mostly irrelevant to the
24 extraction of interesting cosmological dynamics. We should point out, however, that our results here
25 are fully compatible with previous derivations of effective cosmological Friedmann equations from
26 GFT dynamics as, e.g., given in [29]: also in this previous work, effective Friedmann equations can
27 be derived for general states and only depend on expectation values of a few elementary operators,
28 without any dependence on quantum fluctuations or higher moments in general.

29
30 The criterion of semiclassicality, as studied in terms of relative fluctuations in [28, 29], should then
31 be seen as an additional requirement for a meaningful physical interpretation of the resulting effective
32 Friedmann equations (derived from the dynamics of expectation values of relevant operators), even
33 if these equations themselves are unaffected by the magnitude of quantum fluctuations. A quantum
34

35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55 ¹⁵ That gravitational and, more generally, effective field theory couplings like Newton's constant or the cosmological
56 constant are in fact a function of microscopic parameters of the underlying non-spatiotemporal quantum gravity
57 dynamics is something to be expected in any formalism in which spacetime itself is emergent.
58
59
60

cosmology with large quantum fluctuations in macroscopic cosmological observables would not describe the universe we observe, even if such observables happen to satisfy the correct (Friedmann-like) dynamics in mean value. In this respect, it is important to emphasise that if relative quantum fluctuations are required to be small at small curvature ($|\langle \hat{K} \rangle(t) / \langle \hat{K} \rangle(0)| \ll 1$), in our context they turn out to be also small when the curvature is large ($|\langle \hat{K} \rangle(t) / \langle \hat{K} \rangle(0)| \gg 1$). Both these limits are characterised by $|t| \rightarrow \infty$, so that the asymptotic behaviour of the relative fluctuations

$$\sigma_{aa'}^2 = \frac{\Delta(aa')}{\langle \hat{a} \rangle \langle \hat{a}' \rangle} \quad (4.11)$$

is identical¹⁶, as we can see explicitly from (4.8a)–(4.8e). The fact that quantum fluctuations can be relatively small even when curvature is large is a feature shared with LQC models [32, 33]. A crucial property of our approach is that the (averaged) particle number remains constant, as we have emphasised several times above, in sharp contrast to previous results on quantum fluctuations in GFT cosmology [28, 29] where the (average) particle number had a non-trivial relational evolution.¹⁷ In these works, it was shown that the evolution of the particle number crucially affects the evolution of relative quantum fluctuations, suppressing them at large volume (when $N \gg 1$) and possibly enhancing them at small volume if $N \lesssim 1$. This mechanism of suppression and enhancement of fluctuations is of course absent here, which may be one of the reasons why quantum fluctuations can remain small even in a high curvature regime, provided they are small at low curvature.

V. CONCLUSIONS AND DISCUSSION

In this paper we presented a novel way to obtain an effective cosmological dynamics from GFT, motivated by the interpretation of cosmology as quantum gravity hydrodynamics [19, 22]. Contrary to what was done in previous works, this approach does not depend explicitly on a specific choice of states. In order to achieve this, we extended the effective approach developed in [15–17] to

¹⁶ Notice that relative quantum fluctuations are not necessarily small at all times, since they may diverge when the denominator of (4.11) goes to zero. This does not signal that the system is affected by strong quantum fluctuations, but rather that they should not be measured by relative quantities. A more sensible option in this case would be to define a threshold below which fluctuations are considered small, see [53] for a more detailed discussion.

¹⁷ Notice again that these previous results refer to different notions of relational dynamics in GFT. The number operator in [29] is the “time-dependent version” of (3.7a). On the other hand, the (relational) number operator in [25] is the same as in (3.7a), although relational evolution is defined only for a specific choice of states (and with respect to a different relational clock).

1
2
3 describe the behaviour of a finite set of collective GFT observables, assumed to capture macroscopic
4 cosmological physics. That is, we extended this approach to a field-theoretic setting. The dynamics
5 of these observables are governed by a one-body constraint, obtained by integrating the microscopic
6 equation of motion of the model. Similarly to [24] and many subsequent works, this model includes
7 matter in the form of a minimally coupled massless scalar field, and is characterised by negligible
8 interactions. This is reflected also at the level of our algebra of one-body operators of interest, which
9 splits into a geometric two-dimensional subalgebra and a four-dimensional one generated by “matter
10 observables” and the number operator.
11
12

13
14
15
16
17 From the algebra of operators and the one-body constraint we obtained a quantum phase space
18 which can be coordinatised by a finite set of expectation values and moments. We considered
19 only moments up to second order in powers of \hbar , obtaining an effectively 27-dimensional space,
20 described by 6 expectation values and 21 moments. At this order, the quantum constraint produces
21 7 effective constraints, which in turn generate 6 independent gauge flows in the truncated quantum
22 phase space. We fixed 5 of these 6 gauge flows by requiring that the operator \hat{K} that we identify
23 classically with the extrinsic curvature has vanishing moments. The quantum phase space variables
24 related to the other operator generating the geometric subalgebra were then expressed in terms of
25 the remaining variables through the effective constraints. In this way we obtained an effectively
26 reduced phase space parametrised by 15 variables: 14 expectation values and moments related to the
27 four-dimensional “matter” subalgebra and one expectation value $\langle \hat{K} \rangle$. We could use the remaining
28 gauge flow to express the evolution of the 14 non-geometric variables in terms of $\langle \hat{K} \rangle$. Since \hat{K} by
29 construction has vanishing quantum fluctuations (they are gauge-fixed to zero), this amounts to
30 an effective relational description of the evolution of matter degrees of freedom with respect to the
31 extrinsic curvature clock. This is a refreshing change of perspective for cosmological applications of
32 relational dynamics in quantum gravity, where it is usually matter degrees of freedom that are used
33 as physical clocks (and rods) [54]. In the framework we developed, choosing a geometric clock was
34 the most natural choice, since an analogous elimination of matter variables resulting in a reduced
35 quantum phase space of only geometric quantities (and possibly the expectation value of a matter
36 clock) could not be achieved by the methods explained above.
37
38
39
40
41
42
43
44
45
46
47
48
49

50
51 The way relational evolution is obtained in this framework emphasises the idea that in order to
52 define a notion of relational dynamics in an emergent spacetime context as the one we considered
53 here, some prior coarse-graining process is necessary [25]. Here, this coarse-graining is represented by
54 the assumption that the physics of the system is adequately captured by a finite set of macroscopic
55 observables and one collective quantum constraint. This is however still not enough to define a
56
57
58
59
60

1
2
3 “good clock” and consequently a notion of relational dynamics analogous to the classical theory: it is
4 also necessary to be able to find a gauge (the “Zeitgeist” of [16, 17]) in which quantum fluctuations
5 of the clock can be set to zero.
6
7

8 At the level of expectation values, the resulting relational evolution is in agreement with a
9 classical flat FLRW universe. Interestingly, this result does not depend on the magnitude of quantum
10 fluctuations who evolve independently and, if they are required to be relatively small at low curvature,
11 remain small at high curvature. We argue that this feature, shared also with LQC models [32, 33],
12 may be due to the constancy of the number operator. In the framework presented in this paper, the
13 number operator is constant because the quantum constraint is a one-body operator and, as such,
14 trivially commutes with the number operator. In most previous attempts to extract cosmological
15 physics from GFT the particle number is time-dependent, and this time dependence can crucially
16 enhance or suppress quantum fluctuations [28, 29].
17
18
19
20
21
22

23 The methods employed in this work, and thus all the aforementioned results, rely on two important
24 assumptions.
25
26

27 The first one is related to the choice of states. While we remark again that no specific state choice
28 is necessary within this framework, the states we considered are required to satisfy two important
29 conditions: (i) they should be such that the system is characterised by a few macroscopic observables
30 (hydrodynamic condition); and (ii) moments of operators computed in these states should naturally
31 produce a hierarchy characterised by different powers of \hbar (semiclassicality condition). Without the
32 first requirement one would have to take into account the fact that a field theory is characterised by
33 infinitely many operators and “constraints” (the quantum equations of motion), while without the
34 second one one would have to consider the infinitely many moments coordinatising the quantum
35 phase space, even for finite-dimensional systems.
36
37
38
39
40
41

42 The second assumption is related to the choice of the fundamental observables, which depends
43 on the features of the physical system being considered. In our case, we introduced the “curvature
44 operator” \hat{K} by requiring it to satisfy certain commutation relations which classically allow for a
45 cosmological interpretation of the system. However, a microscopic description of such an operator is
46 missing and requires further studies. In fact, we showed that such an operator cannot be represented
47 on a compact group, suggesting that, at least for the type of dynamics we defined here (which is
48 very much simplified compared to the one corresponding to more realistic quantum gravity models),
49 GFT models with non-compact “gravitational gauge group” would be preferred.
50
51
52
53
54

55 Even when these assumptions are made, there are some general limitations to the framework used
56 here. Computations become more and more involved as either the number of fundamental elements
57
58
59
60

of the quantum algebra increase or the commutation relations between the operators become more complicated (e.g., non-linear). These computational problems are the main reason why considering a system of operators and dynamics for which the particle number can vary is complicated. Indeed, in order to do so, one would have to consider an additional n -body constraint ($n > 1$) with non-trivial commutation relations with the number operator. An example would be, of course, a dynamical constraint incorporating the contribution from GFT interactions. This would however force us to include many other operators (with possibly more complicated commutation relations) in order to obtain a closed algebra, which would significantly enlarge the quantum phase space.

Despite these limitations, the method we employed allowed us to (i) obtain effective cosmological dynamics from GFT, (ii) gain insights about the notion of relational dynamics in an emergent quantum gravity scenario, and (iii) study the impact of quantum fluctuations, by making only broad assumptions on the states of interest. The relative simplicity in which the above results were obtained within a well-defined framework illustrates the importance of effective methods for extracting continuum physics from quantum gravity, and in particular suggests that the connection between GFTs (in their hydrodynamic approximation) and cosmological physics may go beyond the choice of specific (condensate) states.

ACKNOWLEDGMENTS

The work of SG was funded by the Royal Society through a University Research Fellowship (UF160622) and a Research Grant for Research Fellows (RGF\R1\180030). AP was supported by the same Research Grant for Research Fellows awarded to SG. LM thanks the University of Pisa and the INFN (section of Pisa) for financial support, and the Ludwig Maximilians-Universität (LMU) Munich for the hospitality.

Appendix A: Quantum Poisson structure

In this section we collect the non-vanishing Poisson brackets of the quantum phase space considered in Section IV. In the formulas below we identify $N = \Pi_0$ and introduce the “fake” variable $\Pi_{-1} = 0$. Unless specified otherwise, $a \in \{N, \Pi_1, \Pi_2, X, \Lambda, K\}$.

Using (2.6) one can verify the following commutation relations which we computed using Wolfram Mathematica,

$$\{\langle \hat{\Lambda} \rangle, \Delta(Ka)\} = (1 + \delta_{aK})\alpha\Delta(Ka), \quad (\text{A1})$$

$$\{\langle \hat{K} \rangle, \Delta(\Lambda a)\} = -(1 + \delta_{a\Lambda})\alpha\Delta(Ka), \quad (A2)$$

$$\{\langle \hat{X} \rangle, \Delta(\Pi_n a)\} = (1 + \delta_{a\Pi_n})n\Delta(\Pi_{n-1}a), \quad a \neq \Pi_{m \neq n}, \quad (A3)$$

$$\{\langle \hat{X} \rangle, \Delta(\Pi_n \Pi_m)\} = n\Delta(\Pi_{n-1}\Pi_m) + m\Delta(\Pi_n \Pi_{m-1}), \quad (A4)$$

$$\{\langle \hat{\Pi}_n \rangle, \Delta(Xa)\} = -(1 + \delta_{aX})n\Delta(\Pi_{n-1}a), \quad (A5)$$

$$\{\Delta(\Lambda X), \Delta(Ka)\} = (1 + \delta_{aK})\alpha\langle \hat{K} \rangle\Delta(Xa), \quad a \neq \Pi_n, \quad (A6)$$

$$\{\Delta(\Lambda X), \Delta(\Pi_n a)\} = (1 + \delta_{a\Pi_n})n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Lambda a), \quad a \neq K, \Pi_{m \neq n}, \quad (A7)$$

$$\{\Delta(\Lambda X), \Delta(K\Pi_n)\} = n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Lambda K) + \alpha\langle \hat{K} \rangle\Delta(X\Pi_n), \quad (A8)$$

$$\{\Delta(\Lambda X), \Delta(\Pi_n \Pi_m)\} = n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Lambda \Pi_m) + m\langle \hat{\Pi}_{m-1} \rangle\Delta(\Lambda \Pi_n), \quad (A9)$$

$$\{\Delta(KX), \Delta(\Lambda a)\} = -(1 + \delta_{a\Lambda})\alpha\langle \hat{K} \rangle\Delta(Xa), \quad a \neq \Pi_n, \quad (A10)$$

$$\{\Delta(KX), \Delta(\Pi_n a)\} = (1 + \delta_{a\Pi_n})n\langle \hat{\Pi}_{n-1} \rangle\Delta(Ka), \quad a \neq \Lambda, \Pi_{m \neq n}, \quad (A11)$$

$$\{\Delta(KX), \Delta(\Lambda \Pi_n)\} = n\langle \hat{\Pi}_{n-1} \rangle\Delta(K\Lambda) - \alpha\langle \hat{K} \rangle\Delta(X\Pi_n), \quad (A12)$$

$$\{\Delta(KX), \Delta(\Pi_n \Pi_m)\} = n\langle \hat{\Pi}_{n-1} \rangle\Delta(K\Pi_m) + m\langle \hat{\Pi}_{m-1} \rangle\Delta(K\Pi_n), \quad (A13)$$

$$\{\Delta(\Lambda \Pi_n), \Delta(Ka)\} = (1 + \delta_{aK})\alpha\langle \hat{K} \rangle\Delta(\Pi_n a), \quad a \neq X, \quad (A14)$$

$$\{\Delta(\Lambda \Pi_n), \Delta(Xa)\} = -(1 + \delta_{aX})n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Lambda a), \quad a \neq K, \quad (A15)$$

$$\{\Delta(\Lambda \Pi_n), \Delta(KX)\} = -n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Lambda a) + \alpha\langle \hat{K} \rangle\Delta(\Pi_n a), \quad (A16)$$

$$\{\Delta(K\Pi_n), \Delta(\Lambda a)\} = -(1 + \delta_{a\Lambda})\alpha\langle \hat{K} \rangle\Delta(\Pi_n a), \quad a \neq X, \quad (A17)$$

$$\{\Delta(K\Pi_n), \Delta(Xa)\} = -(1 + \delta_{aX})n\langle \hat{\Pi}_{n-1} \rangle\Delta(Ka), \quad a \neq \Lambda, \quad (A18)$$

$$\{\Delta(K\Pi_n), \Delta(\Lambda X)\} = -n\langle \hat{\Pi}_{n-1} \rangle\Delta(Ka) - \alpha\langle \hat{K} \rangle\Delta(\Pi_n a), \quad (A19)$$

$$\{\Delta(K^2), \Delta(\Lambda a)\} = -(1 + \delta_{a\Lambda})2\alpha\langle \hat{K} \rangle\Delta(Ka), \quad (A20)$$

$$\{\Delta(\Lambda^2), \Delta(Ka)\} = (1 + \delta_{aK})2\alpha\langle \hat{K} \rangle\Delta(\Lambda a), \quad (A21)$$

$$\{\Delta(\Lambda K), \Delta(Ka)\} = \alpha\langle \hat{K} \rangle\Delta(Ka), \quad a \neq \Lambda, \quad (A22)$$

$$\{\Delta(\Lambda K), \Delta(\Lambda a)\} = -\alpha\langle \hat{K} \rangle\Delta(\Lambda a), \quad a \neq K, \quad (A23)$$

$$\{\Delta(X^2), \Delta(\Pi_n a)\} = (1 + \delta_{a\Pi_n})2n\langle \hat{\Pi}_{n-1} \rangle\Delta(Xa), \quad a \neq \Pi_{m \neq n}, \quad (A24)$$

$$\{\Delta(X^2), \Delta(\Pi_n \Pi_m)\} = 2(n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Pi_m X) + m\langle \hat{\Pi}_{m-1} \rangle\Delta(\Pi_n X)), \quad (A25)$$

$$\{\Delta(X\Pi_n), \Delta(\Pi_m a)\} = (1 + \delta_{a\Pi_m})m\langle \hat{\Pi}_{m-1} \rangle\Delta(\Pi_n a), \quad a \neq X, \Pi_{l \neq m}, \quad (A26)$$

$$\{\Delta(X\Pi_n), \Delta(\Pi_m \Pi_l)\} = m\langle \hat{\Pi}_{m-1} \rangle\Delta(\Pi_n \Pi_l) + l\langle \hat{\Pi}_{l-1} \rangle\Delta(\Pi_n \Pi_m), \quad (A27)$$

$$\{\Delta(X\Pi_n), \Delta(Xa)\} = -(1 + \delta_{aX})n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Pi_n \Pi_l), \quad a \neq \Pi_m, \quad (A28)$$

$$\{\Delta(\Pi_n \Pi_m), \Delta(Xa)\} = -(1 + \delta_{aX})(n\langle \hat{\Pi}_{n-1} \rangle\Delta(\Pi_m a) + m\langle \hat{\Pi}_{m-1} \rangle\Delta(\Pi_n a)). \quad (A29)$$

-
- [1] K.V. Kuchař, *Time and interpretations of quantum gravity*, *International Journal of Modern Physics D* **20** (2011) 3.
- [2] C.J. Isham, *Canonical quantum gravity and the problem of time*, *NATO Sci. Ser. C* **409** (1992) 157 [gr-qc/9210011].
- [3] B. Dittrich, *The Continuum Limit of Loop Quantum Gravity: A Framework for Solving the Theory*, in *Loop Quantum Gravity: The First 30 Years*, A. Ashtekar and J. Pullin, eds., pp. 153–179 (2017) [1409.1450].
- [4] C. Rovelli, *Quantum Gravity*, Cambridge Monographs on Mathematical Physics, Cambridge University Press (2004).
- [5] T. Thiemann, *Modern Canonical Quantum General Relativity*, Cambridge Monographs on Mathematical Physics, Cambridge University Press (2007).
- [6] R.L. Arnowitt, S. Deser and C.W. Misner, *The dynamics of general relativity*, in *Gravitation: an introduction to current research*, L. Witten, ed., Wiley, New York (1962) [gr-qc/0405109].
- [7] C. Rovelli, *What is observable in classical and quantum gravity?*, *Class. Quant. Grav.* **8** (1991) 297.
- [8] C. Rovelli, *Partial observables*, *Phys. Rev. D* **65** (2002) 124013 [gr-qc/0110035].
- [9] J. Tambornino, *Relational Observables in Gravity: a Review*, *SIGMA* **8** (2012) 017 [1109.0740].
- [10] B. Dittrich, *Partial and complete observables for Hamiltonian constrained systems*, *Gen. Rel. Grav.* **39** (2007) 1891 [gr-qc/0411013].
- [11] B. Dittrich, *Partial and complete observables for canonical general relativity*, *Class. Quant. Grav.* **23** (2006) 6155 [gr-qc/0507106].
- [12] D. Oriti, *The Bronstein hypercube of quantum gravity*, in *Beyond Spacetime*, N. Huggett, K. Matsubara and C. Wüthrich, eds., pp. 25–52 (2020) [1803.02577].
- [13] M. Bojowald, B. Sandlhofer, A. Skirzewski and A. Tsobanjan, *Effective Constraints for Quantum Systems*, *Rev. Math. Phys.* **21** (2009) 111 [0804.3365].
- [14] M. Bojowald and A. Skirzewski, *Effective equations of motion for quantum systems*, *Rev. Math. Phys.* **18** (2006) 713 [math-ph/0511043].
- [15] M. Bojowald and A. Tsobanjan, *Effective constraints for relativistic quantum systems*, *Phys. Rev. D* **80** (2009) 125008 [0906.1772].
- [16] M. Bojowald, P.A. Höhn and A. Tsobanjan, *An effective approach to the problem of time*, *Class. Quant. Grav.* **28** (2011) 035006 [1009.5953].
- [17] M. Bojowald, P.A. Höhn and A. Tsobanjan, *Effective approach to the problem of time: General features and examples*, *Phys. Rev. D* **83** (2011) 125023 [1011.3040].
- [18] P.A. Höhn, E. Kubalová and A. Tsobanjan, *Effective relational dynamics of a nonintegrable cosmological model*, *Phys. Rev. D* **86** (2012) 065014 [1111.5193].
- [19] D. Oriti, *The universe as a quantum gravity condensate*, *Comptes Rendus Physique* **18** (2017) 235

- [1612.09521].
- [20] S. Gielen and L. Sindoni, *Quantum Cosmology from Group Field Theory Condensates: a Review*, *SIGMA* **12** (2016) 082 [1602.08104].
- [21] D. Oriti, *Group field theory as the second quantization of loop quantum gravity*, *Class. Quant. Grav.* **33** (2016) 085005 [1310.7786].
- [22] S. Gielen, D. Oriti and L. Sindoni, *Homogeneous cosmologies as group field theory condensates*, *JHEP* **06** (2014) 013 [1311.1238].
- [23] S. Gielen, *Quantum cosmology of (loop) quantum gravity condensates: an example*, *Class. Quant. Grav.* **31** (2014) 155009 [1404.2944].
- [24] D. Oriti, L. Sindoni and E. Wilson-Ewing, *Emergent Friedmann dynamics with a quantum bounce from quantum gravity condensates*, *Class. Quant. Grav.* **33** (2016) 224001 [1602.05881].
- [25] L. Marchetti and D. Oriti, *Effective relational cosmological dynamics from quantum gravity*, *JHEP* **05** (2021) 025 [2008.02774].
- [26] S. Gielen and A. Polaczek, *Hamiltonian group field theory with multiple scalar matter fields*, *Phys. Rev. D* **103** (2021) 086011 [2009.00615].
- [27] E. Wilson-Ewing, *Relational Hamiltonian for group field theory*, *Phys. Rev. D* **99** (2019) 086017 [1810.01259].
- [28] L. Marchetti and D. Oriti, *Quantum Fluctuations in the Effective Relational GFT Cosmology*, *Front. Astron. Space Sci.* **8** (2021) 683649 [2010.09700].
- [29] S. Gielen and A. Polaczek, *Generalised effective cosmology from group field theory*, *Class. Quant. Grav.* **37** (2020) 165004 [1912.06143].
- [30] S. Gielen, *Frozen formalism and canonical quantization in (group) field theory*, 2105.01100.
- [31] A. Ashtekar and T.A. Schilling, *Geometrical Formulation of Quantum Mechanics*, gr-qc/9706069.
- [32] A. Corichi and E. Montoya, *Coherent semiclassical states for loop quantum cosmology*, *Phys. Rev. D* **84** (2011) 044021 [1105.5081].
- [33] C. Rovelli and E. Wilson-Ewing, *Why are the effective equations of loop quantum cosmology so accurate?*, *Phys. Rev. D* **90** (2014) 023538 [1310.8654].
- [34] M. Assanioussi and I. Kotecha, *Thermal quantum gravity condensates in group field theory cosmology*, *Phys. Rev. D* **102** (2020) 044024 [2003.01097].
- [35] P.A. Höhn, *From Classical to Quantum: New Canonical Tools for the Dynamics of Gravity*, Ph.D. thesis, Utrecht U., 2012.
- [36] A. Tsobanjan, *Semiclassical Analysis of Constrained Quantum Systems*, *AIP Conf. Proc.* **1196** (2009) 291 [0911.0648].
- [37] I.M. Gelfand and M.A. Naimark, *On the imbedding of normed rings into the ring of operators on a Hilbert space*, *Matematicheskii Sbornik* **54** (1943) 197.
- [38] I.E. Segal, *Irreducible representations of operator algebras*, *Bulletin of the American Mathematical Society* **53** (1947) 73.

- 1
2
3 [39] M. Bojowald and T. Strobl, *Poisson geometry in constrained systems*, *Rev. Math. Phys.* **15** (2003) 663
4 [hep-th/0112074].
5
6 [40] T. Buchert, *On Average Properties of Inhomogeneous Fluids in General Relativity: Perfect Fluid*
7 *Cosmologies*, *Gen. Rel. Grav.* **33** (2001) 1381 [gr-qc/0102049].
8
9 [41] T. Buchert et al., *Is there proof that backreaction of inhomogeneities is irrelevant in cosmology?*, *Class.*
10 *Quant. Grav.* **32** (2015) 215021 [1505.07800].
11
12 [42] S. Gielen, *Emergence of a low spin phase in group field theory condensates*, *Class. Quant. Grav.* **33**
13 (2016) 224002 [1604.06023].
14
15 [43] M. de Cesare, D. Oriti, A.G.A. Pithis and M. Sakellariadou, *Dynamics of anisotropies close to a*
16 *cosmological bounce in quantum gravity*, *Class. Quant. Grav.* **35** (2018) 015014 [1709.00994].
17
18 [44] J. Ben Geloun and V. Bonzom, *Radiative Corrections in the Boulatov-Ooguri Tensor Model: The*
19 *2-Point Function*, *Int. J. Theor. Phys.* **50** (2011) 2819 [1101.4294].
20
21 [45] S. Carrozza, D. Oriti and V. Rivasseau, *Renormalization of a SU(2) Tensorial Group Field Theory in*
22 *Three Dimensions*, *Commun. Math. Phys.* **330** (2014) 581 [1303.6772].
23
24 [46] M.P. Reisenberger and C. Rovelli, *Spacetime as a Feynman diagram: the connection formulation*, *Class.*
25 *Quant. Grav.* **18** (2001) 121 [gr-qc/0002095].
26
27 [47] M. Bojowald, *Canonical Gravity and Applications*, Cambridge University Press (2010).
28
29 [48] D. Oriti, L. Sindoni and E. Wilson-Ewing, *Bouncing cosmologies from quantum gravity condensates*,
30 *Class. Quant. Grav.* **34** (2017) 04LT01 [1602.08271].
31
32 [49] A. Ashtekar and P. Singh, *Loop quantum cosmology: a status report*, *Class. Quant. Grav.* **28** (2011)
33 213001 [1108.0893].
34
35 [50] G. Calcagni, S. Gielen and D. Oriti, *Group field cosmology: a cosmological field theory of quantum*
36 *geometry*, *Class. Quant. Grav.* **29** (2012) 105005 [1201.4151].
37
38 [51] L. Freidel and S. Speziale, *Twisted geometries: A geometric parametrization of SU(2) phase space*, *Phys.*
39 *Rev. D* **82** (2010) 084040 [1001.2748].
40
41 [52] Y. Li, D. Oriti and M. Zhang, *Group field theory for quantum gravity minimally coupled to a scalar*
42 *field*, *Class. Quant. Grav.* **34** (2017) 195001 [1701.08719].
43
44 [53] A. Ashtekar, L. Bombelli and A. Corichi, *Semiclassical states for constrained systems*, *Phys. Rev. D* **72**
45 (2005) 025008 [gr-qc/0504052].
46
47 [54] K. Giesel and T. Thiemann, *Scalar material reference systems and loop quantum gravity*, *Class. Quant.*
48 *Grav.* **32** (2015) 135015 [1206.3807].
49
50
51
52
53
54
55
56
57
58
59
60