



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/184457/>

Version: Published Version

Article:

Haider, M.S., Youngkong, S., Thavorncharoensap, M. et al. (2022) Priority setting of vaccine introduction in Bangladesh: a multicriteria decision analysis study. *BMJ Open*, 12 (2). e054219. ISSN: 2044-6055

<https://doi.org/10.1136/bmjopen-2021-054219>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

BMJ Open Priority setting of vaccine introduction in Bangladesh: a multicriteria decision analysis study

Mohammad Sabbir Haider ^{1,2} Sitaporn Youngkong ^{1,3}
Montarat Thavorncharoensap ^{1,3} Praveen Thokala ⁴

To cite: Haider MS, Youngkong S, Thavorncharoensap M, *et al.* Priority setting of vaccine introduction in Bangladesh: a multicriteria decision analysis study. *BMJ Open* 2022;**12**:e054219. doi:10.1136/bmjopen-2021-054219

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-054219>).

Received 08 June 2021

Accepted 27 January 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

¹Mahidol University Health Technology Assessment (MUHTA) Graduate Program, Mahidol University, Bangkok, Thailand

²Directorate General of Health Services, Government of Bangladesh Ministry of Health and Family Welfare, Dhaka, Bangladesh

³Social and Administrative Pharmacy Division, Department of Pharmacy, Faculty of Pharmacy, Mahidol University, Bangkok, Thailand

⁴Health Economics and Decision Science, School of Health and Related Research (ScHARR), The University of Sheffield, Sheffield, UK

Correspondence to

Dr Sitaporn Youngkong;
sitaporn.you@mahidol.edu

ABSTRACT

Objective To prioritise vaccines for introduction in Bangladesh.

Methods Multicriteria decision analysis (MCDA) process was used to prioritise potential vaccines for introduction in Bangladesh. A set of criteria were identified, weighted and assigned scores by relevant stakeholders (n=14) during workshop A. The performance matrix of the data of vaccines against the criteria set was constructed and validated with the experts (n=6) in workshop B. The vaccines were ranked and appraised by another group of stakeholders (n=10) in workshop C, and the final workshop D involved the dissemination of the findings to decision-makers (n=28).

Results Five criteria including incidence rate, case fatality rate, vaccine efficacy, size of the population at risk and type of population at risk were used quantitatively to evaluate and to score the vaccines. Two other criteria, cost-effectiveness and outbreak potentiality, were considered qualitatively. On deliberation, the Japanese encephalitis (JE) vaccine was ranked top to be recommended for introduction in Bangladesh.

Conclusions Based on the MCDA results, JE vaccine is planned to be recommended to the decision-makers for introduction into the national vaccine benefit package. The policymakers support the use of systematic evidence-based decision-making processes such as MCDA for vaccine introduction in Bangladesh, and to prioritise health interventions in the country.

INTRODUCTION

Vaccination is the most effective public health measure to prevent infectious diseases.^{1 2} Governments in developing countries prefer to invest in vaccination programmes that can be financially sustainable^{3–5} and while countries often consider cost-effectiveness, this should not be the only criterion for the selection of interventions.^{6 7} Different criteria, such as disease severity, effectiveness, accessibility, quality of care and equity, should be considered during healthcare priority setting.⁸

Decision-making around the introduction of new vaccines in the healthcare benefit package is complex.⁹ There are systematic and evidence-based methods,¹⁰ using priority

Strengths and limitations of this study

- Multicriteria decision analysis (MCDA) process was used to support vaccine introduction decision-making in Bangladesh, contributing to transparency and evidence-informed priority setting.
- Participation of a wide range of stakeholders in this MCDA study ensured the transparency and accountability of decision-making, which is essential for a fair priority setting process.
- Data on the vaccines on the different criteria were gathered from systematic evidence synthesis and validated with experts, and good practice MCDA guidance was followed to elicit the preferences and rank the list of vaccines.
- Different sets of stakeholders took part in the workshops, resulting in a lack of a consistent group of stakeholders (and hence values or preferences) throughout the MCDA process.
- Stakeholders from private sectors and representatives of patient groups were not involved in the process, leading to uncertainty in accountability of the results to those stakeholders.

setting to allocate the scarce resources to meet increasing demand.¹¹ Multicriteria decision analysis (MCDA) is one such approach which evaluates different options considering multiple criteria in an explicit manner,¹² to aid decision-makers to make rational decisions.¹³ MCDA can be a useful approach to support inclusion of health interventions in the benefit package.⁷

Vaccine preventable diseases such as dengue, human papillomavirus (HPV), influenza, Japanese encephalitis (JE) and typhoid are prevalent in Bangladesh.^{14–18} These diseases can be prevented by the introduction of new or underused vaccines by the government of Bangladesh. However, new vaccines have considerable budget impact and need to be prioritised for introduction into the benefit package.¹⁹ In the past, decision-making for vaccine introduction was ad-hoc

but there is increasing interest in prioritisation using systematic evaluation of multiple criteria.¹⁹

As such, we conducted an MCDA study to support prioritisation of vaccines for introduction in the benefit package in Bangladesh. The aims of the study are to support prioritisation of health interventions using an evidence-based systematic process incorporating multiple criteria and involving key relevant stakeholders, and to provide national decision-makers with scientific recommendations on vaccine introduction to better use the limited resources in Bangladesh.

METHODS

We followed the steps outlined in good practice guidelines for the use of MCDA in healthcare.^{20 21} As stakeholder involvement is key, we conducted four workshops (between October 2019 and January 2020) with the relevant stakeholders during the MCDA process. The steps and the workshops are described in further detail below.

Identifying the list of potential vaccines for introduction

The potential vaccines for prioritisation were identified from the recommendations of the WHO, Gavi the vaccine alliance, and Centers for Disease Control and Prevention in the USA (CDC-US). Vaccines which were currently in the expanded programme on immunisation (EPI) programme of the neighbouring countries were also considered. From these sources, vaccines that were not yet introduced in Bangladesh were identified as potential vaccines to be evaluated.

Selecting criteria for vaccine introduction in Bangladesh

A three-step process was used to select criteria for vaccine introduction in Bangladesh. First, a systematic review was conducted to identify all potential criteria for vaccine introduction in Bangladesh, which is described elsewhere in detail.²² Second, from this long list of criteria, a core team of three public health experts of Bangladesh (including the lead author, MSH) excluded criteria that cannot be quantified (eg, political will) and those that were mentioned less frequently.

Finally, the potential criteria list was ranked in workshop A (WS-A) in October 2019, to identify the key criteria to be used for vaccine prioritisation. Stakeholders (n=14) included paediatricians (n=1), public health experts (n=6), virologists (n=2), epidemiologists (n=4) and health economists (n=1). In terms of affiliation, these stakeholders (n=14) were from directorate office (n=4), technical institutes (n=4), non-government organisations (NGOs) (n=3), National Immunization Technical Advisory Group (NITAG) (n=2) and health professional associations (n=1). The criteria, along with their definitions, were presented to the stakeholders (online supplementary A) who were then asked to rank each criterion from '1 to 10', where '1' was the most preferable and '10' was the least preferable criterion. The ranked order of criteria was transformed into ranking weight using the rank order

centroid (ROC) method.²³ Criteria were ranked based on the mean ROC weight, and the stakeholders selected a set of criteria by consensus to be used in the prioritisation of vaccines.

Weighting and scoring

In the same workshop (WS-A), the stakeholders weighted the criteria using direct rating methods. Stakeholders discussed and then agreed by consensus to assign points to each criterion on a scale of 0–100, where '0' depicted the least important, and '100' represented the most important. To calculate the weights, the points assigned for each criterion was normalised (ie, by dividing the points allocated to each criterion by the sum of points of all criteria) using Equation 1.^{24 25}

$$w_i = p_i / \sum p_i \quad (1)$$

where, w_i is the normalised weight of criterion i ; i is the index of criterion; p_i is the points allocated to each criterion.

For scoring, the levels of criteria were identified by the core team from literature review and expert opinion. These were presented to the stakeholders in WS-A, who then assigned scores to the levels in each criterion individually. The stakeholders then deliberated on these individual scores and assigned scores to each level of the criterion by consensus. The range of scores was between 0 and 1, where, '0' depicted the lowest score, and '1' represented the highest score.

Gathering evidence

Data for the potential vaccines were collected from databases and reports from key organisations such as EPI, Communicable Disease Control of Directorate General of Health Services (CDC-DGHS), Institute of Epidemiology, Disease Control and Research and International Centre for Diarrhoeal Disease and Research, Bangladesh. A performance matrix was constructed, which presents data for each vaccine against the set of criteria. Then, workshop B (WS-B) was arranged in November 2019 to validate the data with a group of public health and vaccine experts in the country (n=6), that is, public health experts who were working in the disease surveillance (n=2), DGHS (n=2), Health Economics Unit (n=1) and NITAG (n=1). After reviewing and validation, they signed off on the performance matrix.

Rank ordering the potential vaccines

The scores for the different levels from the WS-A were combined with the validated performance matrix from the WS-B to calculate the scores for each vaccine on the different criteria. Then, using the additive method²¹ (see Equation 2),²⁶ the scores of each vaccine corresponding to the criteria level were multiplied by the weight of each criterion (from WS-A) to calculate the total scores of each potential vaccine. The vaccines were ranked based on the total scores of each vaccine, with the highest total score ranked top, and the next highest total second, and so on.

$$V_j = \sum C_{ij} * W_i \quad (2)$$

where V_j is the total value for alternative j ; C_{ij} is the score of alternative j on criteria i ; W_i is the weight attached to criteria i .

Appraising the rank of vaccines

Workshop C (WS-C) was conducted in December 2019 to appraise the vaccines. Stakeholders included the experts in the area of vaccination (n=10), that is, epidemiologists (n=2), virologists (n=3), infectious disease specialists (n=2), surveillance experts (n=1) and members of the vaccination policy programme (n=2). The performance matrix of the potential vaccines was provided in a paper-based format (online supplementary B) and the stakeholders were asked to assign the rank to the seven potential vaccines individually, where '1' was the most preferable vaccine and '7' was the least preferable vaccine. The mean rank of each vaccine was calculated from the ranks provided by each stakeholder, using the ROC method.²³

The ranking analysis of vaccines retrieved from step 5 (based on findings from WS-A and WS-B) were then presented to the stakeholders, along with the evidence of the cost-effectiveness and outbreak potentiality of each vaccine. Stakeholders then considered all this information and deliberated to reach a consensus on a final ranking of vaccines.

Application of vaccine prioritisation process in Bangladesh health system

A final workshop D was organised in January 2020 with the policymakers (n=28) working in vaccine decision-making, vaccination programme implementation, vaccine-related research and disease surveillance. The stakeholders were representatives from the ministry of health (n=12), the directorate office of health (n=9), development partners (n=2), health professional associations (n=2) and NGOs (n=3). This workshop involved the dissemination of the whole vaccine prioritisation process (including the selection of criteria, identification of vaccines and the MCDA methods), along with the findings.

Patient and public involvement

Patients and the general public were not involved in this study.

RESULTS

The list of potential vaccines for introduction in Bangladesh

WHO recommended 23 vaccines for routine vaccination globally, while the CDC-US recommended 16 vaccines and Gavi the vaccine alliance provided support against 16 infectious diseases.^{27–29} Bangladesh so far introduced 10 vaccines in their benefit package and two additional vaccines for the Haj pilgrimage travellers. Therefore, there were 11 vaccines not included yet in the Bangladesh health benefit package. After discussion among the core team and vaccine experts, four vaccines were excluded:

Table 1 Selecting criteria based on ranking from the WS-A

Criteria	Rank	
	Using the mean of individuals	Consensus after deliberation
Incidence rate of disease*	1	1
Case fatality rate*	2	2
Vaccine efficacy*	3	3
Size of population at risk*	5	4
Type of population at risk*	6	5
Outbreak potentiality	4	6
Cost-effectiveness	7	7
Severity of disease	8	8
Global target	9	9
Equity	10	10

*Criteria selected for vaccine prioritisation in Bangladesh. WS-A, workshop A.

tick-borne encephalitis and yellow fever as Bangladesh lacked incidence data for these diseases, and varicella and hepatitis A virus vaccines as they were not included in the benefit package of the neighbouring countries. Seven vaccines (ie, cholera, dengue, typhoid, HPV, influenza, JE and rotavirus) were then selected for consideration in the priority setting process.

Prioritisation criteria for vaccine introduction in Bangladesh

Sixty-seven criteria were identified in the systematic review, from which the core team identified 10 criteria as being potentially most relevant (table 1). Definitions of these 10 criteria were derived from the literature review.^{30–32}

In the WS-A, stakeholders discussed the importance of each of these 10 criteria and justification for inclusion in the set of prioritisation criteria to be used for vaccine introduction in Bangladesh. Participants ranked individually first and after deliberation, consensus was achieved. Table 1 presents the mean of individual ranking using ROC method and the final consensus ranking. Based on these rankings, stakeholders selected the top five criteria for vaccine prioritisation in Bangladesh (ie, incidence rate, case fatality rate, vaccine efficacy, size of population at risk and type of population at risk). In addition to these five quantitative criteria, stakeholders also decided to include two qualitative criteria: 'outbreak potentiality' and 'cost-effectiveness'. These two criteria were not weighted or scored explicitly, but were used in deliberative discussions.

Performance matrix

The data on the performance of each of seven vaccines against the prioritisation criteria are presented in table 2. The table presents data on the five quantitative criteria used for weighting and scoring, as well as the two qualitative criteria that were used in deliberative discussions.

Table 2 Performance matrix with data of vaccines on the criteria (after validation in WS-B)

Vaccine preventable disease	Incidence rate		Case fatality rate		Vaccine efficacy		Size of population at risk		Type of population at risk		Cost effectiveness*		Outbreak potential†‡
	Number of new cases per 100 000 population per year	Percentage of death among the cases in a year	Effectiveness of vaccine or reduction of diseases provided by vaccine (%)	Number of population at risk of getting infection per year (in millions)	Effectiveness of vaccine or reduction of diseases provided by vaccine (%)	Number of population at risk of getting infection per year (in millions)	Type of population needed to be vaccinated	Type of population at risk	Cost-effectiveness results from published literature				
Cholera ⁵⁶⁻⁵⁹	1640	3.0	53	15.175	53	15.175	Under five children	Cost-effective	High				
Dengue ⁶⁰⁻⁶²	3700	0.16†	66	2.18†	66	2.18†	Adult/high-risk	Very cost-effective	High				
HPV ⁶³⁻⁶⁶	10.6	0.0115	95	1.56	95	1.56	Woman	Highly cost-effective	Low				
Influenza ^{68 67-71}	10 200	0.088	63	15.5	63	15.5	High risk	Cost-effective	Low				
Japanese encephalitis ^{16 68 72-75}	2.7	30.0	96.2	10.77	96.2	10.77	High risk	Very cost-effective	Medium				
Rotavirus ^{68 76-78}	1080	0.0055	43	15.175	43	15.175	Under five children	Very cost-effective	High†				
Typhoid ^{68 79-83}	280	0.30	81.60	15.175	81.60	15.175	Under five children	Cost-effective	Medium				

*Not included in weighting and scoring, used in deliberative discussions in WS-C for final rankings. Judgements on cost-effectiveness were made from conclusions from published literature which evaluated the cost-effectiveness of these vaccines in Bangladesh or similar countries.

†Expert opinion.

‡Not included in weighting and scoring, used in deliberative discussions in WS-C for final rankings. HPV, human papillomavirus; WS-B, workshop B; WS-C, workshop C.

Table 3 Points allocated, and the calculated weights, for the criteria (from WS-A)

Criteria	Points	Weight
Incidence rate	100	0.26
Case fatality rate	85	0.22
Vaccine efficacy	80	0.21
Size of population at risk	75	0.19
Type of population at risk	50	0.13

WS-A, workshop A.

It should be noted that expert opinion (from WS-B) was used when there was no data available from published literature.

As shown in [table 2](#), influenza and dengue fever have the highest incidence among adults or high-risk groups but with relatively low case fatality rates. JE, on the other hand, has a relatively low incidence but with high case fatality rate (almost a third of patients dying from the condition). Among children, cholera and rotavirus seem to be with the highest incidence and cholera with a mortality rate of 3%. Vaccine efficacy seems to be excellent for JE and HPV (both above 90%), quite good for typhoid (above 80%), moderate for dengue and influenza (around 65%) and average for cholera (53%) and rotavirus (43%). All the vaccines seemed to be cost-effective or highly cost-effective. Finally, outbreak potential seems high for dengue, cholera and rotavirus.

Weighting and scoring

The participants of the WS-A consensually assigned 100 points to the criterion of ‘incidence rate’ and four other criteria were assigned points in accordance, with the least important criterion, ‘type of population at risk’ assigned 50 points. The weight of each criterion was calculated by using the normalisation method, and the weight of ‘incidence rate’ was estimated as 0.26, as presented in [table 3](#). ‘Case fatality rate’ and ‘vaccine efficacy’ were weighted similarly (0.22 and 0.21, respectively), ‘size of the population at risk’ had a weight of 0.19 and ‘type of population at risk’ had the lowest weight (0.13).

In the same workshop (WS-A), the stakeholders assigned scores for the different levels of the five criteria by consensus, using direct rating methods. For continuous criteria such as ‘incidence rate’, ‘case fatality rate’, ‘vaccine efficacy’ and ‘size of the population at risk’, the scores were assigned based on the levels of measures (eg, scores of 1, 0.8 and 0.55 for three levels for vaccine efficacy based on whether efficacy is >80%, 60%–80% or <60%), while the scores for categorical criteria such as ‘type of population at risk’ were based on the categories (eg, scores of 1, 0.8, 0.7 and 0.5 for children, high-risk groups, women and adults, respectively). The scores for the different levels of each criterion are presented in [table 4](#).

Table 4 Scores for the levels of criteria (from WS-A)

Criteria	Levels	Score
Incidence rate	Level 1: >1000/100 000	1.0
	Level 2: 100–1000/100 000	0.8
	Level 3: 10–100/100 000	0.5
	Level 4: <10/100 000	0.3
Case fatality rate	Level 1 >10%	1.0
	Level 2: 1%–10%	0.8
	Level 3: <1%	0.4
Vaccine efficacy	Level 1: >80%	1.0
	Level 2: 60%–79%	0.8
	Level 3: <60%	0.55
Size of population at risk	Level 1: >10 million	1.0
	Level 2: 1–10 million	0.8
	Level 3: 100 000–1 million	0.5
	Level 4: <100 000	0.3
Type of population at risk	Level A: children (<5 years)	1.0
	Level C: high-risk group	0.8
	Level B: women	0.7
	Level D: adult	0.5

WS-A, workshop A.

Rank ordering the potential vaccines

After combining the findings from tables 2–4 to estimate the score and weights (ie, the weights from WS-A, and the scores by combining the different levels from WS-A with the data from performance matrix validated in WS-B), the core team performed analysis of seven vaccines and produced the ranking results, as shown in table 5. Cholera vaccine was top-ranked with the highest total score of 0.86 primarily because it affects children, has a high incidence rate, high case fatality rate and with high size of population at risk. Despite having effective vaccines, JE and HPV ranked bottom (with scores of 0.74 and 0.68, respectively) because they have a low incidence rate and low size of population at risk.

Appraising the rank of vaccines

In the WS-C, the stakeholders reviewed the performance matrix and each stakeholder ranked the vaccines individually first. The mean of their individual rankings are presented in table 6. Based on the deliberations of performance matrix, the stakeholders in WS-C ranked HPV, JE and rotavirus, as the first, second and third, respectively. The stakeholders discussed and highlighted the importance of the vaccine for women, which was why HPV was ranked as the first. Then, they gave priority to vaccines with high incidence rate and high case fatality rate; therefore, JE and rotavirus vaccines were ranked next highest. This contrasts with the findings from the quantitative MCDA exercise by the core team (see table 5 using

findings from WS-A and WS-B), which suggested cholera, typhoid and influenza as the top three ranking vaccines.

The results of ranking by the core team (table 5) were then presented to the stakeholders in WS-C, along with the information on the potentiality of outbreak of the diseases and cost-effectiveness (see table 2). After considering all this information, the stakeholders adjusted the ranking by consensus and the final ranking is presented in table 6. HPV, JE and rotavirus still remained top three but the ranking order changed with JE, HPV and rotavirus being first, second and third, respectively.

Application of vaccine prioritisation process in Bangladesh health system

After dissemination of the findings, the policymakers agreed on the importance of appraising new interventions scientifically and supported the use of MCDA in the priority setting process for vaccine introduction decision-making. The key personnel of the ministry of health and family welfare, Bangladesh, stated that *'It is better for Bangladesh at present to have this system to prioritise vaccines in the country. Bangladesh, a lower-middle income country is graduating Gavi funding. So, we have to change our decision-making process from donor influenced decision-making to self-decision-making'*. Based on the MCDA results, JE vaccine is planned to be recommended to the decision-makers for introduction into the national vaccine benefit package. They also highlighted that after the selection of vaccines, the country should prepare for vaccine logistics such as cold-chain capacity and other programmatic issues.

DISCUSSION

Summary of the study

This study represents the first time an explicit priority setting process based on MCDA was used for the prioritisation of vaccines in Bangladesh. Vaccines selected for prioritisation were those which were recommended by the international organisations but not included in health benefit package of Bangladesh. The potential multiple criteria were identified systematically from published literature, and shortlisted in two phases to select five quantitative criteria and two qualitative criteria for the evaluation of the vaccines. Weighting and scoring of the quantitative criteria were explicit and participatory, and the tools used for eliciting scores and weights were user friendly and well understood by the stakeholders. The final ranking of the vaccines was determined after deliberative discussions based on the performance matrix, which considered both quantitative criteria and qualitative criteria.

Statement of the principal findings

Through this explicit MCDA approach, JE vaccine was placed as the top-ranked vaccine and is planned to be recommended to the decision-makers for introduction into the national vaccine benefit package. The policymakers support the use of systematic evidence-based decision-making processes such as MCDA for vaccine

Table 5 Rank order of vaccine using only quantitative criteria (from WS-A and WS-B)

Weight of criteria	Incidence rate				Case fatality rate			Vaccine efficacy			Size of population at risk				Type of population at risk				Total	Rank
	L1	L2	L3	L4	L1	L2	L3	L1	L2	L3	L1	L2	L3	L4	L-A	L-B	L-C	L-D		
Levels	1.0	0.8	0.5	0.3	1.0	0.8	0.4	1.0	0.8	0.55	1.0	0.8	0.5	0.3	1.0	0.8	0.7	0.5	Sum	Rank
Score of levels	1.0	0.8	0.5	0.3	1.0	0.8	0.4	1.0	0.8	0.55	1.0	0.8	0.5	0.3	1.0	0.8	0.7	0.5		
Cholera	(0.26×1.0) 0.26				(0.22×0.8) 0.17			(0.21×0.55) 0.11			(0.19×1.0) 0.19				(0.13×1.0) 0.13				0.86	1
Typhoid	(0.26×0.8) 0.20				(0.22×0.4) 0.09			(0.21×1.0) 0.21			(0.19×1.0) 0.19				(0.13×1.0) 0.13				0.82	2
Influenza	(0.26×1.0) 0.26				(0.22×0.4) 0.09			(0.21×0.8) 0.16			(0.19×1.0) 0.19				(0.13×0.7) 0.09				0.79	3
Rotavirus	(0.26×1.0) 0.26				(0.22×0.4) 0.09			(0.21×0.55) 0.11			(0.19×1.0) 0.19				(0.13×1.0) 0.13				0.78	4
Dengue	(0.26×1.0) 0.26				(0.22×0.4) 0.09			(0.21×0.8) 0.16			(0.19×0.8) 0.15				(0.13×0.7) 0.09				0.75	5
Japanese encephalitis	(0.26×0.3) 0.08				(0.22×1.0) 0.22			(0.21×1.0) 0.21			(0.19×0.8) 0.15				(0.13×0.7) 0.09				0.74	6
HPV	(0.26×0.5) 0.13				(0.22×0.4) 0.09			(0.21×1.0) 0.21			(0.19×0.8) 0.15				(0.13×0.8) 0.10				0.68	7

*Data from performance matrix (table 2) were combined with the scores for different levels (table 4) to estimate the scores for each vaccine. These were then multiplied with weights (table 3) to calculate overall scores, which were then used for ranking.

HPV, human papillomavirus; WS-A, workshop A; WS-B, workshop B.

Table 6 Ranking of vaccines

Vaccine	Ranking from WS-C	Ranking from the analysis of WS-A and WS-B	Final ranking after deliberation in WS-C*
Japanese encephalitis	2	6	1
HPV	1	7	2
Rotavirus	3	4	3
Cholera	5	1	4
Typhoid	4	2	5
Dengue	7	5	6
Influenza	6	3	7

*including consideration of information on cost-effectiveness and outbreak potential.
HPV, human papillomavirus; WS-A, workshop A; WS-B, workshop B; WS-C, workshop C.

introduction in Bangladesh, and to prioritise health interventions in the country.

Strengths of the study, and comparison to findings from other studies

Stakeholder involvement

The MCDA process was supported by different stakeholders. Members of the different decision-making committees (NITAG), implementing bodies (EPI and others) and health professional associations were involved in every step of this study. Stakeholders of implementing agencies—EPI and CDC-DGHS—also participated in the deliberative process and ranking. NITAG members and members of National Committee for Immunization Practices also participated in the final decision-making workshop at the ministry level. Participation of stakeholders in this study ensured the transparency and accountability of decision-making, which is essential for a fair priority setting approach.³³ The importance of involving different stakeholders during the decision-making of vaccine introduction is also highlighted in other countries such as South Korea,³⁴ Oman,³⁵ Indonesia³⁶ and the Netherlands.³⁷

Criteria used in priority setting

Incidence rate of the disease and case fatality rate criteria were weighted highly, indicating that disease burden was considered important for vaccine selection by the stakeholders. This finding is similar to other studies which suggest disease burden as the most common and important criterion considered by other low- and middle-income countries (LMICs) during national decision-making.^{19 38–42} Efficacy of the vaccines was weighted as the next most important criterion suggesting that clinical effectiveness is also important.

Deliberative MCDA

The final ranking in this study was based on deliberation using the performance matrix, where the weights and scores were not explicit. Despite the lack of explicit

weighting and scoring, deliberative discussions are considered to be a very important part of MCDA process as it allows a shared understanding of the data, criteria and priorities. Deliberation among stakeholders followed by simple ranking appears a feasible strategy for the prioritisation of vaccines for introduction in Bangladesh and other LMICs. Kenya and Iran selected vaccines by voting, whereas Oman, India and the Netherlands selected vaccines by expert evaluation which were evidence-based but not systematic.^{35 37 43 44} Korea and Thailand selected vaccines systematically via evidence-based deliberation using DELPHI and MCDA techniques.^{34 45} Recent consensus on the use of MCDA for Health Technology Assessment (HTA),⁴⁶ recommends deliberative MCDA approach over quantitative MCDA. Furthermore, a recent study by WHO encouraged weighting and scoring as they help streamline the deliberative discussions.⁴⁷ The methods used in our study, where the stakeholders deliberated the results from the quantitative MCDA and the performance matrix before finalising the ranking of vaccines, are in line with these recommendations.

Implications for policymakers

While decision-making around vaccines in LMICs has been driven by donor funding, our study shows that it is possible to perform prioritisation systematically using evidence-based MCDA approaches. Based on the results of the MCDA study, the top-ranked JE vaccine is planned to be recommended to the decision-makers for introduction into the national vaccine benefit package. Please note that the ranking of vaccines and the selection of JE vaccine is country specific and may not be applicable to other settings. It is noteworthy that decision-making itself is a dynamic process, and vaccine performance on some criteria is likely to change over time. Therefore, we recommend Bangladesh undertake this priority setting process routinely even though most of the countries evaluate vaccines to be introduced once.^{39 40 43 48–51}

Limitations of the study

Different sets of stakeholders took part in the three workshops, resulting in a lack of a consistent group of stakeholders (and hence values/preferences) throughout the MCDA process. The ranking from quantitative weighting and scoring (from WS-A and WS-B) was different to the ranking by the stakeholders in the WS-C, who ranked the vaccines after a deliberative process. This may be due to the differences in the stakeholder membership between the different workshops and the underlying differences in their preferences.

Furthermore, the vaccine ranking in WS-C was finalised after considering the cost-effectiveness and the outbreak potentiality criteria, as well as the quantitative ranking. Also, the stakeholder preferences were implicit in the WS-C while they were explicitly elicited in the ranking using quantitative weighting and scoring (from WS-A and WS-B). This highlights the importance of ensuring a consistent set of criteria and a consistent preference

elicitation methodology throughout the MCDA process, along with a consistent group of stakeholders. If the membership, the criteria set or the methodology changes between the different workshops, there is a possibility that the ranking may change quite substantially.

Despite the inclusion of a wide variety of stakeholders, our study does not represent all stakeholders' perspectives. Stakeholders from private sectors and representatives of patient groups were not involved in the process leading to uncertainty in accountability of the results to those stakeholders.

Finally, in our study, the cost-effectiveness considerations and data of outbreak potentiality were included as qualitative criteria rather than quantitative criteria with explicit weighting and scoring. It is important to note that cost-effectiveness is not recommended as a criterion in the MCDA,^{52–53} as such, a pragmatic approach was taken to consider this information qualitatively rather than weighting and scoring. While decision-making around vaccines has typically been driven by donor funding assurance, financial considerations are highlighted as being key by stakeholders. Capacity building around economic evaluation and budget impact analysis of vaccines is needed in LMICs such as Bangladesh to support evidence-based priority setting combining MCDA with Value for Money approaches.^{53–55}

CONCLUSIONS

This study presents the first application of MCDA to support vaccine prioritisation in Bangladesh health system. This study involved relevant stakeholders in priority setting process and achieved the objectives of prioritising the vaccines for introduction in Bangladesh in a transparent way, using systematic evidence-based decision-making. JE vaccine was placed as the top-ranked vaccine and is planned to be recommended to the decision-makers for introduction into the national vaccine benefit package. The use of MCDA to prioritise interventions in healthcare should be promoted as the decision-making process can be improved using systematic approaches.

Contributors MSH conceived and designed the study, collected data, analysed the results and drafted the manuscript. SY and PT helped in the study design, data analysis, interpretation of results and reviewed the manuscript. MT contributed to the design of the study and reviewed the manuscript. All authors discussed the findings of the study, edited and approved the manuscript. All authors are responsible for the overall content as guarantors.

Funding International Decision Support Initiative (iDSI) (OPP1087363)

Disclaimer The findings, interpretations and conclusions expressed in this article do not necessarily reflect the views of the funding agencies.

Competing interests The authors declared no potential conflicts of interest with respect to the research and authorship.

Patient consent for publication Not applicable.

Ethics approval Ethical clearance of this study was obtained from the Bangladesh Medical and Research Council (BMRC) and informed written consent was obtained from the stakeholders participating in the workshops.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available from the corresponding author. Email: sitaporn.you@mahidol.edu.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Mohammad Sabbir Haider <http://orcid.org/0000-0003-0013-0051>

Sitaporn Youngkong <http://orcid.org/0000-0002-2448-3954>

Montarat Thavorncharoensap <http://orcid.org/0000-0002-8256-2167>

Praveen Thokala <http://orcid.org/0000-0003-4122-2366>

REFERENCES

- 1 Timmis JK, Black S, Rappuoli R. Improving accountability in vaccine decision-making. *Expert Rev Vaccines* 2017;16:1057–66.
- 2 Brenzel L, Wolfson LJ, Fox-Rushby J. Vaccine-Preventable Diseases—Chapter 20. *Disease Control Priorities in Developing Countries*:389–411.
- 3 Ehreth J. The value of vaccination: a global perspective. *Vaccine* 2003;21:4105–17.
- 4 Ehreth J. The global value of vaccination. *Vaccine* 2003;21:596–600.
- 5 Mauskopf J, Standaert B, Connolly MP, et al. Economic analysis of vaccination programs: an ISPOR good practices for outcomes research Task force report. *Value Health* 2018;21:1133–49.
- 6 Baltussen R, Jansen MP, Mikkelsen E, et al. Priority setting for universal health coverage: we need evidence-informed Deliberative processes, not just more evidence on cost-effectiveness. *Int J Health Policy Manag* 2016;5:615–8.
- 7 Norheim OF. Ethical priority setting for universal health coverage: challenges in deciding upon fair distribution of health services. *BMC Med* 2016;14:75.
- 8 Youngkong S, Kapiriri L, Baltussen R. Setting priorities for health interventions in developing countries: a review of empirical studies. *Trop Med Int Health* 2009;14:930–9.
- 9 Munira SL, Fritzen SA. What influences government adoption of vaccines in developing countries? A policy process analysis. *Soc Sci Med* 2007;65:1751–64.
- 10 Baltussen R, Niessen L. Priority setting of health interventions: the need for multi-criteria decision analysis. *Cost Eff Resour Alloc* 2006;4:14.
- 11 Mitton C, Donaldson C. Health care priority setting: principles, practice and challenges. *Cost Eff Resour Alloc* 2004;2:3.
- 12 Thokala P, Duenas A. Multiple criteria decision analysis for health technology assessment. *Value Health* 2012;15:1172–81.
- 13 Marsh K, Goetghebuer M, Thokala P. *Multi-Criteria decision analysis to support healthcare decisions*. Springer, 2017.
- 14 Haque F, Hossain MJ, Kundu SK. Cholera Outbreaks in Urban Bangladesh. In: Sunnivaale, ed. 2011. *Epidemiology*. 3, 2013.
- 15 Hossain MJ, Gurley ES, Montgomery S, et al. Hospital-Based surveillance for Japanese encephalitis at four sites in Bangladesh, 2003–2005. *Am J Trop Med Hyg* 2010;82:344–9.
- 16 Paul RC, Rahman M, Gurley ES, et al. A novel low-cost approach to estimate the incidence of Japanese encephalitis in the catchment area of three hospitals in Bangladesh. *Am J Trop Med Hyg* 2011;85:379–85.
- 17 Hasan AZ, Saha S, Saha SK, et al. Using pneumococcal and rotavirus surveillance in vaccine decision-making: a series of case studies in Bangladesh, Armenia and the Gambia. *Vaccine* 2018;36:4939–43.
- 18 Dhar-Chowdhury P, Paul KK, Haque CE, Chowdhury PDP KK, Hossain S, et al. Dengue seroprevalence, seroconversion and risk factors in Dhaka, Bangladesh. *PLoS Negl Trop Dis* 2017;11:e0005475.

- 19 Uddin J, Sarma H, Bari TI, *et al.* Introduction of new vaccines: decision-making process in Bangladesh. *J Health Popul Nutr* 2013;31:211–7.
- 20 Marsh K, Jzerman M, Thokala P, *et al.* Multiple Criteria Decision Analysis for Health Care Decision Making--Emerging Good Practices: Report 2 of the ISPOR MCDA Emerging Good Practices Task Force. *Value Health* 2016;19:125–37.
- 21 Thokala P, Devlin N, Marsh K, *et al.* Multiple Criteria Decision Analysis for Health Care Decision Making--An Introduction: Report 1 of the ISPOR MCDA Emerging Good Practices Task Force. *Value Health* 2016;19:1–13.
- 22 Haider MS. Multi-Criteria Decision Analysis for Priority Setting of Vaccine Introduction in Bangladesh [Ph.D. Thesis] 2020:Mahidol University.
- 23 Roszkowska E. Rank ordering criteria weighting methods--a comparative overview; 2013.
- 24 Podvezko A, Podvezko V. Influence of data transformation on multicriteria evaluation result. *Procedia Engineering* 2015;122:151–7.
- 25 *Normalization techniques for multi-criteria decision making: analytical hierarchy process case study.* 2016, Springer
- 26 Marsh K, Thokala P, Mühlbacher A. Incorporating Preferences and Priorities into MCDA: Selecting an Appropriate Scoring and Weighting Technique. In: Marsh K, Goetghebeur M, Thokala P, eds. *Multi-Criteria decision analysis to support healthcare decisions*. Cham: Springer International Publishing, 2017: 47–66.
- 27 World Health Organization. WHO recommendations for routine immunization - summary tables 2019 [updated 26 April 2019; cited 2019 21 November 2019], 2019. Available: https://www.who.int/immunization/policy/immunization_tables/en/ [Accessed 21 November, 2019].
- 28 Gavi The Vaccine Alliance. Gavi, The Vaccine Alliance-Vaccine Support 2019 [cited 2019 21 November 2019]. Available: <https://www.gavi.org/programmes-impact/types-support/vaccine-support>
- 29 Centers for Disease Control and Prevention. Centers for disease control and prevention 2019, 2019. Available: <https://www.cdc.gov/vaccines/> [Accessed 21 November 2019].
- 30 Rashid K. Hyder's Textbook of community Medicine and public Health. Dhaka RHM publishers; 2008.
- 31 World Health Organization. World Health Statistics 2012 - Indicator compendium: World Health Organization; 2012.
- 32 World Health Organization. Global reference list of 100 core health indicators: World Health organization; 2015.
- 33 Daniels N. Accountability for Reasonableness. *BMJ* 2000;321:1300–1.
- 34 Choe YJ, Han OP, Cho H, *et al.* Prioritization of the introduction of new vaccines to the National immunization program in the Republic of Korea. *Vaccine* 2014;32:6049–53.
- 35 Al Awaidy S. The National Committee for vaccines regulation and surveillance of vaccine-preventable diseases in the Sultanate of Oman: evidence-based approach and consensus decision-making. *Vaccine* 2010;28 Suppl 1:A39–41.
- 36 Hadisoemarto PF, Reich MR, Castro MC. Introduction of pentavalent vaccine in Indonesia: a policy analysis. *Health Policy Plan* 2016;31:1079–88.
- 37 van der Putten IM, Paulus ATG, Hilgsmann M, *et al.* Evidence-Informed vaccine decision making: the introduction of human papilloma virus (HPV) vaccination in the Netherlands. *Health Policy* 2019;123:260–266.
- 38 Burchett HED, Mounier-Jack S, Griffiths UK, *et al.* National decision-making on adopting new vaccines: a systematic review. *Health Policy Plan* 2012;27 Suppl 2:ii62–76.
- 39 Hadisoemarto PF, Reich MR, Castro MC. Introduction of pentavalent vaccine in Indonesia: a policy analysis. *Health Policy Plan* 2016;31:1079–88.
- 40 Kant L. NTAGI Subcommittee recommendations on Haemophilus influenzae type B (Hib) vaccine introduction in India. *Indian Pediatr* 2009;46:945–54.
- 41 Makinen M, Kaddar M, Moldrem V, *et al.* New vaccine adoption in lower-middle-income countries. *Health Policy Plan* 2012;27 Suppl 2:ii39–49.
- 42 van der Putten IM, Evers SMAA, Deogaonkar R, *et al.* Stakeholders' perception on including broader economic impact of vaccines in economic evaluations in low and middle income countries: a mixed methods study. *BMC Public Health* 2015;15:356.
- 43 Dawa J, Chaves SS, Ba Nguz A, *et al.* Developing a seasonal influenza vaccine recommendation in Kenya: process and challenges faced by the National immunization technical Advisory group (NITAG). *Vaccine* 2019;37:464–72.
- 44 Muliylil JP, Bhan MK, Bhattacharya SK, *et al.* NTAGI Subcommittee recommendations on Haemophilus influenzae type B (Hib) vaccine introduction in India. *Indian Pediatr* 2009;46:945–54.
- 45 Pooripussarakul S, Riewpaiboon A, Bishai D, *et al.* What criteria do decision makers in Thailand use to set priorities for vaccine introduction? *BMC Public Health* 2016;16:684.
- 46 Baltussen R, Marsh K, Thokala P, *et al.* Multicriteria decision analysis to support health technology assessment agencies: benefits, limitations, and the way forward. *Value Health* 2019;22:1283–8.
- 47 Botwright S, Giersing BK, Meltzer MI, *et al.* The CAPACITI Decision-Support tool for national immunization programs. *Value Health* 2021;24:1150–7.
- 48 Dorji T, Tshomo U, Phuntsho S, *et al.* Introduction of a national HPV vaccination program into Bhutan. *Vaccine* 2015;33:3726–30.
- 49 Fletcher MA, Tetelboum R, Fritzell B. Time to recommend pneumococcal vaccination for all children in Europe: experience in France. *Eur J Pediatr* 2002;161 Suppl 2:S132–4.
- 50 Garpenholt O, Fredlund H, Timпка T. Immunization against Haemophilus influenzae type b in Sweden--a study of the introduction process. *Scand J Public Health* 2001;29:271–8.
- 51 Ngcobo NJ, Cameron NA. The decision making process on new vaccines introduction in South Africa. *Vaccine* 2012;30 Suppl 3:C9–13.
- 52 Marsh KD, Sculpher M, Caro JJ, *et al.* The use of MCDA in HTa: great potential, but more effort needed. *Value Health* 2018;21:394–7.
- 53 Marsh K, Thokala P, Youngkong S, *et al.* Incorporating MCDA into HTa: challenges and potential solutions, with a focus on lower income settings. *Cost Eff Resour Alloc* 2018;16:43.
- 54 Wilson ECF, Peacock SJ, Ruta D. Priority setting in practice: what is the best way to compare costs and benefits? *Health Econ* 2009;18:467–78.
- 55 Airoldi M, Morton A, Smith J. Healthcare prioritisation at the local level: a socio-technical approach 2011.
- 56 Khan AI, Levin A, Chao DL, *et al.* The impact and cost-effectiveness of controlling cholera through the use of oral cholera vaccines in urban Bangladesh: a disease modeling and economic analysis. *PLoS Negl Trop Dis* 2018;12:e0006652.
- 57 World Health Organization. Cholera, 2014. *Wkly Epidemiol Rec* 2015;90:517–44.
- 58 Hossain MA, AHMM L, Rayhan MI. *Population projection of Bangladesh: dynamics and trends 2011-2061*. Dhaka, Bangladesh: Bangladesh Bureau of Statistics, Statistics and Information Division, Ministry of Planning, 2015.
- 59 Eisenberg MC, Robertson SL, Tien JH. Identifiability and estimation of multiple transmission pathways in cholera and waterborne disease. *J Theor Biol* 2013;324:84–102.
- 60 Haider S, Chaikledkaew U, Thavorncharoensap M. *Cost-Effectiveness of dengue vaccine introduction in Dhaka City, Bangladesh. HTAsiaLink*. Thailand: Chiang Mai, 2018.
- 61 World Health Organization. Dengue vaccine: who position paper – July 2016. *Wkly Epidemiol Rec* 2016;91:349–64.
- 62 World Health Organization. Dengue vaccine: WHO position paper, July 2016 - recommendations. *Vaccine* 2017;35:1200–1.
- 63 Mahumud RA, Gow J, Alam K, *et al.* Cost-Effectiveness of the introduction of two-dose bi-valent (Cervarix) and quadrivalent (Gardasil) HPV vaccination for adolescent girls in Bangladesh. *Vaccine* 2020;38:165–72.
- 64 Services DGoH. National strategy for cervical cancer prevention and control Bangladesh (2017-2022): Directorate General of health services; 2017.
- 65 Jit M, Brisson M, Portnoy A, *et al.* Cost-Effectiveness of female human papillomavirus vaccination in 179 countries: a prime modelling study. *Lancet Glob Health* 2014;2:e406–14.
- 66 Riesen M, Garcia V, Low N, *et al.* Modeling the consequences of regional heterogeneity in human papillomavirus (HPV) vaccination uptake on transmission in Switzerland. *Vaccine* 2017;35:7312–21.
- 67 Ahmed M, Aleem MA, Roguski K, *et al.* Estimates of seasonal influenza-associated mortality in Bangladesh, 2010–2012. *Influenza Other Respir Viruses* 2018;12:65–71.
- 68 Brooks WA, Goswami D, Rahman M, *et al.* Influenza is a major contributor to childhood pneumonia in a tropical developing country. *Pediatr Infect Dis J* 2010;29:216–21.
- 69 Zaman K, Roy E, Arifeen SE, *et al.* Effectiveness of maternal influenza immunization in mothers and infants. *N Engl J Med* 2008;359:1555–64.
- 70 Orenstein EW, Orenstein LAV, Diarra K, *et al.* Cost-Effectiveness of maternal influenza immunization in Bamako, Mali: a decision analysis. *PLoS One* 2017;12:e0171499.
- 71 van den Driessche P. Reproduction numbers of infectious disease models. *Infect Dis Model* 2017;2:288–303.
- 72 Touch S, Suraratdecha C, Samnang C, *et al.* A cost-effectiveness analysis of Japanese encephalitis vaccine in Cambodia. *Vaccine* 2010;28:4593–9.



- 73 World Health Organization. Japanese encephalitis vaccines: who position paper. *Weekly Epidemiological Record= Relevé épidémiologique hebdomadaire* 2015;90:69–88.
- 74 Tandan JB, Ohrr H, Sohn YM, *et al.* Single dose of SA 14-14-2 vaccine provides long-term protection against Japanese encephalitis: a case-control study in Nepalese children 5 years after immunization. drjbtandan@yahoo.com. *Vaccine* 2007;25:5041–5.
- 75 . Estimation of parameters and basic reproductive ratio for Japanese encephalitis transmission in the Philippines using a sequential Monte Carlo filter. *2017 IEEE Conference on Control Technology and Applications (CCTA)*. IEEE, 2017.
- 76 Sarker AR, Sultana M, Mahumud RA, *et al.* Cost-Effectiveness analysis of introducing universal childhood rotavirus vaccination in Bangladesh. *Hum Vaccin Immunother* 2018;14:189–98.
- 77 Troeger C, Khalil IA, Rao PC, *et al.* Rotavirus vaccination and the global burden of rotavirus diarrhea among children younger than 5 years. *JAMA Pediatr* 2018;172:958–65.
- 78 Zaman K, Dang DA, Victor JC, *et al.* Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in Asia: a randomised, double-blind, placebo-controlled trial. *Lancet* 2010;376:615–23.
- 79 Antillón M, Bilcke J, Paltiel AD, *et al.* Cost-Effectiveness analysis of typhoid conjugate vaccines in five endemic low- and middle-income settings. *Vaccine* 2017;35:3506–14.
- 80 Marchello CS, Hong CY, Crump JA. Global typhoid fever incidence: a systematic review and meta-analysis. *Clin Infect Dis* 2019;68:S105–16.
- 81 Yu AT, Amin N, Rahman MW, *et al.* Case-Fatality ratio of blood culture-confirmed typhoid fever in Dhaka, Bangladesh. *J Infect Dis* 2018;218:S222–6.
- 82 Shakya M, Colin-Jones R, Theiss-Nyland K, *et al.* Phase 3 efficacy analysis of a typhoid conjugate vaccine trial in Nepal. *N Engl J Med* 2019;381:2209–18.
- 83 Pitzer VE, Bowles CC, Baker S, *et al.* Predicting the impact of vaccination on the transmission dynamics of typhoid in South Asia: a mathematical modeling study. *PLoS Negl Trop Dis* 2014;8:e2642.

SUPPLEMENTARY**Supplementary A: Data collection instrument of Workshop A - Ranking of criteria**

Name of the participants: _____

Designation: _____

Organization: _____

Note: Pls. make rank of the following criteria - 1 to 10 (Where is the most important and 10 is the less important)

CRITERIA	DEFINITION	RANK
Criteria of Disease		
Case fatality rate	Percentage of death among the cases	
Incidence rate of disease	Number of new cases per 100,000 population per year	
Outbreak potentiality	Potentiality of the disease to be epidemic Potentiality will be measured by the reproduction of the disease	
Severity of disease	Symptoms of the disease; how severe the disease are in the most of the cases	
Size of population at risk	Size of the population at risk or the target population for vaccination	
Type of Target population/ Demographic consideration	Demographic consideration or Target population for the vaccination against the disease (e.g. children or female or adult)	
Criteria of Vaccine		
Cost-effectiveness	Cost-effectiveness of vaccine; Incremental cost-effectiveness ration (ICER) will be \$/QALY gained or \$/DALY avoided if the vaccine introduced in comparison to the No vaccination	
Vaccine efficacy	Effectiveness of vaccine or the percentage reduction of diseases provided by vaccine	
Other Criteria		
Equity	Disease occur more in economically poor people or disadvantaged population	
Global Target	Global agenda of eradication/ elimination/ control target	

Supplementary B: Data collection instrument of Workshop C - Ranking of vaccines

Name of the participants: _____

Designation: _____

Organization: _____

Please rank vaccines from 1-7, where 1 is most favourable and 7 is least favourable

	Criteria	Rotavirus	HPV	Cholera	JE	Typhoid	Influenza	Dengue
D I S E A S E	1. Incidence rate Number of new cases per 100,000 population per year	1080/100,000	24.3/100,000	210/100,000	2.7/100,000	280/100,000	10,000/ 100,000 person year (2008) 6600/100,000 person year (2009) 17000/100,000 person year (2010)	1340-5780/ 100,000 person- season
	2. Case fatality rate Percentage of death among the cases	0.03% 12.42/100,000 among < 5 years of age (Rotavirus gastroenteritis mortality) (1.24%)	1.8% 50%	1.5%	10-30% (25%)	0.3% (1%)	0.08%	2.5%
V A C C I N E	Type of vaccine	RV5; Live attenuated	Human Papillomaviruses Nanovalent	Shanchol	SA14-14-2 JE Vaccine	Typhoid Conjugate vaccine	Influenza trivalent vaccine: Single dose	Dengvaxia live attenuated, recombinant tetravalent vaccine
	Dosage	3 dosages	2 dosages	2 dosages	Single dose	Single dose	Single dose	3 dosages
	3. Vaccine efficacy Effectiveness of vaccine or the percentage reduction of diseases provided by vaccine	55% (40-85%)	90-100%	50-60%	95% >85%	50-72%	40% 40-60%	66%
P O P U L A T I O N	4. Type of Target population Demographic consideration or Target population for the vaccination against the disease (e.g. children or female or adult)	Under 5 population	Girls of 10 years of age or class 5 student 1.54 million	1-5years of population Urban and high risk population	1-15 years of children; Routine immunization 9- 12 months	Under 5 population	High risk group	Dhaka City population
	5. Size of population at risk (million) Number of population need to be vaccinated or size of the population at risk or the target population for vaccination	15.17	9.17 1.54	13.3	7.4 (1-15 years of children)	15.17	15.47	2.18
	RANK							