

This is a repository copy of Differences in older adults walking football initiation and maintenance influences across respondent characteristics: a cross-sectional survey.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/184924/</u>

Version: Accepted Version

Article:

Cholerton, R., Butt, J., Quirk, H. orcid.org/0000-0003-2716-4681 et al. (1 more author) (2022) Differences in older adults walking football initiation and maintenance influences across respondent characteristics: a cross-sectional survey. Journal of Aging and Physical Activity, 30 (6). pp. 936-949. ISSN 1063-8652

https://doi.org/10.1123/japa.2021-0305

Accepted author manuscript version reprinted, by permission, from Journal of Aging and Physical Activity, 2022, https://doi.org/10.1123/japa.2021-0305. © Human Kinetics, Inc.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

1 MANUSCRIPT

- 2
- 3 Running title: Differences in walking football initiation and maintenance influences across
- 4 respondent characteristics: a cross-sectional survey in 50-75 year -old adults

Abstract

2 Despite health benefits gained from physical activity and sport participation, older adults are 3 less likely to be active. This study investigates what influences 50–75-year-olds (N=439) to 4 initiate and maintain walking football, across gender, socioeconomic status, number of health 5 conditions and physical activity level. It also considers relationships between participant 6 characteristics and influences, and intentions to play after a forced break (COVID-19). 7 Results of a UK online cross-sectional survey found those with two or more health conditions 8 rated social influences significantly higher in initiation and maintenance, than participants 9 with no health conditions. Multiple regression analysis found a positive walking football 10 culture and perceived use of maintenance resources contributed significantly to intentions to 11 return to play after COVID-19 restrictions eased. Practitioners should consider providing 12 opportunities for social connection, foster a positive walking football culture, and encourage 13 players to utilise maintenance resources (e.g., scheduling sessions) in older adult walking 14 football sessions.

15

16 Keywords: behaviour change, soccer, physical activity, survey research, older adults17

Introduction

2 Research has shown that physical activity (PA) improves physiological and psychological 3 health markers in older adults (Chapman et al., 2013; Rezende et al., 2014), however reports 4 have suggested that older adults in the UK are more likely to lead inactive lives compared 5 with the general population (Sport England, 2018). Furthermore, research has found that 6 older adults are less likely to maintain PA in the long-term (Kendrick et al., 2018). Sport 7 participation, as a particular form of PA, offers additional psychological and physical benefits 8 in older age. These benefits include less total sedentary behaviour, quicker reaction times, 9 and better self-reported health related quality of life than those who do not play 10 sport(Gayman et al., 2017; Pesce & Audiffren, 2011). Despite this, research has reported a 11 number of barriers to sport participation in older age, including a lack of accessible sport 12 programmes (including inappropriate facilities or lack of senior competitions) (Jenkin et al., 13 2018). Furthermore, a low socio-economic status (SES), identifying as a woman, and poor 14 health have been reported widely as barriers to PA and sport participation (Dhalwani et al., 15 2016; Fox et al., 2011; Hirvensalo & Lintunen, 2011; Jenkin et al., 2018). In order to 16 increase numbers of older adults who benefit from sport participation, it is important to 17 identify determinants and factors which promote long-term participation. As a result, sport 18 programmes can be tailored and optimised to promote and encourage physical activity 19 behaviour among a diverse range of older adult populations.

20

Walking football (soccer) is a form of adapted sport becoming popular in the UK, where
football rules are adapted for players to play the game at a walking pace (Lloyd, 2019).
Researchers have suggested adapted sport as a form of sport participation which may mitigate
some barriers expressed by older adults, for example health status (Jenkin et al., 2018).
Studies investigating the effectiveness of walking football on older adult health have shown

1 promising findings relating to improving physiological health, and the sport being a 2 sustainable form of activity for older adults (McEwan et al., 2019; Reddy et al., 2017). 3 Qualitative methods that have been used to explore the experiences of older adults who 4 participated in a walking football intervention found that perceived benefits included the 5 enjoyment of social interactions and walking football (Reddy et al., 2017). However, these 6 studies tended to investigate the overall impact of walking football and the benefits 7 associated with playing the sport. Research within the field of behaviour change has 8 suggested that differences to physiological and mental health are seen after sustaining PA 9 longer than six months (Almeida et al., 2014; Laitakari et al., 1996; Liu-Ambrose, 2010) and 10 further differences are seen between those who are starting (initiating) and maintaining a 11 behaviour long term (Rothman, 2000; van Stralen et al., 2009, 2010). It is therefore important 12 to investigate the influences involved in different phases of behaviour change, so relevant 13 approaches to helping older adults sustain walking football play long term can be employed. 14

15 Recent qualitative research into walking football has furthered understanding of psychosocial 16 and behavioural aspects of initiating and maintaining participation among older adults 17 (Cholerton et al., 2020, 2021). This research found multiple individual, social and 18 environmental influences present across initiation and maintenance (e.g., values relating to 19 health and sport, team connections). Further evidence suggests that some influences also 20 differ across phases, for example walking football culture (e.g. the values a club and sporting 21 body holds) being influential at maintenance of walking football, but not initiation. There is 22 also research evidence that influences differ across varying participant characteristics, such as 23 SES and gender (e.g., different forms of walking football awareness across varying SES). 24 Previous research has also found differing levels of PA and sport participation among older 25 adults with varying characteristics, for example across gender and SES (Kamphuis et al.,

1 2009; Murtagh et al., 2015) as well as varying experiences of PA and sport participation 2 among participants with different numbers of health conditions and weekly PA levels 3 (Dhalwani et al., 2016; Jenkin et al., 2017; Stenner et al., 2020). As it is evident that varying 4 PA and sport experiences are seen across a number of participant characteristics, it is 5 important to understand whether there are any differences in influences relating to walking 6 football initiation and maintenance, across these characteristics. This knowledge would allow 7 clubs and sporting bodies to make informed and tailored recommendations relating to the 8 promotion of the game and session delivery, with the aim of aiding sustained participation in 9 the sport for varying audiences.

10

11 Despite the known benefits of long-term PA and sport participation, research has suggested 12 that older people have reduced chances of maintaining PA. Kendrick and colleagues (2018) 13 found only 48% of participants achieved the stated PA target (150 minutes) in the most 14 successful exercise programme group assessed, 24-months after the exercise programme had 15 ended. The Coronavirus (COVID-19) pandemic also had a direct effect on walking football 16 clubs, due to strict health and safety restrictions imposed (The FA, 2020). As breaks in play 17 due to illness, injury and holidays are discussed throughout previous walking football 18 qualitative research (Cholerton et al., 2020, 2021) and relapse prevention becoming a key 19 aspect of PA maintenance strategies (Sherwood et al., 2008), understanding key influences which are more likely to aid players to return to playing walking football (following enforced 20 21 cessation) is important to investigate.

22

The aims of the present study were to: (a) to investigate differences in influences of walking
football initiation across SES, gender, health conditions and weekly PA, b) to investigate
differences in influences of walking football maintenance across SES, gender, health

conditions and weekly PA, and c) to examine which characteristics, initiation and
 maintenance influences contribute most to the intention to continue playing walking football
 after COVID-19 restrictions ease.

- 4
- 5

Methods

6 **Respondents and recruitment**

7 The reporting of the study methods adheres to the Checklist for Reporting Results of Internet 8 E-Surveys (CHERRIES) guidelines (Eysenbach, 2004). Respondents were 439 older adults 9 (392 male, 47 female) aged between 50-75 years. They were recruited from UK walking 10 football clubs and purposive sampling was applied as part of the process. Recruitment was 11 conducted solely online due to the COVID-19 pandemic, using social media, and contacting 12 clubs via email, to advertise the survey to their players. Further snowball sampling was used, 13 with players being asked at the end of the survey to advertise to other players. To complete 14 the survey, respondents were required to have at least six months' experience playing 15 walking football, to ensure respondents satisfied the minimum term of maintenance according to previous literature (Lally et al., 2010; van Stralen et al., 2009). Due to COVID-16 19 restrictions coming into effect on 23rd March 2020 in the UK, respondents were therefore 17 18 recruited if they had started playing walking football on or before the 23rd of September 19 2019. The respondents were also required to be of UK residence, so postcodes for SES data 20 could be recorded. Respondents were also asked to provide information around their lifetime 21 sport experience (types of sport played, years played and days per week played) and how 22 they became aware of walking football to gather further information regarding their 23 awareness of walking football.

24

25 Sample size

1 G*Power 3 was used to calculate a sample size for the cross-sectional survey (Faul et al., 2 2007), based on a medium effect size ($f^2 = 0.15$), estimated power of 0.90 and alpha level of 3 0.05. Sufficient sample sizes calculated were N=116 for study aims one and two (Hotelling's 4 T2 and MANOVA analyses), and N=89 for aim three (multiple regression analysis), however 5 the research team aimed to recruit at least 300 participants, similar to papers with similar 6 purposes (Ashford et al., 1993; Ashton et al., 2017; Ryu et al., 2018), who used sample sizes 7 of 339, 282, and 153 participants, respectively. A sample size target of 300 was chosen in 8 order to account for dropout and to produce scalable findings.

9

10 Survey development

11 An empirically grounded survey was designed to achieve the study aims. The findings from 12 the previous qualitative studies conducted by the research team (Cholerton et al., 2020, 2021) 13 directly informed the themes and content of the survey. Due to the lack of existing 14 instruments to assess walking football participation, this approach was employed so the new 15 survey was context-specific to walking football, and key concepts from the qualitative phase 16 could be suitably measured within a larger sample. This approach is consistent with similar 17 research in the field of sport and exercise, and health psychology (Gould et al., 2002; Walker 18 et al., 2021; Wallace et al., 2012).

19

The survey development involved three rounds. The first round of development involved forming survey subscales, using the lower order themes contained in each higher order theme from the research team's previous studies (Cholerton et al., 2020, 2021). The second stage of survey development involved adapting previous questions from suitable existing surveys within the PA literature, or developing new questions, observing similar language used in previous questionnaires and surveys. This follows similar research which has used combinations of previously published surveys and developed questions to answer the
research aims (Ashford et al., 1993; Crowther et al., 2017; Hwang et al., 2020; Rice et al.,
2019). The third round of survey development involved formatting questions within the
online survey software Qualtrics (Provo, UT, 2020). Eighteen pages or web screens of items
were shown to participants. The survey was developed so respondents were able to review or
change their answers, via a back button on the survey.

7

8 Measures

9 Independent variables

10 Sixty-six items in total were used in the survey. Two items relating to the respondent's age 11 and length of walking football play were used to ensure respondents were screened correctly. Four items measured respondent characteristics. Gender (male, female, prefer not to say) was 12 13 collected, and SES was measured via Indices of Multiple Deprivation (Ministry of Housing 14 Communities and Local Government, 2019; Northern Ireland Statistical Research Agency, 15 2017; Scottish Government, 2016; StatsWales, 2019). Information on number of health 16 conditions was reported using Potter and colleagues' (2017) long term conditions 17 questionnaire. Weekly PA level measured by a single item questionnaire, developed by 18 Milton and colleagues (Milton et al., 2011). A further three items measured respondents' 19 change in PA since COVID-19 restrictions were implemented, lifelong sport experience, and 20 how respondents became aware of walking football, measured by defined categorical options, 21 as well as an open-ended response for 'Other'.

22

23 Dependent variables

Eighteen items measured influences on walking football initiation and 35 items were used tomeasure influences on walking football maintenance. The items were adapted from a range of

1 surveys, or developed by the researcher based on quotes from the research team's previous 2 qualitative research (Anderson, 2004; Anderson et al., 2016; Barrett-Lennard, 1962; 3 Cholerton et al., 2020; 2021; Devereaux Melillo et al., 1997; Estabrooks & Carron, 2000; 4 Jowett & Ntoumanis, 2004; Markland & Tobin, 2004; Myers et al., 2006; Newton et al., 5 2007; Pelletier et al., 2013). For each statement, respondents were expected to answer each of 6 the items on a 5-point Likert scale, ranging from 'not true of me at all' to 'extremely true of me'. Examples of statements include 'The location of the walking football club was suitable 7 8 for me' and 'I have good friends in this walking football group'. Six items formed the 9 intention to continue section of the survey. Two items were developed by the researcher to 10 ascertain whether clubs had resumed at the time of answering the survey, and if the 11 respondent had returned to walking football. A further two items then measured the intention 12 to play once walking football clubs had reopened, and wording was adapted from a previous 13 survey administered to older adults (Li et al., 2019).

14

15 **Procedure**

16 Following university ethics approval for the study (Ethics ID: ER25811320), a pilot survey 17 was administered to walking football players in August 2020. The pilot survey was 18 conducted with 12 respondents (10 male, 2 female). A further survey section containing 19 questions related to the survey process were asked at the piloting stage. The median 20 completion time for the pilot was 17 minutes. Minor changes were made to the survey after 21 piloting, such as formatting some questions for clarification. This included emboldening text 22 to clarify questions, for example "In the past week, on how many days have you done a total 23 of 30 minutes or more of physical activity, which was enough to raise your breathing rate?". 24 When asked if the time to complete the survey was acceptable, all respondent selected 'yes'.

1 The main survey data collection took place online between September and November 2020. 2 The web link to the survey contained a welcome page with a participation information sheet 3 and consent form for participants to provide informed consent, and the survey was in an open 4 format, meaning that it was not password protected. After completing the survey, respondents 5 were shown a debrief form. Upon completion of the study, respondents were offered the 6 choice to enter a raffle to win one of two gift vouchers, with research suggesting that 7 incentives can aid response rates (Singer & Ye, 2013). The process adhered to university 8 guidelines (Sheffield Hallam University, 2015).

9

10 Data analysis

11 Before data analysis took place, data files were cleaned to identify missing, invalid or 12 insufficient data (Fritchoff Davis, 2012). Duplicated survey responses, identified through IP 13 address and postcode, were removed before analysis. A criterion for including partially 14 completed surveys was devised by the research team prior to analysing the data, to ensure 15 sufficient completion for each section was included, as composite scores were needed for coding and data analysis. Any survey response which did not meet 67% completion 16 17 (participants not completing the initiation questionnaire) was discounted, due to data 18 insufficiency in relation to answering the study aims. Out of 514 returned surveys, 75 survey 19 responses were excluded from analysis due to insufficient response completion, duplication 20 of respondents, or not meeting the inclusion criteria. Four hundred and thirty nine responses 21 were used to analyse descriptive statistics and initiation analysis. Two responses were 22 discounted from the maintenance analysis and a further four responses were discounted from 23 the intention to return to play analysis due to insufficient completion. Gender was assigned 24 (male or female), and SES was determined via converting postcodes into IMD (Index of 25 Multiple Deprivation) rankings, which are used in the UK to measure relative deprivation in

1 small areas (Department for Communities and Local Government, 2015; Northern Ireland 2 Statistical Research Agency, 2017; Scottish Government, 2020; StatsWales, 2019). These 3 ranks range from one to 32844, one being the most deprived rank and 32844 being the least 4 deprived rank. These ranks were then divided into two groups, 'high' and 'low' SES. Number 5 of health conditions were calculated and split into three groups, representing those with no 6 health conditions, those with one health condition, and those with two or more health 7 conditions, the latter group defined as 'multimorbidity' (Barnett et al., 2012; Dhalwani et al., 8 2016). Weekly PA was split into three groups, representing those who did PA on 0-1 days, 9 those who did PA on 2-4 days, and those who did PA on 5+ days (Sport England, 2020a). For 10 the intention to return to play analysis, the responses for perceived change in PA over 11 COVID-19 restrictions were split into three groups, 'more', 'same' or 'less', in line with the 12 Sport England report on COVID-19 behaviours (Sport England, 2020b). 13

14 Cronbach's alpha tests were run for each factor, in order to assess how much the items on the 15 scales devised were measuring the same underlying construct (Field, 2013). Any scales with 16 a reliability score (Cronbach's alpha) over 0.6 was considered acceptable, as when measuring 17 psychological constructs in exploratory studies, scores below 0.7 can be expected due to the 18 diversity of the constructs measured (Field, 2013).

19

For the first and second research aims (differences in initation and maintenance influences between participant characteristics), differences according to gender and SES were examined using Hotelling's T² test, and Bonferroni-adjusted alpha levels were used to correct for the increase in Type I errors from the multiple individual tests (Field, 2013). Differences according to health conditions and weekly PA levels were investigated using a one-way multivariate analysis of variance. Where significant results occurred, post hoc tests (univariate ANOVAs and Tukey's Honest Significant Difference) were run. For the third
research aim (i.e., which variables contributed most to the intention to continue play),
multiple regression analysis was performed and 13 independent variables relating to the
characteristics and influences were examined, with the mean for the two intention to play
items as the dependent variable (Field, 2013). Effect sizes for significant results were
interpreted according to Cohen (1988), using Cohen's d for Hotelling's T and adjusted R²
values for multiple regression.

- 8
- 9

Results

10 Table 1 displays the characteristics of the respondents. The sample consisted of 439 older 11 adults who had played walking football for over six months where the majority of 12 respondents (89.3%) were male compared to female respondents (10.7%). The majority of 13 respondents were categorised as being from high SES areas (67.7%) and a third of 14 respondents were from low SES areas (32.3%). Most respondents were from England 15 (91.3%). The number of health conditions varied; a third of respondents respondents reported one health condition (34.4%) and 22.6 percent of respondents reported two or more health 16 17 conditions, with 43.1 percent of respondents reporting no health conditions. Weekly PA levels also varied, with 46 respondents (10.5%) reporting low active levels (30+ minutes on 18 19 0-1 days per week), 235 participants (53.5%) reporting 2-4 days per week of PA, and 158 20 participants (36%) reporting five or more days a week of activity. Regarding perceived 21 change following the COVID-19 UK restrictions, the majority of respondents recorded they had taken part in 'Less' PA (41.5%), 28 percent of respondents recorded the 'same', and 30.4 22 23 percent of respondents recorded 'more' PA. Regarding lifetime sport experience, the

24 respondents' mean days per week playing sport tended to decrease over the lifespan. When

1	respondents were asked about how they became aware of walking football, being made aware
2	via a friend was the most common amongst the overall sample of respondents (33.7%).
3 4	[Insert Table 1 here]
5	
6	Survey internal consistency
7	For both initiation and maintenance sections of the survey, subscales were deemed acceptable
8	for the purposes of this study (Field, 2013). Table two displays individual Cronbach's Alpha
9	scores for each subscale.
10	
11	[Insert Table 2 here]
12	
13	Main effects of gender, SES, number of health conditions and weekly PA level on
14	walking football initiation influences
15	Descriptive statistics for differences in initiation influences on walking football play for
16	gender, SES, health conditions and weekly PA are shown below (Table 3 and 4). Means,
17	standard deviations and F scores are shown for each dependent variable. Male players rated
18	social influences higher than female players, whereas female players rated psychological
19	influences and walking football environment higher. The differences between gender on the
20	combined dependent variables for initiation (psychological influences, social interactions,
21	and walking football environment) was not statistically significant (F(3, 435) = 1.94 , $p = .12$,
22	Wilks' $\Lambda = .99$; partial $\eta 2 = .01$). When assessing SES, low SES respondents reported
23	psychological influences, social interactions and walking football environment as being more
24	influential on walking football initiation than high SES counterparts. The differences between
25	high and low SES respondents on the combined dependent variables was not statistically
26	significant (F(3, 435) = 1.10, p = .35, Wilks' Λ = .99; partial η 2 = .01).

2 [Insert Table 3 here]

- 3 [Insert Table 4 here]
- 4

5 Regarding weekly PA, those who participated in weekly PA for 5 or more days per week 6 rated the highest on psychological influences and walking football environment, and those 7 engaging in PA for 2-4 days per week rated highest on social interactions. The differences 8 between the level of weekly PA on the combined dependent variables was not statistically 9 significant (F(6, 868) = 1.09, p = .37, Wilks' $\Lambda = .99$; partial $\eta 2 = .01$). When assessing the 10 number of health conditions, those with no health conditions rated all initiation influences 11 lower than those with one health condition, or two or more health conditions. The differences 12 between the number of health conditions on the combined dependent variables was statistically significant (F(6, 868) = 2.56, p = .018; Wilks' Λ = .97; partial η 2 = .02; d = .21). 13 14 Follow-up univariate ANOVAs showed that social interactions were statistically significantly 15 different between respondents with different numbers of health conditions (F= 4.882, p =16 .008), using a Bonferroni adjusted α level of .017. Tukey post-hoc tests found that for the 17 influence of social interactions, respondents with two or more health conditions had 18 statistically significantly higher mean scores than respondents with no health conditions (p = 19 .005). No significant differences were observed for either psychological influences or 20 walking football environment. 21 22 [insert Figure 1 here] 23 24 Main effects of gender, SES, number of health conditions and weekly PA level on

25 walking football maintenance influences

- Descriptive statistics for differences in maintenance influences on walking football play for
 gender, SES, health conditions and weekly PA are shown below. Means, standard deviations
 and F scores are shown for each dependent variable (see tables 5 and 6).
- 4

5 Male players rated psychological influences higher than female players, where female players 6 rated social influences, walking football culture, session specific factors and maintenance 7 resources higher. The differences between male and female players on the combined 8 dependent variables in maintenance (psychological influences, social-level influences, 9 walking football culture, walking football session specific factors, and availability of 10 maintenance resources) was not statistically significant (F(5, 431) = 1.70, p = .13, Wilks' $\Lambda =$ 11 .98; partial $\eta 2 = .02$). When assessing SES, low SES respondents reported psychological 12 influences, social influences, walking football culture and session specific factors as being 13 more influential on walking football maintenance than high SES counterparts, but those from 14 high SES areas rated maintenance resources (e.g., scheduling session, sticking with the 15 decision to attend) as being more of an influence in maintenance than low SES counterparts. 16 The differences between high and low SES respondents on the combined dependent variables was not statistically significant (F(5, 431) = 0.64, p = .67, Wilks' $\Lambda = .99$; partial $\eta 2 = .01$). 17 18

19 [insert table 5 and 6]

20

Regarding weekly PA, those who participated in weekly PA for 5 or more days per week rated the highest on psychological influences, walking football culture, walking football session specific factors and maintenance resources. Those who engaged in weekly PA for 2-4 days per week rated the highest on social influences. The differences between the level of weekly PA on the combined dependent variables was not statistically significant (F(10, 860)

1	= 1.04, p = .41, Wilks' Λ = .98; partial η 2 = .012). When assessing the number of health
2	conditions, those with two or more health conditions rated the highest on social influences,
3	walking football culture and maintenance resources, where those with one health condition
4	rated the highest on psychological influences and session specific factors. The differences
5	between the number of health conditions on the combined dependent variables was
6	statistically significant (F(10, 860) = 2.58, p = .004; Wilks' Λ = .94; partial η 2 = .029;
7	d=0.21). Follow-up univariate ANOVAs showed that social interactions were statistically
8	significantly different between respondents with different numbers of health conditions (F =
9	6.014, $p = .003$), using a Bonferroni adjusted α level of .01. Tukey post-hoc tests showed that
10	for social influences, respondents with two or more health conditions had statistically
11	significantly higher mean scores than respondents with no health conditions ($p = .003$). No
12	significant differences were observed for either psychological influences or walking football
13	environment.

15 [insert Figure 2]

16

17 **Contributors to intention to return to play**

18 For the regression analysis conducted, bivariate correlations and regression coefficients are 19 presented below (see table 7 and 8). The intention to play (variable 1) was significantly and 20 positively associated with one initiation variable, Psychological Influences (3), and all 21 maintenance variables, including Psychological Influences (6), Social Influences (7), 22 Walking Football Culture (8), Walking Football Session Structure (9), and Maintenance Resources (10). The multiple regression model statistically significantly predicted intention 23 24 $(F(13, 419) = 10.94, p < 0.001, adj. R^2 = .23)$. The model accounted for 23% of the variance, which is considered a medium effect size (Cohen, 1988). Two variables from maintenance, 25

1	walking football culture (8), and availability of maintenance resources (10), added
2	statistically significantly to the regression ($p \le .01$). The regression coefficient for walking
3	football culture indicates that a point increase in the walking football culture mean score was
4	associated with an increase of 0.100 in intention to play walking football after COVID-19
5	restrictions had eased. The regression coefficient relating to maintenance resources indicates
6	that a point increase in availability of maintenance resources mean score was associated with
7	an increase of .399, in intention to return to play walking football.
8	
9	[Insert table 7 here]
10	[Insert table 8 here]
11	
12	Discussion
13	This is the first study of its kind to assess influences for initiating and maintaining walking
14	football play in older adults. Furthermore, this study builds on previous literature by
15	assessing influences by group characteristics, including SES, gender, number of health
16	conditions and weekly PA, and which influences contribute most to the intention to play
17	walking football after COVID-19 restrictions eased. This study adds to the behaviour change
18	literature, providing evidence for the presence of different influences at different stages of
19	walking football participation, especially in specific group characteristics.
20	
21	The respondent sample was mostly male players from high SES areas, which represents a
22	similar ratio of male and female players participating in walking football in the general
23	population (Walking Football Association, 2020). This supports previous research
24	highlighting gender differences seen in sport participation, with players more likely to be
25	male (Breuer et al., 2011). Further reports concerning SES also highlights that those from

1 higher SES areas tended to take part in more PA per week (Sport England, 2020a).

2 Percentages of those reporting none, one, and two or more health conditions was in line with 3 previous research, which have examined numbers of health conditions in older adults 4 (Martinez-Gomez et al., 2017). Whilst weekly PA levels within the sample varied, it is 5 important to note that over 40% of respondents perceived a decrease in PA over the COVID-6 19 pandemic, consistent with recent reports from (Sport England, 2021). Lastly, perception of 7 sport participation levels from adolescence to middle age is consistent with previous reports 8 and literature, which have stated a similar decline in both PA and sport participation (McPhee 9 et al., 2016).

10

11 Differences in walking football initiation and maintenance influences across gender,

12 SES, number of health conditions and weekly PA level

13 The significant results surrounding number of health conditions and social influences suggest 14 that those with a higher number of health conditions rate social interactions as being more 15 influential on the initiation of walking football (see figure 1). Regarding SES, gender and 16 weekly PA results, this contradicts previous research into sport participation in older adults, 17 specifically concerning gender differences and SES differences which have previously been 18 seen amongst those participating in older adulthood (Faß & Schlesinger, 2019). Possible 19 explanations for the inconsistency between previous research and this study may lie in the group characteristics, the sport context assessed, and the imbalance between male and female 20 21 respondents. Firstly, within this population, influences of walking football initiation may not 22 be affected by gender or SES level, as it may be in a wider sport participation context (Faß & 23 Schlesinger, 2019). Secondly, walking football is a recently developed adapted sport, and it is 24 possible that sport type and intensity may provide less variation in the influences assessed 25 among these groups. This may be especially prominent when assessing influences by the

level of weekly PA performed, and due to the walking nature of the sport, walking football
 may be more accessible to those of varying PA levels, compared to other sporting contexts
 (Jenkin et al., 2017, 2018).

4

5 Post-hoc tests found that those with two or more reported health conditions rated social 6 interactions higher than those with no health conditions, suggesting that those with more 7 health conditions find social interactions a more important influence upon initiation of the 8 sport. The findings support previous literature, which suggests that social support from both 9 significant others and group members have been considered important in initiation of PA, and 10 sport is important in fostering social connections (Jenkin et al., 2017; van Stralen et al., 11 2009). Furthermore, recent intervention research into walking football with older adults 12 highlights the importance of social interactions amongst other players (Lamont et al., 2017; 13 McEwan et al., 2019; Reddy et al., 2017). This study's results also support qualitative 14 research where participants highlighted the importance of social connections in initiation 15 when managing a health condition, feeling less isolated due to being surrounded with people who have similar health conditions (Cholerton et al., 2020). The current study's findings, 16 17 coupled with previous research, highlights the importance of social interactions during 18 initiation for those with a larger number of health conditions, and further research may 19 consider investigating the use of walking football in improving social connections within 20 those with health conditions. Despite these significant results, however, it is important to 21 consider the small effect size, and in practice, social interactions may be one of many 22 influences involved in initiation for those with a larger number of health conditions. 23

When investigating the second aim of the study - assessing differences in maintenance
influences - no significant results were found between SES, gender and weekly PA levels

1 when investigating maintenance influences. During analysis of the number of health 2 conditions, however, significant differences were found for social influences, and those with 3 two or more reported health conditions rated social influences higher than those with no 4 health conditions. This, as with initiation, suggests that those with more health conditions 5 find social influences more important during maintenance of walking football. This also 6 supports previous walking football research which found social connections were important 7 for those managing mental health conditions (Lamont et al., 2017). In a PA context, social 8 support from group members has been found to have a positive effect on maintenance in 9 older adults (van Stralen et al., 2009). As the findings suggest that those with more health 10 conditions tend to find social interactions as more important than those without, 11 considerations should be made as to how peer support can be utilised for those who are 12 managing health conditions whilst playing walking football.

13

14 For the third aim of the study, multiple regression results indicate that there are two 15 contributors to the intention to return to play once COVID-19 restrictions had eased; positive 16 walking football culture (e.g., players respecting each other's abilities), and higher use of 17 maintenance resources (e.g., behaviours, such as scheduling in sessions on a weekly basis). 18 The results suggest that the more positive the perceived walking football culture, the more 19 likely players intend to return to walking football. Results also suggest that the more 20 maintenance resources a player perceives themselves to use, the more likely they are to 21 intend to play after COVID-19 restrictions ease. Little research has directly assessed the 22 importance of a caring climate in older adults, within a sport context, however, results 23 regarding a positive walking football culture and 'caring climate' have been found in youth 24 sport participation, with caring climates having an effect on motivational outcomes and 25 enjoyment of the sport (Gerabinis et al., 2018). Furthermore, previous research has

1 highlighted the importance of maintenance resources (e.g., action planning, coping planning) 2 in long-term behaviour change in a general population (Kwasnicka et al., 2016; Peels et al., 3 2020). Whilst direct parallels cannot be drawn to other breaks in play such as injury or 4 holidays, the ceasing of walking football sessions due to COVID-19 restrictions provides a 5 similar context where there was a forced break in play for all walking football players in the 6 UK. It is important to note that whilst no differences in gender, SES, number of health 7 conditions and weekly PA were seen as influencing intention to play in these data, 8 characteristics may need to be controlled for in future research. This may provide further 9 understanding into factors that predict intention to play after a forced break.

10

11 Implications for future research

12 This study can help inform future research within the walking football context and help to 13 examine the mechanisms involved in long-term behaviour change within this population. The 14 research highlights the importance of social connections on those with health conditions in 15 both initiation and maintenance phases, as well as a positive walking football culture and use of maintenance resources (e.g., scheduling sessions, redefining PA expectations) on the 16 17 intention to continue after a forced break. Future research may consider generating hypotheses relating to the differences in influences between characteristics, and investigating 18 19 the types of social influences which are most likely to aid initiation and maintenance in 20 walking football players. Furthermore, assessing whether type of health condition (e.g. either 21 mental or physical health conditions) affects influences at initiation or maintenance may be 22 considered, to further support for specific health conditions.

23

Refining the survey questions used in this survey and running psychometric analysis (e.g.,
factor analysis) would assess the reliability of the constructs further. Research focusing on

1 the effects of a positive walking football culture on older adult sport maintenance may be 2 considered, to further research based on the multiple regression results. The same applies to 3 the use of maintenance resources, and further research including investigation of the types of 4 maintenance resources used with older adult populations may be useful in understanding the 5 mechanisms of maintenance in more detail, so older adults can be adequately supported in 6 positive PA behaviour change. Research exploring the use of maintenance resources, and 7 assessing fit with theories such as Self Determination Theory (Ryan & Deci, 2000), and 8 frameworks such as COM-B and the behaviour change wheel (Michie et al., 2011) may be 9 warranted, so robust intervention methodology can be developed to investigate the use of 10 maintenance resources and the effect on initiation and maintenance of walking football. 11 Techniques effective in positively changing behaviour can then be isolated, tested and 12 encouraged among players.

13

14 Implications for policy and practice

15 The findings can inform future policy surrounding walking football, and initiatives to support 16 older players in maintaining the sport. Policies around peer support systems, for example 17 buddy schemes (pairing a new player with an experienced player), could be implemented, as 18 highlighted in previous research as having an effect on PA levels in older adults (Lindsay 19 Smith et al., 2017) and also may be particularly beneficial to those living with health 20 conditions. In some cases, providing funding to set up clubs for those who manage certain 21 health conditions may allow players to benefit from appropriate levels of play, and enhanced 22 peer support. It is also important to consider the effects of a caring climate and a positive 23 culture in the intention to continue playing walking football, which seems to be evident in 24 older adults playing the sport. Previous research within organisational psychology has 25 highlighted that positive perceptions of an organisation's culture has been linked with

employee retention (Anitha & Begum, 2016). This may be similar when assessing
commitment and success in sporting contexts (Wagstaff & Burton-Wylie, 2018). Local
Authorities, National Governing Bodies and coaches may consider encouraging a caring
culture through marketing campaigns conveying messaging around understanding players'
ability levels, and introducing a code fostering an inclusive playing environment, to
encourage positive club culture.

7

8 Practice implications include considerations for the coach and individual clubs to consider. 9 Within engagement and maintenance, this study highlights the need to understand the social 10 requirements of those with health conditions. As other studies have highlighted the positive 11 effects of social connections on PA participation in older adults (Maula et al., 2019), 12 encouraging social events and sessions with those who have similar health conditions may 13 promote longer-term engagement in the sport for this population. Furthermore, the COVID-14 19 pandemic has provided a unique environment to analyse PA patterns of older adults when 15 going through a forced break from structured PA sessions. Overall, this study found that selfreported PA levels did not rise (70.5% reporting the 'same' or 'less' PA) during COVID-19 16 17 restrictions being implemented. This is in line with previous research stating a negligible rise 18 in self-reported PA levels (Richardson et al., 2021). In light of previous research stating 19 importance of understanding the lapse and relapse in maintenance of PA (Kahn et al., 2002), 20 this research highlights the need for a positive sporting culture, and supporting players to 21 increase maintenance resources. Regarding club practice, tailoring club environments (e.g. 22 appropriate warm ups and sessions with reduced intensity) and actively fostering a positive 23 culture (e.g. through fair refereeing and accepting all abilities into the club), may provide 24 opportunities for long-term continuation in walking football. Regarding coach practice, 25 training courses focusing on the key principles of creating a caring and positive culture,

1 alongside working with players to use maintenance resources, is important to consider. 2 Previous research highlights the importance of coping strategies on a positive exercise 3 outcome in long-term exercisers (Stetson et al., 2005), therefore providing tangible 4 maintenance support to players may aid long-term continuation, for example by encouraging 5 players to schedule sessions in advance and encouraging players to attend, despite adverse 6 weather. Extra training for coaches in understanding behaviour change and how to encourage 7 players to continue walking football may be beneficial, to support players in maintaining the 8 sport.

9

10 Strengths and limitations

11 This is the first known survey assessing influences of walking football initiation and 12 maintenance in older adults. This work strengthens the current literature by providing a 13 context-specific analysis of influences on behaviour change in a specific age group. The 14 online survey was developed using the qualitative data from the research team's previous 15 qualitative research (Cholerton et al., 2020, 2021), assessing multiple influences that were 16 discussed by participants directly. The qualitative work provided direct participant 17 experiences that were a base for designing this study, and as limited research has explored 18 experiences in this group and sport, provides a unique survey to assess behaviour change 19 influences in walking football. Another strength of the study includes responses from a large 20 sample (n=439) of walking football players between 50-75 years old, and sample size 21 calculations suggest that the sample is adequately powered. Nevertheless, the sample of 22 female players was low. Despite the sample being, to the researcher's knowledge, 23 representative of the known population of walking football players in the UK (Walking 24 Football Association, 2020), further research may look to increase the sample of female 25 players in analyses, which may yield different findings. Furthermore, the respondents were

1 overwhelmingly based in England, compared with Wales, Northern Ireland and Scotland, 2 despite researchers contacting organisations from all countries. Further representation of 50-3 75-year-old players in different countries within the UK could be achieved by working 4 closely with national walking football organisations to promote research, and increasing 5 accessibility to the survey, (e.g., providing bothonline and paper surveys) to improve 6 response rates in this age group across the UK. It is also important to note the cross-sectional 7 nature of the survey, assessing those who are actively maintaining, and also dichotomised 8 SES scores may affect the detail of the SES data collected. Therefore, longitudinal or 9 intervention research would be useful to conduct, as well as performing analysis with a wider 10 range of SES levels to determine any smaller differences seen in this characteristic. Self-11 reporting and retrospective recall were also expected of the respondents, especially due to 12 walking football clubs not playing due to the COVID-19 pandemic during survey 13 distribution. Future research should attempt to survey those that are currently initiating the 14 sport, in order to capture live initiation data.

15

16 Conclusion

17 This cross-sectional survey is the first to assess initiation and maintenance influences, across 18 respondent characteristics, in 50-75 year old adults playing walking football. The survey 19 results revealed that those with two or more health conditions rated social interactions as 20 being more important to both initiation and maintenance, than those with no health 21 conditions. Furthermore, walking football culture and the availability of maintenance 22 resources were found to be significant contributors related to the intention to continue 23 playing. Walking football clubs and bodies should consider the importance of delivering a 24 positive and inclusive walking football culture. Further consideration should be taken 25 concerning the encouragement of players using maintenance strategies to continue play, as

- 1 well as facilitating social connections in those with health conditions. Further research is
- 2 needed to examine which factors in the walking football environment contribute to a higher
- 3 motivation to maintain the sport, and test these within an intervention setting.

1	References
2	Almeida, O. P., Khan, K. M., Hankey, G. J., Yeap, B. B., Golledge, J., & Flicker, L. (2014).
3	150 minutes of vigorous physical activity per week predicts survival and successful
4	ageing: a population-based 11-year longitudinal study of 12 201 older Australian
5	men. British Journal of Sports Medicine, 48(3), 220–225.
6	https://doi.org/10.1136/bjsports-2013-092814
7	Anderson. (2004). Athletic Identity and Its Relation to Exercise Behavior: Scale
8	Development and Initial Validation. Journal of Sport and Exercise Psychology, 26(1),
9	39-56. https://doi.org/10.1123/jsep.26.1.39
10	Anderson, Bovard, R. S., Wang, Z., Beebe, T. J., & Murad, M. H. (2016). A survey of social
11	support for exercise and its relationship to health behaviours and health status among
12	endurance Nordic skiers. BMJ Open, 6(6), 1-6. https://doi.org/10.1136/bmjopen-
13	2015-010259
14	Anitha, J., & Begum, N. F. (2016). Role of Organisational Culture and Employee
15	Commitment in Employee Retention. ASBM Journal of Management, 9(1), 17-28.
16	Ashford, B., Biddle, S., & Goudas, M. (1993). Participation in community sports centres:
17	Motives and predictors of enjoyment. Journal of Sports Sciences, 11(3), 249-256.
18	https://doi.org/10.1080/02640419308729992
19	Ashton, L. M., Hutchesson, M. J., Rollo, M. E., Morgan, P. J., & Collins, C. E. (2017).
20	Motivators and Barriers to Engaging in Healthy Eating and Physical Activity: A
21	Cross-Sectional Survey in Young Adult Men. American Journal of Men's Health,
22	11(2), 330-343. https://doi.org/10.1177/1557988316680936
23	Barnett, K., Mercer, S. W., Norbury, M., Watt, G., Wyke, S., & Guthrie, B. (2012).
24	Epidemiology of multimorbidity and implications for health care, research, and

1	medical education: A cross-sectional study. The Lancet, 380(9836), 37-43.
2	https://doi.org/10.1016/S0140-6736(12)60240-2
3	Barrett-Lennard, G. T. (1962). Dimensions of therapist response as causal factors in
4	therapeutic change. Psychological Monographs: General and Applied, 76(43), 1–36.
5	https://doi.org/10.1037/h0093918
6	Breuer, C., Hallmann, K., & Wicker, P. (2011). Determinants of sport participation in
7	different sports. Managing Leisure, 16(4), 269-286.
8	https://doi.org/10.1080/13606719.2011.613625
9	Chapman, S. B., Aslan, S., Spence, J. S., DeFina, L. F., Keebler, M. W., Didehbani, N., &
10	Lu, H. (2013). Shorter term aerobic exercise improves brain, cognition, and
11	cardiovascular fitness in aging. Frontiers in Aging Neuroscience, 5(NOV), 1-9.
12	https://doi.org/10.3389/fnagi.2013.00075
13	Cholerton, R., Breckon, J., Butt, J., & Quirk, H. (2020). Experiences Influencing Walking
14	Football Initiation in 55- to 75-Year-Old Adults: A Qualitative Study. Journal of
15	Aging and Physical Activity, 28(4), 521-533. https://doi.org/10.1123/japa.2019-0123
16	Cholerton, R., Quirk, H., Breckon, J., & Butt, J. (2021). Experiences and Strategies
17	Influencing Older Adults to Continue Playing Walking Football. Journal of Aging
18	and Physical Activity, AOP, 1-13. https://doi.org/10.1123/japa.2020-0058
19	Cohen, J. (1988). Statistical power analysis for the behavioral sciences (2nd ed.). Routledge.
20	Crowther, F., Sealey, R., Crowe, M., Edwards, A., & Halson, S. (2017). Team sport athletes'
21	perceptions and use of recovery strategies: A mixed-methods survey study. BMC
22	Sports Science, Medicine and Rehabilitation, 9(6), 1–10.
23	https://doi.org/10.1186/s13102-017-0071-3

1	Department for Communities and Local Government. (2015). The English indices of
2	deprivation 2015 statistical release.
3	https://dclgapps.communities.gov.uk/imd/idmap.html
4	Devereaux Melillo, K., Williamson, E., Futrell, M., & Chamberlain, C. (1997). A self-
5	assessment tool to measure older adults' perceptions regarding physical fitness and
6	exercise activity. Journal of Advanced Nursing, 25(6), 1220-1226.
7	https://doi.org/10.1046/j.1365-2648.1997.19970251220.x
8	Dhalwani, N. N., O'Donovan, G., Zaccardi, F., Hamer, M., Yates, T., Davies, M., & Khunti,
9	K. (2016). Long terms trends of multimorbidity and association with physical activity
10	in older English population. International Journal of Behavioral Nutrition and
11	Physical Activity, 13(1), 1-9. https://doi.org/10.1186/s12966-016-0330-9
12	Estabrooks, P. A., & Carron, A. V. (2000). The Physical Activity Group Environment
13	Questionnaire: An instrument for the assessment of cohesion in exercise classes.
14	Group Dynamics, 4(3), 230-243. https://doi.org/10.1037/1089-2699.4.3.230
15	Eysenbach, G. (2004). Improving the quality of web surveys: The Checklist for Reporting
16	Results of Internet E-Surveys (CHERRIES). Journal of Medical Internet Research,
17	6(3), 1–6. https://doi.org/10.2196/jmir.6.3.e34
18	Faß, E., & Schlesinger, T. (2019). The role of individual resources, health behaviour and age
19	perception as determinants of sports participation in older age. Ageing and Society, 1-
20	27. https://doi.org/10.1017/S0144686X19001260
21	Faul, F., Erdfelder, E., Lang, A., & Buchner, A. (2007). G*Power 3: A flexible statistical
22	power analysis program for the social, behavioral, and biomedical sciences. Behavior
23	Research Methods, 39(2), 175–191. https://doi.org/10.3758/BF03193146
24	Field, A. (2013). Discovering Statistics Using IBM SPSS Statistics (A. Field (ed.); 4th ed.).
25	SAGE Publications.

1	Fox, K. R., Hillsdon, M., Sharp, D., Cooper, A. R., Coulson, J. C., Davis, M., Harris, R.,
2	McKenna, J., Narici, M., Stathi, A., & Thompson, J. L. (2011). Neighbourhood
3	deprivation and physical activity in UK older adults. Health and Place, 17(2), 633-
4	640. https://doi.org/10.1016/j.healthplace.2011.01.002
5	Fritchoff Davis, M. (2012). Data Cleaning. In N. J. Salkind (Ed.), Encyclopedia of Research
6	Design (pp. 326–328). SAGE Publications Ltd.
7	https://doi.org/10.2307/j.ctv16755wv.14
8	Gayman, A. M., Fraser-Thomas, J., Spinney, J. E. L., Stone, R. C., & Baker, J. (2017).
9	Leisure-time Physical Activity and Sedentary Behaviour in Older People: The
10	Influence of Sport Involvement on Behaviour Patterns in Later Life. AIMS Public
11	Health, 4(2), 171–188. https://doi.org/10.3934/publichealth.2017.2.171
12	Gerabinis, P., Hatzigeorgiadis, A., Theodorakis, Y., & Goudas, M. (2018). Sport Climate ,
13	Developmental Experiences and Motivational Outcomes in Youth Sport. Journal of
14	Education and Human Development, 7(3), 58–65.
15	https://doi.org/10.15640/jehd.v7n3a7
16	Gould, D., Guinan, D., Greenleaf, C., & Chung, Y. (2002). A survey of U.S. Olympic
17	coaches: Variables perceived to have influenced athlete performances and coach
18	effectiveness. Sport Psychologist, 16(3), 229–250.
19	https://doi.org/10.1123/tsp.16.3.229
20	Hirvensalo, M., & Lintunen, T. (2011). Life-course perspective for physical activity and
21	sports participation. European Review of Aging and Physical Activity, 8(1), 13-22.
22	https://doi.org/10.1007/s11556-010-0076-3
23	Hwang, J., Li, W., Stough, L. M., Lee, C., & Turnbull, K. (2020). People with disabilities'
24	perceptions of autonomous vehicles as a viable transportation option to improve
25	mobility: An exploratory study using mixed methods. International Journal of

2	https://doi.org/10.1080/15568318.2020.1833115
3	Jenkin, C. R., Eime, R. M., Westerbeek, H., O'Sullivan, G., & Van Uffelen, J. G. Z. (2017).
4	Sport and ageing: A systematic review of the determinants and trends of participation
5	in sport for older adults. BMC Public Health, 17(1). https://doi.org/10.1186/s12889-
6	017-4970-8
7	Jenkin, C. R., Eime, R. M., Westerbeek, H., & Van Uffelen, J. G. Z. (2018). Sport for adults
8	aged 50+ years: Participation benefits and barriers. Journal of Aging and Physical
9	Activity, 26(3), 363-371. https://doi.org/10.1123/japa.2017-0092
10	Jowett, S., & Ntoumanis, N. (2004). The Coach–Athlete Relationship Questionnaire (CART-
11	Q): development and initial validation. Scandinavian Journal of Medicine and
12	Science in Sports, 14, 245-257. https://doi.org/10.1046/j.1600-0838.2003.00338.x
13	Kahn, E. B., Ramsey, L. T., Brownson, R. C., Heath, G. W., Howze, E. H., Powell, K. E.,
14	Stone, E. J., Rajab, M. W., & Corso, P. (2002). The effectiveness of interventions to
15	increase physical activity: A systematic review. American Journal of Preventive
16	Medicine, 22(4 SUPPL. 1), 73-107. https://doi.org/10.1016/S0749-3797(02)00434-8
17	Kamphuis, C. B. M., van Lenthe, F. J., Giskes, K., Huisman, M., Brug, J., & Mackenbach, J.
18	P. (2009). Socioeconomic differences in lack of recreational walking among older
19	adults: The role of neighbourhood and individual factors. International Journal of
20	Behavioral Nutrition and Physical Activity, 6, 1–11. https://doi.org/10.1186/1479-
21	5868-6-1
22	Kendrick, D., Orton, E., Lafond, N., Audsley, S., Maula, A., Morris, R., Vedhara, K., &
23	Iliffe, S. (2018). Keeping active: maintenance of physical activity after exercise
24	programmes for older adults. Public Health, 164, 118-127.
25	https://doi.org/10.1016/j.puhe.2018.08.003

1	Kwasnicka, D., Dombrowski, S. U., White, M., & Sniehotta, F. (2016). Theoretical
2	explanations for maintenance of behavior change : a systematic review of behavior
3	theories Theoretical explanations for maintenance of behaviour change : a systematic
4	review of behaviour theories. Health Psychology Review, 10(3), 277–296.
5	https://doi.org/10.1080/17437199.2016.1151372
6	Laitakari, J., Vuori, I., & Oja, P. (1996). Is long-term maintenance of health-related physical
7	activity possible? An analysis of concepts and evidence. Health Education Research,
8	11(4), 463–477. https://doi.org/10.1093/her/11.4.463
9	Lally, P., Van Jaarsveld, C. H. M., Potts, H. W. W., & Wardle, J. (2010). How are habits
10	formed: Modelling habit formation in the real world. European Journal of Social
11	Psychology, 40(6), 998-1009. https://doi.org/10.1002/ejsp.674
12	Lamont, E., Harris, J., McDonald, G., Kerin, T., & Dickens, G. L. (2017). Qualitative
13	investigation of the role of collaborative football and walking football groups in
14	mental health recovery. Mental Health and Physical Activity, 12, 116-123.
15	https://doi.org/10.1016/j.mhpa.2017.03.003
16	Li, J., Hsu, C. C., & Lin, C. T. (2019). Leisure participation behavior and psychological well-
17	being of elderly adults: An empirical study of Tai Chi Chuan in China. International
18	Journal of Environmental Research and Public Health, 16(18).
19	https://doi.org/10.3390/ijerph16183387
20	Lindsay Smith, G., Banting, L., Eime, R., O'Sullivan, G., & van Uffelen, J. G. Z. (2017). The
21	association between social support and physical activity in older adults: A systematic
22	review. International Journal of Behavioral Nutrition and Physical Activity, 14(1), 1-
23	21. https://doi.org/10.1186/s12966-017-0509-8

1	Liu-Ambrose, T. (2010). Resistance Training and Executive Functions: A 12-Month
2	Randomized Controlled Trial. Archives of Internal Medicine, 170(2), 170.
3	https://doi.org/10.1001/archinternmed.2009.494
4	Lloyd, M. (2019). Walking football "has helped save lives." https://www.bbc.co.uk/news/uk-
5	wales-50267287
6	Markland, D., & Tobin, V. (2004). A modification to the behavioural regulation in exercise
7	questionnaire to include an assessment of amotivation. Journal of Sport and Exercise
8	Psychology, 26(2), 191-196. https://doi.org/10.1123/jsep.26.2.191
9	Martinez-Gomez, D., Guallar-Castillon, P., Garcia-Esquinas, E., Bandinelli, S., & Rodríguez-
10	Artalejo, F. (2017). Physical Activity and the Effect of Multimorbidity on All-Cause
11	Mortality in Older Adults. Mayo Clinic Proceedings, 92(3), 376–382.
12	https://doi.org/10.1016/j.mayocp.2016.12.004
13	Maula, A., LaFond, N., Orton, E., Iliffe, S., Audsley, S., Vedhara, K., & Kendrick, D. (2019).
14	Use it or lose it: a qualitative study of the maintenance of physical activity in older
15	adults. BMC Geriatrics, 19(1), 349-349. https://doi.org/10.1186/s12877-019-1366-x
16	McEwan, G., Buchan, D., Cowan, D., Arthur, R., Sanderson, M., & Macrae, E. (2019).
17	Recruiting Older Men to Walking Football: A Pilot Feasibility Study. Explore, 15(3),
18	206–214. https://doi.org/10.1016/j.explore.2018.12.001
19	McPhee, J. S., French, D. P., Jackson, D., Nazroo, J., Pendleton, N., & Degens, H. (2016).
20	Physical activity in older age: perspectives for healthy ageing and frailty.
21	Biogerontology, 17(3), 567-580. https://doi.org/10.1007/s10522-016-9641-0
22	Michie, S., van Stralen, M. M., & West, R. (2011). The behaviour change wheel: A new
23	method for characterising and designing behaviour change interventions.
24	Implementation Science, 6(1), 42-42. https://doi.org/10.1186/1748-5908-6-42

1	Milton, K., Bull, F. C., & Bauman, A. (2011). Reliability and validity testing of a single-item
2	physical activity measure. British Journal of Sports Medicine, 45(3), 203–208.
3	https://doi.org/10.1136/bjsm.2009.068395
4	Ministry of Housing Communities and Local Government. (2019). English indices of
5	deprivation 2019. https://www.gov.uk/government/statistics/english-indices-of-
6	deprivation-2019
7	Murtagh, E. M., Murphy, M. H., Murphy, N. M., & Woods, C. (2015). Prevalence and
8	Correlates of Physical Inactivity in Community-Dwelling Older Adults in Ireland.
9	PLoS ONE, 10(2), 1–11. https://doi.org/10.1371/journal.pone.0118293
10	Myers, N. D., Feltz, D. L., Maier, K. S., Wolfe, E. W., & Reckase, M. D. (2006). Athletes'
11	evaluations of their head coach's coaching competency. Research Quarterly for
12	<i>Exercise and Sport</i> , 77(1), 111–121.
13	https://doi.org/10.1080/02701367.2006.10599337
14	Newton, M., Fry, M., Watson, D., Gano-Overway, L., Kim, MS., Magyar, M., &
15	Guivernau, M. (2007). Psychometric properties of the caring climate scale in a
16	physical activity setting. Revista de Psicología Del Deporte, 16(1), 67-84.
17	Northern Ireland Statistical Research Agency. (2017). Northern Ireland Multiple Deprivation
18	Measure 2017 (NIMDM2017).
19	https://www.nisra.gov.uk/statistics/deprivation/northern-ireland-multiple-deprivation-
20	measure-2017-nimdm2017
21	Peels, D. A., Verboon, P., van Stralen, M. M., Bolman, C., Golsteijn, R. H. J., Mudde, A. N.,
22	de Vries, H., & Lechner, L. (2020). Motivational factors for initiating and maintaining
23	physical activity among adults aged over fifty targeted by a tailored intervention.
24	Psychology and Health, 35(10), 1184–1206.
25	https://doi.org/10.1080/08870446.2020.1734202

1	Pelletier, L. G., Rocchi, M. A., Vallerand, R. J., Deci, E. L., & Ryan, R. M. (2013).
2	Validation of the revised sport motivation scale (SMS-II). Psychology of Sport and
3	Exercise, 14(3), 329-341. https://doi.org/10.1016/j.psychsport.2012.12.002
4	Pesce, C., & Audiffren, M. (2011). Does acute exercise switch off switch costs? A study with
5	younger and older athletes. Journal of Sport and Exercise Psychology, 33(5), 609-
6	626. https://doi.org/10.1123/jsep.33.5.609
7	Potter, C. M., Batchelder, L., A'Court, C., Geneen, L., Kelly, L., Fox, D., Baker, M.,
8	Bostock, J., Coulter, A., Fitzpatrick, R., Forder, J. E., Gibbons, E., Jenkinson, C.,
9	Jones, K., & Peters, M. (2017). Long-Term Conditions Questionnaire (LTCQ): Initial
10	validation survey among primary care patients and social care recipients in England.
11	BMJ Open, 7(11), 1-12. https://doi.org/10.1136/bmjopen-2017-019235
12	Reddy, P., Dias, I., Holland, C., Campbell, N., Nagar, I., Connolly, L., Krustrup, P., &
13	Hubball, H. (2017). Walking football as sustainable exercise for older adults-A pilot
14	investigation. European Journal of Sport Science, 17(5), 638-645.
15	https://doi.org/10.1080/17461391.2017.1298671
16	Rezende, L. F. M. d., Rey-López, J. P., Matsudo, V. K, & Carmo Luiz, O (2014).
17	Sedentary behavior and health outcomes among older adults: a systematic review.
18	BMC Public Health, 14, 333-333. https://doi.org/10.1186/1471-2458-14-333
19	Rice, W. S., Turan, B., Fletcher, F. E., Nápoles, T. M., Walcott, M., Batchelder, A., Kempf,
20	M. C., Konkle-Parker, D. J., Wilson, T. E., Tien, P. C., Wingood, G. M., Neilands, T.
21	B., Johnson, M. O., Weiser, S. D., & Turan, J. M. (2019). A Mixed Methods Study of
22	Anticipated and Experienced Stigma in Health Care Settings among Women Living
23	with HIV in the United States. AIDS Patient Care and STDs, 33(4), 184–195.
24	https://doi.org/10.1089/apc.2018.0282

1	Richardson, D. L., Duncan, M. J., Clarke, N. D., Myers, T. D., & Tallis, J. (2021). The
2	influence of COVID-19 measures in the United Kingdom on physical activity levels,
3	perceived physical function and mood in older adults: A survey-based observational
4	study. Journal of Sports Sciences, 39(8), 887-899.
5	https://doi.org/10.1080/02640414.2020.1850984
6	Rothman, A. J. (2000). Toward a theory-based analysis of behavioral maintenance. Health
7	Psychology, 19(1 SUPPL.), 64-69. https://doi.org/10.1037//0278-6133.19.suppl1.64
8	Ryan, R. M., & Deci, E. L. (2000). Self-determination theory and the facilitation of intrinsic
9	motivation, social development, and well-being. American Psychologist, 55(1), 68-
10	78. https://doi.org/10.1037/0003-066X.55.1.68
11	Ryu, J., Yang, H., Kim, A. C. H., Kim, K. M., & Heo, J. (2018). Understanding pickleball as
12	a new leisure pursuit among older adults. Educational Gerontology, 44(2-3), 128-
13	138. https://doi.org/10.1080/03601277.2018.1424507
14	Scottish Government. (2016). Scottish Index of Multiple Deprivation (SIMD) 2016.
15	https://data.gov.uk/dataset/a448dd2a-9197-4ea0-8357-c2c9b3c29591/scottish-index-
16	of-multiple-deprivation-simd-2016
17	Scottish Government. (2020). Scottish index of multiple deprivation 2020.
18	https://www.gov.scot/collections/scottish-index-of-multiple-deprivation-2020/
19	Sheffield Hallam University. (2015). Incentives to Research Participants.
20	https://www.shu.ac.uk/~/media/home/research/files/ethics/03-research-
21	incentives.pdf?la=en
22	Sherwood, N. E., Martinson, B. C., Crain, A. L., Hayes, M. G., Pronk, N. P., & O'Connor, P.
23	J. (2008). A new approach to physical activity maintenance: Rationale, design, and
24	baseline data from the Keep Active Minnesota trial. BMC Geriatrics, 8, 1–10.
25	https://doi.org/10.1186/1471-2318-8-17

1	Singer, E., & Ye, C. (2013). The Use and Effects of Incentives in Surveys. The Annals of the
2	American Academy of Political and Social Science, 645, 112–141.
3	Sport England. (2018). Active Lives Adult Survey (pp. 8-8).
4	https://www.sportengland.org/media/13530/spotlight-on-older-adults.pdf
5	Sport England. (2020a). Active Lives Adult Survey May 19/20 Report (Issue October, pp. 1-
6	30). https://sportengland-production-files.s3.eu-west-2.amazonaws.com/s3fs-
7	public/2020-10/Active Lives Adult May 19-20
8	Report.pdf?AYzBswpBmlh9cNcH8TFctPI38v4Ok2JD
9	Sport England. (2020b). COVID-19 Briefing: Exploring attitudes and behaviours in England
10	during the COVID-19 pandemic (pp. 98-98). https://www.sportengland.org/know-
11	your-audience/demographic-knowledge/coronavirus#the_story_so_far
12	Sport England. (2021). Understanding the impact of of Covid-19 (pp. 1–28).
13	StatsWales. (2019). WIMD 2019. https://statswales.gov.wales/Catalogue/Community-Safety-
14	and-Social-Inclusion/Welsh-Index-of-Multiple-Deprivation/WIMD-2019#:~:text=The
15	Welsh Index of Multiple, several different types of deprivation.
16	Stenner, B. J., Buckley, J. D., & Mosewich, A. D. (2020). Reasons why older adults play
17	sport: A systematic review. Journal of Sport and Health Science.
18	https://doi.org/10.1016/j.jshs.2019.11.003
19	Stetson, B. A., Beacham, A. O., Frommelt, S. J., Boutelle, K. N., Cole, J. D., Ziegler, C. H.,
20	& Looney, S. W. (2005). Exercise slips in high-risk situations and activity patterns in
21	long-term exercisers: An application of the relapse prevention model. Annals of
22	Behavioral Medicine, 30(1), 25-35. https://doi.org/10.1207/s15324796abm3001_4
23	The FA. (2020). Grassroots football in England has been suspended.
24	https://www.thefa.com/news/2020/mar/16/grassroots-football-suspended-160320

1	van Stralen, M., de Vries, H., Mudde, A. N., Bolman, C., & Lechner, L. (2009). Determinants
2	of initiation and maintenance of physical activity among older adults: A literature
3	review. Health Psychology Review, 3(2), 147–207.
4	https://doi.org/10.1080/17437190903229462
5	van Stralen, M., Lechner, L., Mudde, A. N., De Vries, H., & Bolman, C. (2010).
6	Determinants of awareness, initiation and maintenance of physical activity among the
7	over-fifties: A Delphi study. Health Education Research, 25(2), 233-247.
8	https://doi.org/10.1093/her/cyn045
9	Wagstaff, C. R. D., & Burton-Wylie, S. (2018). Organizational culture in sport: A
10	conceptual, definitional, and methodological review. Sport and Exercise Psychology
11	Review, 14(2), 32–52.
12	Walker, R., Limbert, C., & Smith, P. M. (2021). Exploring the Perceived Barriers and
13	Benefits of Physical Activity Among Wounded, Injured, and/or Sick Military
14	Veterans. Journal of Social, Behavioral, and Health Sciences, 15(1).
15	https://doi.org/10.5590/JSBHS.2021.15.1.11
16	Walking Football Association. (2020). The state of the game: Player and club survey (Issue
17	September, pp. 1–16). https://thewfa.co.uk/walking-football-survey-september-2020/
18	Wallace, S., Clark, M., & White, J. (2012). "It's on my iPhone": Attitudes to the use of
19	mobile computing devices in medical education, a mixed-methods study. BMJ Open,
20	2(4), 1–7. https://doi.org/10.1136/bmjopen-2012-001099
21	
22	

Table 1

Respondent characteristics.

Participant Characteristic	Classification	Frequency	Percent
Gender	Male	392	89.3
Gender	Female	47	10.7
	50-54	21	4.8
	55-59	72	16.4
Age Range	60-64	133	30.3
	65-69	131	29.8
	70-75	82	18.7
SES	High	297	67.7
525	Low	142	32.3
	England	401	91.3
	Scotland	35	8.0
UK country of residence	Wales	1	0.2
	NI	2	0.5
	0 HCs	189	43.1
Health Conditions	1 HC	151	34.4
	2+ HCs	99	22.6
	0-1 day	46	10.5
Weekly PA (days)	2-4 days	235	53.5
weekiy FA (uays)	5+ days	158	36.0
	Less	182	41.5
	Same	123	28.0
Weekly PA (days) erceived change in PA COVID- 19	More	134	30.5
	Local flyer/poster	57	13.0
	Local/national newspaper	50	11.4
	Half time during a game	4	0.9
Awareness of walking football	Social media	56	12.8
	Radio	2	0.5
	Friend told me about the sessions	148	33.7
	Other	122	27.8
		Mean	SD
Mean years playing s	oort between ages 12-18	5.58	1.18
Mean days per week playi	ng sport between ages 12-18	3.68	1.58
Mean years playing s	oort between ages 19-34	12.44	3.92
Mean days per week playi	ng sport between ages 19-34	2.86	1.32
Mean years playing s	oort between ages 35-54	13.97	6.01
Mean days per week plavi	ng sport between ages 35-54	2.47	1.30

Table 2 *Cronbac*

2 Cronbach's Alpha scores for survey subscales.

Subscale	Cronbach's Alpha (α) score
Initiation Variables	
Psychological Influences	0.84
Social interactions	0.69
Walking football environment	0.65
Maintenance Variables	
Psychological influences	0.81
Social-level influences	0.88
Walking football culture	0.81
Valking football session specific factors	0.70
Availability of maintenance resources	0.71
Intention to play	0.81

7 Descriptive statistics on initiation influences across gender and SES.

Initiation Influences			Gender			SES							
	Ма	le	Female			Hig	h	Lo	w				
	Mean	SD	Mean	SD	F	Mean	SD	Mean	SD	F			
Psychological Influences	3.69	0.74	3.74	0.87	0.18	3.67	0.78	3.75	0.70	1.26			
Social Interactions	3.23	0.85	3.17	0.82	0.20	3.17	0.86	3.32	0.79	2.97			
Walking football environment	3.97	0.70	4.16	0.68	3.28	3.98	0.71	4.02	0.70	0.37			

*p<0.05

8

9 Table 4

10 Descriptive statistics on initiation influences across number of health conditions and weekly PA level.

Initiation Influences			Healt	h cond	itions			Weekly PA								
	0 HCs		1 HC		2+ HCs			0-1 Days		2-4 Days		5+ Days				
	Mean	SD	Mean	SD	Mean	SD	F	Mean	SD	Mean	SD	Mean	SD	F		
Psychological Influences	3.64	0.86	3.77	0.66	3.70	0.67	1.25	3.53	0.70	3.67	0.72	3.78	0.81	2.11		
Social Interactions	3.11	0.84	3.23	0.84	3.43	0.82	4.88*	3.21	0.80	3.23	0.83	3.21	0.89	0.42		
Walking football environment	3.95	0.71	4.05	0.73	3.98	0.65	0.84	3.86	0.75	4.00	0.70	4.02	0.70	0.86		

*p<0.05

11

17 Descriptive statistics on maintenance influences across gender and SES.

Maintenance Influences			Gender		SES						
	Ма	le	Fem	ale		High		Low			
	Mean	SD	Mean	SD	F	Mean	SD	Mean	SD	F	
Psychological Influences	4.24	0.59	4.18	0.75	0.41	4.21	0.63	4.27	0.57	0.73	
Social Influences	3.09	0.86	3.19	0.82	0.55	3.07	0.87	3.16	0.83	0.87	
Walking football culture	4.20	0.83	4.46	0.79	3.89	4.24	0.85	4.21	0.77	0.09	
Walking football session specific factors	3.89	0.75	3.97	0.76	0.45	3.88	0.75	3.94	0.74	0.70	
Maintenance resources	4.44	0.52	4.53	0.56	0.27	4.45	0.53	4.45	0.51	0.25	

^{*}p<0.05

21 Descriptive statistics on maintenance influences across number of health conditions and weekly PA level.

Maintenance Influences			Health	1 condi	tions			Weekly PA								
	0 HCs		1 HC		2+ HCs			0-1 Days		2-4 Days		5+ Days				
	Mean	SD	Mean	SD	Mean	SD	F	Mean	SD	Mean	SD	Mean	SD	F		
Psychological Influences	4.20	0.66	4.29	0.53	4.20	0.62	1.24	4.11	0.64	4.21	0.60	4.29	0.61	1.80		
Social Influences	2.95	0.82	3.15	0.88	3.30	0.83	6.01*	3.04	0.82	3.11	0.83	3.10	0.90	0.13		
Walking football culture	4.24	0.83	4.19	0.87	4.26	0.76	0.26	4.03	0.83	4.24	0.79	4.26	0.87	1.45		
Walking football session specific factors	3.87	0.79	3.96	0.71	3.87	0.73	0.72	3.69	0.84	3.90	0.73	3.96	0.75	2.31		
Maintenance resources	4.43	0.54	4.45	0.50	4.49	0.53	0.37	4.44	0.52	4.44	0.51	4.47	0.56	0.10		

*p<0.05

Table 6

25 Multiple regression coefficients and standard errors for intention to play.

Intention to play			onfidence al for B		Standardized Coefficients	R²	ΔR^2
	В	Lower Bound	Upper Bound	Std. Error	β		
Model						0.25	0.23
(Constant)	3.28	2.77	3.80	0.20			
Gender	-0.06	-0.22	0.09	0.06	-0.05		
SES	-0.04	-0.14	0.06	0.04	-0.04		
Number of health conditions	-0.01	-0.08	0.05	0.02	-0.03		
Weekly PA	0.00	-0.07	0.08	0.03	0.00		
Perceived change in PA over COVID	0.04	-0.02	0.10	0.02	0.08		
Initiation: Psychological Influences	-0.07	-0.17	0.02	0.04	-0.13		
Initiation: Social Interactions	-0.05	-0.13	0.03	0.03	-0.10		
Initiation: Walking Football environment	-0.06	-0.15	0.04	0.04	-0.09		
Maintenance: Psychological influences	0.02	-0.10	0.14	0.05	0.03		
Maintenance: Social influences	0.02	-0.06	0.10	0.03	0.03		
Maintenance: Walking football culture	0.10***	0.03	0.17	0.03	0.20***		
Maintenance: Walking football session specific factors	-0.01	-0.10	0.08	0.03	-0.02		
Maintenance: Maintenance resources	0.40***	0.29	0.51	0.04	0.50***		

Note: Model = 'Enter' method in SPSS statistics; B = unstandardized regression coefficient; CI = confidence interval; R² = coefficient of determination; ΔR^2 = adjusted R². ***p<0.001.

29 Descriptive statistics and bivariate correlations among study variables.

	Mean	SD	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1. Intention to Play	4.86	0.42	1													
2. Gender 3. SES	1.11 1.32	0.31 0.47	-0.01 -0.07	1 -0.01	1											
4. Number of health conditions 5. Weekly PA	1.80 2.26	0.79 0.63	-0.04 0.04	-0.04 -0.05	0.137** -0.04	1 -0.09	1									
6. Perceived change in PA over COVID-19 7. Initiation - Psychological	1.89	0.84	0.09	0.05	-0.04	-0.10	0.25	1								
influences 8. Initiation - Social interactions	3.70 3.22	0.76 0.84	0.13** 0.05	0.02 -0.02	0.06 0.09	0.04 0.143**	0.11 0.01	0.02 -0.10	1 0.47**	1						
9. Initiation -Walking football environment 10. Maintenance - Psychological	3.99	0.71	0.09	0.09	0.04	0.02	0.06	-0.01	0.46**	0.51**	1					
influences	4.23	0.61	0.216**	-0.03	0.04	0.02	0.09	-0.03	0.71**	0.43**	0.45**	1				
11. Maintenance - Social influences 12. Maintenance - Walking football	3.10	0.86	0.129**	0.04	0.05	0.17	0.02	-0.01	0.45**	0.63**	0.50**	0.43**	1			
culture	4.23	0.83	0.233**	0.10	-0.01	0.00	0.06	0.00	0.34**	0.40**	0.56**	0.37**	0.50**	1		
13. Maintenance - WF session specific factors 14. Maintenance - Maintenance	3.90	0.75	0.14**	0.03	0.04	0.01	0.09	0.06	0.47**	0.44**	0.64**	0.46**	0.56**	0.48**	1	
Resources	4.46	0.52	0.439**	0.05	-0.01	0.04	0.03	-0.01	0.53**	0.34**	0.36**	0.59**	0.35**	0.32**	0.40**	1

**p<0.01