



This is a repository copy of *Low exposure long-baseline neutrino oscillation sensitivity of the DUNE experiment*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/185511/>

Version: Submitted Version

Article:

Collaboration, DUNE, Abud, AA, Abi, B et al. (1154 more authors) (Submitted: 2021) Low exposure long-baseline neutrino oscillation sensitivity of the DUNE experiment. arXiv. (Submitted)

© 2021 The Authors. For reuse permissions, please contact the Author(s).

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Low exposure long-baseline neutrino oscillation sensitivity of the DUNE experiment

A. Abed Abud,^{126,25} B. Abi,¹⁵² R. Acciarri,⁶⁶ M. A. Acero,¹¹ M. R. Adames,¹⁸⁷ G. Adamov,⁷¹
D. Adams,²¹ M. Adinolfi,²⁰ A. Aduszkiewicz,⁸⁰ J. Aguilar,¹²⁵ Z. Ahmad,¹⁹⁸ J. Ahmed,²⁰¹ B. Aimard,⁵²
B. Ali-Mohammadzadeh,^{86,37} T. Alion,¹⁸⁵ K. Allison,⁴⁵ S. Alonso Monsalve,^{25,58} M. AlRashed,¹¹⁹ C. Alt,⁵⁸
A. Alton,¹² P. Amedo,⁸⁴ J. Anderson,⁷ C. Andreopoulos,^{172,126} M. Andreotti,^{87,67} M. P. Andrews,⁶⁶ F. Andrianala,⁵
S. Andringa,¹²³ N. Anfimov,¹¹¹ A. Ankowski,¹⁷³ M. Antoniaassi,¹⁸⁷ M. Antonova,⁸³ A. Antoshkin,¹¹¹ S. Antusch,¹⁴
A. Aranda-Fernandez,⁴⁴ L. O. Arnold,⁴⁷ M. A. Arroyave,⁵⁷ J. Asaadi,¹⁹⁰ L. Asquith,¹⁸⁵ A. Aurisano,⁴²
V. Aushiev,¹²² D. Autiero,⁹⁷ M. Ayala-Torres,⁴³ F. Azfar,¹⁵² A. Back,¹⁰⁵ H. Back,¹⁵³ J. J. Back,²⁰¹ C. Backhouse,¹⁹⁶
I. Bagaturia,⁷¹ L. Bagby,⁶⁶ N. Balashov,¹¹¹ S. Balasubramanian,⁶⁶ P. Baldi,³⁰ B. Baller,⁶⁶ B. Bambah,⁸¹
F. Barao,^{123,99} G. Barenboim,⁸³ G. J. Barker,²⁰¹ W. Barkhouse,¹⁴⁵ C. Barnes,¹³⁵ G. Barr,¹⁵² J. Barranco
Monarca,⁷⁶ A. Barros,¹⁸⁷ N. Barros,^{123,60} J. L. Barrow,¹³² A. Basharina-Freshville,¹⁹⁶ A. Bashyal,⁷ V. Basque,¹³¹
E. Belchior,³⁶ J.B.R. Battat,²⁰² F. Battisti,¹⁵² F. Bay,⁴ J. L. Bazo Alba,¹⁶⁴ J. F. Beacom,¹⁵⁰ E. Bechetoille,⁹⁷
B. Behera,⁴⁶ L. Bellantoni,⁶⁶ G. Bellettini,¹⁶² V. Bellini,^{86,37} O. Beltramello,²⁵ N. Benekos,²⁵ C. Benitez
Montiel,⁹ F. Bento Neves,¹²³ J. Berger,⁴⁶ S. Berkman,⁶⁶ P. Bernardini,^{89,175} R. M. Berner,¹⁵ S. Bertolucci,^{85,18}
M. Betancourt,⁶⁶ A. Betancur Rodríguez,⁵⁷ A. Bevan,¹⁶⁷ Y. Bezawada,²⁹ T.J.C. Bezerra,¹⁸⁵ A. Bhardwaj,¹²⁸
V. Bhatnagar,¹⁵⁵ M. Bhattacharjee,¹⁰³ S. Bhuller,²⁰ B. Bhuyan,¹⁰³ S. Biagi,⁹⁵ J. Bian,³⁰ M. Biassoni,⁹⁰
K. Biery,⁶⁶ B. Bilki,^{16,108} M. Bishai,²¹ A. Bitadze,¹³¹ A. Blake,¹²⁴ F. D. M. Blaszczyk,⁶⁶ G. C. Blazey,¹⁴⁶
E. Blucher,⁴⁰ J. Boissevain,¹²⁷ S. Bolognesi,²⁴ T. Bolton,¹¹⁹ L. Bomben,^{90,107} M. Bonesini,^{90,137} M. Bongrand,¹⁵⁶
C. Bonilla-Diaz,³⁸ F. Bonini,²¹ A. Booth,¹⁶⁷ F. Boran,¹⁶ S. Bordini,²⁵ A. Borkum,¹⁸⁵ N. Bostan,¹⁴⁸ P. Bour,⁴⁹
C. Bourgeois,¹⁵⁶ D. Boyden,¹⁴⁶ J. Bracinik,¹⁷ D. Braga,⁶⁶ D. Brailsford,¹²⁴ A. Branca,⁹⁰ A. Brandt,¹⁹⁰ J. Bremer,²⁵
C. Brew,¹⁷² S. J. Brice,⁶⁶ C. Brizzolari,^{90,137} C. Bromberg,¹³⁶ J. Brooke,²⁰ A. Bross,⁶⁶ G. Brunetti,^{90,137}
M. Brunetti,²⁰¹ N. Buchanan,⁴⁶ H. Budd,¹⁶⁹ I. Butorov,¹¹¹ I. Cagnoli,^{85,18} D. Caiulo,⁹⁷ R. Calabrese,^{87,67}
P. Calafiura,¹²⁵ J. Calcutt,¹³⁶ M. Calin,²² S. Calvez,⁴⁶ E. Calvo,²⁶ A. Caminata,⁸⁸ M. Campanelli,¹⁹⁶ D. Caratelli,⁶⁶
G. Carini,²¹ B. Carlus,⁹⁷ M. F. Carneiro,²¹ P. Carniti,⁹⁰ I. Caro Terrazas,⁴⁶ H. Carranza,¹⁹⁰ T. Carroll,²⁰⁵
J. F. Castaño Forero,⁶ A. Castillo,¹⁷⁷ C. Castromonte,¹⁰⁶ E. Catano-Mur,²⁰⁴ C. Cattadori,⁹⁰ F. Cavalier,¹⁵⁶
F. Cavanna,⁶⁶ S. Centro,¹⁵⁴ G. Cerati,⁶⁶ A. Cervelli,⁸⁵ A. Cervera Villanueva,⁸³ M. Chalifour,²⁵ A. Chappell,²⁰¹
E. Chardonnet,¹⁵⁷ N. Charitonidis,²⁵ A. Chatterjee,¹⁶³ S. Chattopadhyay,¹⁹⁸ H. Chen,²¹ M. Chen,³⁰ Y. Chen,¹⁵
Z. Chen,¹⁸³ Y. Cheon,¹⁹⁴ D. Cherdack,⁸⁰ C. Chi,⁴⁷ S. Childress,⁶⁶ A. Chiriacescu,²² G. Chisnall,¹⁸⁵ K. Cho,¹¹⁷
S. Choate,¹⁴⁶ D. Chokheli,⁷¹ P. S. Chong,¹⁵⁹ A. Christensen,⁴⁶ D. Christian,⁶⁶ G. Christodoulou,²⁵ A. Chukanov,¹¹¹
M. Chung,¹⁹⁴ E. Church,¹⁵³ V. Cicero,^{85,18} P. Clarke,⁵⁹ T. E. Coan,¹⁸² A. G. Cocco,⁹² J. A. B. Coelho,¹⁵⁷
N. Colton,⁴⁶ E. Conley,⁵⁵ R. Conley,¹⁷³ J. M. Conrad,¹³² M. Convery,¹⁷³ S. Copello,⁸⁸ L. Cremaldi,¹⁴¹
L. Cremonesi,¹⁶⁷ J. I. Crespo-Anadón,²⁶ M. Crisler,⁶⁶ E. Cristaldo,⁹ R. Cross,¹²⁴ A. Cudd,⁴⁵ C. Cuesta,²⁶ Y. Cui,³²
D. Cussans,²⁰ O. Dalager,³⁰ H. da Motta,²³ L. Da Silva Peres,⁶⁵ C. David,^{208,66} Q. David,⁹⁷ G. S. Davies,¹⁴¹
S. Davini,⁸⁸ J. Dawson,¹⁵⁷ K. De,¹⁹⁰ P. Debbs,¹⁰⁸ I. De Bonis,⁵² M. P. Decowski,^{144,3} A. de Gouvêa,¹⁴⁷ P. C. De
Holanda,³⁶ I. L. De Icaza Astiz,¹⁸⁵ A. Deisting,¹⁷⁰ P. De Jong,^{144,3} A. Delbart,²⁴ D. Delepine,⁷⁶ M. Delgado,⁶
A. Dell'Acqua,²⁵ P. De Lurgio,⁷ J. R. T. de Mello Neto,⁶⁵ D. M. DeMuth,¹⁹⁷ S. Dennis,³⁵ C. Densham,¹⁷²
G. W. Deptuch,²¹ A. De Roeck,²⁵ V. De Romeri,⁸³ G. De Souza,³⁶ R. Devi,¹¹² R. Dharmapalan,⁷⁹ M. Dias,¹⁹⁵
F. Diaz,¹⁶⁴ J. S. Díaz,¹⁰⁵ S. Di Domizio,^{88,70} L. Di Giulio,²⁵ P. Ding,⁶⁶ L. Di Noto,^{88,70} C. Distefano,⁹⁵
R. Diurba,¹⁴⁰ M. Diwan,²¹ Z. Djurcic,⁷ D. Doering,¹⁷³ S. Dolan,²⁵ F. Dolek,¹⁶ M. J. Dolinski,⁵⁴ L. Domine,¹⁷³
D. Douglas,¹³⁶ D. Douillet,¹⁵⁶ G. Drake,⁶⁶ F. Drielsma,¹⁷³ L. Duarte,¹⁹⁵ D. Duchesneau,⁵² K. Duffy,⁶⁶
P. Dunne,¹⁰² H. Duyang,¹⁸¹ O. Dvornikov,⁷⁹ D. A. Dwyer,¹²⁵ A. S. Dyshkant,¹⁴⁶ M. Eads,¹⁴⁶ A. Earle,¹⁸⁵
D. Edmunds,¹³⁶ J. Eisch,⁶⁶ L. Emberger,^{131,133} S. Emery,²⁴ A. Ereditato,²⁰⁶ T. Erjavec,²⁹ C. O. Escobar,⁶⁶
G. Eurin,²⁴ J. J. Evans,¹³¹ E. Ewart,¹⁰⁵ A. C. Ezeribe,¹⁷⁸ K. Fahey,⁶⁶ A. Falcone,^{90,137} M. Fani,¹²⁷ C. Farnese,⁹³
Y. Farzan,⁹⁸ D. Fedoseev,¹¹¹ J. Felix,⁷⁶ Y. Feng,¹⁰⁹ E. Fernandez-Martinez,¹³⁰ P. Fernandez Menendez,⁸³
M. Fernandez Morales,⁸⁴ F. Ferraro,^{88,70} L. Fields,¹⁴⁸ P. Filip,⁴⁸ F. Filthaut,^{144,168} A. Fiorentini,¹⁷⁹ M. Fiorini,^{87,67}
R. S. Fitzpatrick,¹³⁵ W. Flanagan,⁵¹ B. Fleming,²⁰⁶ R. Flight,¹⁶⁹ S. Fogarty,⁴⁶ W. Foreman,¹⁰¹ D. V. Forero,¹³⁴
J. Fowler,⁵⁵ W. Fox,¹⁰⁵ J. Franc,⁴⁹ K. Francis,¹⁴⁶ D. Franco,²⁰⁶ J. Freeman,⁶⁶ J. Freestone,¹³¹ J. Fried,²¹
A. Friedland,¹⁷³ F. Fuentes Robayo,²⁰ S. Fuess,⁶⁶ I. K. Furic,⁶⁸ A. P. Furmanski,¹⁴⁰ A. Gabrielli,⁸⁵ A. Gago,¹⁶⁴
H. Gallagher,¹⁹³ A. Gallas,¹⁵⁶ A. Gallego-Ros,²⁶ N. Gallice,^{91,138} V. Galymov,⁹⁷ E. Gamberini,²⁵ T. Gamble,¹⁷⁸
F. Ganacim,¹⁸⁷ R. Gandhi,⁷⁷ R. Gandrajula,¹³⁶ F. Gao,¹⁶³ S. Gao,²¹ D. Garcia-Gamez,⁷⁴ M. Á. García-Peris,⁸³
S. Gardiner,⁶⁶ D. Gastler,¹⁹ J. Gauvreau,¹⁴⁹ G. Ge,⁴⁷ N. Geffroy,⁵² B. Gelli,³⁶ A. Gendotti,⁵⁸ S. Gent,¹⁸⁰
Z. Ghorbani-Moghaddam,⁸⁸ P. Giammaria,³⁶ T. Giammaria,^{87,67} D. Gibin,¹⁵⁴ I. Gil-Botella,²⁶ S. Gilligan,¹⁵¹
C. Girerd,⁹⁷ A. K. Giri,¹⁰⁴ D. Gnani,¹²⁵ O. Gogota,¹²² M. Gold,¹⁴² S. Gollapinni,¹²⁷ K. Gollwitzer,⁶⁶ R. A. Gomes,⁶²

L. V. Gomez Bermeo,¹⁷⁷ L. S. Gomez Fajardo,¹⁷⁷ F. Gonnella,¹⁷ J. A. Gonzalez-Cuevas,⁹ D. Gonzalez-Diaz,⁸⁴
M. Gonzalez-Lopez,¹³⁰ M. C. Goodman,⁷ O. Goodwin,¹³¹ S. Goswami,¹⁶¹ C. Gotti,⁹⁰ E. Goudzovski,¹⁷ C. Grace,¹²⁵
R. Gran,¹³⁹ E. Granados,⁷⁶ P. Granger,²⁴ A. Grant,⁵³ C. Grant,¹⁹ D. Gratieri,⁶⁹ P. Green,¹³¹ L. Greenler,²⁰⁵
J. Greer,²⁰ J. Grenard,²⁵ W. C. Griffith,¹⁸⁵ M. Groh,⁴⁶ J. Grudzinski,⁷ K. Grzelak,²⁰⁰ W. Gu,²¹ E. Guardincerri,¹²⁷
V. Guarino,⁷ M. Guarise,^{87, 67} R. Guenette,⁷⁸ E. Guerard,¹⁵⁶ M. Guerzoni,⁸⁵ D. Guffanti,⁹¹ A. Guglielmi,⁹³
B. Guo,¹⁸¹ V. Gupta,¹⁴⁴ K. K. Guthikonda,¹¹⁸ R. Gutierrez,⁶ P. Guzowski,¹³¹ M. M. Guzzo,³⁶ S. Gwon,⁴¹ C. Ha,⁴¹
A. Habig,¹³⁹ H. Hadavand,¹⁹⁰ R. Haenni,¹⁵ A. Hahn,⁶⁶ J. Haiston,¹⁷⁹ P. Hamacher-Baumann,¹⁵² T. Hamernik,⁶⁶
P. Hamilton,¹⁰² J. Han,¹⁶³ D. A. Harris,^{208, 66} J. Hartnell,¹⁸⁵ T. Hartnett,¹⁷² J. Harton,⁴⁶ T. Hasegawa,¹¹⁶
C. Hasnip,¹⁵² R. Hatcher,⁶⁶ K. W. Hatfield,³⁰ A. Hatzikoutelis,¹⁷⁶ C. Hayes,¹⁰⁵ K. Hayrapetyan,¹⁶⁷ J. Hays,¹⁶⁷
E. Hazen,¹⁹ M. He,⁸⁰ A. Heavey,⁶⁶ K. M. Heeger,²⁰⁶ J. Heise,¹⁷⁴ S. Henry,¹⁶⁹ M. A. Hernandez Morquecho,¹⁰¹
K. Herner,⁶⁶ J. Hewes,⁴² T. Hill,¹⁰⁰ S. J. Hillier,¹⁷ A. Himmel,⁶⁶ E. Hinkle,⁴⁰ L.R. Hirsch,¹⁸⁷ J. Ho,⁷⁸ J. Hoff,⁶⁶
A. Holin,¹⁷² E. Hoppe,¹⁵³ G. A. Horton-Smith,¹¹⁹ M. Hostert,¹⁴⁰ A. Hourlier,¹³² B. Howard,⁶⁶ R. Howell,¹⁶⁹
I. Hristova,¹⁷² M. S. Hronek,⁶⁶ J. Huang,²⁹ G. Iles,¹⁰² N. Ilic,¹⁹² A. M. Ilescu,⁸⁵ R. Illingworth,⁶⁶ G. Ingratta,^{85, 18}
A. Ioannisian,²⁰⁷ B. Irwin,¹⁴⁰ L. Isenhower,¹ R. Itay,¹⁷³ C.M. Jackson,¹⁵³ V. Jain,² E. James,⁶⁶ W. Jang,¹⁹⁰
B. Jargowsky,³⁰ F. Jediny,⁴⁹ D. Jena,⁶⁶ Y. S. Jeong,^{41, 108} C. Jesús-Valls,⁸² X. Ji,²¹ L. Jiang,¹⁹⁹ S. Jiménez,²⁶
A. Jipa,²² R. Johnson,⁴² N. Johnston,¹⁰⁵ B. Jones,¹⁹⁰ S. B. Jones,¹⁹⁶ M. Judah,¹⁶³ C. K. Jung,¹⁸³ T. Junk,⁶⁶
Y. Jwa,⁴⁷ M. Kabirnezhad,¹⁵² A. Kaboth,^{170, 172} I. Kadenko,¹²² D. Kaira,⁴⁷ I. Kakorin,¹¹¹ A. Kalitkina,¹¹¹
F. Kamiya,⁶⁴ N. Kaneshige,³³ G. Karagiorgi,⁴⁷ G. Karaman,¹⁰⁸ A. Karcher,¹²⁵ M. Karolak,²⁴ Y. Karyotakis,⁵²
S. Kasai,¹²¹ S. P. Kasetti,¹²⁸ L. Kashur,⁴⁶ N. Kazaryan,²⁰⁷ E. Kearns,¹⁹ P. Keener,¹⁵⁹ K.J. Kelly,⁶⁶ E. Kemp,³⁶
O. Kemularia,⁷¹ W. Ketchum,⁶⁶ S. H. Kettell,²¹ M. Khabibullin,⁹⁶ A. Khotjantsev,⁹⁶ A. Khvedelidze,⁷¹ D. Kim,¹⁸⁸
B. King,⁶⁶ B. Kirby,⁴⁷ M. Kirby,⁶⁶ J. Klein,¹⁵⁹ K. Koehler,²⁰⁵ L. W. Koerner,⁸⁰ D. H. Koh,¹⁷³ S. Kohn,^{28, 125}
P. P. Koller,¹⁵ L. Kolupaeva,¹¹¹ D. Korablev,¹¹¹ M. Kordosky,²⁰⁴ T. Kosc,⁹⁷ U. Kose,²⁵ V. A. Kostelecký,¹⁰⁵
K. Kothekar,²⁰ L. Kreczko,²⁰ F. Krennrich,¹⁰⁹ I. Kreslo,¹⁵ W. Kropp,³⁰ Y. Kudenko,⁹⁶ V. A. Kudryavtsev,¹⁷⁸
S. Kulagin,⁹⁶ J. Kumar,⁷⁹ P. Kumar,¹⁷⁸ P. Kunze,⁵² N. Kurita,¹⁷³ C. Kuruppu,¹⁸¹ V. Kus,⁴⁹ T. Kutter,¹²⁸
J. Kvasnicka,⁴⁸ D. Kwak,¹⁹⁴ A. Lambert,¹²⁵ B. J. Land,¹⁵⁹ C. E. Lane,⁵⁴ K. Lang,¹⁹¹ T. Langford,²⁰⁶
M. Langstaff,¹³¹ J. Larkin,²¹ P. Lasorak,¹⁸⁵ D. Last,¹⁵⁹ C. Lastoria,²⁶ A. Laudrie,²⁰⁵ G. Laurenti,⁸⁵ A. Lawrence,¹²⁵
I. Lazanu,²² R. LaZur,⁴⁶ M. Lazzaroni,^{91, 138} T. Le,¹⁹³ S. Leardini,⁸⁴ J. Learned,⁷⁹ P. LeBrun,⁹⁷ T. LeCompte,⁷
C. Lee,⁶⁶ S. Y. Lee,¹¹⁴ G. Lehmann Miotto,²⁵ R. Lehnert,¹⁰⁵ M. A. Leigui de Oliveira,⁶⁴ M. Leitner,¹²⁵
L. M. Lepin,¹³¹ S. W. Li,¹⁷³ T. Li,⁵⁹ Y. Li,²¹ H. Liao,¹¹⁹ C. S. Lin,¹²⁵ Q. Lin,¹⁷³ S. Lin,¹²⁸ R. A. Lineros,³⁸
J. Ling,¹⁸⁴ A. Lister,²⁰⁵ B. R. Littlejohn,¹⁰¹ J. Liu,³⁰ S. Lockwitz,⁶⁶ T. Loew,¹²⁵ M. Lokajicek,⁴⁸ I. Lomidze,⁷¹
K. Long,¹⁰² T. Lord,²⁰¹ J. M. LoSecco,¹⁴⁸ W. C. Louis,¹²⁷ X.-G. Lu,¹⁵² K.B. Luk,^{28, 125} B. Lunday,¹⁵⁹ X. Luo,³³
E. Luppi,^{87, 67} T. Lux,⁸² V. P. Luzio,⁶⁴ D. MacFarlane,¹⁷³ A. A. Machado,³⁶ P. Machado,⁶⁶ C. T. Macias,¹⁰⁵
J. R. Macier,⁶⁶ A. Maddalena,⁷³ A. Madera,²⁵ P. Madigan,^{28, 125} S. Magill,⁷ K. Mahn,¹³⁶ A. Maio,^{123, 60}
A. Major,⁵⁵ J. A. Maloney,⁵⁰ G. Mandrioli,⁸⁵ R. C. Mandujano,³⁰ J. Maneira,^{123, 60} L. Manenti,¹⁹⁶ S. Manly,¹⁶⁹
A. Mann,¹⁹³ K. Manolopoulos,¹⁷² M. Manrique Plata,¹⁰⁵ V. N. Manyam,²¹ L. Manzanillas,¹⁵⁶ M. Marchan,⁶⁶
A. Marchionni,⁶⁶ W. Marciano,²¹ D. Marfatia,⁷⁹ C. Mariani,¹⁹⁹ J. Maricic,⁷⁹ R. Marie,¹⁵⁶ F. Marinho,⁶³
A. D. Marino,⁴⁵ D. Marsden,¹³¹ M. Marshak,¹⁴⁰ C. M. Marshall,¹⁶⁹ J. Marshall,²⁰¹ J. Marteau,⁹⁷ J. Martin-Albo,⁸³
N. Martinez,¹¹⁹ D.A. Martinez Caicedo,¹⁷⁹ P. Martínez Miravé,⁸³ S. Martynenko,¹⁸³ V. Mascagna,^{90, 107}
K. Mason,¹⁹³ A. Mastbaum,¹⁷¹ F. Matichard,¹²⁵ S. Matsuno,⁷⁹ J. Matthews,¹²⁸ C. Mauger,¹⁵⁹ N. Mauri,^{85, 18}
K. Mavrokoridis,¹²⁶ I. Mawby,²⁰¹ R. Mazza,⁹⁰ A. Mazzacane,⁶⁶ E. Mazzucato,²⁴ T. McAskill,²⁰² E. McCluskey,⁶⁶
N. McConkey,¹³¹ K. S. McFarland,¹⁶⁹ C. McGrew,¹⁸³ A. McNab,¹³¹ A. Mefodiev,⁹⁶ P. Mehta,¹¹³ P. Melas,¹⁰
O. Mena,⁸³ H. Mendez,¹⁶⁵ P. Mendez,²⁵ D. P. Méndez,²¹ A. Menegolli,^{94, 158} G. Meng,⁹³ M. D. Messier,¹⁰⁵
W. Metcalf,¹²⁸ T. Mettler,¹⁵ M. Mewes,¹⁰⁵ H. Meyer,²⁰³ T. Miao,⁶⁶ G. Michna,¹⁸⁰ T. Miedema,^{144, 168} V. Mikola,¹⁹⁶
R. Milincic,⁷⁹ G. Miller,¹³¹ W. Miller,¹⁴⁰ J. Mills,¹⁹³ C. Milne,¹⁰⁰ O. Mineev,⁹⁶ A. Minotti,^{91, 137} O. G. Miranda,⁴³
S. Miryala,²¹ C. S. Mishra,⁶⁶ S. R. Mishra,¹⁸¹ A. Mislivec,¹⁴⁰ D. Mladenov,²⁵ I. Mocioiu,¹⁶⁰ K. Moffat,⁵⁶
N. Moggi,^{85, 18} R. Mohanta,⁸¹ T. A. Mohayai,⁶⁶ N. Mokhov,⁶⁶ J. Molina,⁹ L. Molina Bueno,⁸³ E. Montagna,^{85, 18}
A. Montanari,⁸⁵ C. Montanari,^{94, 66, 158} D. Montanari,⁶⁶ L. M. Montano Zetina,⁴³ J. Moon,¹³² S. H. Moon,¹⁹⁴
M. Mooney,⁴⁶ A. F. Moor,³⁵ D. Moreno,⁶ C. Morris,⁸⁰ C. Mossey,⁶⁶ E. Motuk,¹⁹⁶ C. A. Moura,⁶⁴ J. Mousseau,¹³⁵
G. Moustier,¹²⁴ W. Mu,⁶⁶ L. Mualem,³⁴ J. Mueller,⁴⁶ M. Muether,²⁰³ S. Mufson,¹⁰⁵ F. Muheim,⁵⁹ A. Muir,⁵³
M. Mulhearn,²⁹ D. Munford,⁸⁰ H. Muramatsu,¹⁴⁰ S. Murphy,⁵⁸ J. Musser,¹⁰⁵ J. Nachtman,¹⁰⁸ S. Nagu,¹²⁹
M. Nalbandyan,²⁰⁷ R. Nandakumar,¹⁷² D. Naples,¹⁶³ S. Narita,¹¹⁰ A. Nath,¹⁰³ A. Navrer-Agasson,¹³¹ N. Nayak,³⁰
M. Nebot-Guinot,⁵⁹ K. Negishi,¹¹⁰ J. K. Nelson,²⁰⁴ J. Nesbit,²⁰⁵ M. Nessi,²⁵ D. Newbold,¹⁷² M. Newcomer,¹⁵⁹
D. Newhart,⁶⁶ H. Newton,⁵³ R. Nichol,¹⁹⁶ F. Nicolas-Arnaldos,⁷⁴ E. Niner,⁶⁶ K. Nishimura,⁷⁹ A. Norman,⁶⁶

A. Norrick,⁶⁶ R. Northrop,⁴⁰ P. Novella,⁸³ J. A. Nowak,¹²⁴ M. Oberling,⁷ J. P. Ochoa-Ricoux,³⁰ A. Olivier,¹⁶⁹
 A. Olshevskiy,¹¹¹ Y. Onel,¹⁰⁸ Y. Onishchuk,¹²² J. Ott,³⁰ L. Pagani,²⁹ S. Pakvasa,⁷⁹ G. Palacio,⁵⁷ O. Palamara,⁶⁶
 S. Palestini,²⁵ J. M. Paley,⁶⁶ M. Pallavicini,^{88,70} C. Palomares,²⁶ J. L. Palomino-Gallo,¹⁸³ W. Panduro Vazquez,¹⁷⁰
 E. Pantic,²⁹ V. Paolone,¹⁶³ V. Papadimitriou,⁶⁶ R. Papaleo,⁹⁵ A. Papanestis,¹⁷² S. Paramesvaran,²⁰ S. Parke,⁶⁶
 E. Parozzi,^{90,137} Z. Parsa,²¹ M. Parvu,²² S. Pascoli,^{56,18} L. Pasqualini,^{85,18} J. Pasternak,¹⁰² J. Pater,¹³¹
 C. Patrick,¹⁹⁶ L. Patrizii,⁸⁵ R. B. Patterson,³⁴ S. J. Patton,¹²⁵ T. Patzak,¹⁵⁷ A. Paudel,⁶⁶ B. Paulos,²⁰⁵
 L. Paulucci,⁶⁴ Z. Pavlovic,⁶⁶ G. Pawloski,¹⁴⁰ D. Payne,¹²⁶ V. Pec,¹⁷⁸ S. J. M. Peeters,¹⁸⁵ E. Pennacchio,⁹⁷
 A. Penzo,¹⁰⁸ O. L. G. Peres,³⁶ J. Perry,⁵⁹ D. Pershey,⁵⁵ G. Pessina,⁹⁰ G. Petrillo,¹⁷³ C. Petta,^{86,37}
 R. Petti,¹⁸¹ V. Pia,^{85,18} F. Piastra,¹⁵ L. Pickering,¹³⁶ F. Pietropaolo,^{25,93} R. Plunkett,⁶⁶ R. Poling,¹⁴⁰
 X. Pons,²⁵ N. Poonthottathil,¹⁰⁹ F. Poppi,^{85,18} S. Pordes,⁶⁶ J. Porter,¹⁸⁵ M. Potekhin,²¹ R. Potenza,^{86,37}
 B. V. K. S. Potukuchi,¹¹² J. Pozimski,¹⁰² M. Pozzato,^{85,18} S. Prakash,³⁶ T. Prakash,¹²⁵ M. Prest,⁹⁰ S. Prince,⁷⁸
 F. Psihas,⁶⁶ D. Pugnere,⁹⁷ X. Qian,²¹ J. L. Raaf,⁶⁶ V. Radeka,²¹ J. Rademacker,²⁰ B. Radics,⁵⁸ A. Rafique,⁷
 E. Raguzin,²¹ M. Rai,²⁰¹ M. Rajaoalisoa,⁴² I. Rakhno,⁶⁶ A. Rakotonandrasana,⁵ L. Rakotondravohitra,⁵
 Y. A. Ramachers,²⁰¹ R. Rameika,⁶⁶ M. A. Ramirez Delgado,¹⁵⁹ B. Ramson,⁶⁶ A. Rappoldi,^{94,158} G. Raselli,^{94,158}
 P. Ratoff,¹²⁴ S. Raut,¹⁸³ R. F. Razakamiandra,⁵ E. Rea,¹⁴⁰ J.S. Real,⁷⁵ B. Rebel,^{205,66} M. Reggiani-Guzzo,¹³¹
 T. Rehak,⁵⁴ J. Reichenbacher,¹⁷⁹ S. D. Reitzner,⁶⁶ H. Rejeb Sfar,²⁵ A. Renshaw,⁸⁰ S. Rescia,²¹ F. Resnati,²⁵
 A. Reynolds,¹⁵² M. Ribas,¹⁸⁷ S. Riboldi,⁹¹ C. Riccio,¹⁸³ G. Riccobene,⁹⁵ L. C. J. Rice,¹⁶³ J. Ricol,⁷⁵ A. Rigamonti,²⁵
 Y. Rigaut,⁵⁸ D. Rivera,¹⁵⁹ A. Robert,⁷⁵ L. Rochester,¹⁷³ M. Roda,¹²⁶ P. Rodrigues,¹⁵² M. J. Rodriguez
 Alonso,²⁵ E. Rodriguez Bonilla,⁶ J. Rodriguez Rondon,¹⁷⁹ S. Rosauro-Alcaraz,¹³⁰ M. Rosenberg,¹⁶³ P. Rosier,¹⁵⁶
 B. Roskovec,³⁰ M. Rossella,^{94,158} M. Rossi,²⁵ J. Rout,¹¹³ P. Roy,²⁰³ A. Rubbia,⁵⁸ C. Rubbia,⁷² B. Russell,¹²⁵
 D. Ruterbories,¹⁶⁹ A. Rybnikov,¹¹¹ A. Saa-Hernandez,⁸⁴ R. Saakyan,¹⁹⁶ S. Sacerdoti,¹⁵⁷ T. Safford,¹³⁶ N. Sahu,¹⁰⁴
 P. Sala,^{91,25} N. Samios,²¹ O. Samoylov,¹¹¹ M. C. Sanchez,¹⁰⁹ V. Sandberg,¹²⁷ D. A. Sanders,¹⁴¹ D. Sankey,¹⁷²
 S. Santana,¹⁶⁵ M. Santos-Maldonado,¹⁶⁵ N. Saoulidou,¹⁰ P. Sapienza,⁹⁵ C. Sarasty,⁴² I. Sarcevic,⁸ G. Savage,⁶⁶
 V. Savinov,¹⁶³ A. Scaramelli,⁹⁴ A. Scarff,¹⁷⁸ A. Scarpelli,²¹ H. Schellman,^{151,66} S. Schifano,^{87,67} P. Schlabach,⁶⁶
 D. Schmitz,⁴⁰ K. Scholberg,⁵⁵ A. Schukraft,⁶⁶ E. Segreto,³⁶ A. Selyunin,¹¹¹ C. R. Senise,¹⁹⁵ J. Sensenig,¹⁵⁹
 M. Seoane,⁸⁴ A. Sergi,¹⁷ D. Sgalaberna,⁵⁸ M. H. Shaevitz,⁴⁷ S. Shafaq,¹¹³ M. Shamma,³² R. Sharankova,¹⁹³
 H. R. Sharma,¹¹² R. Sharma,²¹ R. Kumar,¹⁶⁶ T. Shaw,⁶⁶ C. Shepherd-Themistocleous,¹⁷² A. Sheshukov,¹¹¹
 S. Shin,¹¹⁴ I. Shoemaker,¹⁹⁹ D. Shoultz,¹³⁶ R. Shrock,¹⁸³ H. Siegel,⁴⁷ L. Simard,¹⁵⁶ F. Simon,^{66,133} J. Sinclair,¹⁵
 G. Sinev,¹⁷⁹ Jaydip Singh,¹²⁹ J. Singh,¹²⁹ L. Singh,²⁷ V. Singh,^{27,13} R. Sipos,²⁵ F. W. Sippach,⁴⁷ G. Sirri,⁸⁵
 A. Sitraka,¹⁷⁹ K. Siyeon,⁴¹ K. Skarpaas,¹⁷³ A. Smith,³⁵ E. Smith,¹⁰⁵ P. Smith,¹⁰⁵ J. Smolik,⁴⁹ M. Smy,³⁰
 E.L. Snider,⁶⁶ P. Snopok,¹⁰¹ D. Snowden-Ifft,¹⁴⁹ M. Soares Nunes,¹⁸⁶ H. Sobel,³⁰ M. Soderberg,¹⁸⁶ S. Sokolov,¹¹¹
 C. J. Solano Salinas,¹⁰⁶ S. Söldner-Rembold,¹³¹ S.R. Soleti,¹²⁵ N. Solomey,²⁰³ V. Solovov,¹²³ W. E. Sondheim,¹²⁷
 M. Sorel,⁸³ A. Sotnikov,¹¹¹ J. Soto-Oton,²⁶ A. Sousa,⁴² K. Soustruznik,³⁹ F. Spaggiardi,¹⁵² M. Spanu,^{90,137}
 J. Spitz,¹³⁵ N. J. C. Spooner,¹⁷⁸ K. Spurgeon,¹⁸⁶ M. Stancari,⁶⁶ L. Stanco,^{93,154} R. Stein,²⁰ H. M. Steiner,¹²⁵
 A. F. Steklain Lisboá,¹⁸⁷ J. Stewart,²¹ B. Stillwell,⁴⁰ J. Stock,¹⁷⁹ F. Stocker,²⁵ T. Stokes,¹²⁸ M. Strait,¹⁴⁰
 T. Strauss,⁶⁶ A. Stuart,⁴⁴ J. G. Suarez,⁵⁷ H. Sullivan,¹⁹⁰ D. Summers,¹⁴¹ A. Surdo,⁸⁹ V. Susic,¹⁴ L. Suter,⁶⁶
 C. M. Suter,^{86,37} R. Svoboda,²⁹ B. Szczerbinska,¹⁸⁹ A. M. Szelc,⁵⁹ H. A. Tanaka,¹⁷³ B. Tapia Oregui,¹⁹¹
 A. Tapper,¹⁰² S. Tariq,⁶⁶ E. Tatar,¹⁰⁰ R. Tayloe,¹⁰⁵ A. M. Teklu,¹⁸³ M. Tenti,⁸⁵ K. Terao,¹⁷³ C. A. Ternes,⁸³
 F. Terranova,^{90,137} G. Testera,⁸⁸ T. Thakore,⁴² A. Thea,¹⁷² J. L. Thompson,¹⁷⁸ C. Thorn,²¹ S. C. Timm,⁶⁶
 V. Tishchenko,²¹ L. Tomassetti,^{87,67} A. Tonazzo,¹⁵⁷ D. Torbunov,¹⁴⁰ M. Torti,^{90,137} M. Tortola,⁸³
 F. Tortorici,^{86,37} N. Tosi,⁸⁵ D. Totani,³³ M. Toups,⁶⁶ C. Touramanis,¹²⁶ R. Travaglini,⁸⁵ J. Trevor,³⁴
 S. Trilov,²⁰ W. H. Trzaska,¹¹⁵ Y. Tsai,⁶⁶ Y.-T. Tsai,¹⁷³ Z. Tsamalaidze,⁷¹ K. V. Tsang,¹⁷³ N. Tsverava,⁷¹
 S. Tufanli,²⁵ C. Tull,¹²⁵ E. Tyley,¹⁷⁸ M. Tzanov,¹²⁸ L. Uboldi,²⁵ M. A. Uchida,³⁵ J. Urheim,¹⁰⁵ T. Usher,¹⁷³
 S. Uzunyan,¹⁴⁶ M. R. Vagins,¹²⁰ P. Vahle,²⁰⁴ G. A. Valdivieso,⁶¹ R. Valentim,¹⁹⁵ Z. Vallari,³⁴ E. Vallazza,⁹⁰
 J. W. F. Valle,⁸³ S. Vallecorsa,²⁵ R. Van Berg,¹⁵⁹ R. G. Van de Water,¹²⁷ F. Varanini,⁹³ D. Vargas,⁸² G. Varner,⁷⁹
 J. Vasel,¹⁰⁵ S. Vasina,¹¹¹ G. Vasseur,²⁴ N. Vaughan,¹⁵¹ K. Vaziri,⁶⁶ S. Ventura,⁹³ A. Verdugo,²⁶ S. Vergani,³⁵
 M. A. Vermeulen,¹⁴⁴ M. Verzocchi,⁶⁶ M. Vicenzi,^{88,70} H. Vieira de Souza,^{36,90} C. Vignoli,⁷³ C. Vilela,²⁵
 B. Viren,²¹ T. Vrba,⁴⁹ T. Wachala,¹⁴³ A. V. Waldron,¹⁰² M. Wallbank,⁴² C. Wallis,⁴⁶ H. Wang,³¹ J. Wang,¹⁷⁹
 L. Wang,¹²⁵ M.H.L.S. Wang,⁶⁶ Y. Wang,³¹ Y. Wang,¹⁸³ K. Warburton,¹⁰⁹ D. Warner,⁴⁶ M.O. Wascko,¹⁰²
 D. Waters,¹⁹⁶ A. Watson,¹⁷ P. Weatherly,⁵⁴ A. Weber,^{172,152} M. Weber,¹⁵ H. Wei,²¹ A. Weinstein,¹⁰⁹
 D. Wenman,²⁰⁵ M. Wetstein,¹⁰⁹ A. White,¹⁹⁰ L. H. Whitehead,³⁵ D. Whittington,¹⁸⁶ M. J. Wilking,¹⁸³
 C. Wilkinson,^{125,*} Z. Williams,¹⁹⁰ F. Wilson,¹⁷² R. J. Wilson,⁴⁶ W. Wisniewski,¹⁷³ J. Wolcott,¹⁹³ T. Wongjirad,¹⁹³
 A. Wood,⁸⁰ K. Wood,¹²⁵ E. Worcester,²¹ M. Worcester,²¹ C. Wret,¹⁶⁹ W. Wu,⁶⁶ W. Wu,³⁰ Y. Xiao,³⁰

F. Xie,¹⁸⁵ E. Yandel,³³ G. Yang,¹⁸³ K. Yang,¹⁵² T. Yang,⁶⁶ A. Yankelevich,³⁰ N. Yershov,⁹⁶ K. Yonehara,⁶⁶
 T. Young,¹⁴⁵ B. Yu,²¹ H. Yu,²¹ H. Yu,¹⁸⁴ J. Yu,¹⁹⁰ W. Yuan,⁵⁹ R. Zaki,²⁰⁸ J. Zalesak,⁴⁸ L. Zambelli,⁵²
 B. Zamorano,⁷⁴ A. Zani,⁹¹ L. Zazueta,²⁰⁴ G. P. Zeller,⁶⁶ J. Zennamo,⁶⁶ K. Zeug,²⁰⁵ C. Zhang,²¹ M. Zhao,²¹
 E. Zhivun,²¹ G. Zhu,¹⁵⁰ E. D. Zimmerman,⁴⁵ S. Zucchelli,^{85,18} J. Zuklin,⁴⁸ V. Zutshi,¹⁴⁶ and R. Zwaska⁶⁶

(The DUNE Collaboration)

- ¹Abilene Christian University, Abilene, TX 79601, USA
²University of Albany, SUNY, Albany, NY 12222, USA
³University of Amsterdam, NL-1098 XG Amsterdam, The Netherlands
⁴Antalya Bilim University, 07190 Döşemealtı/Antalya, Turkey
⁵University of Antananarivo, Antananarivo 101, Madagascar
⁶Universidad Antonio Nariño, Bogotá, Colombia
⁷Argonne National Laboratory, Argonne, IL 60439, USA
⁸University of Arizona, Tucson, AZ 85721, USA
⁹Universidad Nacional de Asunción, San Lorenzo, Paraguay
¹⁰University of Athens, Zografou GR 157 84, Greece
¹¹Universidad del Atlántico, Barranquilla, Atlántico, Colombia
¹²Augustana University, Sioux Falls, SD 57197, USA
¹³Banaras Hindu University, Varanasi - 221 005, India
¹⁴University of Basel, CH-4056 Basel, Switzerland
¹⁵University of Bern, CH-3012 Bern, Switzerland
¹⁶Beykent University, Istanbul, Turkey
¹⁷University of Birmingham, Birmingham B15 2TT, United Kingdom
¹⁸Università del Bologna, 40127 Bologna, Italy
¹⁹Boston University, Boston, MA 02215, USA
²⁰University of Bristol, Bristol BS8 1TL, United Kingdom
²¹Brookhaven National Laboratory, Upton, NY 11973, USA
²²University of Bucharest, Bucharest, Romania
²³Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, RJ 22290-180, Brazil
²⁴IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France
²⁵CERN, The European Organization for Nuclear Research, 1211 Meyrin, Switzerland
²⁶CIEMAT, Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas, E-28040 Madrid, Spain
²⁷Central University of South Bihar, Gaya, 824236, India
²⁸University of California Berkeley, Berkeley, CA 94720, USA
²⁹University of California Davis, Davis, CA 95616, USA
³⁰University of California Irvine, Irvine, CA 92697, USA
³¹University of California Los Angeles, Los Angeles, CA 90095, USA
³²University of California Riverside, Riverside CA 92521, USA
³³University of California Santa Barbara, Santa Barbara, California 93106 USA
³⁴California Institute of Technology, Pasadena, CA 91125, USA
³⁵University of Cambridge, Cambridge CB3 0HE, United Kingdom
³⁶Universidade Estadual de Campinas, Campinas - SP, 13083-970, Brazil
³⁷Università di Catania, 2 - 95131 Catania, Italy
³⁸Universidad Católica del Norte, Antofagasta, Chile
³⁹Institute of Particle and Nuclear Physics of the Faculty of Mathematics and Physics of the Charles University, 180 00 Prague 8, Czech Republic
⁴⁰University of Chicago, Chicago, IL 60637, USA
⁴¹Chung-Ang University, Seoul 06974, South Korea
⁴²University of Cincinnati, Cincinnati, OH 45221, USA
⁴³Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional (Cinvestav), Mexico City, Mexico
⁴⁴Universidad de Colima, Colima, Mexico
⁴⁵University of Colorado Boulder, Boulder, CO 80309, USA
⁴⁶Colorado State University, Fort Collins, CO 80523, USA
⁴⁷Columbia University, New York, NY 10027, USA
⁴⁸Institute of Physics, Czech Academy of Sciences, 182 00 Prague 8, Czech Republic
⁴⁹Czech Technical University, 115 19 Prague 1, Czech Republic
⁵⁰Dakota State University, Madison, SD 57042, USA
⁵¹University of Dallas, Irving, TX 75062-4736, USA
⁵²Laboratoire d'Annecy de Physique des Particules, Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LAPP-IN2P3, 74000 Annecy, France
⁵³Daresbury Laboratory, Cheshire WA4 4AD, United Kingdom
⁵⁴Drexel University, Philadelphia, PA 19104, USA
⁵⁵Duke University, Durham, NC 27708, USA

- ⁵⁶ *Durham University, Durham DH1 3LE, United Kingdom*
- ⁵⁷ *Universidad EIA, Envigado, Antioquia, Colombia*
- ⁵⁸ *ETH Zurich, Zurich, Switzerland*
- ⁵⁹ *University of Edinburgh, Edinburgh EH8 9YL, United Kingdom*
- ⁶⁰ *Faculdade de Ciências da Universidade de Lisboa - FCUL, 1749-016 Lisboa, Portugal*
- ⁶¹ *Universidade Federal de Alfenas, Poços de Caldas - MG, 37715-400, Brazil*
- ⁶² *Universidade Federal de Goiás, Goiania, GO 74690-900, Brazil*
- ⁶³ *Universidade Federal de São Carlos, Araras - SP, 13604-900, Brazil*
- ⁶⁴ *Universidade Federal do ABC, Santo André - SP, 09210-580, Brazil*
- ⁶⁵ *Universidade Federal do Rio de Janeiro, Rio de Janeiro - RJ, 21941-901, Brazil*
- ⁶⁶ *Fermi National Accelerator Laboratory, Batavia, IL 60510, USA*
- ⁶⁷ *University of Ferrara, Ferrara, Italy*
- ⁶⁸ *University of Florida, Gainesville, FL 32611-8440, USA*
- ⁶⁹ *Fluminense Federal University, 9 Icaraí Niterói - RJ, 24220-900, Brazil*
- ⁷⁰ *Università degli Studi di Genova, Genova, Italy*
- ⁷¹ *Georgian Technical University, Tbilisi, Georgia*
- ⁷² *Gran Sasso Science Institute, L'Aquila, Italy*
- ⁷³ *Laboratori Nazionali del Gran Sasso, L'Aquila AQ, Italy*
- ⁷⁴ *University of Granada & CAFPE, 18002 Granada, Spain*
- ⁷⁵ *University Grenoble Alpes, CNRS, Grenoble INP, LPSC-IN2P3, 38000 Grenoble, France*
- ⁷⁶ *Universidad de Guanajuato, Guanajuato, C.P. 37000, Mexico*
- ⁷⁷ *Harish-Chandra Research Institute, Jhansi, Allahabad 211 019, India*
- ⁷⁸ *Harvard University, Cambridge, MA 02138, USA*
- ⁷⁹ *University of Hawaii, Honolulu, HI 96822, USA*
- ⁸⁰ *University of Houston, Houston, TX 77204, USA*
- ⁸¹ *University of Hyderabad, Gachibowli, Hyderabad - 500 046, India*
- ⁸² *Institut de Física d'Altes Energies (IFAE)—Barcelona Institute of Science and Technology (BIST), Barcelona, Spain*
- ⁸³ *Instituto de Física Corpuscular, CSIC and Universitat de València, 46980 Paterna, Valencia, Spain*
- ⁸⁴ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, E-15782, Galicia, Spain*
- ⁸⁵ *Istituto Nazionale di Fisica Nucleare Sezione di Bologna, 40127 Bologna BO, Italy*
- ⁸⁶ *Istituto Nazionale di Fisica Nucleare Sezione di Catania, I-95123 Catania, Italy*
- ⁸⁷ *Istituto Nazionale di Fisica Nucleare Sezione di Ferrara, I-44122 Ferrara, Italy*
- ⁸⁸ *Istituto Nazionale di Fisica Nucleare Sezione di Genova, 16146 Genova GE, Italy*
- ⁸⁹ *Istituto Nazionale di Fisica Nucleare Sezione di Lecce, 73100 - Lecce, Italy*
- ⁹⁰ *Istituto Nazionale di Fisica Nucleare Sezione di Milano Bicocca, 3 - I-20126 Milano, Italy*
- ⁹¹ *Istituto Nazionale di Fisica Nucleare Sezione di Milano, 20133 Milano, Italy*
- ⁹² *Istituto Nazionale di Fisica Nucleare Sezione di Napoli, I-80126 Napoli, Italy*
- ⁹³ *Istituto Nazionale di Fisica Nucleare Sezione di Padova, 35131 Padova, Italy*
- ⁹⁴ *Istituto Nazionale di Fisica Nucleare Sezione di Pavia, I-27100 Pavia, Italy*
- ⁹⁵ *Istituto Nazionale di Fisica Nucleare Laboratori Nazionali del Sud, 95123 Catania, Italy*
- ⁹⁶ *Institute for Nuclear Research of the Russian Academy of Sciences, Moscow 117312, Russia*
- ⁹⁷ *Institut de Physique des 2 Infinis de Lyon, 69622 Villeurbanne, France*
- ⁹⁸ *Institute for Research in Fundamental Sciences, Tehran, Iran*
- ⁹⁹ *Instituto Superior Técnico - IST, Universidade de Lisboa, Portugal*
- ¹⁰⁰ *Idaho State University, Pocatello, ID 83209, USA*
- ¹⁰¹ *Illinois Institute of Technology, Chicago, IL 60616, USA*
- ¹⁰² *Imperial College of Science Technology and Medicine, London SW7 2BZ, United Kingdom*
- ¹⁰³ *Indian Institute of Technology Guwahati, Guwahati, 781 039, India*
- ¹⁰⁴ *Indian Institute of Technology Hyderabad, Hyderabad, 502285, India*
- ¹⁰⁵ *Indiana University, Bloomington, IN 47405, USA*
- ¹⁰⁶ *Universidad Nacional de Ingeniería, Lima 25, Perú*
- ¹⁰⁷ *University of Insubria, Via Ravasi, 2, 21100 Varese VA, Italy*
- ¹⁰⁸ *University of Iowa, Iowa City, IA 52242, USA*
- ¹⁰⁹ *Iowa State University, Ames, Iowa 50011, USA*
- ¹¹⁰ *Iwate University, Morioka, Iwate 020-8551, Japan*
- ¹¹¹ *Joint Institute for Nuclear Research, Dzhelapov Laboratory of Nuclear Problems 6 Joliot-Curie, Dubna, Moscow Region, 141980 RU*
- ¹¹² *University of Jammu, Jammu-180006, India*
- ¹¹³ *Jawaharlal Nehru University, New Delhi 110067, India*
- ¹¹⁴ *Jeonbuk National University, Jeonrabuk-do 54896, South Korea*
- ¹¹⁵ *University of Jyväskylä, FI-40014, Finland*
- ¹¹⁶ *High Energy Accelerator Research Organization (KEK), Ibaraki, 305-0801, Japan*
- ¹¹⁷ *Korea Institute of Science and Technology Information, Daejeon, 34141, South Korea*
- ¹¹⁸ *K L University, Vaddeswaram, Andhra Pradesh 522502, India*

- ¹¹⁹ *Kansas State University, Manhattan, KS 66506, USA*
- ¹²⁰ *Kavli Institute for the Physics and Mathematics of the Universe, Kashiwa, Chiba 277-8583, Japan*
- ¹²¹ *National Institute of Technology, Kure College, Hiroshima, 737-8506, Japan*
- ¹²² *Taras Shevchenko National University of Kyiv, 01601 Kyiv, Ukraine*
- ¹²³ *Laboratório de Instrumentação e Física Experimental de Partículas, 1649-003 Lisboa and 3004-516 Coimbra, Portugal*
- ¹²⁴ *Lancaster University, Lancaster LA1 4YB, United Kingdom*
- ¹²⁵ *Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*
- ¹²⁶ *University of Liverpool, L69 7ZE, Liverpool, United Kingdom*
- ¹²⁷ *Los Alamos National Laboratory, Los Alamos, NM 87545, USA*
- ¹²⁸ *Louisiana State University, Baton Rouge, LA 70803, USA*
- ¹²⁹ *University of Lucknow, Uttar Pradesh 226007, India*
- ¹³⁰ *Madrid Autonoma University and IFT UAM/CSIC, 28049 Madrid, Spain*
- ¹³¹ *University of Manchester, Manchester M13 9PL, United Kingdom*
- ¹³² *Massachusetts Institute of Technology, Cambridge, MA 02139, USA*
- ¹³³ *Max-Planck-Institut, Munich, 80805, Germany*
- ¹³⁴ *University of Medellín, Medellín, 050026 Colombia*
- ¹³⁵ *University of Michigan, Ann Arbor, MI 48109, USA*
- ¹³⁶ *Michigan State University, East Lansing, MI 48824, USA*
- ¹³⁷ *Università del Milano-Bicocca, 20126 Milano, Italy*
- ¹³⁸ *Università degli Studi di Milano, I-20133 Milano, Italy*
- ¹³⁹ *University of Minnesota Duluth, Duluth, MN 55812, USA*
- ¹⁴⁰ *University of Minnesota Twin Cities, Minneapolis, MN 55455, USA*
- ¹⁴¹ *University of Mississippi, University, MS 38677 USA*
- ¹⁴² *University of New Mexico, Albuquerque, NM 87131, USA*
- ¹⁴³ *H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland*
- ¹⁴⁴ *Nikhef National Institute of Subatomic Physics, 1098 XG Amsterdam, Netherlands*
- ¹⁴⁵ *University of North Dakota, Grand Forks, ND 58202-8357, USA*
- ¹⁴⁶ *Northern Illinois University, DeKalb, IL 60115, USA*
- ¹⁴⁷ *Northwestern University, Evanston, IL 60208, USA*
- ¹⁴⁸ *University of Notre Dame, Notre Dame, IN 46556, USA*
- ¹⁴⁹ *Occidental College, Los Angeles, CA 90041*
- ¹⁵⁰ *Ohio State University, Columbus, OH 43210, USA*
- ¹⁵¹ *Oregon State University, Corvallis, OR 97331, USA*
- ¹⁵² *University of Oxford, Oxford, OX1 3RH, United Kingdom*
- ¹⁵³ *Pacific Northwest National Laboratory, Richland, WA 99352, USA*
- ¹⁵⁴ *Università degli Studi di Padova, I-35131 Padova, Italy*
- ¹⁵⁵ *Panjab University, Chandigarh, 160014 U.T., India*
- ¹⁵⁶ *Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France*
- ¹⁵⁷ *Université de Paris, CNRS, Astroparticule et Cosmologie, F-75006, Paris, France*
- ¹⁵⁸ *Università degli Studi di Pavia, 27100 Pavia PV, Italy*
- ¹⁵⁹ *University of Pennsylvania, Philadelphia, PA 19104, USA*
- ¹⁶⁰ *Pennsylvania State University, University Park, PA 16802, USA*
- ¹⁶¹ *Physical Research Laboratory, Ahmedabad 380 009, India*
- ¹⁶² *Università di Pisa, I-56127 Pisa, Italy*
- ¹⁶³ *University of Pittsburgh, Pittsburgh, PA 15260, USA*
- ¹⁶⁴ *Pontificia Universidad Católica del Perú, Lima, Perú*
- ¹⁶⁵ *University of Puerto Rico, Mayaguez 00681, Puerto Rico, USA*
- ¹⁶⁶ *Punjab Agricultural University, Ludhiana 141004, India*
- ¹⁶⁷ *Queen Mary University of London, London E1 4NS, United Kingdom*
- ¹⁶⁸ *Radboud University, NL-6525 AJ Nijmegen, Netherlands*
- ¹⁶⁹ *University of Rochester, Rochester, NY 14627, USA*
- ¹⁷⁰ *Royal Holloway College London, TW20 0EX, United Kingdom*
- ¹⁷¹ *Rutgers University, Piscataway, NJ, 08854, USA*
- ¹⁷² *STFC Rutherford Appleton Laboratory, Didcot OX11 0QX, United Kingdom*
- ¹⁷³ *SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA*
- ¹⁷⁴ *Sanford Underground Research Facility, Lead, SD, 57754, USA*
- ¹⁷⁵ *Università del Salento, 73100 Lecce, Italy*
- ¹⁷⁶ *San Jose State University, San José, CA 95192-0106, USA*
- ¹⁷⁷ *Universidad Sergio Arboleda, 11022 Bogotá, Colombia*
- ¹⁷⁸ *University of Sheffield, Sheffield S3 7RH, United Kingdom*
- ¹⁷⁹ *South Dakota School of Mines and Technology, Rapid City, SD 57701, USA*
- ¹⁸⁰ *South Dakota State University, Brookings, SD 57007, USA*
- ¹⁸¹ *University of South Carolina, Columbia, SC 29208, USA*
- ¹⁸² *Southern Methodist University, Dallas, TX 75275, USA*

- ¹⁸³ *Stony Brook University, SUNY, Stony Brook, NY 11794, USA*
¹⁸⁴ *Sun Yat-Sen University, Guangzhou, 510275*
¹⁸⁵ *University of Sussex, Brighton, BN1 9RH, United Kingdom*
¹⁸⁶ *Syracuse University, Syracuse, NY 13244, USA*
¹⁸⁷ *Universidade Tecnológica Federal do Paraná, Curitiba, Brazil*
¹⁸⁸ *Texas A&M University, College Station, Texas 77840*
¹⁸⁹ *Texas A&M University - Corpus Christi, Corpus Christi, TX 78412, USA*
¹⁹⁰ *University of Texas at Arlington, Arlington, TX 76019, USA*
¹⁹¹ *University of Texas at Austin, Austin, TX 78712, USA*
¹⁹² *University of Toronto, Toronto, Ontario M5S 1A1, Canada*
¹⁹³ *Tufts University, Medford, MA 02155, USA*
¹⁹⁴ *Ulsan National Institute of Science and Technology, Ulsan 689-798, South Korea*
¹⁹⁵ *Universidade Federal de São Paulo, 09913-030, São Paulo, Brazil*
¹⁹⁶ *University College London, London, WC1E 6BT, United Kingdom*
¹⁹⁷ *Valley City State University, Valley City, ND 58072, USA*
¹⁹⁸ *Variable Energy Cyclotron Centre, 700 064 West Bengal, India*
¹⁹⁹ *Virginia Tech, Blacksburg, VA 24060, USA*
²⁰⁰ *University of Warsaw, 02-093 Warsaw, Poland*
²⁰¹ *University of Warwick, Coventry CV4 7AL, United Kingdom*
²⁰² *Wellesley College, Wellesley, MA 02481, USA*
²⁰³ *Wichita State University, Wichita, KS 67260, USA*
²⁰⁴ *William and Mary, Williamsburg, VA 23187, USA*
²⁰⁵ *University of Wisconsin Madison, Madison, WI 53706, USA*
²⁰⁶ *Yale University, New Haven, CT 06520, USA*
²⁰⁷ *Yerevan Institute for Theoretical Physics and Modeling, Yerevan 0036, Armenia*
²⁰⁸ *York University, Toronto M3J 1P3, Canada*
- (Dated: September 6, 2021)

The Deep Underground Neutrino Experiment (DUNE) will produce world-leading neutrino oscillation measurements over the lifetime of the experiment. In this work, we explore DUNE’s sensitivity to observe charge-parity violation (CPV) in the neutrino sector, and to resolve the mass ordering, for exposures of up to 100 kiloton-megawatt-years (kt-MW-yr). The analysis includes detailed uncertainties on the flux prediction, the neutrino interaction model, and detector effects. We demonstrate that DUNE will be able to unambiguously resolve the neutrino mass ordering at a 3σ (5σ) level, with a 66 (100) kt-MW-yr far detector exposure, and has the ability to make strong statements at significantly shorter exposures depending on the true value of other oscillation parameters. We also show that DUNE has the potential to make a robust measurement of CPV at a 3σ level with a 100 kt-MW-yr exposure for the maximally CP-violating values $\delta_{\text{CP}} = \pm\pi/2$. Additionally, the dependence of DUNE’s sensitivity on the exposure taken in neutrino-enhanced and antineutrino-enhanced running is discussed. An equal fraction of exposure taken in each beam mode is found to be close to optimal when considered over the entire space of interest.

I. INTRODUCTION

The Deep Underground Neutrino Experiment (DUNE) [1] is a next-generation long-baseline neutrino oscillation experiment which will utilize high-intensity ν_μ and $\bar{\nu}_\mu$ beams with peak neutrino energies of ≈ 2.5 GeV over a 1285 km baseline to carry out a detailed study of neutrino mixing. Some of DUNE’s key scientific goals are the definitive determination of the neutrino mass ordering, the definitive observation of charge-parity symmetry violation (CPV) for more than 50% of possible true values of the charge-parity violating phase, δ_{CP} , and the precise measurement of other three-neutrino oscillation parameters. These measurements will help guide theory in understanding if there are new symmetries in the neutrino sector and whether there is

a relationship between the generational structure of quarks and leptons [2]. Observation of CPV in neutrinos would be an important step in understanding the origin of the baryon asymmetry of the universe [3, 4]. DUNE has a rich physics program beyond the three-neutrino oscillation accelerator neutrino program described here. These include beyond standard model searches [5], supernova neutrino detection [6], and solar neutrino detection [7]. Additional physics possibilities with DUNE are discussed in Refs. [8] and [9].

Neutrino oscillation experiments have so far measured five of the three-neutrino mixing parameters [10–12]: the three mixing angles θ_{12} , θ_{23} , and θ_{13} ; and the two squared-mass differences Δm_{21}^2 and $|\Delta m_{31}^2|$, where $\Delta m_{ij}^2 = m_i^2 - m_j^2$ is the difference between the squares of the neutrino masses. The neutrino mass ordering (the sign of Δm_{31}^2) is not currently known, though recent results show a weak preference for the normal ordering (NO), where $\Delta m_{31}^2 > 0$, over the inverted ordering

* Corresponding author: cwilkinson@lbl.gov

(IO) [13–15]. The value of δ_{CP} is not well known, though neutrino oscillation data are beginning to provide some information on its value [13, 16].

The oscillation probability of $(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$ through matter in the standard three-flavor model and a constant matter density approximation can be written as [17]:

$$\begin{aligned}
P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) &\simeq \sin^2 \theta_{23} \sin^2 2\theta_{13} \frac{\sin^2(\Delta_{31} - aL)}{(\Delta_{31} - aL)^2} \Delta_{31}^2 \\
&+ \sin 2\theta_{23} \sin 2\theta_{13} \sin 2\theta_{12} \frac{\sin(\Delta_{31} - aL)}{(\Delta_{31} - aL)} \Delta_{31} \\
&\times \frac{\sin(aL)}{(aL)} \Delta_{21} \cos(\Delta_{31} \pm \delta_{\text{CP}}) \\
&+ \cos^2 \theta_{23} \sin^2 2\theta_{12} \frac{\sin^2(aL)}{(aL)^2} \Delta_{21}^2,
\end{aligned} \tag{1}$$

where

$$a = \pm \frac{G_{\text{F}} N_e}{\sqrt{2}} \approx \pm \frac{1}{3500 \text{ km}} \left(\frac{\rho}{3.0 \text{ g/cm}^3} \right),$$

$aL \approx 0.367$, G_{F} is the Fermi constant, N_e is the number density of electrons in the Earth’s crust, $\Delta_{ij} = 1.267 \Delta m_{ij}^2 L / E_\nu$, Δm_{ij}^2 is in eV^2 , L is the baseline in km, and E_ν is the neutrino energy in GeV. Both δ_{CP} and a terms are positive (negative) for $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) oscillations. The matter effect asymmetry encapsulated in the a terms arises from the presence of electrons and absence of positrons in the Earth’s crust [18, 19]. In the analysis described here, the oscillation probabilities are calculated exactly [20].

DUNE has published sensitivity estimates [21] to CPV and the neutrino mass ordering, as well as other oscillation parameters, for large exposures of up to 1104 kiloton-megawatt-years (kt-MW-yr), which show the ultimate sensitivity of the experiment. Sophisticated studies with a detailed treatment of systematic uncertainties were carried out only at large exposures. In this work, DUNE’s sensitivity at low exposures is explored further, with a detailed systematics treatment, including an investigation into how the run plan may be optimized to enhance sensitivity to CPV and/or mass ordering. It is shown that DUNE will produce world-leading results at relatively short exposures, which highlights the need for a high-performance near detector complex from the beginning of the experiment.

The DUNE far detector (FD) will ultimately consist of four modules, each with a 17 kt total mass. The neutrino beam-line has an initial design intensity of 1.2 MW, with a planned upgrade to 2.4 MW. We assume a combined yearly Fermilab accelerator and neutrino beam-line uptime of 57% [8], and include this in our calculation of calendar years in this work. As the FD deployment schedule and beam power scenarios are both subject to change, the results shown in this work are consistently given in terms of exposure in units of kt-MW-yr, which is agnostic to the exact staging scenario, but can easily be expressed

in terms of experiment years for any desired scenario. For example, with two FD modules, assuming a fiducial mass of 10 kt and a beam intensity of 1.2 MW, exposure would accumulate at a rate of 24 kt-MW-yr per calendar year, although a ramp up in beam power is expected before reaching the design intensity in early running. In this work, the single-phase horizontal drift technology is assumed for all FD modules (see Section IID), which is a necessary simplification, but alternative technologies which may have slightly different performance are under investigation for some FD modules.

The analysis framework is described in Section II, including a description of the flux, neutrino interaction and detector models and associated uncertainties. A study on the dependence of the sensitivity to CPV and mass ordering on the fraction of data collected in neutrino-enhanced or antineutrino-enhanced running is given in Section III. A detailed study on the CPV and mass ordering sensitivities at low exposures are described in Sections IV and V, respectively. Finally, conclusions are presented in Section VI.

II. ANALYSIS FRAMEWORK

This work uses the flux, neutrino interaction and detector model described in detail in Ref. [21], implemented in the CAFAna framework [22]. This section provides an overview of the key analysis features. Further details on all aspects can be found in Ref. [21].

A. Neutrino flux

DUNE will operate with two different beam modes, which depend on the polarity of the electromagnetic horns used to focus secondary particles produced after protons from the primary beam line interact in the target. Forward horn current (FHC) corresponds to neutrino-enhanced running, and reverse horn current (RHC) corresponds to antineutrino-enhanced running. In both FHC and RHC there are significant contributions from neutrinos with energies between 0.5–6 GeV, with a flux peak at ≈ 2.5 GeV. The neutrino flux prediction is generated with G4LBNF [8, 23], using the LBNF optimized beam design [8]. Flux uncertainties are due to the production rates and kinematic distributions of hadrons produced in the target and the parameters of the beam line, such as horn currents and horn and target positioning (“focusing uncertainties”) [8]. They are evaluated using current measurements of hadron production and estimates of alignment tolerances, giving flux uncertainties of approximately 8% at the first oscillation maximum, which are highly correlated across energy bins and neutrino flavors. A flux covariance as a function of neutrino energy, beam mode, detector, and neutrino species is generated with a “toy throw” approach, which is built using variations (“throws”) of the systematics propagated through

the full beam line simulation. To reduce the number of parameters used in the fit, the covariance matrix is diagonalized, and each principal component is treated as an uncorrelated nuisance parameter. Only the first ≈ 30 principal components (out of 108) were found to have a significant effect in the analysis and were included. The shapes of the unoscillated fluxes at the ND and FD are similar, and the differences between them are understood at the percent level.

B. Neutrino interaction model

The interaction model used is based on GENIE v2.12.10 [24, 25], although the combination of models used is much closer to some of the physics tunes available with GENIE v3.00.06, including a number of uncertainties beyond those provided by either GENIE version. These are motivated by data, although the available (anti)neutrino data taken on argon targets is sparse, leading to an uncertainty model that relies in a number of places on light target (mostly hydrocarbon) data. Variations in the cross sections are implemented either using GENIE reweighting parameters, or with *ad hoc* weights of events designed to parameterize uncertainties or cross-section corrections currently not implemented within GENIE. The latter were developed using alternative generators or GENIE configurations, or custom weightings using the NUISANCE package [26]. Further details about the uncertainties used can be found in Ref. [21] (Section 3).

The nuclear model which describes the initial state of nucleons in the nucleus is the Bodek-Ritchie global Fermi gas model [27], which includes empirical modifications to the nucleon momentum distribution to account for short-range correlation effects. The quasi-elastic model uses the Llewellyn-Smith formalism [28] with a simple dipole axial form factor, and BBBA05 vector form factors [29]. Nuclear screening effects and uncertainties are included based on the T2K 2017/8 parameterization [30] of the Valencia group’s [31, 32] Random Phase Approximation model. The Valencia model of the multi-nucleon, 2p2h, contribution to the cross section [31, 32] is used, as described in Ref. [33]. Both MINERvA [34] and NOvA [35] have shown that this model underpredicts observed event rates on carbon at relevant neutrino energies for DUNE. Modifications to the model are constructed to produce agreement with MINERvA CC-inclusive data [34], which are used in the analysis to introduce additional uncertainties on the 2p2h contribution, with energy dependent uncertainties, and extra freedom between neutrinos and antineutrinos. Uncertainties are added on scaling the 2p2h prediction from carbon to argon on electron-scattering measurements of short-range correlated (SRC) pairs taken on multiple targets [36], separately for neutrinos and antineutrinos. GENIE uses a modified version of the Rein-Sehgal (R-S) model for pion production [37]. A data-driven modification to the

GENIE model is included based on reanalyzed neutrino–deuterium bubble chamber data [38, 39]. The Deep Inelastic Scattering (DIS) model implemented in GENIE uses the Bodek-Yang parametrization [40], and GRV98 parton distribution functions [41]. Hadronization is described by the AKGY model [42], which uses the KNO scaling model [43] for invariant masses $W \leq 2.3$ GeV and PYTHIA6 [44] for invariant masses $W \geq 3$ GeV, with a smooth transition between the two models for intermediate invariant masses. Additional uncertainties developed by the NOvA Collaboration [45] to describe their resonance to DIS transition region data are also included. The final state interaction model and uncertainties available in GENIE are retained [46–48].

The cross sections include terms proportional to the lepton mass, which are significant at low energies where quasielastic processes dominate. Some of the form factors in these terms have significant uncertainties in the nuclear environment. Separate (and anticorrelated) uncertainties on the cross section ratio σ_μ/σ_e for neutrinos and antineutrinos are adopted from Ref. [49]. Additionally, some ν_e charged-current (CC) interactions occur at four-momentum transfers where ν_μ CC interactions are kinematically forbidden, and so cannot be constrained by ν_μ cross-section measurements. To reflect this, a 100% uncertainty is applied in the phase space present for ν_e but absent for ν_μ .

C. Near detector simulation and reconstruction

The near detector (ND) hall will be located 574 m downstream of the proton target and ≈ 60 m underground. The reference design for the DUNE ND system is fully described in Ref. [9], and consists of a liquid argon (LAr) time projection chamber (TPC) referred to as ND-LAr, a magnetized high-pressure gaseous argon TPC (ND-GAr), and an on-axis beam monitor (SAND). Additionally, ND-LAr and ND-GAr are designed to move perpendicular to the beam axis in order to take data at various off-axis angles (the DUNE-PRISM technique). ND-LAr is a modular detector based on the ArgonCube design [50–52], with a total active LAr volume of 105 m^3 (a LAr mass of 147 tons). ND-GAr is implemented in this analysis as a cylindrical TPC filled with a 90/10 mixture of argon and CH_4 at 10 bar, surrounded by a granular, high-performance electromagnetic calorimeter (ECal). ND-GAr sits immediately downstream of the LAr cryostat and serves as a muon spectrometer for ND-LAr [9].

Neutrino interactions are simulated in the active volume of ND-LAr. The propagation of neutrino interaction products through the ND-LAr and ND-GAr detector volumes is simulated using a Geant4-based program [53]. As pattern recognition and reconstruction software has not yet been fully developed for the ND, this analysis uses a parameterized reconstruction based on the Geant4 simulated energy deposits in active detector volumes.

Only CC-inclusive interactions originating in the LAr are considered in this analysis, with a fiducial volume (FV) which excludes 50 cm from the sides and upstream edge, and 150 cm from the downstream edge of the active region, containing a total fiducial mass of ≈ 50 t. Most muons with kinetic energies greater than 1 GeV exit ND-LAr. Energetic forward-going muons pass into ND-GAr, where their momentum and charge are reconstructed by curvature. Muon energy is reconstructed by range for tracks that stop in the LAr, and the charge cannot be determined event-by-event. Events with muons that exit the LAr active volume and do not match to a track in ND-GAr are rejected, as the muon momentum is not well reconstructed. For FHC beam running, the wrong-sign background is small and the charge is assumed to be negative for all LAr-contained muons. For RHC beam running, a Michel electron is required at the end of these stopped tracks to suppress the wrong-sign μ^- by a factor of four.

All generated muons and charged pions are evaluated as potential muon candidates. Tracks are classified as muons if their length is at least 1 m, and their mean energy deposit is less than 3 MeV/cm. The minimum length requirement imposes an effective threshold on the true muon kinetic energy of about 200 MeV, but greatly suppresses potential neutral current (NC) backgrounds with low-energy, non-interacting charged pions. Charged-current events are required to have exactly one muon candidate, and if the charge is reconstructed by curvature, it must be of the appropriate sign. Hadronic energy in the ND is reconstructed by summing all charge deposits in the LAr active volume that are not associated with the muon. To remove events where the hadronic energy is badly reconstructed due to charged particles exiting the detector, a veto region is defined as the outer 30 cm of the active volume on all sides, and events with more than 30 MeV total energy deposited in the veto region are rejected. Only a fraction of neutron kinetic energy is typically observed (24% on average with large fluctuations), resulting in poor energy reconstruction of events with energetic neutrons. The reconstructed neutrino energy, $E_\nu^{\text{rec}} = E_\mu^{\text{rec}} + E_{\text{had}}^{\text{rec}}$, is the sum of the reconstructed hadronic energy, $E_{\text{had}}^{\text{rec}}$, and the reconstructed muon energy, E_μ^{rec} . The reconstructed inelasticity, $y_{\text{rec}} = 1 - E_\mu^{\text{rec}}/E_\nu^{\text{rec}}$, is the fraction of the neutrino energy that is carried by hadrons.

D. Far detector simulation and reconstruction

The DUNE FD design consists of four separate LArTPC detector modules, each with a total LAr mass of 17 kt, installed ≈ 1.5 km underground at the Sanford Underground Research Facility (SURF) [54]. The technologies to be deployed for the four modules and their order of construction are still under investigation, so in this analysis, only the single-phase design with a horizontal drift [55] is used. In this design, signals from drift

electrons in the $13.3 \times 12.0 \times 57.5$ m³ active volume are read out by ≈ 5 mm spaced wires in anode readout planes. Scintillation light produced at the time of the neutrino interaction is detected and used to reconstruct the start time of the electron drift. We have developed a full simulation chain, which generates neutrino events in a Geant4 model of the FD geometry and simulates the electronics readout. We have developed a reconstruction package to calculate efficiencies and reconstructed neutrino energy estimators for the four CC-inclusive FD samples used in the analysis (ν_μ -like FHC, ν_e -like FHC, $\bar{\nu}_\mu$ -like RHC and $\bar{\nu}_e$ -like RHC).

The electronics response to the ionization electrons and scintillation light is simulated in the wire planes and photon detectors, respectively. Algorithms are applied to remove the impact of the LArTPC electric field and electronics response from the raw detector signal to identify hits, and to cluster hits that may be grouped together due to proximity in time and space. Clusters from different wire planes are matched to form high-level objects such as tracks and showers using the Pandora toolkit [56, 57]. Event classification is carried out through image recognition techniques using a convolutional neural network [58] which classifies events as $\langle \bar{\nu}_\mu \rangle$ -CC, $\langle \bar{\nu}_e \rangle$ -CC, $\langle \bar{\nu}_\tau \rangle$ -CC, and NC. The $\langle \bar{\nu}_e \rangle$ and $\langle \bar{\nu}_\mu \rangle$ efficiencies in both beam modes exceed 90% in the flux peak.

The neutrino energy for $\langle \bar{\nu}_\mu \rangle$ -CC ($\langle \bar{\nu}_e \rangle$ -CC) events is estimated by the sum of the energy of the longest reconstructed track (highest energy reconstructed electromagnetic shower) and the hadronic energy. For both event types, the hadronic energy is estimated from the charge of reconstructed hits that are not in the primary track or shower, and corrections are applied to each hit charge for recombination and electron lifetime effects. For $\langle \bar{\nu}_\mu \rangle$ -CC events, the energy of the longest track is estimated by range if the track is contained or by multiple Coulomb scattering if it is exiting. For 0.5–4 GeV neutrino energies, the observed neutrino energy resolution is 15–20%. The muon energy resolution is 4% for contained tracks and 18% for exiting tracks. The electron energy resolution is approximately $4\% \oplus 9\%/\sqrt{E}$. The hadronic energy resolution is 34%.

E. Detector systematics

Detector effects impact the event selection efficiency as well as the reconstruction of neutrino energy and inelasticity (the variables used in the oscillation fits). The main sources of detector systematic uncertainties are limitations of the expected calibration and modeling of particles in the detector. Important differences between the ND and FD LArTPC design, size, detector environment, and calibration strategy, lead to uncertainties that do not fully correlate between the two detectors. The degree of correlation is under active study, but in this analysis they are treated as being completely uncorrelated. Detailed simulations of detector effects are under devel-

opment. In this analysis, uncertainties on the energy scale, energy resolution, particle responses, and detector acceptance are included to encapsulate these effects. The absolute scale uncertainties shift the reconstructed energy distributions, while the resolution uncertainties narrow or broaden them.

An uncertainty on the overall energy scale is included in the analysis presented here, as well as particle energy scale and resolution uncertainties that are separate and uncorrelated between four particle classes: muons, charged hadrons, neutrons, and electromagnetic showers. In the ND, muons reconstructed by range in LAr and by curvature in the ND-GAr are treated separately and assigned uncorrelated uncertainties. For each class of particle, uncertainties on the energy scale are introduced as a function of the reconstructed particle energy, E , with a constant term, a term proportional to \sqrt{E} , and a term proportional to $1/\sqrt{E}$. A 10% uncertainty on the energy resolution is also included, and treated as uncorrelated between the four particle classes. The parameters produce a shift to the kinematic variables in an event, as opposed to simply assigning a weight to each simulated event. The scale of the uncertainties is motivated by what has been achieved in recent experiments, including calorimetric based approaches (NOvA, MINERvA) and LArTPCs (LArIAT, MicroBooNE, ArgoNeut).

In addition to impacting energy reconstruction, the E-field model also affects the definition of the FD FV, which is sensitive to electron drift. An additional 1% uncertainty is therefore included on the total fiducial mass, which is conservatively treated as uncorrelated between the $\langle \bar{\nu}_\mu \rangle$ and $\langle \bar{\nu}_e \rangle$ samples due to the potential distortion caused by large electromagnetic showers in the electron sample.

The FD is sufficiently large that acceptance is not expected to vary significantly as a function of event kinematics. However, the ND acceptance does vary as a function of both muon and hadronic kinematics due to various containment criteria. Uncertainties are evaluated on the muon and hadron acceptance at the ND based on the change in the acceptance as a function of muon kinematics and true hadronic energy.

F. Sensitivity Methods

Systematics are implemented in the analysis using one-dimensional response functions for each analysis bin, and oscillation weights are calculated exactly, in fine (50 MeV) bins of true neutrino energy. For a given set of inputs — flux, oscillation parameters, cross sections, detector energy response matrices, and detector efficiency — an expected event rate can be produced. Minimization is performed using the MINUIT [59] package.

Oscillation sensitivities are obtained by simultaneously fitting the ν_μ -like FHC, ν_e -like FHC, $\bar{\nu}_\mu$ -like RHC and $\bar{\nu}_e$ -like RHC FD spectra along with the ν_μ FHC and $\bar{\nu}_\mu$ RHC samples from the ND. In the studies, all oscillation

Parameter	Central value	Relative uncertainty
θ_{12}	0.5903	2.3%
θ_{23} (NO)	0.866	4.1%
θ_{23} (IO)	0.869	4.0%
θ_{13} (NO)	0.150	1.5%
θ_{13} (IO)	0.151	1.5%
Δm_{21}^2	7.39×10^{-5} eV ²	2.8%
Δm_{32}^2 (NO)	2.451×10^{-3} eV ²	1.3%
Δm_{32}^2 (IO)	-2.512×10^{-3} eV ²	1.3%
ρ	2.848 g cm ⁻³	2%
δ_{CP} (NO)	-2.53 (rad.)	—
δ_{CP} (IO)	-1.33 (rad.)	—

TABLE I. Central value and relative uncertainty of neutrino oscillation parameters from a global fit [60, 61] to neutrino oscillation data. The matter density is taken from Ref. [62]. Because the probability distributions are somewhat non-Gaussian (particularly for θ_{23}), the relative uncertainty is computed using 1/6 of the 3σ allowed range from the fit, rather than 1/2 of the 1σ range. For θ_{23} , θ_{13} , and Δm_{31}^2 , the best-fit values and uncertainties depend on whether NO or IO is assumed. The best fit for δ_{CP} is used as a test point in the analysis, but no uncertainty is assigned.

parameters shown in Table I are allowed to vary. Gaussian penalty terms (taken from Table I) are applied to θ_{12} , Δm_{21}^2 , and the matter density, ρ , of the Earth along the DUNE baseline [62]. Some studies presented in this work include a Gaussian penalty term on θ_{13} (also taken from NuFIT 4.0, given in Table I), which is precisely measured by experiments sensitive to reactor antineutrino disappearance [63–65]. The remaining parameters, $\sin^2 \theta_{23}$, Δm_{32}^2 , and δ_{CP} are allowed to vary freely, with no penalty terms. The penalty terms are treated as uncorrelated with each other, and uncorrelated with other parameters.

Flux, cross-section, and FD detector parameters are allowed to vary in the fit, but are constrained by a penalty term corresponding to the prior uncertainty. ND detector uncertainties are included via a covariance matrix based on the shape difference between ND prediction and the “data” (which comes from the simulation in this sensitivity study). The covariance matrix is constructed with a throwing technique. For each “throw”, all ND energy scale, resolution, and acceptance parameters are simultaneously thrown according to their respective uncertainties, and the modified prediction is produced by varying the relevant quantities away from the nominal prediction according to the thrown parameter values. The bin-to-bin covariance is determined by comparing the resulting spectra with the nominal prediction, in the same binning as is used in the oscillation sensitivity analysis.

The compatibility of a particular oscillation hypothesis with both ND and FD data is evaluated using the

Parameter	Prior	Range
$\sin^2 \theta_{23}$	Uniform	[0.4; 0.6]
$ \Delta m_{32}^2 $ ($\times 10^{-3}$ eV ²)	Uniform	[2.3; 2.7]
δ_{CP} (π)	Uniform	[-1; 1]
θ_{13} (NO)	Gaussian	0.1503 ± 0.0023 (rad.)
θ_{13} (IO)	Gaussian	0.1510 ± 0.0023 (rad.)

TABLE II. Treatment of the oscillation parameters for the simulated data set studies. The value and uncertainty for θ_{13} in both NO and IO used in the analysis come from NuFIT 4.0 [60, 61].

standard Poisson log-likelihood ratio [66]:

$$\begin{aligned}
\chi^2(\vec{\vartheta}, \vec{x}) &= -2 \log \mathcal{L}(\vec{\vartheta}, \vec{x}) \\
&= 2 \sum_i^{N_{\text{bins}}} \left[M_i(\vec{\vartheta}, \vec{x}) - D_i + D_i \ln \left(\frac{D_i}{M_i(\vec{\vartheta}, \vec{x})} \right) \right] \\
&+ \sum_j^{N_{\text{sys}}} \left[\frac{\Delta x_j}{\sigma_j} \right]^2 \\
&+ \sum_k^{N_{\text{bins}}^{\text{ND}}} \sum_l^{N_{\text{bins}}^{\text{ND}}} (M_k(\vec{x}) - D_k) V_{kl}^{-1} (M_l(\vec{x}) - D_l),
\end{aligned} \tag{2}$$

where $\vec{\vartheta}$ and \vec{x} are the vector of oscillation parameter and nuisance parameter values, respectively; $M_i(\vec{\vartheta}, \vec{x})$ and D_i are the MC expectation and fake data in the i th reconstructed bin (summed over all selected samples), with the oscillation parameters neglected for the ND; Δx_j and σ_j are the difference between the nominal and current value, and the prior uncertainty on the j th nuisance parameter; and V_{kl} is the covariance matrix between ND bins described previously. To protect against false minima, all fits are repeated starting at four different δ_{CP} values ($-\pi$, $-\pi/2$, 0 , $\pi/2$), in both mass orderings, and in both $\sin^2 \theta_{23}$ octants, and the lowest obtained χ^2 value is taken as the true minimum.

Two approaches are used for the sensitivity studies presented in this work. Asimov studies [67] are carried out (in Section III) in which the fake (Asimov) dataset is the same as the nominal MC. In these, the true value of all systematic uncertainties and oscillation parameters are set to their nominal value (see Table I) except the parameters of interest, which are set to a test point. Then a fit is carried out in which all parameters can vary, constrained by their prior uncertainty where applicable. For the smallest exposures investigated with an Asimov study in this work, all samples have at least 100 events, satisfying the Gaussian approximation inherent in the Asimov method. Toy throw studies are performed (in Sections IV and V) in which an ensemble of systematic, oscillation parameter and statistical throws are made. Systematic throws are made according to their prior Gaussian uncertainties, oscillation parameters are randomly chosen as described in Table II, and Poisson fluctuations are then


applied to all analysis bins, based on the mean event count for each bin after the systematic adjustments have been applied. For each throw in the ensemble, the test statistic is minimized, and the best-fit value of all parameters is determined. The expected resolution for parameters of interest are then determined from the spread in the distribution of their post-fit values.

Asimov studies are computationally efficient, and for Gaussian parameters and uncertainties, give a good sense of the median sensitivity of an experiment. Toy throwing studies are computationally expensive, fully explore the parameter space, and make fewer assumptions about the behavior of parameters and uncertainties.


G. Near and far detector samples and statistics

In this work, the sensitivity as a function of FD exposure is explored and results are reported in terms of kt-MW-yr, which does not assume any specific FD or beam intensity staging scenario. However, the ND used in this analysis (ND-LAr with a downstream muon spectrometer) is assumed not to be staged, and as such the ND sample size corresponding to a particular FD exposure will vary based on the staging scenario. The nominal staging scenario from Ref. [21] is therefore retained for the purpose of normalizing the ND samples at each FD exposure. In that scenario, a 7 year exposure corresponds to 336 kt-MW-yr at the FD, and 480 t-MW-yr at the ND, summed over both beam modes. The ND statistics used in this analysis are scaled assuming this ratio throughout, using the same fraction of exposure in each beam mode as used at the FD. The ND samples used in this analysis are relatively quickly systematics limited in both beam modes, and so these approximations are unlikely to have a significant impact on the results.

The oscillation analysis presented here includes two CC-inclusive samples originating in the ND-LAr FV, an FHC ν_μ and an RHC $\bar{\nu}_\mu$ sample. These samples are both binned in two dimensions, as a function of reconstructed neutrino energy and inelasticity, $y_{\text{rec}} = 1 - E_\mu^{\text{rec}}/E_\nu^{\text{rec}}$. The sample distributions for both FHC and RHC are shown in Figure 1 for an exposure of 105 t-MW-yr, corresponding to 100 kt-MW-yr at the far detector with the assumptions stated above. The size of the systematic uncertainty bands from all of the flux, cross-section and ND detector systematics used in the analysis and described above are shown, as well as the postfit uncertainty bands obtained by performing an Asimov fit to the ND data. It is clear that even after a relatively small exposure of 105 t-MW-yr, the ND samples are very high statistics, and are systematics limited in the binning used in the analysis. Backgrounds in the $(\bar{\nu}_\mu)$ -CC samples are also shown in Figure 1. NC backgrounds are predominantly from NC π^\pm production where the pion leaves a long track and does not shower. Wrong-sign contamination in the beam is a background where the charge of the muon is not reconstructed, which particularly affects low reconstructed




(a) FHC




(b) RHC

FIG. 1. ND samples in both FHC and RHC, shown in the reconstructed neutrino energy and reconstructed inelasticity binning (y_{rec}) used in the analysis, for a 105 t-MW-yr exposure (equivalent to a 100 kt-MW-yr exposure at the FD), with an equal split between FHC and RHC. The size of the systematic uncertainty bands from all of the flux, cross-section and ND detector systematics used in the analysis are shown, as well as the postfit uncertainty bands obtained by performing an Asimov fit to the ND data. NC backgrounds and wrong-sign contributions to the total event rate are also shown. Statistical uncertainties are too small to be visible on this plot scale.

neutrino energies in RHC. The wrong-sign background is also larger at high reconstructed inelasticity, y_{rec} , due to the kinematics of neutrino and antineutrino scattering.



(a) FHC



(b) RHC

FIG. 2. Reconstructed energy distribution of selected CC $\bar{\nu}_e$ -like events in the FD, for a 50 kt-MW-yr exposure in both FHC and RHC beam modes, for a total 100 kt-MW-yr exposure. The plots are shown for NO, all other oscillation parameters are set to their NuFIT 4.0 best-fit values (see Table I). The size of the systematic uncertainty bands from all of the flux, cross-section and FD detector systematics used in the analysis are shown, as well as the postfit uncertainty bands with parameters constrained by ND data. Backgrounds are also shown, the largest contribution comes from intrinsic $\bar{\nu}_e$ contamination in the beam, although NC and other flavors, $\bar{\nu}_\mu + \bar{\nu}_\tau$, also contribute.

The expected FD FHC ν_e and RHC $\bar{\nu}_e$ samples are shown in Figure 2 for a 100 kt-MW-yr total FD exposure, split equally between FHC and RHC beam modes. The systematic uncertainty bands with and without the ND constraint applied are shown, as well as the back-

ground contributions. There are contributions from both ν_e and $\bar{\nu}_e$ in both beam modes. The NC, intrinsic beam $\bar{\nu}_e$, and wrong flavor contamination is also shown; the largest background comes from the intrinsic $\bar{\nu}_e$ beam contribution in both modes. After a 50 kt-MW-yr exposure in FHC, the ν_e sample statistical uncertainty is close to the systematic uncertainty before the ND constraint, although is still clearly statistics limited when the ND constraint is applied. The $\bar{\nu}_e$ sample is still strongly statistics limited after 50 kt-MW-yr exposure in RHC. The difference is largely due to the difference in the ν_e and $\bar{\nu}_e$ cross sections.


The expected FD FHC ν_μ and RHC $\bar{\nu}_\mu$ samples are shown in Figure 3 for a 100 kt-MW-yr total FD exposure, split equally between FHC and RHC beam modes. The systematic uncertainty bands with and without the ND constraint applied are shown, as well as the background contributions. Although the wrong-sign ν_μ contribution to the RHC $\bar{\nu}_\mu$ sample is shown separately, it still provides useful information for constraining the oscillation parameters and is included in the analysis. The statistics are much higher than in Figure 2; the statistical uncertainty on the ν_μ FHC sample is smaller than the systematic uncertainty band for some regions of phase space, even after the ND constraint is applied, although the statistical uncertainty is larger than the constrained systematic uncertainty in the “dip” region, around 2.5 GeV, which is likely to have the most impact on the analysis. The statistical uncertainty on the $\bar{\nu}_\mu$ RHC sample is larger, again due to the smaller $\bar{\nu}_\mu$ (than ν_μ) cross section and lower fluxes in RHC running. The statistical uncertainty around the 2.5 GeV dip region is significantly larger than the systematic uncertainty band, although as for the FHC ν_μ sample, the statistical uncertainty is smaller than the systematics for some regions of the parameter space.

Events with a reconstructed neutrino energy of less than 0.5 GeV (which are shown in Figures 1, 2 and 3) or a reconstructed neutrino energy greater than 10 GeV are not included in the analysis for any of the FD or ND samples.


III. RUN PLAN OPTIMIZATION

In previous DUNE sensitivity studies [21], equal running times in FHC and RHC were assumed, based on early sensitivity estimates for different scenarios. In this section, the dependence of the median CPV and mass ordering significances are studied, for different fractions of time spent in each beam mode, using the full analysis framework described in Section II.

Figure 4 shows DUNE’s Asimov sensitivity to CPV for a total 100 kt-MW-yr far detector exposure, with different fractions of FHC and RHC running, at the NuFIT 4.0 best fit value in both NO and IO (see Table I), shown with and without a penalty on θ_{13} applied. For each point tested, all oscillation and nuisance param-



(a) FHC



(b) RHC

FIG. 3. Reconstructed energy distribution of selected CC $(\bar{\nu}_\mu)$ -like events in the FD, for 50 kt-MW-yr exposure in both FHC and RHC beam modes, for a total 100 kt-MW-yr exposure. The plots are shown for NO, all other oscillation parameters are set to their NuFIT 4.0 best-fit values (see Table I). The size of the systematic uncertainty bands from all of the flux, cross-section and ND detector systematics used in the analysis are shown, as well as the postfit uncertainty bands with parameters constrained by ND data. NC backgrounds and wrong-sign contributions to the event rate are also shown.

ters are allowed to vary, and three fits are carried out, two where δ_{CP} is set to the CP-conserving values $\delta_{\text{CP}} = 0$ and $\delta_{\text{CP}} = \pm\pi$, the minimum of which is the CP-conserving best-fit value, and another where δ_{CP} is allowed to vary. The difference in the best-fit χ^2 values is calculated:

$$\Delta\chi_{\text{CPV}}^2 = \min \{ \chi_{\delta_{\text{CP}}=0}^2, \chi_{\delta_{\text{CP}}=\pm\pi}^2 \} - \chi_{\text{CPV}}^2, \quad (3)$$

and the square root of the difference is used as the figure of merit on the y-axis in Figure 4. There are some caveats associated with this figure of merit, which are discussed in

Section IV. A 100 kt-MW-yr exposure is shown as it was identified in Ref [21] as the exposure at which DUNE's median CPV significance exceeds 3σ at $\delta_{\text{CP}} = \pm\pi/2$, an important milestone in DUNE's physics program (with equal beam mode running).

Figure 4 shows that when the reactor constraint on θ_{13} is included, the sensitivity to CPV can be increased in some regions of δ_{CP} parameter space with more FHC than RHC running. However, this degrades the sensitivity in other regions, most notably for $\delta_{\text{CP}} > 0$ regardless of the true mass ordering. This is due to a degeneracy between δ_{CP} and the octant of $\sin^2 2\theta_{23}$ because both parameters impact the rate of ν_e appearance. The degeneracy is resolved by including antineutrino data; the octant of $\sin^2 2\theta_{23}$ affects the rate of ν_e and $\bar{\nu}_e$ appearance in the same way, but the effect of δ_{CP} is reversed for antineutrinos.

For regions of phase space where the octant degeneracy does not affect the result (e.g., $\sin^2 \theta_{23} \approx 0.5$), there is no degradation in the sensitivity, and enhanced FHC running increases the sensitivity for all values of δ_{CP} . Increasing the fraction of RHC decreases the sensitivity for the entire δ_{CP} range when the reactor θ_{13} constraint is included, relative to equal beam mode running. This is due to the lower statistics of the $\bar{\nu}_e$ sample (see Figure 2) because of the reduced antineutrino flux and cross section. For short exposures, DUNE will not have a competitive independent measurement of θ_{13} , so the main analysis will include the reactor θ_{13} constraint. Nonetheless, it is instructive to look at the results without the penalty applied. In this case, the sensitivity is severely degraded (as expected) for 100% running in either beam mode.

Figure 5 shows DUNE's Asimov sensitivity to the mass ordering for a total 24 kt-MW-yr far detector exposure, with different fractions of FHC and RHC running, and the same four true oscillation parameter sets. A 24 kt-MW-yr exposure is used in Figure 5 as it is around the exposure at which DUNE's median mass ordering significance exceeds 5σ for some values of δ_{CP} [21]. For each point tested, all oscillation and nuisance parameters are allowed to vary, and two fits are carried out, one using each ordering. The difference in the best-fit χ^2 values is calculated:

$$\Delta\chi_{\text{MO}}^2 = \chi_{\text{IO}}^2 - \chi_{\text{NO}}^2, \quad (4)$$

and the square root of the difference is used as the figure of merit on the y-axis in Figure 5. There are some caveats associated with this figure of merit, which are discussed in Section V.

It is clear from Figure 5 that the mass ordering sensitivity has a strong dependence on the fraction of running in each beam mode. As in the CPV case, the effect is very different with and without the reactor θ_{13} constraint included. If the true ordering is normal and the reactor θ_{13} penalty is applied, the sensitivity increases significantly with increasing FHC running, with a full 1σ increase in the sensitivity between equal beam running and 100% FHC for most values of δ_{CP} . Conversely, if the ordering




FIG. 4. The Asimov CPV sensitivity as a function of the true value of δ_{CP} , for a total exposure of 100 kt-MW-yr with different fractions of FHC and RHC running, with and without a θ_{13} penalty applied in the fit. Results are shown for both true normal and inverted ordering, with the true oscillation parameter values set to the NuFit 4.0 best fit point in each ordering (see Table I).

is inverted, 100% FHC running would degrade the sensitivity by $\geq 1\sigma$ for all values of δ_{CP} at the NuFIT 4.0 best fit point. Overall, the sensitivity to the inverted ordering is improved by a more equal split between the beam modes. It is clear that 100% RHC running gives poor sensitivity for all values tested.

Without the reactor θ_{13} constraint, the greatest sensitivity is obtained with close to an equal split of FHC and RHC running, and the sensitivity is significantly reduced with 100% FHC running. This is because of a degeneracy between the effect of θ_{13} and the mass ordering on the rate of ν_e appearance in FHC mode. If the mass ordering is normal, the ν_e rate in FHC will be enhanced; without

the reactor constraint, this excess can be accommodated by increasing the value of θ_{13} .

For comparison, Figure 6 shows the Asimov CPV and mass ordering sensitivities, with and without the reactor θ_{13} constraint included, for true normal ordering only, for a large exposure of 336 kt-MW-yr, with different fractions of FHC and RHC running. At large exposures, running with strongly enhanced FHC no longer improves the sensitivity over equal beam mode running, with or without the θ_{13} penalty applied, for either CPV or mass ordering determination. This can be understood because the enhancement to the statistics that enhanced FHC brings is no longer as important to the sensitivity, and




FIG. 5. The Asimov mass ordering sensitivity as a function of the true value of δ_{CP} , for a total exposure of 24 kt-MW-yr with different fractions of FHC and RHC running, with and without a θ_{13} penalty applied in the fit. Results are shown for both true normal and inverted ordering, with the true oscillation parameter values set to the NuFIT 4.0 best fit point in each ordering (see Table I).

DUNE is able to place a constraint on the value of θ_{13} with its own data.

Overall, the sensitivity to CPV and the mass ordering is dependent on the division of running time between FHC and RHC, but a choice that increases the sensitivity in some region of parameter space can severely decrease the sensitivity in other regions. If there is strong reason to favor, for example, normal over inverted ordering when DUNE starts to take data, Figure 5 shows that this could be more rapidly verified by running with more FHC data than RHC data, as the reactor θ_{13} constraint will be used in the main low exposure analysis. However, if this choice is wrong, this might cause DUNE to take longer

to reach the same significance. Clearly this is an important consideration which should be revisited shortly before DUNE begins to collect data. Similarly, the CPV sensitivity shown in Figure 4 might be optimized if there is a strong reason to favor gaining sensitivity for $\delta_{CP} > 0$ or $\delta_{CP} < 0$, at a cost of reducing the sensitivity to CPV if δ_{CP} has the other sign. But, it is clear from Figures 4 and 5 that equal running in FHC and RHC gives a close to optimal sensitivity across all of the parameter space, and as such is a reasonable *a priori* choice of run plan for studies of the DUNE sensitivity. Additionally, it is clear from Figure 6 that the improvement in the sensitivity with unequal beam running is a feature at low exposures,




FIG. 6. The Asimov CPV and mass ordering sensitivities as a function of the true value of δ_{CP} , for a total exposure of 336 kt-MW-yr with different fractions of FHC and RHC running, with and without a θ_{13} penalty applied in the fit. Results are shown for both true normal ordering only, with the true oscillation parameter values set to the NuFIT 4.0 NO best fit point (see Table I).

but not at high exposures, particularly because at high exposures when DUNE is able to constrain all the oscillation parameters with precision [21], there is a stronger motivation to run a DUNE-only analysis, without relying on the reactor θ_{13} measurement.

IV. CP VIOLATION SENSITIVITY

In this section, CPV sensitivity results are presented. For simplicity, only true NO will be shown unless explicitly stated. In all cases, a joint ND+FD fit is performed, and a θ_{13} penalty is always applied to incorporate the re-

actor measurement, as described in Section II. An equal split between FHC and RHC running is assumed based on the results obtained in Section III. Asimov sensitivities, as shown in Section III, are instructive but do not give information on how the expected sensitivity may vary with statistical or systematic uncertainties, or for variations in the other oscillation parameters of interest.

Figure 7 shows the significance with which CPV ($\delta_{CP} \neq \{0, \pm\pi\}$) can be observed for both NO and IO, for exposures of 66 and 100 kt-MW-yr. The sensitivity metric used is the square root of the difference between the best fit χ^2 values obtained for a CP-conserving fit and one where δ_{CP} is allowed to vary, as shown in Equ-




FIG. 7. Significance of the DUNE determination of CP-violation ($\delta_{\text{CP}} \neq \{0, \pm\pi\}$) as a function of the true value of δ_{CP} , for 66 kt-MW-yr (blue) and 100 kt-MW-yr (orange) exposures, for normal (top) and inverted (bottom) orderings. The width of the transparent bands cover 68% of fits in which random throws are used to simulate systematic, oscillation parameter and statistical variations, with independent fits performed for each throw constrained by prior uncertainties. The solid lines show the median significance.

tion 3, which is calculated for each throw of the systematic, other oscillation parameters and statistics.

The sensitivity shown in Figure 7 has a characteristic double peak structure because the significance of a CPV measurement decreases around CP-conserving values. The systematic and statistical variations mean that all throws have $\Delta\chi_{\text{CPV}}^2 \geq 0$, and therefore neither the median significance nor the band showing the central 68% of throws reach exactly 0 at CP-conserving values. This

is entirely expected, it simply means that random variations in the data will cause us to obtain a 1σ measurement of CPV $\approx 32\%$ of the time for CP-conserving values. Median significances are slightly higher for IO than for NO, and by exposures of 100 kt-MW-yr, the median significance exceeds 3σ for the maximal CP-violating values of $\pm\pi/2$. This presentation of the CPV sensitivity was followed in Ref. [21], and is very informative at high exposures. Around CP-conserving values ($\delta_{\text{CP}} = \{0, \pm\pi\}$), the distribution of the sensitivity metric $\sqrt{\Delta\chi_{\text{CPV}}^2}$ is non-Gaussian for all exposures. Additionally, at lower exposures, as shown in Figure 7, the distribution of $\sqrt{\Delta\chi_{\text{CPV}}^2}$ around maximally CP-violating values of $\delta_{\text{CP}} = \pm\pi/2$ is increasingly non-Gaussian, making the spread in sensitivity harder to interpret with this presentation.

The CPV significance in Figure 7 (and previously in Ref. [21]) is calculated using constant $\Delta\chi^2$ critical values, where $\Delta\chi_{\text{CPV}}^2 \leq 1, 4, 9$ corresponds to a significance of 1, 2 and 3σ for one degree of freedom. This assumption holds when Wilks' theorem can be applied [68], but can lead to incorrect coverage where it cannot. It is known to break down for low-statistics samples, around physical boundaries, in the case of cyclic parameters, and where there are significant degeneracies. It is likely that a constant $\Delta\chi^2$ treatment will break down for δ_{CP} , where all of these issues apply, as has indeed been shown by the T2K Collaboration [13].

The Feldman-Cousins method [69] is a brute force numerical method to calculate confidence intervals with correct coverage. A large number of toy experiments are produced, where the parameter(s) of interest (here δ_{CP}) is set to a desired true value, all other systematic and oscillation parameters are thrown, as described in Section II, and a statistical throw is made, for the two ND samples and four FD sample used in the analysis. Then two fits are performed, one where the parameter(s) of interest are fixed to the true value, and another where the test statistic is minimized with respect to the parameter(s) of interest. In both fits, all other parameters are allowed to vary. For each throw, the profile likelihood ratio $\Delta\chi_{\text{FC}}^2$ is calculated using the minimum χ^2 values for those two fits, as in Equation 5.

$$\Delta\chi_{\text{FC}}^2 = \chi^2(\theta_{\text{true}}) - \min_{\theta} \chi^2(\theta) \quad (5)$$

The distribution of these throws is used to calculate the $\Delta\chi_{\text{FC}}^2$ value that gives the desired coverage, with the appropriate fraction of toys above/below the calculated value. These are labelled *critical values*, and are denoted $\Delta\chi_c^2$. A distribution of $\Delta\chi_{\text{FC}}^2$ values is shown in Figure 8 for an example ND+FD analysis with a 100kt-MW-yr exposure at the far detector, equal FHC and RHC run fractions, and the reactor θ_{13} constraint applied. In Figure 8, the $\Delta\chi_c^2$ values corresponding to for 68.27% (1σ), 90%, 95.45% (2σ) and 99.73% (3σ) of the throws are indicated. The $\Delta\chi_c^2$ values were only calculated up to the 3σ level due to the very large number of throws required for higher confidence levels.




FIG. 8. Distribution of $\Delta\chi_{\text{FC}}^2$ values, calculated using Equation 5, for a large number of throws with true $\delta_{\text{CP}} = 0$, and a 100 kt-MW-yr exposure. The $\Delta\chi_{\text{FC}}^2$ values (vertical lines) obtained using the Feldman-Cousins method show the $\Delta\chi_{\text{FC}}^2$ value below which 68.27% (1σ), 90%, 95.45% (2σ) and 99.73% (3σ) of throws reside, with the calculated values given in the legend. The number of throws used is also given.

An uncertainty on the value of $\Delta\chi_c^2$ obtained from the toy throw distribution (e.g., Figure 8), is obtained using a bootstrap rethrowing method [70]. The empirical PDF obtained from the throws is treated as the true PDF, and B independent samples of size n are drawn from it, where n is the total number of throws used to build the empirical PDF. Each throw can be drawn multiple times in this method, so the ensemble of throws is different in each sample. Then, the standard deviation $s_{\hat{\vartheta}}$, on the $\Delta\chi_c^2$ values of interest, ϑ , are calculated for each of the B samples using:

$$s_{\hat{\vartheta}} = \sqrt{\frac{1}{B-1} \sum_{i=0}^B (\vartheta_i^* - \bar{\vartheta}^*)^2}, \quad (6)$$

where ϑ_i^* denotes the calculated $\Delta\chi_c^2$ value of interest for each of the samples, and $\bar{\vartheta}^*$ is their average value.

Figure 9 shows the evolution of the $\Delta\chi_c^2$ values as a function of exposure for $\delta_{\text{CP}} = 0$, the relevant value for CPV sensitivity, for an ND+FD analysis with equal FHC and RHC running and the reactor θ_{13} constraint applied. For all significance levels tested, the $\Delta\chi_c^2$ rise quickly as a function of exposure, and stabilize at values slightly higher than those suggested by the constant $\Delta\chi^2$ method by exposures of ≈ 100 kt-MW-yr. The initial rise in the $\Delta\chi_c^2$ values is due to the low statistics at those exposures. Overall, this implies that the CPV significance is slightly weaker than what would be inferred from $\sigma = \sqrt{\Delta\chi_{\text{CPV}}^2}$, as used for example in Figure 7. Crucially, there is no constant increase in the $\Delta\chi_c^2$ values over time as has been reported by the T2K Collaboration [13]. Details on the number of toy throws used at each point of Figure 9 are




FIG. 9. The $\Delta\chi_c^2$ values corresponding to 68.27% (1σ), 90%, 95.45% (2σ) and 99.73% (3σ) of throws, shown for true $\delta_{\text{CP}} = 0$, as a function of exposure. A linear x -axis scale is used to highlight the stability of $\Delta\chi_c^2$ values for large exposures. The uncertainty on the $\Delta\chi_c^2$ values is obtained using Equation 6, and is indicated as the shaded line. To guide the eye, horizontal dashed lines are included which indicate the 1σ , 90%, 2σ and 3σ $\Delta\chi^2$ values assumed using the constant- $\Delta\chi^2$ method, with one degree of freedom. The distribution of throws used produced to calculate the $\Delta\chi_c^2$ values shown are given in Figure 19.

given in the Appendix, and the toy throw distributions from which the $\Delta\chi_c^2$ values and their uncertainties were calculated are shown in Figure 19.

As δ_{CP} is a cyclical parameter, with physical boundaries at $\pm\pi$, it is interesting to see how the $\Delta\chi_c^2$ values evolve as a function of it. Figure 10 shows the $\Delta\chi_c^2$ as a function of true δ_{CP} , for an ND+FD analysis with equal FHC and RHC running including the reactor θ_{13} constraint, for both 100 kt-MW-yr and 336 kt-MW-yr exposures. There is a noticeable, although not large, depression in the $\Delta\chi_c^2$ values at $\delta_{\text{CP}} = \pm\pi/2$ for all significance levels considered. This effect is larger at the lower, 100 kt-MW-yr, exposure, and is larger at higher significance levels. It is also clear from Figure 10 that the $\Delta\chi_c^2$ behaviour is very similar at $\delta_{\text{CP}} = \pm\pi/2$ as at $\delta_{\text{CP}} = 0$. Although the $\Delta\chi_c^2$ values are relevant for CPV sensitivity, this evolution of the $\Delta\chi_c^2$ values with δ_{CP} will be important for estimating DUNE's δ_{CP} resolution. Details on the number of toy throws used at each point of Figure 10 are given in the Appendix, and the toy throw distributions used to calculate the $\Delta\chi_c^2$ values and uncertainties are shown for the 100 kt-MW-yr (336 kt-MW-yr) test points in Figure 20 (Figure 21).

DUNE's CPV sensitivity is calculated using Equation 3 from an ensemble of throws of all systematic, other oscillation parameters and statistics. Figure 11 shows the fraction of throws for which DUNE would observe a CPV significance above a discrete threshold, as a function of




FIG. 10. The $\Delta\chi^2_c$ values corresponding to 68.27% (1σ), 90%, 95.45% (2σ) and 99.73% (3σ) of throws, shown as a function of true δ_{CP} , for exposures of 100 kt-MW-yr and 336 kt-MW-yr. The uncertainty on the $\Delta\chi^2_c$ values is obtained using Equation 6, and is indicated as the shaded line. To guide the eye, horizontal dashed lines are included which indicate the 1σ , 90%, 2σ and 3σ $\Delta\chi^2_{CPV}$ values assumed using the constant- $\Delta\chi^2$ method, with one degree of freedom. The distribution of throws used produced to calculate the $\Delta\chi^2_c$ values shown are given in Figure 20 (Figure 21) for 100 kt-MW-yr (336 kt-MW-yr).

the true value of δ_{CP} , for $1-3\sigma$ significances and for a variety of exposures. The shaded histograms show the complete treatment including FC, while the dashed histograms show the constant- $\Delta\chi^2$ treatment, to show the deviation from Wilks' theorem, and to facilitate comparison at higher significances where the FC treatment becomes computationally prohibitive.

The point at which the median significance (50% of throws) passes different significance thresholds can be

easily read from the figures, and can be compared with those shown in Figure 7. The same double peak structure seen in Figure 7 can be observed. The median significance for measuring CPV exceeds 3σ after ≈ 100 kt-MW-yr, but a significant fraction of throws exceed 3σ at shorter exposures. For a 336 kt-MW-yr exposure, the fraction of throws for which the significance is less than 3σ at maximal values of δ_{CP} is very small. In general, the effect of the Feldman-Cousins correction is to reduce the fraction of toy throws that cross each significance threshold (with respect to the constant- $\Delta\chi^2$ result), by a maximum of $\approx 10\%$, but the exact fraction changes as a function of true δ_{CP} value and exposure. An exception to this general trend is the 3σ behaviour at 24 kt-MW-yr, the lowest exposure shown, where the significance increases. This is due to fall in the 3σ $\Delta\chi^2_c$ value towards the lowest exposures observed in Figure 9. The number of throws carried out at each exposure is indicated on each plot. The number of throws decreases as a function of exposure because fixed computing resources were used for each configuration, and the time for the ensemble of fits carried out for each throw to complete increases slightly with exposure. The final 336 kt-MW-yr exposure has more throws because it was generated for the analysis presented in Ref. [21], where more than one projection was considered — requiring more throws to sample the space.

Figure 12 shows the fraction of throws which exceed different significance thresholds at the maximal δ_{CP} violation value of $\delta_{CP} = -\pi/2$, and for 50% of δ_{CP} values as a function of exposure, with and without FC corrections, for $1-3\sigma$ significance values. Figure 12 was produced using the same throws used for Figure 11, with additional points from higher exposures used in Ref. [21], but not shown in Figure 11 (646 kt-MW-yr and 1104 kt-MW-yr). After ≈ 200 kt-MW-yr, the median significance (including FC correction) for 50% of the δ_{CP} range is greater than 3σ . It is clear from Figure 12 that the effect of the FC correction is not large, and $\approx 10\%$ longer exposures are required for the median expected significance to cross each threshold than without correction, at both $\delta_{CP} = -\pi/2$ and for the 50% range of δ_{CP} values.

Calculating $\Delta\chi^2_c$ values above 3σ using the FC method is challenging due to the large number of throws to explore the tails of the $\Delta\chi^2_{FC}$ distribution and prohibitive computational cost. In Figure 13, the fraction of throws that exceed $1-5\sigma$ significance, calculated only with the constant- $\Delta\chi^2$ method are shown in order to explore DUNE's sensitivity at higher significance levels. All the caveats described above relating to the constant- $\Delta\chi^2$ method still apply. Figure 13 shows that although the median significance to CPV does not exceed 5σ until ≈ 336 kt-MW-yr, there are significant fractions of throws at lower exposures which reach 5σ significance. Figure 14 shows the fraction of throws which exceed different significance thresholds at the maximal CP-violating value of $\delta_{CP} = -\pi/2$, and for 50% of all δ_{CP} values, as a function of exposure. By ≈ 200 kt-MW-yr, where the median




FIG. 11. Fraction of throws for which significance of DUNE's CP-violation test ($\delta_{CP} \neq \{0, \pm\pi\}$) exceeds $1-3\sigma$, calculated using both the FC (shaded histograms) and constant- $\Delta\chi^2$ (dashed lines) methods, as a function of the true value of δ_{CP} . Shown for NO, for a number of different exposures. The number of throws used to make each figure is also shown.




FIG. 12. Fraction of throws for which the significance of DUNE's CP-violation test ($\delta_{CP} \neq \{0, \pm\pi\}$) exceeds $1-3\sigma$, for $\delta_{CP} = -\pi/2$ and for 50% of δ_{CP} values, calculated with the FC (solid lines) and constant- $\Delta\chi^2$ (dashed lines) methods, as a function of exposure.




FIG. 13. Fraction of throws for which the significance of DUNE's CP-violation test ($\delta_{CP} \neq \{0, \pm\pi\}$) exceeds 1–5 σ , as a function of the true value of δ_{CP} . Shown for NO, for a number of different exposures. The number of throws used to make each figure is also shown.




FIG. 14. Fraction of throws for which the significance of DUNE's CP-violation test ($\delta_{CP} \neq \{0, \pm\pi\}$) exceeds 1–5 σ , both assuming $\delta_{CP} = -\pi/2$, and for 50% of δ_{CP} values, shown as a function of exposure, for NO.

significance for 50% of the δ_{CP} range is greater than 3σ , the sensitivity at $\delta_{\text{CP}} = -\pi/2$ exceeds 4σ .

V. NEUTRINO MASS ORDERING SENSITIVITY

In this section, the toy-throwing approach described in Section II is used to explore the neutrino mass ordering sensitivity as a function of exposure in detail. In all cases, a joint ND+FD fit is performed, and the reactor θ_{13} constraint is always applied, as described in Section II. An equal split between FHC and RHC running is assumed based on the results obtained in Section III.

Figure 15 shows the significance with which the neutrino mass ordering can be determined for both true NO and IO, for exposures of 66 and 100 kt-MW-yr. The sensitivity metric used is the square root of the difference between the best fit χ^2 value obtained using each ordering, as shown in Equation 4, which is calculated for each throw of the systematics, other oscillation parameters and statistics. The characteristic shape of the MH sensitivity in Figure 15 results from near degeneracy between matter and CPV effects that occurs near $\delta_{\text{CP}} = \pi/2$ ($\delta_{\text{CP}} = -\pi/2$) for true normal (inverted) ordering. Dedicated studies have shown that special attention must be paid to the statistical interpretation of neutrino mass ordering sensitivities [71–73] because the $\Delta\chi_{\text{MO}}^2$ metric does not follow the expected chi-square distribution for one degree of freedom, so the interpretation of the $\sqrt{\Delta\chi_{\text{MO}}^2}$ as the sensitivity is complicated.

Given the complications with the interpretation of significance for mass ordering determination, it is instructive to look at the distribution of the test-statistic (Equation 4), which gives more information than the 68% central band and median throw shown in Figure 15. Figure 16 shows the distribution of $\Delta\chi_{\text{MO}}^2$ obtained for a large ensemble of throws, for both true normal and inverted orderings, for a number of different exposures. There is a uniform distribution of true δ_{CP} used in the throws at each exposure. The change in shape at higher exposures in Figure 16 is due to the degeneracy between δ_{CP} and the effect of the mass ordering, and as might be expected from Figure 15, the separation between hierarchies is greater for some true values of δ_{CP} than others. This additional structure starts to become obvious from a ≈ 66 kt-MW-yr exposure, at which point the CPV sensitivity is not very strong (see Section IV). For all exposures, the shape of the throw distribution is highly non-Gaussian, which makes it difficult to apply simple corrections to the sensitivity of the sort described in Ref. [73]. As a result alternatives to $\sqrt{\Delta\chi_{\text{MO}}^2}$ as a sensitivity metric are not explored, the full information is given in Figure 16.

Figure 16 also indicates the probability for the test statistic $\Delta\chi_{\text{MO}}^2$ to be less (more) than zero from the toy throws for true normal (inverted) orderings at each exposure. This information is summarized in Figure 17.




FIG. 15. Significance of the DUNE determination of the neutrino mass ordering, as a function of the true value of δ_{CP} , for 66 kt-MW-yr (blue) and 100 kt-MW-yr (orange) exposures. The width of the transparent bands cover 68% of fits in which random throws are used to simulate systematic, oscillation parameter and statistical variations, with independent fits performed for each throw constrained by prior uncertainties. The solid lines show the median significance.

This marks the proportion of toys which appear more like the incorrect ordering than the true ordering for the toy, and gives a sense of the ambiguity between the hierarchies, although it is not easily converted to a single number sensitivity. It is clear from Figures 16 and 17 that DUNE is sensitive to the mass ordering even from very low (≈ 12 kt-MW-yr) exposures, with a small probability for preferring the incorrect ordering. By exposures of 66 kt-MW-yr, the overlap between the orderings is very small with $\approx 1\%$ of toy throws which appear more like




FIG. 16. The distribution of $\Delta\chi^2_{MO} = \chi^2_{IO} - \chi^2_{NO}$ values shown for both true normal (red) and true inverted (blue) hierarchies built using random throws of the systematic parameters, the oscillation parameters and with statistical variations. In each case, the χ^2 values are separately minimized with respect to all variable parameters before calculating the test statistic. The fraction of throws for which the value of $\Delta\chi^2_{MO}$ is greater than (less than) 0 is also given for inverted (normal) hierarchies. For each ordering and exposure, approximately 100,000 throws were used.

the incorrect ordering than the true ordering.

Figure 18 shows an alternative way to present the result of the throws as a function of δ_{CP} , which is complementary to Figure 15. The fraction of throws for which the simple figure of merit (the square-root of Equation 4) exceeds different confidence levels are shown, for 1–5 σ significances, and a variety of exposures, all for true NO. The same throws are used as in Figures 16. Despite the caveats regarding the interpretation of $\sqrt{\Delta\chi^2_{MO}}$ as units of σ , the general trend is clear, and provides more information about the expected DUNE sensitivity at low exposures. As with Figures 11 and 13, the point at which the median significance (50% of throws) passes different significance thresholds can be easily read from the figures, and can be compared with those shown in Figure 15. The same general shape as a function of δ_{CP} as was observed in Figure 15 can be seen. The general trend would be very similar in IO, reflected in the line $\delta_{CP} = 0$, although

a slightly longer exposure is required to reach the same sensitivity. The median significance for $\delta_{CP} = -\pi/2$ exceeds 5 σ for 24 kt-MW-yr, at which point the fraction of throws for which the significance is 3 σ or smaller is only $\approx 2\%$. By 66 kt-MW-yr, 100% of the throws exceed 5 σ at $\delta_{CP} = -\pi/2$. By 100 kt-MW-yr exposures, the median significance approaches 5 σ for all true values of δ_{CP} . At long exposures of 336 kt-MW-yr, almost 100% of the throws exceed 5 σ for all values of δ_{CP} .

VI. CONCLUSION

In this work a detailed exploration of DUNE’s sensitivity to CPV and the mass ordering at low exposures has been presented. The analysis uses the same framework, flux, cross section and detector models and selections as were used in Ref. [21], which showed the ultimate DUNE




FIG. 17. The probability for preferring the wrong neutrino mass ordering as a function of exposure, shown for both true NO and IO.

sensitivity to CPV, the neutrino mass ordering and other oscillation parameters, with large statistics samples after long exposures.

The effect of operating with different run plans, involving different ratios of FHC and RHC beam modes, on the mass ordering and CPV Asimov sensitivities was explored. It was found that for low exposures, the sensitivity to both CPV and the mass ordering can be increased for certain regions of parameter space, but at a cost to the sensitivity in other regions. This sensitivity increase is in part produced by leveraging the strong θ_{13} constraint available from reactor experiments. If there is a strong reason to favor the exploration of a given region of parameter space when DUNE begins to take data, this issue should be re-visited. However, with no strong motivation to focus on a given ordering or region of δ_{CP} parameter space, equal FHC and RHC beam running provides a close to optimal sensitivity across all of the parameter space, so was used for the subsequent detailed sensitivity studies. The increase in sensitivity for unequal beam running is also a feature of low exposure running, and degrades the sensitivity almost uniformly across the parameter space investigated for large exposures, with and without a θ_{13} constraint applied.

The studies presented here demonstrate that a full treatment of DUNE's sensitivity at low exposures supports the conclusions made in Refs. [21] and [8] using Asimov studies. In particular, the median CPV sensitivity is $\approx 3\sigma$ for $\delta_{CP} = \pm\pi/2$ after approximately a 100 kt-MW-yr FD exposure. Variations in the expected sensitivity around the median value were also explored. Additionally, it was shown that the CPV sensitivity is not significantly degraded when Feldman-Cousins corrections are included, leading to $\approx 10\%$ longer exposures to reach a given significance level. Crucially, it was found that after an initial low-exposure rise, the Feldman-Cousins

$\Delta\chi_c^2$ do not change as a function of exposure, unlike the rise with exposure which has been observed by the T2K experiment [13].

It has also been shown that strong statements on the mass ordering can be expected with very short exposures of ≈ 12 kt-MW-yr, which supports the results shown in Refs. [21] and [8] with a more complete treatment of the systematic uncertainty.

Although the analysis used here makes no assumptions about the FD staging scenario, and results are given as a function of exposure only, the results are dependent on having a performant ND complex from the start of the experiment. In particular, the low-exposures necessary to make world-leading statements about the mass ordering can only be given with confidence with ND samples included in the fit. Additional samples of events from detectors other than ND-LAr in the DUNE ND complex are not explicitly included in this analysis, but there is an assumption that it will be possible to control the uncertainties to the level used in the analysis, and it should be understood that that implicitly relies on having a highly capable ND.

ACKNOWLEDGMENTS

This document was prepared by the DUNE collaboration using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by Fermi Research Alliance, LLC (FRA), acting under Contract No. DE-AC02-07CH11359. This work was supported by CNPq, FAPERJ, FAPEG and FAPESP, Brazil; CFI, IPP and NSERC, Canada; CERN; MŠMT, Czech Republic; ERDF, H2020-EU and MSCA, European Union; CNRS/IN2P3 and CEA, France; INFN, Italy; FCT, Portugal; NRF, South Korea; CAM, Fundación “La Caixa”, Junta de Andalucía-FEDER, and MICINN, Spain; SERI and SNSF, Switzerland; TÜBİTAK, Turkey; The Royal Society and UKRI/STFC, United Kingdom; DOE and NSF, United States of America. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility operated under Contract No. DE-AC02-05CH11231.




FIG. 18. Fraction of throws for which the DUNE sensitivity to the mass ordering exceeds 1–5 σ significance, as a function of the true value of δ_{CP} . Shown for NO, for a number of different exposures. The number of throws used to make each figure is also shown.

-
- [1] B. Abi *et al.* (DUNE), JINST **15**, T08008 (2020), arXiv:2002.02967 [physics.ins-det].
- [2] X. Qian and P. Vogel, Prog. Part. Nucl. Phys. **83**, 1 (2015), arXiv:1505.01891 [hep-ex].
- [3] M. Fukugita and T. Yanagida, Phys. Lett. **B174**, 45 (1986).
- [4] S. Davidson, E. Nardi, and Y. Nir, Phys. Rept. **466**, 105 (2008), arXiv:0802.2962 [hep-ph].
- [5] B. Abi *et al.* (DUNE), Eur. Phys. J. C **81**, 322 (2021), arXiv:2008.12769 [hep-ex].
- [6] B. Abi *et al.* (DUNE), Eur. Phys. J. C **81**, 423 (2021), arXiv:2008.06647 [hep-ex].
- [7] F. Capozzi, S. W. Li, G. Zhu, and J. F. Beacom, Phys. Rev. Lett. **123**, 131803 (2019), arXiv:1808.08232 [hep-ph].
- [8] B. Abi *et al.* (DUNE), (2020), arXiv:2002.03005 [hep-ex].
- [9] A. Abed Abud *et al.*, (2021), arXiv:2103.13910 [physics.ins-det].
- [10] F. Capozzi *et al.*, Phys. Rev. D **95**, 096014 (2017), [Addendum: Phys.Rev.D 101, 116013 (2020)], arXiv:2003.08511 [hep-ph].
- [11] P. F. de Salas, D. V. Forero, S. Gariazzo, P. Martínez-Miravé, O. Mena, C. A. Ternes, M. Tórtola, and J. W. F. Valle, JHEP **02**, 071 (2021), arXiv:2006.11237 [hep-ph].
- [12] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, and A. Zhou, JHEP **09**, 178 (2020), arXiv:2007.14792 [hep-ph].
- [13] K. Abe *et al.* (T2K), Phys. Rev. D **103**, 112008 (2021), arXiv:2101.03779 [hep-ex].
- [14] K. Abe *et al.* (Super-Kamiokande), Phys. Rev. **D97**, 072001 (2018), arXiv:1710.09126 [hep-ex].
- [15] M. A. Acero *et al.* (NOvA), Phys. Rev. Lett. **123**, 151803 (2019), arXiv:1906.04907 [hep-ex].
- [16] K. Abe *et al.* (T2K), Nature **580**, 339 (2020), arXiv:1910.03887 [hep-ex].
- [17] H. Nunokawa, S. J. Parke, and J. W. Valle, Prog. Part. Nucl. Phys. **60**, 338 (2008), arXiv:0710.0554 [hep-ph].
- [18] L. Wolfenstein, Phys. Rev. **D17**, 2369 (1978).

- [19] S. Mikheev and A. Y. Smirnov, *Sov. J. Nucl. Phys.* **42**, 913 (1985).
- [20] J. Kopp, *Int. J. Mod. Phys. C* **19**, 523 (2008), arXiv:physics/0610206.
- [21] B. Abi *et al.* (DUNE), *Eur. Phys. J.* **C80**, 978 (2020), arXiv:2006.16043 [hep-ex].
- [22] NOvA Collaboration, ed., “NOvA-ART,” (Redmine, 2019) Chap. CAFAna overview, https://cdcvns.fnal.gov/redmine/projects/novaart/wiki/CAFAAna_overview.
- [23] L. Aliaga *et al.* (MINERvA), *Phys. Rev.* **D94**, 092005 (2016), [Addendum: *Phys. Rev. D* **95**, no.3, 039903 (2017)], arXiv:1607.00704 [hep-ex].
- [24] C. Andreopoulos *et al.*, *Nucl. Instrum. Meth.* **A614**, 87 (2010), arXiv:0905.2517 [hep-ph].
- [25] C. Andreopoulos *et al.*, (2015), arXiv:1510.05494 [hep-ph].
- [26] P. Stowell *et al.*, *JINST* **12**, P01016 (2017), arXiv:1612.07393 [hep-ex].
- [27] A. Bodek and J. L. Ritchie, *Phys. Rev. D* **23**, 1070 (1981).
- [28] C. Llewellyn Smith, *Phys. Rept.* **3**, 261 (1972).
- [29] R. Bradford, A. Bodek, H. Budd, and J. Arrington, *Nuclear Physics B - Proceedings Supplements* **159**, 127 (2006).
- [30] K. Abe *et al.* (T2K), *Phys. Rev. Lett.* **121**, 171802 (2018), arXiv:1807.07891 [hep-ex].
- [31] J. Nieves, I. Ruiz Simo, and M. J. Vicente Vacas, *Phys. Rev.* **C83**, 045501 (2011), arXiv:1102.2777 [hep-ph].
- [32] R. Gran, J. Nieves, F. Sanchez, and M. J. Vicente Vacas, *Phys. Rev.* **D88**, 113007 (2013), arXiv:1307.8105 [hep-ph].
- [33] J. Schwehr, D. Cherdack, and R. Gran, (2016), arXiv:1601.02038 [hep-ph].
- [34] P. A. Rodrigues *et al.* (MINERvA), *Phys. Rev. Lett.* **116**, 071802 (2016), arXiv:1511.05944 [hep-ex].
- [35] M. A. Acero *et al.* (NOvA), *Phys. Rev.* **D98**, 032012 (2018), arXiv:1806.00096 [hep-ex].
- [36] C. Colle *et al.*, *Phys. Rev.* **C92**, 024604 (2015), arXiv:1503.06050 [nucl-th].
- [37] D. Rein and L. M. Sehgal, *Annals Phys.* **133**, 79 (1981).
- [38] C. Wilkinson, P. Rodrigues, S. Cartwright, L. Thompson, and K. McFarland, *Phys. Rev.* **D90**, 112017 (2014), arXiv:1411.4482 [hep-ex].
- [39] P. Rodrigues, C. Wilkinson, and K. McFarland, *Eur. Phys. J.* **C76**, 474 (2016), arXiv:1601.01888 [hep-ex].
- [40] A. Bodek and U. Yang, *J. Phys. G* **29**, 1899 (2003), arXiv:hep-ex/0210024.
- [41] M. Glück, E. Reya, and A. Vogt, *Eur. Phys. J. C* **5**, 461 (1998), arXiv:hep-ph/9806404.
- [42] T. Yang, C. Andreopoulos, H. Gallagher, K. Hoffmann, and P. Kehayias, *Eur. Phys. J.* **C63**, 1 (2009), arXiv:0904.4043 [hep-ph].
- [43] Z. Koba, H. B. Nielsen, and P. Olesen, *Nucl. Phys.* **B40**, 317 (1972).
- [44] T. Sjostrand, S. Mrenna, and P. Z. Skands, *JHEP* **05**, 026 (2006), arXiv:hep-ph/0603175 [hep-ph].
- [45] M. Sanchez, in *XXVIII International Conference on Neutrino Physics and Astrophysics (Neutrino 2018)* (Zenodo, 2018) <https://zenodo.org/record/1286758>.
- [46] S. Dytman and A. Meyer, *AIP Conf. Proc.* **1405**, 213 (2011).
- [47] S. Dytman, *Proceedings, Workshop on Neutrino Interactions, Systematic uncertainties and near detector physics: Session of CETUP* 2014: Lead/Dead Wood, South Dakota, USA, July 22-31, 2014*, *AIP Conf. Proc.* **1680**, 020005 (2015).
- [48] S. Dytman, *Acta Phys. Polon.* **B40**, 2445 (2009).
- [49] M. Day and K. S. McFarland, *Phys. Rev.* **D86**, 053003 (2012), arXiv:1206.6745 [hep-ph].
- [50] C. Amsler *et al.* (ArgonCube), *ArgonCube: a novel, fully-modular approach for the realization of large-mass liquid argon TPC neutrino detectors*, Tech. Rep. CERN-SPSC-2015-009. SPSC-I-243 (CERN, Geneva, 2015).
- [51] D. A. Dwyer *et al.*, *JINST* **13**, P10007 (2018), arXiv:1808.02969 [physics.ins-det].
- [52] M. Auger, Y. Chen, A. Ereditato, D. Goeldi, I. Kreslo, D. Lorca, M. Luethi, T. Mettler, J. R. Sinclair, and M. S. Weber, *Instruments* **2**, 3 (2018), arXiv:1711.11409 [physics.ins-det].
- [53] S. Agostinelli *et al.* (GEANT4), *Nucl. Instrum. Meth. A* **506**, 250 (2003).
- [54] B. Abi *et al.* (DUNE), (2018), arXiv:1807.10334 [physics.ins-det].
- [55] B. Abi *et al.* (DUNE), *JINST* **15**, T08010 (2020), arXiv:2002.03010 [physics.ins-det].
- [56] J. S. Marshall and M. A. Thomson, *Eur. Phys. J.* **C75**, 439 (2015), arXiv:1506.05348 [physics.data-an].
- [57] R. Acciarri *et al.* (MicroBooNE), *Eur. Phys. J.* **C78**, 82 (2018), arXiv:1708.03135 [hep-ex].
- [58] B. Abi *et al.* (DUNE), *Phys. Rev.* **D102**, 092003 (2020), arXiv:2006.15052 [physics.ins-det].
- [59] F. James, “MINUIT Function Minimization and Error Analysis: Reference Manual Version 94.1,” (1994), CERN-D-506, CERN-D506.
- [60] I. Esteban, M. C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni, and T. Schwetz, *JHEP* **01**, 106 (2019), arXiv:1811.05487 [hep-ph].
- [61] I. Esteban, M. C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni, and T. Schwetz, “Nufit4.0,” (2018).
- [62] B. Roe, *Phys. Rev.* **D95**, 113004 (2017), arXiv:1707.02322 [hep-ex].
- [63] T. Abrahão *et al.* (Double Chooz), *JHEP* **01**, 190 (2021), arXiv:2007.13431 [hep-ex].
- [64] D. Adey *et al.* (Daya Bay), *Phys. Rev. Lett.* **121**, 241805 (2018), arXiv:1809.02261 [hep-ex].
- [65] G. Bak *et al.* (RENO), *Phys. Rev. Lett.* **121**, 201801 (2018), arXiv:1806.00248 [hep-ex].
- [66] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev.* **D98**, 030001 (2018).
- [67] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, *Eur. Phys. J. C* **71**, 1554 (2011), [Erratum: *Eur. Phys. J. C* **73**, 2501(2013)], arXiv:1007.1727 [physics.data-an].
- [68] S. S. Wilks, *Ann. Math. Stat.* **9**, 60 (1938).
- [69] G. J. Feldman and R. D. Cousins, *Phys. Rev. D* **57**, 3873 (1998), arXiv:physics/9711021.
- [70] J. A. Rice, *Mathematical Statistics and Data Analysis.*, 3rd ed. (Belmont, CA: Duxbury Press., 2006).
- [71] E. Ciuffoli, J. Evslin, and X. Zhang, *JHEP* **01**, 095 (2014), arXiv:1305.5150 [hep-ph].
- [72] X. Qian *et al.*, *Phys. Rev.* **D86**, 113011 (2012), arXiv:1210.3651 [hep-ph].
- [73] M. Blennow, P. Coloma, P. Huber, and T. Schwetz, *JHEP* **1403**, 028 (2014), arXiv:1311.1822 [hep-ph].

Appendix: Feldman-Cousins throw distributions

The distribution of throws used to calculate the $\Delta\chi_c^2$ values for Figure 9 for nine different exposures with $\delta_{CP} = 0$ are shown in Figure 19; for Figure 10a for nine different values of δ_{CP} with an exposure of 100 kt-MW-yr in Figure 20; and for Figure 10b for nine different values

of δ_{CP} with an exposure of 336 kt-MW-yr in Figure 21. For each distribution shown in Figures 19, 20 and 21, the calculated $\Delta\chi_c^2$ values corresponding to for 68.27% (1σ), 90%, 95.45% (2σ) and 99.73% (3σ) of the throws are given and indicated with a vertical line. The number of throws used is also given. The $\Delta\chi_c^2$ values were only calculated up to the 3σ level due to the very large number of throws required for higher confidence levels.




FIG. 19. Distribution of $\Delta\chi^2$ values, calculated using Equation 5, for a large number of throws with true $\delta_{CP} = 0$, for a variety of exposures. The $\Delta\chi_c^2$ values (vertical lines) obtained using the Feldman-Cousins method show the $\Delta\chi_{FC}^2$ value below which 68.27% (1 σ), 90%, 95.45% (2 σ) and 99.73% (3 σ) of throws reside, with the calculated values given in the legend. The number of throws used is also given.




FIG. 20. Distribution of $\Delta\chi^2$ values, calculated using Equation 5, for a large number of throws for 9 different values of true δ_{CP} , for a 100 kt-MW-yr exposure. The $\Delta\chi^2_{FC}$ values (vertical lines) obtained using the Feldman-Cousins method show the $\Delta\chi^2_{FC}$ value below which 68.27% (1σ), 90%, 95.45% (2σ) and 99.73% (3σ) of throws reside, with the calculated values given in the legend. The number of throws used is also given.




FIG. 21. Distribution of $\Delta\chi^2$ values, calculated using Equation 5, for a large number of throws for 9 different values of true δ_{CP} , for a 336 kt-MW-yr exposure. The $\Delta\chi^2_{FC}$ values (vertical lines) obtained using the Feldman-Cousins method show the $\Delta\chi^2_{FC}$ value below which 68.27% (1 σ), 90%, 95.45% (2 σ) and 99.73% (3 σ) of throws reside, with the calculated values given in the legend. The number of throws used is also given.