

This is a repository copy of *The importance of eco-evolutionary dynamics for predicting and managing insect range shifts*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/189463/>

Version: Published Version

Article:

Wellenreuther, Maren, Dudaniec, Rachael Y, Neu, Anika et al. (10 more authors) (2022) The importance of eco-evolutionary dynamics for predicting and managing insect range shifts. *Current Opinion in Insect Science*. 100939. ISSN 2214-5753

<https://doi.org/10.1016/j.cois.2022.100939>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



ELSEVIER



The importance of eco-evolutionary dynamics for predicting and managing insect range shifts

Maren Wellenreuther^{1,2}, Rachael Y Dudaniec³, Anika Neu⁴, Jean-Philippe Lessard⁵, Jon Bridle⁶, José A Carbonell^{7,13}, Sarah E Diamond⁸, Katie E Marshall⁹, Camille Parmesan^{10,11,12}, Michael C Singer^{10,11}, Janne Swaegers¹³, Chris D Thomas¹⁴ and Lesley T Lancaster¹⁵

Evolutionary change impacts the rate at which insect pests, pollinators, or disease vectors expand or contract their geographic ranges. Although evolutionary changes, and their ecological feedbacks, strongly affect these risks and associated ecological and economic consequences, they are often underappreciated in management efforts. Greater rigor and scope in study design, coupled with innovative technologies and approaches, facilitates our understanding of the causes and consequences of eco-evolutionary dynamics in insect range shifts. Future efforts need to ensure that forecasts allow for demographic and evolutionary change and that management strategies will maximize (or minimize) the adaptive potential of range-shifting insects, with benefits for biodiversity and ecosystem services.

Addresses

¹ The New Zealand Institute for Plant & Food Research Ltd, Nelson, New Zealand

² School of Biological Sciences, The University of Auckland, Auckland, New Zealand

³ School of Natural Sciences, Macquarie University, Sydney, NSW 2109, Australia

⁴ Zoological Institute and Museum, University of Greifswald, Greifswald, Germany

⁵ Department of Biology, Concordia University, Montreal, Canada

⁶ Department of Genetics, Evolution and Environment, University College London, UK

⁷ Department of Zoology, Faculty of Biology, University of Seville, Seville, Spain

⁸ Department of Biology, Case Western Reserve University, Cleveland, OH 44106, USA

⁹ Department of Zoology, University of British Columbia, Vancouver, BC, Canada

¹⁰ Station d'Écologie Théorique et Expérimentale (SETE), CNRS, 2 route du CNRS, 09200 Moulis, France

¹¹ Biological and Marine Sciences, University of Plymouth, Plymouth PL4 8AA, UK

¹² Dept of Geological Sciences, University of Texas at Austin, Austin, Texas 78712

¹³ Laboratory of Evolutionary Stress Ecology and Ecotoxicology, University of Leuven, Charles Deberiotstraat 32, Leuven B-3000, Belgium

¹⁴ Leverhulme Centre for Anthropocene Biodiversity, University of York, Wentworth Way, York YO10 5DD, UK

¹⁵ School of Biological Sciences, University of Aberdeen, Aberdeen UK AB24 2TZ

Corresponding author: Lesley T Lancaster (lesleylancaster@abdn.ac.uk)

Current Opinion in Insect Science 2022, 52:100939

This review comes from a themed issue on **Global change biology**

Edited by **Robby Stoks** and **Nedim Tüzün**

<https://doi.org/10.1016/j.cois.2022.100939>

2214-5745/© 2022 Elsevier Ltd. All rights reserved.

Introduction

Insect range expansions and contractions in response to human activities are widespread and are reshaping entire food webs and altering ecosystem functions and services [1,2]. Practical consequences of range shifts include the rescue of species from degraded habitats, more robust populations with expanded ranges, and increases in local biodiversity. While such consequences can often be favorable, negative impacts are also observed, if range-shifting species include agricultural pests and vectors of human and agricultural diseases, or disrupt native communities and ecosystem services [3,4]. Because both ecological and economic consequences of insect range shifts are important, it is a priority to understand what drives some insects, but not others, to undergo rapid range shifts, and to further understand drivers of variation. Until recently, however, forecasting models have mostly considered ecological drivers such as colonization, biotic interactions, and abundance shifts, without taking into account rapid evolution associated with range shifting, which can affect all of these ecological parameters. Rapid evolution commonly occurs during range shifts; this phenomenon was first shown in insects [5], and is clearly a dominant phenomenon in this group, facilitated by their sensitivities to environmental change and by their short life cycles and large population sizes. These characteristics can facilitate rapid transgenerational genetic or epigenetic change [6,7]. Strong life-history trade-offs in insects can further help to maintain

evolutionary variability, and therefore, perhaps counter-intuitively, can increase fitness when encountering novel environments during range expansions [8].

Range shifts are ecological processes, so when they involve evolution they invoke eco-evolutionary dynamics, i.e. interactions between evolutionary change and ecological outcomes. For example, insect range expansions can both drive evolution of dietary niche breadth [9•] and be facilitated by it [5•,10]. There is well-established theoretical understanding of eco-evolutionary dynamic feedbacks between microevolutionary change and demographic shifts during colonization of novel environments [11]. However, empirical evidence lags behind theoretical developments. This omission is largely due to a lack of established approaches for discovering, diagnosing and monitoring eco-evolutionary processes in wild systems. Lack of data to test models therefore makes it difficult to predict evolutionary responses and their relationship to ecological patterns and processes. This is unfortunate, given that conservation and management actions depend on a robust prediction of range shift trajectories. To increase awareness of this gap between theory and data, we (1) highlight opportunities that insects offer for identifying the eco-evolutionary basis of range shifts and their applied relevance; (2) discuss recent advances in our understanding of eco-evolutionary dynamics of insect range shifts; and (3) identify critical avenues for robustly assessing the temporal and spatial scale of those dynamics to improve predictions of range shifts and their impacts on biodiversity.

The importance of studying eco-evolutionary dynamics during insect range expansions

Insects are among the most rapidly range-expanding of all terrestrial taxa [12•]. Range-shifting insects have a large impact on the assembly of resident biotas [13], on pollinator networks [4] and other ecosystem services [14•]. Life histories of range-expanding insects are often distinct from resident communities in that they exhibit increased resilience to stressors, ecological generalism, or competitive ability [15]. These observations have been used to suggest that colonizing species are those already with ecological traits that enable rapid spread. However, this conclusion rests on the assumption that ecological and evolutionary processes occur on different time scales, where the evolution of the traits that confer the capacity for ecological disruption is thought to occur in advance of geographical expansion. However, we argue that this assumption is often unlikely to be true, in part because attempts to discover traits predisposing range expansions have failed or detected only weak effects (reviewed in [16•]). Instead, we suggest that traits conferring colonization ability most likely evolve *during* range expansions (see also: [17•]; Figure 1). Conversely, traits and processes likely to arise during range

contraction may potentially confer enhanced sensitivity to further declines, although evolutionary processes during range contraction are even less well characterized than those associated with expansion.

Insects further offer ideal opportunities for detailed studies of range shifts. Data on insect distributional shifts is increasing in volume owing to technological advances in rapid insect surveying methods [18], such as remote sensing and techniques that exploit frequencies of solar radiation to track individuals and their microclimates [19•]. These advances offer more in-depth insight into the densities, distributions and dispersal behavior of flying insects. However, such data are highly biased towards terrestrial insect communities, which are unlikely to be representative of aquatic species and the important ecosystem services they provide. Increasing research on the eco-evolutionary dynamics of range shifting pest species and disease vectors [20] is particularly needed, given their relevance for food sustainability and human health.


Eco-evolutionary dynamics during insect range shifts

Many studies of evolution during range shifts in insects have focused on changes in physiological traits, such as thermal tolerances, shaped under ecological selection in the new part of the range (e.g. [21–23]). However, novel demographic conditions arising from range expansion can also impose selection on characteristics such as enhanced dispersal or resistance to inbreeding, and on life history traits such as reduced generation time, voltinism and diapause that can speed up the expansion itself [24]. In addition, gene flow from the historic range and genetic drift further shape genetic variation and, thus alter evolutionary dynamics [25]. Neutral, rather than adaptive processes, can dominate evolutionary changes during colonization events, making it difficult to predict range shift outcomes. Nonetheless, certain anticipatory responses may also evolve, for example, serial colonization at the leading edge of range shifts can select for indiscriminate individuals that are more likely to accept marginal habitat conditions [26].

Recent studies highlight the importance of trait plasticity for insect adaptation during range shifts [23,27]. In particular, increased learning ability may be important for insects to cope with environmental heterogeneity and unpredictability during colonization events [28,29]. Evolutionary increases in learning ability, as in any form of plasticity, can further facilitate environmental adaptation by allowing persistence in a novel environment until genetic accommodation has occurred [30].

Dispersal evolution is often part of eco-evolutionary feedback loops, both as a driver and response [31]. In addition to being favored during serial colonization [24],

Figure 1



Mechanistic, simplified representation of eco-evolutionary dynamics during insect range shifts and their importance for (applied) research. The gear wheels can move in the direction indicated by the arrows, such that the exemplary selection of demographic, environmental and social drivers (D) (see color legend on bottom right) may individually, or in combination, (1) evoke evolutionary responses (e.g. epigenetic, or genetic) (R) in range expanding insect populations on contemporary time scales (2). Understanding the rates and strengths of these processes will be crucial for theoretical advancements and practical implications of insect range shifts (see boxes on bottom left). Moreover, responses can induce new driving forces (3), resulting in positive or negative eco-evolutionary feedbacks and feedback loops, for example, an increase in a driving process A induces a genetic response that further increases versus decreases trait A, thus accelerating or dampening the dynamic and resulting range shift trajectory.

high or low population densities at the range limit may further facilitate changes in dispersal propensity [32]. Evolutionary impacts of range shift-induced dispersal traits may not evolve independently from other traits, meaning that changes in dispersal will impact correlated traits, further modifying eco-evolutionary dynamics [33•,34•]. Furthermore, insect dispersal traits are often temperature-sensitive, suggesting that the interplay between shifts in environmental plasticity and dispersal needs to be considered together to predict likely evolutionary dynamics [35].

Social dynamics also evolve readily during range shifts, influencing adaptive potential to both expand and to resist contraction. This may occur for instance if sexual or other socially-mediated selection differs between the range core and the range edge [36], due to changes in fragmentation, relatedness, or density conditions at fluctuating range edges that impact reproductive and competitive opportunities. Novel social dynamics at range margins can feed back to impact individual phenotypes and fitness, for example, via invoking generalized stress responses that also confer fitness in novel environments [37]. Another consequence of range shifts may be skewed sex ratios, particularly where sexes differ in dispersal, meaning the more dispersive sex reaches the range limit in higher numbers. The resulting skew can impact mating systems or the strength of sexual selection (e.g. in damselflies: [7•]).

An understudied area is the complex, interactive effect of eco-evolutionary change on community dynamics during insect range shifts. For example, novel plant-insect interactions caused by unequal rates of range shifting likely impact both plant and insect biodiversity, population sizes, and evolutionary potential, with shifts in pollinator and pest communities impacting food security. In particular, encountering novel resources during range shifts can trigger diet evolution (e.g. increase or reduction in Lepidopteran host breadth: [9•,38•,39•]). Moreover, shifts in plasticity, physiology, life history and dispersal associated with range shifts can further impact future community dynamics: colonizing lineages, which have already adapted to spatial variability, may then have an evolutionary advantage over existing residents when future environments become more variable or less predictable [40]. To take into account the evolution of interactions among species and environments, an approach that includes multiple drivers and responses is needed to detect how biotic interactions affect and are impacted by eco-evolutionary processes during range shifts (Figure 1).

Advances in the study of eco-evolutionary dynamics

To advance our understanding of eco-evolutionary dynamics, a temporal approach is needed to quantify the

nature, rate and magnitude of population changes along range expansion trajectories. However, due to practical constraints, most studies assess range shift dynamics across only one or a few generations. Improving temporal resolution may be accomplished via judicious use of historical records or genetic inference of past demography and adaptation, to make links between previous or ongoing population processes and resulting demographic characteristics or trait values [7•,39]. Statistical approaches that account for abiotic and biotic factors that can bias range estimates are also developing rapidly to enable more accurate modeling of past and current range dynamics [41••].

Recent advances in genomic sequencing approaches and computational improvements further present the opportunity to track historical eco-evolutionary processes using a greatly expanded set of ‘omic’ markers. Spatial genomic datasets allow for high-powered tests for dispersal and demographic processes that influence genetic connectivity, and provide insight into spatial and temporal dynamics of neutral and adaptive genomic variation [42,43]. Likewise, transcriptomic approaches can test the relationships between environment, gene expression, and phenotype in insect populations impacted by range shifting, providing insight into how the genetic basis of traits, and their plasticities, may evolve [44,45•]. Epigenomics, the study of DNA modifications that impact gene expression, is another growing field that identifies how insects can rapidly respond to a changing climate [6], with accumulating evidence suggesting that modifications can have significant transgenerational impact [46]. One interesting idea is that the evolution of ‘epigenetic potential’ (i.e. proportion of the genome available for new epigenetic modifications) may be an important mechanism of plasticity evolution during range shifts [47], a hypothesis that remains untested in insects.

Effective use of these techniques depends on careful experimental design to allow for generalizations about the role of eco-evolutionary dynamics in the likelihood, rate, and outcomes of range shifts. For instance, effective design may include careful sampling over time within the range as well as at the edge, repeated sampling across multiple range shift transects within a species, or comparing outcomes across range-shifting versus nonshifting species along a shared spatial gradient [48]. Meta-analyses adopting this latter approach have recently revealed that *only range-shifting insects* exhibit classical latitudinal gradients in dietary and thermal niche breadth. This work suggests that rapid niche evolution during range shifts is an important cause of macroecological patterns [9•,49]. In addition, replicated assessment of trait or distribution outcomes across independent expansion trajectories within a species, such as in [38], are crucial to understand what patterns are due to stochastic versus general responses of populations.

Appropriate application of ‘omic approaches often requires a mechanistic understanding of how genetic variation links to relevant traits, which in turn impact demography and adaptation. We therefore advocate for combining wild studies with laboratory or field experiments to robustly corroborate these links. Experimental approaches may involve evolution of replicate laboratory populations under range shift conditions, and adopting an ‘evolve and resequence’ approach to track how genetic and phenotypic (co)variation respond to experimentally altered patterns of demography and environmental variation [50]. Alternatively, common garden or reciprocal transplant studies can be used to establish causality. Such common garden studies can also be combined with more complex breeding designs coupled to sequencing, to experimentally disentangle and quantify genetic and environmental factors [51].

Increased collaborative efforts are also essential. Extending the temporal and spatial scales of study requires investment and prioritization at multi-institutional, and in many cases, multi-national levels. A multidisciplinary approach is also critical for integrating data from environmental, physiological, demographic, genetic, and epigenetic data sources. Effectively combining this information will be essential for modeling evolutionary trajectories and feedbacks that facilitate or hinder range shifts. Such models are still in their infancy. However, recent work suggests that incorporation of data on genetic variation improves species distribution models and forecasts [52], and the development of genetically explicit process-based models is increasing at a rapid pace [53,54]. Inverse fitting of process-based forecasting models parameterized from rigorous empirical demographic and genomic data represents the next stage in forecasting sophistication [55].

Conclusions

Accurate forecasting of which insect species will or will not range-shift or adapt is urgently needed. Insect range shifts are already impacting biodiversity, human disease risk, agriculture, and forestry, with severe economic consequences [3]. Providing the needed empirical data to understand and model the eco-evolutionary processes that underpin range shifts demands multidisciplinary insights e.g. from ‘omic’ approaches, and combinations of wild and laboratory studies at the appropriate temporal and ecological scale. Once we understand the biological processes limiting or enhancing insect range shifts, species and communities identified as being at risk can be targeted to support their ecological and adaptive potentials.

Conflict of interest statement

The authors declare no conflict of interest.

Acknowledgements

We would like to thank Robby Stoks for inviting us to participate in this special issue.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Halsch CA, Shapiro AM, Fordyce JA, Nice CC, Thorne JH, Waetjen DP, Forister ML: **Insects and recent climate change**. *Proc Natl Acad Sci U S A* 2021, **118**:2002543117.

This long-term study infers that climate change has a dramatic influence on insect range dynamics, even in situations where land-use change has been minimal.

2. Lehmann P, Ammunet T, Barton M, Battisti A, Eigenbrode SD, Jepsen JU, Kalinkat G, Neuvonen S, Niemelä P, Terblanche JS, et al.: **Complex responses of global insect pests to climate warming**. *Front Ecol Environ* 2020, **18**:141-150.

This study evaluates responses of 31, globally distributed pest species to warming, and concludes that species respond in highly individual ways, with responses depending on physiological, phenological, demographic traits and evolutionary changes. The authors suggest that a multidimensional approach and consideration of multiple interacting traits and trade-offs are needed when predicting insect climate response.

3. IPCC: **Climate Change 2022: Impacts, Adaptation, and Vulnerability**. Cambridge University Press; 2022.
4. Vasiliev D, Greenwood S: **The role of climate change in pollinator decline across the Northern Hemisphere is underestimated**. *Sci Total Environ* 2021, **775**:145788.
5. Thomas CD, Bodsworth EJ, Wilson RJ, Simmons AD, Davies ZG, Musche M, Conradt L: **Ecological and evolutionary processes at expanding range margins**. *Nature* 2001, **411**:577-581.

This review highlights the importance of considering multiple aspects of diversity when estimating responses to climate change. Beta diversity and genetic diversity, in particular, may be strongly affected.

6. McCaw BA, Stevenson TJ, Lancaster LT: **Epigenetic responses to temperature and climate**. *Integr Comp Biol* 2020, **60**:1469-1480.
7. Dudaniec RY, Carey AR, Svensson EI, Hansson B, Yong CJ, Lancaster LT: **Latitudinal clines in sexual selection, sexual size dimorphism and sex-specific genetic dispersal during a poleward range expansion**. *J Anim Ecol* 2021, <https://doi.org/10.1111/1365-2656.13488>.

This landscape genomic study finds that the demographic change and increasing rate of male-biased dispersal during an insect range expansion alters the strength and direction of sexual selection.

8. Parmesan C, Singer MC: **Mosaics of climatic stress across species' ranges: tradeoffs cause adaptive evolution to limits of climatic tolerance**. *Philos Trans R Soc B* 2022, **377**.
9. Lancaster LT: **Host use diversification during range shifts shapes global variation in Lepidopteran dietary breadth**. *Nat Ecol Evol* 2020, .

This study revisits the idea that niche breadth often increases with latitude, and reports that niche breadth is better related to range position than it is to latitude, suggesting that intraspecific dynamics are more important than latitudinal variation in environmental selection in shaping macroecological distributions of niche breadth.

10. Neu A, Lötters S, Nörenberg L, Wiemers M, Fischer K: **Reduced host-plant specialization is associated with the rapid range expansion of a Mediterranean butterfly**. *J Biogeogr* 2021, **48**:3016-3031.
11. Polechová J: **Is the sky the limit? On the expansion threshold of a species' range**. *PLoS Biol* 2018, **16**:1-18.
12. Lenoir J, Bertrand R, Comte L, Bourgeaud L, Hattab T, Murielle J, Grenouillet G: **Species better track climate warming in the oceans than on land**. *Nat Ecol Evol* 2020, **4**:1044-1059.

This synthesis study provides a comprehensive database on published range shift estimates.

13. Fitt RNL, Lancaster LT: **Range shifting species reduce phylogenetic diversity in high latitude communities via competition.** *J Anim Ecol* 2017, **86**:543-555.
14. Wallingford PD, Morelli TL, Allen JM, Beaury EM, Blumenthal DM, Bradley BA, Dukes JS, Early R, Fusco EJ, Goldberg DE, et al.: **Adjusting the lens of invasion biology to focus on the impacts of climate-driven range shifts.** *Nat Clim Chang* 2020 **10**:398-405.
- This review highlights the fact that range shifts can have similar consequences to invasions, and thus calls upon the community to assess and mitigate risks to communities from range shifts.
15. Ghisbain G, Gérard M, Wood TJ, Hines HM, Michez D: **Expanding insect pollinators in the Anthropocene.** *Biol Rev* 2021, **96**:2755-2770.
16. Beissinger SR, Riddell EA: **Why are species' traits weak predictors of range shifts?** *Annu Rev Ecol Syst* 2021 **2021**, **52**:47-66.
- This review highlights the importance of considering (1) intraspecific variation, (2) plasticity, (3) ecological context and traits that determine rates of environmental exposure, (4) a larger range of ecological drivers, (5) evolutionary history, (6) trait interactions, (7) estimation inaccuracies, and (8) non-linear patterns, when attempting trait-based forecasting of range shifts.
17. Lancaster LT: **On the macroecological significance of eco-evolutionary dynamics: the range shift-niche breadth hypothesis.** *Philos Trans R Soc B* 2022, **377**.
- This review compares the main hypotheses for why range size and niche breadth covary with each other and with latitude, and introduces a new hypothesis based on the observation that niche breadth is particularly likely to evolve during range expansions (to higher latitudes). Evolutionary mechanisms for why this is likely to occur are presented, with evidence from theory and empirical studies.
18. Karlsson D, Hartop E, Forshage M, Jaschhof M, Ronquist F: **The Swedish Malaise Trap Project: A 15 Year Retrospective on a Countrywide Insect Inventory.** *Biodivers Data J* **8** e47255 2020, **8**:e47255.
19. Walter T, Degen J, Pfeiffer K, Stöckl A, Montenegro S, Degen T: **A new innovative real-time tracking method for flying insects applicable under natural conditions.** *BMC Zool* 2021 **6**:1-11.
- This study develops a new protocol to tag individual insects with photoluminescent dots, in order to track short-range movements using an infrared detection system. The approach has strong promise to advance study of individual insect behaviours, particularly for flying insects
20. Mushegian AA, Tougeron K: **Animal-microbe interactions in the context of diapause.** *Biol Bull* 2019, **237**:180-191.
21. Carbonell JA, Wang YJ, Stoks R: **Evolution of cold tolerance and thermal plasticity in life history, behaviour and physiology during a poleward range expansion.** *J Anim Ecol* 2021, <https://doi.org/10.1111/1365-2656.13482>
22. Neu A, Fischer K: **Indications for rapid evolution of trait means and thermal plasticity in range-expanding populations of a butterfly.** *J Evol Biol* 2022, **35**:124-133.
23. Swaegers J, Sánchez-Guillén RA, Carbonell JA, Stoks R: **Convergence of life history and physiology during range expansion toward the phenotype of the native sister species.** *Sci Total Environ* 2021, <https://doi.org/10.1016/J.SCITOTENV.2021.151530>
24. Phillips BL, Brown GP, Shine R: **Life-history evolution in range-shifting populations.** *Ecology* 2010, **91**:1617-1627.
25. Miller TEX, Angert AL, Brown CD, Lee-Yaw JA, Lewis M, Lutscher F, Marculis NG, Melbourne BA, Shaw AK, Szűcs M, et al.: **Eco-evolutionary dynamics of range expansion.** *Ecology* 2020, **101**:1-14.
26. Martin Y, Titeux N, Van Dyck H: **Range expansion, habitat use, and choosiness in a butterfly under climate change: Marginality and tolerance of oviposition site selection.** *Ecol Evol* 2021, **11**:2336-2345.
27. Hällfors MH, Pöyry J, Heliölä J, Kohonen I, Kuussaari M, Leinonen R, Schmucki R, Sihvonen P, Saastamoinen M: **Combining range and phenology shifts offers a winning strategy for boreal Lepidoptera.** *Ecol Lett* 2021, **24**:1619-1632.
28. Nieberding CM, Van Dyck H, Chittka L: **Adaptive learning in non-social insects: from theory to field work, and back.** *Curr Opin Insect Sci* 2018, **27**:75-81.
29. Braem S, Turlure C, Nieberding C, Van Dyck H: **Oviposition site selection and learning in a butterfly under niche expansion: an experimental test.** *Anim Behav* 2021, **180**:101-110.
30. Diamond SE, Martin RA: **Buying time: plasticity and population persistence.** *Phenotypic Plast Evol* 2021, <https://doi.org/10.1201/9780429343001-11>
31. Hanski IA: **Eco-evolutionary spatial dynamics in the Glanville fritillary butterfly.** *Proc Natl Acad Sci U S A* 2011, **108**:14397-14404.
32. Travis JMJ, Mustin K, Benton TG, Dytham C: **Accelerating invasion rates result from the evolution of density-dependent dispersal.** *J Theor Biol* 2009, **259**:151-158.
33. Taylor-Cox ED, Macgregor CJ, Corthine A, Hill JK, Hodgson JA, Saccheri IJ: **Wing morphological responses to latitude and colonisation in a range expanding butterfly.** *PeerJ* 2020, **8**:e10352.
- These authors investigate factors affecting wing characteristics in a range-expanding butterfly, and highlight that range shifts can be most rapid when the direction of increase in plasticity and selection act in the same direction along the range expansion gradient.
34. Ochocki BM, Saltz JB, Miller TEX: **Demography-dispersal trait correlations modify the eco-evolutionary dynamics of range expansion.** *Am Nat* 2020, **195**:231-246.
- Using experimental evolution and modeling, this study highlights that genetic or maternal correlation among dispersal and demography can enhance or repress the rate of range shifting.
35. Jourdan J, Baranov V, Wagner R, Plath M, Haase P: **Elevated temperatures translate into reduced dispersal abilities in a natural population of an aquatic insect.** *J Anim Ecol* 2019, **88**:1498-1509.
36. Svensson EI, Willink B, Duryea MC, Lancaster LT: **Temperature drives pre-reproductive selection and shapes the biogeography of a female polymorphism.** *Ecol Lett* 2020, **23**:149-159.
37. Wood C, Fitt RNL, Lancaster LT: **Evolving social dynamics prime thermal tolerance during a poleward range shift.** *Biol J Linn Soc* 2019, **126**.
38. Freedman MG, Jason C, Ramirez SR, Strauss SY: **Host plant adaptation during contemporary range expansion in the monarch butterfly.** *Evolution (N Y)* 2020, **74**:377-391.
- Thus study capitalized on historical colonization records and common garden experiments to discover replicate loss of generalization in derived, island populations of the Monarch butterfly, with observed patterns of local adaptation likely due to underlying demographic changes, rather than by generalist-specialist trade-offs.
39. Singer MC, Parmesan C: **Colonizations cause diversification of host preferences: a mechanism explaining increased generalization at range boundaries expanding under climate change.** *Glob Chang Biol* 2021, <https://doi.org/10.1111/gcb.15656>.
- Leveraging long-term population and genetic data, this study concludes that younger populations are more likely to exhibit a wider range of host preferences than older populations, despite generally exhibiting lower genetic diversity. Implications for range dynamics are discussed.
40. Lancaster LT, Morrison G, Fitt RN: **Life history trade-offs, the intensity of competition, and coexistence in novel and evolving communities under climate change.** *Philos Trans R Soc B Biol Sci* 2017, **372**.
41. Clare JDJ, Townsend PA, Zuckerberg B: **Generalized model-based solutions to false-positive error in species detection/ nondetection data.** *Ecology* 2021, **102**:e03241.
- This study presents an approach to deal with bias in occupancy records that are used to infer ranges and range changes. The authors highlight that estimates of range dynamics from erroneous occurrence patterns can be severely biased, even if the underlying observations exhibit only minor bias. This highlights the need to carefully control for false positives, in particular when estimating range changes

42. Dudaniec RY, Yong CJ, Lancaster LT, Svensson EI, Hansson B: **Signatures of local adaptation along environmental gradients in a range-expanding damselfly (*Ischnura elegans*)**. *Mol Ecol* 2018, **27**:2576-2593.
43. Swaegers J, Mergeay J, Van Geystelen A, Therry L, Larmuseau MHD, Stoks R: **Neutral and adaptive genomic signatures of rapid poleward range expansion**. *Mol Ecol* 2015, **24**:6163-6176.
44. Lancaster LT, Dudaniec RY, Chauhan P, Wellenreuther M, Svensson EI, Hansson B: **Gene expression under thermal stress varies across a geographical range expansion front**. *Mol Ecol* 2016, **25**:1141-1156.
45. Swaegers J, Spanier KI, Stoks R: **Genetic compensation rather than genetic assimilation drives the evolution of plasticity in response to mild warming across latitudes in a damselfly**. *Mol Ecol* 2020, **29**:4823-4834.
- Despite the fact that range shifts are predicted to be most rapid when plasticity acts in the same direction as selection (see note for reference 33), this study finds that gene expression patterns for range-expanding species are more likely to exhibit signals of genetic compensation, that is, range expanding populations showed increased plasticity in response to ancestral conditions than to the derived conditions at the range front.
46. Anastasiadi D, Venney CJ, Bernatchez L, Wellenreuther M, Wellenreuther M: **Epigenetic inheritance and reproductive mode in plants and animals**. *Trends Ecol Evol* 2021, **2021**.
47. Hanson HE, Koussayer B, Kilvitis HJ, Schrey AW, Dylan Maddox J, Martin LB: **Epigenetic potential in native and introduced populations of house sparrows (*Passer domesticus*)**. *Integr Comp Biol* 2020, **60**:1458-1468.
48. Janion-Scheepers C, Phillips L, Sgrò CM, Duffy GA, Hallas R, Chown SL: **Basal resistance enhances warming tolerance of alien over indigenous species across latitude**. *Proc Natl Acad Sci U S A* 2018, **115**:145-150.
49. Lancaster LT: **Widespread range expansions shape latitudinal variation in insect thermal limits**. *Nat Clim Chang* 2016, **6**:618-621.
50. Larsen CD, Hargreaves AL: **Miniaturizing landscapes to understand species distributions**. *Ecography ((Cop))* 2020, **43**:1625-1638.
51. Oomen RA, Hutchings JA: **Genomic reaction norms inform predictions of plastic and adaptive responses to climate change**. *J Anim Ecol* 2022,, <https://doi.org/10.1111/1365-2656.13707>
52. Razgour O, Forester B, Taggart JB, Bekaert M, Juste J, Ibáñez C, Puechmaille SJ, Novella-Fernandez R, Alberdi A, Manel S: **Considering adaptive genetic variation in climate change vulnerability assessment reduces species range loss projections**. *Proc Natl Acad Sci U S A* 2019, **116**:10418-10423.
53. Currat M, Arenas M, Quilodràn CS, Excoffier L, Ray N: **SPLATCHE3: simulation of serial genetic data under spatially explicit evolutionary scenarios including long-distance dispersal**. *Bioinformatics* 2019, **35**:4480.
54. Landguth EL, Forester BR, Eckert AJ, Shirk AJ, Menon M, Whipple A, Day CC, Cushman SA: **Modelling multilocus selection in an individual-based, spatially-explicit landscape genetics framework**. *Mol Ecol Resour* 2020, **20**:605-615.
55. Waldvogel A, Feldmeyer B, Rolshausen G, Exposito-Alonso M, Rellstab C, Kofler R, Mock T, Schmid K, Schmitt I, Bataillon T, et al.: **Evolutionary genomics can improve prediction of species' responses to climate change**. *Evol Lett* 2020, **4**:4-18.