



This is a repository copy of *Kinetic resolution of 2-Aryl-4-methylenepiperidines toward enantioenriched functionalizable piperidine fragments*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/190517/>

Version: Published Version

---

**Article:**

Choi, A., Meijer, A.J.H.M. [orcid.org/0000-0003-4803-3488](https://orcid.org/0000-0003-4803-3488), Silvestri, I.P. et al. (1 more author) (2022) Kinetic resolution of 2-Aryl-4-methylenepiperidines toward enantioenriched functionalizable piperidine fragments. *The Journal of Organic Chemistry*, 87 (13). pp. 8819-8823. ISSN 0022-3263

<https://doi.org/10.1021/acs.joc.2c00862>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Kinetic Resolution of 2-Aryl-4-methylenepiperidines toward Enantioenriched Functionalizable Piperidine Fragments

Anthony Choi, Anthony J. H. M. Meijer, Ilaria Proietti Silvestri, and Iain Coldham\*



Cite This: *J. Org. Chem.* 2022, 87, 8819–8823



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

**ABSTRACT:** The base *n*-BuLi with sparteine allows a kinetic resolution of *N*-Boc-2-aryl-4-methylenepiperidines. The 2,2-disubstituted products and recovered starting materials were isolated with high enantiomeric ratios. From VT-NMR spectroscopy and DFT studies, the rate of rotation of the *N*-Boc group is fast. Lithiation and trapping of the enantioenriched starting materials gave 2,2-disubstituted piperidines with retention of stereochemistry. Functionalization of the 4-methylene group led to a variety of 2,4-disubstituted piperidines without loss of enantiopurity that could be useful building blocks for drug discovery.



Piperidines are important molecular motifs found in many pharmaceutical drugs and natural products.<sup>1,2</sup> Examples include glasdegib, which is a drug that has been used in the treatment of acute myeloid leukemia,<sup>3</sup> and LNP023, which has been developed as a serine protease factor B inhibitor (Figure 1).<sup>4</sup> Within these structures, an aryl group is present in the 2-





Figure 1. Structures of glasdegib and LNP023.

position of the piperidine core along with an additional functional group in the 4-position. In other piperidine-based drugs, different functional groups extend from the piperidine core resulting in the presence of various stereogenic centers.<sup>5</sup> The ability to control the stereochemistry of these stereocenters within a substituted piperidine remains an area of interest in organic synthesis to allow the development of novel, structurally diverse 3D molecules. These molecules are particularly important in the expansion of fragment libraries that are used in screening processes when developing new drugs.<sup>5</sup> To support the synthesis of these compounds, the use of functionalizable 3D fragments that possess substituents and synthetic handles prove useful as they would allow a large array

of compounds to be synthesized in a more timely and cost-efficient manner.

Previously, we have reported the synthesis of enantiomerically enriched 2-arylpiperidines via kinetic resolution using the chiral base system of *n*-BuLi and (–)-sparteine, which selectively deprotonates one enantiomer of the starting material (Scheme 1, Boc = CO<sub>2</sub>tBu).<sup>6</sup> The resolved 2-arylpiperidines can then be treated with *n*-BuLi and an electrophile to give enantioenriched 2,2-disubstituted compounds without a loss of selectivity. To expand on this research, we wanted to investigate the lithiation of substituted 2-arylpiperidines containing functionalizable groups in the 4-

## Scheme 1. Previous Work on Kinetic Resolution of 2-Arylpiperidines



Received: April 12, 2022

Published: June 14, 2022




position.<sup>7</sup> There are limited examples of kinetic resolution of disubstituted piperidines<sup>8,9</sup> and only one method as far as we are aware for the asymmetric synthesis of 2,4-dialkyl-substituted piperidines by kinetic resolution.<sup>9</sup> This approach uses a chiral hydroxamic acid to obtain good selectivities, although it was important that the starting diastereoisomeric 2,4-disubstituted piperidine was the *trans*-isomer and the chiral hydroxamic acid needed for the reaction is not available commercially.

To apply our chemistry to the formation of a variety of derivatives it was important to ensure the functional group installed in the 4-position was not only inert toward lithiation but also able to undergo further functional group interconversions. We chose to install a methylene group in the 4-position as it meets these criteria. In this paper, we report that *N*-Boc-2-aryl-4-methylenepiperidines are viable substrates for highly selective kinetic resolution by asymmetric deprotonation and can be converted to a selection of 2,4-disubstituted piperidines.

Accessing the 2-aryl-4-methylenepiperidines required for our study was achieved from the enones **1a–h**, which could be synthesized using known methods.<sup>10</sup> Reduction of the enone using *L*-Selectride gave the 2-aryl-4-piperidones **2a–h** which were converted to the desired 4-methylene derivatives **3a–h** using a Wittig olefination (Scheme 2).

### Scheme 2. Preparation of *N*-Boc-2-aryl-4-methylenepiperidines




<sup>a</sup>Yields quoted over two steps.

Applying knowledge from previous work,<sup>11</sup> it was pleasing that lithiation of piperidine **3a** occurred smoothly at  $-40\text{ }^{\circ}\text{C}$  in THF. The organolithium intermediate could be trapped with a variety of electrophiles to give 2,2-disubstituted compounds **4a–9a** in 70–90% yields (Scheme 3). Electrophiles screened included methyl chloroformate (to give **4a**), alkyl or allyl halides (to give **5a–7a**),  $\text{Bu}_3\text{SnCl}$ , and  $\text{Me}_3\text{SiCl}$  (to give **8a–9a**).


The successful formation of products **4a–9a** suggests that rotation of the Boc group is rapid under the conditions of the reaction. The base *n*-BuLi is known to coordinate to the carbonyl group prior to lithiation.<sup>12</sup> Lithiation occurs only from the rotamer in which the carbonyl group is directed toward the benzylic proton. The rate of rotation of the carbonyl was determined by VT-NMR spectroscopy (see the Supporting Information). Coalescence of the signals for the benzylic proton in the <sup>1</sup>H NMR spectrum at  $\sim 5.5$  ppm occurred at about 253 K. Line shape analysis of the mixture of rotamers (ratio  $\sim 1.2:1$ ) showed that the activation parameters

### Scheme 3. Initial Lithiation–Trapping Studies



for rotation of the Boc group were  $\Delta H^\ddagger \approx 50$  kJ/mol and  $\Delta S^\ddagger \approx -10$  J/K/mol, and hence, the half-life for rotation  $t_{1/2} \approx 10$  s at 195 K ( $\Delta G^\ddagger \approx 52$  kJ mol<sup>-1</sup>; see the Supporting Information).

DFT calculations on the rotamers of piperidine **3a** were also carried out based on previous work using the B3LYP-D3BJ functional with the def2-TZVP basis set (B3LYP-D3BJ//def2-TZVP).<sup>13–15</sup> The minimal energy structures for the rotamers of piperidine **3a** were found to be when the phenyl group occupied an axial position (Figure 2a,c). Of these two




**Figure 2.** Optimized geometries of **3a** in THF solution. (a,c) Minimum energy structures with the phenyl group in the axial position. (b) Lowest energy transition state with the phenyl group in the equatorial position.

structures, the rotamer in which the carbonyl group is pointing toward the benzylic proton was lower in thermal energy by  $\sim 170$  Jmol<sup>-1</sup>. Transition-state calculations indicated rotation of the Boc group was most likely to occur through the lowest energy equatorial transition state (Figure 2b). The Gibbs energy of activation was calculated to be  $\sim 55$  kJ/mol at 195 K (with  $\Delta H \approx 43$  kJ/mol and  $\Delta S \approx -63$  J/K/mol). This value matches well with the results obtained from VT-NMR and indicates that the rate of ring flipping between the axial and equatorial positions of the phenyl group was fast along with rotation of the Boc group under the reaction conditions.

Having obtained the knowledge that rotation of the Boc group is fast and that racemic lithiation–trapping is successful, we began investigating the kinetic resolution of piperidine **3a** using the chiral base system *n*-BuLi/(+)-sparteine. Optimization of the reaction conditions (see the Supporting Information) led to the best results when 0.8 equiv of *n*-

BuLi was added to a mixture containing the piperidine **3a** and 0.9 equiv of (+)-sparteine in toluene at  $-78\text{ }^{\circ}\text{C}$  (Scheme 4).


#### Scheme 4. Optimized Kinetic Resolution of **3a**



Quenching the reaction after 1 h with MeOCOCl gave the recovered (S)-**3a** in 41% yield with an excellent enantiomer ratio (er 97:3) together with the substituted product (R)-**4a** (58%, er 73:27). This equates to a selectivity factor  $s \approx 16$ .

These initial optimized conditions for piperidine **3a** were then applied to other 2-aryl-4-methylenepiperidines to obtain similar results (Scheme 5). The kinetic resolution reaction

#### Scheme 5. Scope of Kinetic Resolution




<sup>a</sup>Using 1.0 equiv of *n*-BuLi, 1.1 equiv of (+)-sparteine (and 2 h for **3e** and **3h**). <sup>b</sup>Using (–)-sparteine.

tolerated a variety of functional groups in the *meta* or *para* position of the 2-aryl ring including electron-donating and -withdrawing groups (Me, OMe, F, Cl, CF<sub>3</sub>). However, limitations of the reaction could be observed with the methoxy-substituted substrates (**3e**, **3g–h**) where more equivalents of *n*-BuLi and (+)-sparteine were required (1.0 and 1.1 equiv, respectively) to achieve good enantiomer ratios of the recovered 2-aryl-4-methylenepiperidines. A possible explanation for this is that some of the chiral base may coordinate to the oxygen atom in the methoxy group; hence, more equivalents of the base were required to ensure adequate deprotonation. By applying this revised method, and sometimes by increasing the lithiation time to 2 h, the recovered

(S)-**3e**, **3g–h** were isolated with good enantiomer ratios although typically with slightly lower yields.

In addition to changing the aryl group in the 2-position, the scope of the kinetic resolution reaction could be expanded further. By using the chiral ligand (–)-sparteine the enantioselectivity of the reaction could be inverted. This allowed formation of the recovered starting material (R)-**3a** with good selectivity, along with the substituted product (S)-**4a** (Scheme 5). The optimized reaction conditions for the kinetic resolution of piperidines **3a** and **3b** were tested at larger (gram) scales (Scheme 6). In both reactions, by applying an


#### Scheme 6. Scale-up of the Kinetic Resolution



acid–base wash on the crude products, the chiral ligand (+)-sparteine could be recovered in good yields. After column chromatography, the recovered starting materials (S)-**3a** and (S)-**3b** were isolated in good yields and selectivities along with their corresponding quenched products.

Having investigated the scope of the kinetic resolution reaction with various *N*-Boc-2-aryl-4-methylenepiperidines we sought to illustrate their use as potential fragments in organic synthesis. To explore this possibility, several reactions were carried out on piperidine (S)-**3a** (er 99:1) to generate a diverse range of compounds without loss of enantiopurity. Lithiation of piperidine (S)-**3a** with *n*-BuLi in THF at  $-78\text{ }^{\circ}\text{C}$  followed by trapping with MeOCOCl gave ester (S)-**4a** and maintained the high enantioenrichment (er 99:1) (Scheme 7) due to the


#### Scheme 7. Lithiation–Trapping of (S)-**3a** and Removal of Boc




configurationally stable intermediate organolithium species.<sup>11</sup> Removal of the Boc group in piperidine (S)-**3a** was achieved using HCl in dioxane, which gave piperidine (S)-**10** as the hydrochloride salt. The same reaction on piperidine (S)-**3b** (er 94:6) gave (S)-**11** as the hydrochloride salt which was recrystallized. Subsequent reattachment of the Boc group to give (S)-**3b** followed by chiral stationary-phase HPLC confirmed an improvement in the enantiomeric ratio (er 99:1) (Scheme 8).

In addition to these transformations, functionalization of the 4-methylene group in piperidine (S)-**3a** was investigated. Hydroboration of the alkene was carried out using BH<sub>3</sub>·THF, and the intermediate organoborane was oxidized to give alcohol **11** as a single diastereoisomer (dr 99:1) in 67% yield (Scheme 9). This was subsequently reacted with *p*-bromobenzoyl chloride in the presence of triethylamine and

## Scheme 8. Improving the Enantiomeric Ratio of (S)-3b



## Scheme 9. Alkene Functionalization of (S)-3a



DMAP, and the absolute configuration of the ester product was determined by single-crystal X-ray analysis (Figure 3).




Figure 3. X-ray analysis of *p*-bromobenzoate **12** (ellipsoids at 50% probability).

This confirmed the *cis*-stereochemistry of **12** and that the piperidine **3a** had the 2*S* configuration, as expected based on the use of (+)-sparteine.<sup>16,17</sup> Alternatively, ozonolysis of the 4-methylene group in piperidine (S)-**3a** was carried out at  $-78$  °C, and subsequent reductive workup using Me<sub>2</sub>S gave the enantioenriched 2-aryl-4-piperidone (S)-**2a** in high yield. This could be reacted with DAST to give difluoride (S)-**14** without any loss in enantiopurity. Furthermore, this compound was found to be stable toward lithiation using *n*-BuLi in THF at  $-78$  °C. Trapping the organolithium intermediate with methyl chloroformate gave the ester (S)-**15** without loss of enantiopurity.

In summary, kinetic resolution by deprotonation of 2-arylpiperidines can be extended to their 4-methylene derivatives. High enantiomer ratios can be obtained by proton abstraction preferentially from one enantiomer of the substrates using the base *n*-BuLi and the chiral ligand sparteine (either enantiomer). The *N*-Boc group was found to rotate rapidly under the reaction conditions, and therefore, the presence of both rotamers is not detrimental to the lithiation chemistry. Functional group interconversion of the 4-methylene group provides access to a selection of different

substituted piperidines. The ability to achieve an effective kinetic resolution allows the asymmetric synthesis of 2,4-disubstituted and 2,2,4-trisubstituted piperidines.

## ■ ASSOCIATED CONTENT

## Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.2c00862>.

Experimental procedures; characterization data; NMR spectra; HPLC traces; X-ray crystallographic data; VT-NMR spectra and analysis; DFT analyses with Cartesian coordinates (PDF)

## Accession Codes

CCDC 2155914 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

## ■ AUTHOR INFORMATION

## Corresponding Author

Iain Coldham – Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, U.K.; [orcid.org/0000-0003-4602-6292](https://orcid.org/0000-0003-4602-6292); Email: [i.coldham@sheffield.ac.uk](mailto:i.coldham@sheffield.ac.uk)

## Authors

Anthony Choi – Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, U.K.

Anthony J. H. M. Meijer – Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, U.K.; [orcid.org/0000-0003-4803-3488](https://orcid.org/0000-0003-4803-3488)

Ilaria Proietti Silvestri – Liverpool ChiroChem, Department of Chemistry, University of Liverpool, Liverpool L69 7ZD, U.K.; [orcid.org/0000-0002-4951-3072](https://orcid.org/0000-0002-4951-3072)

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.joc.2c00862>

## Notes

For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising.

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We acknowledge support for this research from the EPSRC (Grant No. EP/R024294/1), the Royal Society (Short Industry Fellowship SIF\R2\202031), and the University of Sheffield. We thank Craig Robertson for the single-crystal X-ray analysis and Khalid Douidin for help with NMR spectra. We acknowledge the Faculty of Science mass spectrometry service at the University of Sheffield.

## ■ REFERENCES

- (1) Bari, A.; Iqbal, A.; Khan, Z. A.; Shahzad, S. A.; Yar, M. Synthetic Approaches toward Piperidine Related Structures: A Review. *Synth. Commun.* **2020**, *50*, 2572–2589.
- (2) Srivastava, N.; Macha, L.; Ha, H.-J. Stereoselective Synthesis of 2,6-Disubstituted Piperidine Alkaloids. *Org. Biomol. Chem.* **2020**, *18*, 5493–5512.
- (3) Munchhof, M. J.; Li, Q.; Shavnya, A.; Borzillo, G. V.; Boyden, T. L.; Jones, C. S.; LaGreca, S. D.; Martinez-Alsina, L.; Patel, N.;

Pelletier, K.; Reiter, L. A.; Robbins, M. D.; Tkalcovic, G. T. Discovery of PF-04449913, a Potent and Orally Bioavailable Inhibitor of Smoothed. *ACS Med. Chem. Lett.* **2012**, *3*, 106–111.

(4) Mainolfi, N.; Ehara, T.; Karki, R. G.; Anderson, K.; Mac Sweeney, A.; Liao, S.-M.; Argikar, U. A.; Jendza, K.; Zhang, C.; Powers, J.; Klosowski, D. W.; Crowley, M.; Kawanami, T.; Ding, J.; April, M.; Forster, C.; Serrano-Wu, M.; Capparelli, M.; Ramqaj, R.; Solovay, C.; Cumin, F.; Smith, T. M.; Ferrara, L.; Lee, W.; Long, D.; Prentiss, M.; De Erkenez, A.; Yang, L.; Liu, F.; Sellner, H.; Sirockin, F.; Valeur, E.; Erbel, P.; Ostermeier, D.; Ramage, P.; Gerhartz, B.; Schubart, A.; Flohr, S.; Gradoux, N.; Feifel, R.; Vogg, B.; Wiesmann, C.; Maibaum, J.; Eder, J.; Sedrani, R.; Harrison, R. A.; Mogi, M.; Jaffee, B. D.; Adams, C. M. Discovery of 4-((2*S*,4*S*)-4-Ethoxy-1-((5-methoxy-7-methyl-1*H*-indol-4-yl)methyl)piperidin-2-yl)benzoic Acid (LNP023), a Factor B Inhibitor Specifically Designed To Be Applicable to Treating a Diverse Array of Complement Mediated Diseases. *J. Med. Chem.* **2020**, *63*, 5697–5722.

(5) (a) For examples of piperidines in fragment libraries, see: Wang, G.; Chen, L.; Xian, T.; Liang, Y.; Zhang, X.; Yang, Z.; Luo, M. Discovery and SAR Study of Piperidine-Based Derivatives as Novel Influenza Virus Inhibitors. *Org. Biomol. Chem.* **2014**, *12*, 8048–8060. (b) Johnson, J. A.; Nicolaou, C. A.; Kirberger, S. E.; Pandey, A. K.; Hu, H.; Pomerantz, W. C. K. Evaluating the Advantages of Using 3D-Enriched Fragments for Targeting BET Bromodomains. *ACS Med. Chem. Lett.* **2019**, *10*, 1648–1654. (c) Downes, T. D.; Jones, S. P.; Klein, H. F.; Wheldon, M. C.; Atobe, M.; Bond, P. S.; Firth, J. D.; Chan, N. S.; Waddelove, L.; Hubbard, R. E.; Blakemore, D. C.; De Fusco, C.; Roughley, S. D.; Vidler, L. R.; Whatton, M. A.; Woolford, A. J.-A.; Wrigley, G. L.; O'Brien, P. Design and Synthesis of 56 Shape-Diverse 3D Fragments. *Chem.—Eur. J.* **2020**, *26*, 8969–8975.

(6) Cochrane, E. J.; Leonori, D.; Hassall, L. A.; Coldham, I. Synthesis and Kinetic Resolution of *N*-Boc-2-arylpiperidines. *Chem. Commun.* **2014**, *50*, 9910–9913.

(7) (a) For other examples of lithiated/arylated piperidines, see: Mitchell, E. A.; Peschiulli, A.; Lefevre, N.; Meerpoel, L.; Maes, B. U. W. Direct  $\alpha$ -Functionalization of Saturated Cyclic Amines. *Chem.—Eur. J.* **2012**, *18*, 10092–10142. (b) Seel, S.; Thaler, T.; Takatsu, K.; Zhang, C.; Zipse, H.; Straub, B. F.; Mayer, P.; Knochel, P. Highly Diastereoselective Arylations of Substituted Piperidines. *J. Am. Chem. Soc.* **2011**, *133*, 4774–4777. (c) Millet, A.; Larini, P.; Clot, E.; Baudoin, O. Ligand-controlled  $\beta$ -selective C(sp<sup>3</sup>)-H arylation of *N*-Boc-piperidines. *Chem. Sci.* **2013**, *4*, 2241–2247. (d) Tait, M. B.; Butterworth, S.; Clayden, J. 2,2- and 2,6-Diarylpiperidines by Aryl Migration within Lithiated Urea Derivatives of Tetrahydropyridines. *Org. Lett.* **2015**, *17*, 1236–1239. (e) Beng, T. K.; Takeuchi, H.; Weber, M.; Sarpong, R. Stereocontrolled synthesis of vicinally functionalized piperidines by nucleophilic  $\beta$ -addition of alkylolithiums to  $\alpha$ -aryl substituted piperidine enecarbamates. *Chem. Commun.* **2015**, *51*, 7653–7656.

(8) (a) Li, W.; Yang, H.; Li, R.; Lv, H.; Zhang, X. Kinetic Resolution of Racemic 3,4-Disubstituted 1,4,5,6-Tetrahydropyridine and 3,4-Disubstituted 1,4-Dihydropyridines via Rh-Catalyzed Asymmetric Hydrogenation. *ACS Catal.* **2020**, *10*, 2603–2608. (b) Wang, T.; Du, L.-D.; Wan, D.; Li, X.; Chen, X.-Z.; Wu, G.-F. Use of Lipase Catalytic Resolution in the Preparation of Ethyl (2*S*,5*R*)-5-((Benzyloxy)-amino)piperidine-2-carboxylate, a Key Intermediate of the  $\beta$ -Lactamase Inhibitor Avibactam. *Org. Process Res. Dev.* **2018**, *22*, 1738–1744. (c) France, S. P.; Hussain, S.; Hill, A. M.; Hepworth, L. J.; Howard, R. M.; Mulholland, K. R.; Flitsch, S. L.; Turner, N. J. One-Pot Cascade Synthesis of Mono- and Disubstituted Piperidines and Pyrrolidines using Carboxylic Acid Reductase (CAR),  $\omega$ -Transaminase ( $\omega$ -TA), and Imine Reductase (IRED) Biocatalysts. *ACS Catal.* **2016**, *6*, 3753–3759. (d) Lei, B.-L.; Zhang, Q.-S.; Yu, W.-H.; Ding, Q.-P.; Ding, C.-H.; Hou, X.-L. Kinetic Resolution of 2-Substituted 2,3-Dihydro-4-pyridones by Palladium-Catalyzed Asymmetric Allylic Alkylation: Catalytic Asymmetric Total Synthesis of Indolizidine (–)-209l. *Org. Lett.* **2014**, *16*, 1944–1947. (e) Solymár, M.; Forró, E.; Fülöp, F. Enzyme-catalyzed kinetic resolution of piperidine hydroxy esters. *Tetrahedron: Asym.* **2004**, *15*, 3281–3287.

(9) (a) Wanner, B.; Kreituss, I.; Gutierrez, O.; Kozłowski, M. C.; Bode, J. W. Catalytic Kinetic Resolution of Disubstituted Piperidines by Enantioselective Acylation: Synthetic Utility and Mechanistic Insights. *J. Am. Chem. Soc.* **2015**, *137*, 11491–11497. See also: (b) Dooley, C. J.; Burtea, A.; Mitilian, C.; Dao, W. T.; Qu, B.; Salzameda, N. T.; Rychnovsky, S. D. Using the Competing Enantioselective Conversion Method to Assign the Absolute Configuration of Cyclic Amines with Bode's Acylation Reagents. *J. Org. Chem.* **2020**, *85*, 10750–10759.

(10) Guo, F.; Dhakal, R. C.; Dieter, R. K. Conjugate Addition Reactions of *N*-Carbamoyl-4-pyridones and 2,3-Dihydropyridones with Grignard Reagents in the Absence of Cu(I) Salts. *J. Org. Chem.* **2013**, *78*, 8451–8464.

(11) Sheikh, N. S.; Leonori, D.; Barker, G.; Firth, J. D.; Campos, K. R.; Meijer, A. J. H. M.; O'Brien, P.; Coldham, I. An Experimental and in Situ IR Spectroscopic Study of the Lithiation–Substitution of *N*-Boc-2-Phenylpyrrolidine and -Piperidine: Controlling the Formation of Quaternary Stereocenters. *J. Am. Chem. Soc.* **2012**, *134*, 5300–5308.

(12) Bertini Gross, K. M.; Beak, P. Complex-Induced Proximity Effects: The Effect of Varying Directing-Group Orientation on Carbamate-Directed Lithiation Reactions. *J. Am. Chem. Soc.* **2001**, *123*, 315–321.

(13) (a) Carter, N.; Li, X.; Reavey, L.; Meijer, A. J. H. M.; Coldham, I. Synthesis and Kinetic Resolution of Substituted Tetrahydroquinolines by Lithiation then Electrophilic Quench. *Chem. Sci.* **2018**, *9*, 1352–1357. (b) Choi, A.; El-Tunsi, A.; Wang, Y.; Meijer, A. J. H. M.; Li, J.; Li, X.; Proietti Silvestri, I.; Coldham, I. Asymmetric synthesis of 2-arylidolines and 2,2-disubstituted indolines by kinetic resolution. *Chem.—Eur. J.* **2021**, *27*, 11670–11675. (c) El-Tunsi, A.; Carter, N.; Yeo, S.-H.; Priest, J. D.; Choi, A.; Kobras, C. M.; Ndlovu, S.; Proietti Silvestri, I.; Fenton, A. K.; Coldham, I. Kinetic Resolution by Lithiation: Highly Enantioselective Synthesis of Substituted Dihydrobenzoxazines and Tetrahydroquinoxalines. *Synthesis* **2022**, *54*, 355–368.

(14) Becke, A. D. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652.

(15) Weigend, F.; Ahlrichs, R. Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297–3305.

(16) Kasten, K.; Selig, N.; O'Brien, P. Enantioselective Lithiation–Substitution of Nitrogen-Containing Heterocycles. *Org. Reactions* **2019**, *100*, 255–328.

(17) Bailey, W. F.; Beak, P.; Kerrick, S. T.; Ma, S.; Wiberg, K. B. An Experimental and Computational Investigation of the Enantioselective Deprotonation of Boc-piperidine. *J. Am. Chem. Soc.* **2002**, *124*, 1889–1896.