

This is a repository copy of *“Everything the hujur tells is very educative but if I cannot apply those in my own life then there is no meaning”*:A mixed-methods process evaluation of a smoke-free homes intervention in Bangladesh.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/191743/>

Version: Accepted Version

Article:

Jackson, Catherine, Azdi, Zunayed, Kellar, Ian et al. (9 more authors) (2022) *“Everything the hujur tells is very educative but if I cannot apply those in my own life then there is no meaning”*:A mixed-methods process evaluation of a smoke-free homes intervention in Bangladesh. BMC Public Health. 1889. ISSN 1471-2458

<https://doi.org/10.1186/s12889-022-14283-6>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 ***“Everything the hujur tells is very educative but if I cannot apply those in my own***
2 ***life then there is no meaning”***: A mixed-methods process evaluation of a smoke-
3 **free homes intervention in Bangladesh.**

4

5 Cath Jackson^{1,2}, Zunayed Al Azdi³, Ian Kellar⁴, Noreen Dadirai Mdege¹, Caroline Fairhurst¹, Tarana Ferdous³,
6 Catherine Hewitt¹, Rumana Huque³, Anna-Marie Marshall¹, Sean Semple⁵, Aziz Sheikh⁶, Kamran Siddiqi¹,
7 MCLASS II trial team*

8

9 Corresponding author: Cath Jackson

10

11 1. Department of Health Sciences, University of York, York, UK

12 2. Valid Research Ltd, Wetherby, UK

13 3. ARK Foundation, Dhaka, Bangladesh

14 4. School of Psychology, University of Leeds, Leeds, UK

15 5. Institute for Social Marketing and Health, University of Stirling, Stirling, Scotland, UK

16 6. Usher Institute, University of Edinburgh, Edinburgh, Scotland, UK

17

18 **ABSTRACT**

19 **Background**

20 Second-hand smoke exposure from tobacco significantly contributes to morbidity and mortality worldwide.
21 A cluster RCT in Bangladesh compared a community-based smoke-free home (SFH) intervention delivered
22 in mosques, with or without indoor air quality (IAQ) feedback to households to no intervention. Neither
23 was effective nor cost-effective compared to no intervention using an objective measure of second-hand
24 smoke. This paper presents the process evaluation embedded within the trial and seeks to understand this.

25
26 **Methods**

27 A mixed method process evaluation comprising interviews with 30 household leads and six imams (prayer
28 leader in mosque), brief questionnaire completed by 900 household leads (75% response), fidelity
29 assessment of intervention delivery in six (20%) mosques and research team records. Data were
30 triangulated using meta-themes informed by three process evaluation functions: implementation,
31 mechanisms of impact and context.

32
33 **Results**

34 IMPLEMENTATION: Frequency of SFH intervention delivery was judged moderate to good. However there
35 were mixed levels of intervention fidelity and poor reach. Linked Ayahs (verses of the Qur'an) with health
36 messages targeting SHS attitudes were most often fully implemented and had greatest reach (along with
37 those targeting social norms). Frequency and reach of the IAQ feedback were good. MECHANISMS OF
38 IMPACT: Both interventions had good acceptability. However, views on usefulness of the interventions in
39 creating a SFH were mixed. Individual drivers to behaviour change were new SFH knowledge with
40 corresponding positive attitudes, social norms and intentions. Individual barriers were a lack of self-efficacy
41 and plans. CONTEXT: Social context drivers to SFH intervention implementation in mosques were in place
42 and important. No context barriers to implementation were reported. Social context drivers to SHS
43 behaviour change were children's requests. Barriers were women's reluctance to ask men to smoke outside

44 alongside general reluctance to request this of visitors. (Not) having somewhere to smoke outside was a
45 physical context (barrier) and driver.

46

47 **Conclusions**

48 Despite detailed development and adaption work with relevant stakeholders, the SFH intervention and IAQ
49 feedback became educational interventions that were motivational but insufficient to overcome significant
50 context barriers to reduce objectively measured SHS exposure in the home. Future interventions could
51 usefully incorporate practical support for SFH behaviour change. Moreover, embedding these into
52 community wide strategies that include practical cessation support and enforcement of SFH legislation is
53 needed.

54

55 **Study Registration**

56 Current Controlled Trials [ISRCTN49975452](https://www.clinicaltrials.gov/ct2/show/study?term=ISRCTN49975452)

57

58 **Key words:** tobacco, second-hand smoke, smoke free homes, faith, mosque, intervention, process
59 evaluation, Bangladesh

60

61

62 **BACKGROUND**

63 Exposure to second-hand tobacco smoke (SHS) is estimated to cause 1.2 million deaths and loss of 11
64 million disability-adjusted life years worldwide every year [1]. Our focus was Bangladesh and SHS exposure
65 in homes. In a recent study of 1746 households in Mirpur, Dhaka, over half (55%) self-reported that
66 smoking by household members and visitors was permitted inside the home [2]. Unfortunately, evidence of
67 effective interventions in South Asia to reduce SHS exposure in the home is lacking [3-5]. Moreover, poor
68 reporting means that the intervention elements with greatest efficacy are difficult to identify [3-5].

69
70 International literature shows an association between religious faith and reducing or eliminating smoking
71 behaviours [6-12] with proposed mechanisms including the idea of leading a “puritanical” life, having
72 spiritual strength to resist temptations for future benefit, and being part of a social network of people who
73 lead healthy lives. Relatedly, religious leaders are often highly respected and trusted by their communities
74 [7-12]. Together, these suggest that religious teachings, settings and leaders offer potential to deliver
75 tobacco control interventions.

76
77 In Bangladesh, 89% of the population is Muslim [13]. Islamic teachings focus on principles of minimising
78 harm to individuals and society; and maximising opportunities for individual and collective well-being [9].
79 As such, smoking is discouraged, although whether it is decreed as *mukrooh* (discouraged) or *haram*
80 (prohibited) varies [9]. To date, very few evaluations of Islamic faith-based interventions targeting smoking
81 behaviours have been undertaken [11,14,15].

82
83 A 2018 Cochrane review of interventions to promote smoke-free homes (SFH) reported that 24 of 78
84 included studies found statistically significant reductions in children’s SHS exposure [3]. No one
85 intervention strategy was identified as the gold standard. Successful strategies included motivational
86 interviewing, brief counselling, nicotine replacement therapy for smoking cessation for parents who smoke,
87 and feedback on markers of SHS exposure including the use of indoor air quality (IAQ) feedback. IAQ
88 feedback offers participants objectively measured information on the impact that smoking has on

89 concentrations of air pollutants in their homes to motivate them to reduce or stop smoking inside. This has
90 been effective in reducing SHS in homes and/or children's biomarkers of SHS exposure in several trials
91 across settings and formats, including immediate and delayed feedback [16-22].

92
93 We conducted a three-arm cluster randomised controlled trial, MCLASS (Muslim Communities Learning
94 About SHS) II, in 45 mosques from the Mirpur area of Dhaka, to evaluate effectiveness and cost-
95 effectiveness of a community-based SFH intervention delivered in mosques with (n=16) or without (n=14)
96 IAQ feedback in reducing exposure to SHS in the home [23,24]. Both interventions are described in Table 1.
97 Mosques in the control arm (n=15) received no intervention. We found that at 3- and 12-months post
98 randomisation there were no significant differences on mean 24-hour household airborne fine particulate
99 matter (<2.5 microns in diameter [PM2.5]) concentration between the SFH intervention, with or without
100 IAQ feedback, and no intervention. The interventions were also not cost-effective when compared to no
101 intervention. We therefore concluded that these interventions could not be recommended for Bangladesh
102 [24]. In this paper, we present the findings from our embedded process evaluation [25], to understand their
103 lack of influence on trial outcomes.

104

105 **Table 1:** Description of the content and delivery of SFH and IAQ feedback interventions

<p>SFH intervention</p> <p>CONTENT: A set of 12 health messages relating to smoking and SHS exposure, each supported by at least one verse (Ayah) from the Qur'an, or an Islamic faith-based decree. The messages were developed through a set of iterative workshops involving Islamic scholars, public health professionals and behavioural scientists [26]. They addressed key barriers and drivers of smoking behaviours (attitudes, self-efficacy, social norms, intention formation, action and coping planning, see Figure 1 and Additional file 1).</p> <p>DELIVERY: Imams and khatibs were trained in a half-day session on the intervention and its delivery including detailed guidance on linking the messages and Ayahs. They then delivered the messages and Ayahs in the form of Khutbah (formal sermon preached by the imam in Arabic) to those attending Friday Jumu'ah prayer over 12 weeks</p>
--

(one linked Ayah-message per week). They also distributed copies of a short SFH booklet to their congregation in any way they saw best. The booklet contained a brief description of the 12 linked Ayahs-messages.

IAQ feedback

CONTENT: A two-page personalised leaflet designed in consultation with community members. It contained feedback on the air quality (PM_{2.5} concentration) measured within a home at baseline using the Dylos DC 1700 (Dylos, California, USA), an optical particle counter validated for use in domestic settings. Specifically feedback comprised a comparison of the 24-hour mean PM_{2.5} concentration measured in the home to the World Health Organization (WHO) guidance limit of 25 µg/m³ [27], the total time the IAQ was above this guidance limit, and the maximum concentration measured during the 24-hour measurement period. It included graphical information on how smoking activity impacted on IAQ over the 24-hour measurement period (with classifications: hazardous if >150 µg/m³, unhealthy if 36-150 µg/m³, moderate if 12-35 µg/m³, and good if <12 µg/m³), information about the adverse effects of SHS exposure, recommendations to reduce SHS exposure in the home, and a target that was achievable by implementing SFH rules within the home.

DELIVERY: Trial field investigators delivered and discussed the personalised IAQ feedback with members of the households in person (in their homes) in approximately 10 minutes.

Both the SFH manual and IAQ feedback leaflet are available here [Muslim Communities Learning About Second-hand Smoke in Bangladesh \(MCLASS II\) - Health Sciences, University of York.](#)

106

107 **METHODS**

108 **Overview of study design**

109 This was a mixed method process evaluation conducted November 2018 to January 2019. It comprised
110 interviews with household leads (trial participants) and imams (prayer leader in mosque), a brief
111 questionnaire administered to household leads, fidelity assessment of intervention delivery and research
112 team records. Findings from the different data sets were triangulated using meta-themes [28] based on the
113 UK Medical Research Council's [25] three process evaluation functions:

- 114
- Implementation – what is delivered (frequency, fidelity, reach)?

- 115 • Mechanisms of impact – how does the delivered intervention produce change? (intervention
 116 acceptability and usefulness, individual barriers and drivers to SHS behaviour change)
- 117 • Context – how does context affect implementation and outcomes? (social and physical context barriers
 118 and drivers to intervention implementation, and to SHS behaviour change)
- 119 SHS behaviour change included smokers not smoking inside the home and non-smokers requesting
 120 residents and visitors to smoke outside.

121

122 **Interviews**

123 *Participants*

124 Semi-structured interviews were conducted post-intervention (at 3-month follow-up) with a sample of 30
 125 household leads (14 in SFH arm, 16 in SFH+IAQ arm). Household leads were the nominated trial participant
 126 for participating households (n=1801: 560 SFH, 640 SFH+IAQ, 601 control) where at least one adult resident
 127 was smoking regularly, at least one adult resident was a non-smoker and at least one resident attended a
 128 participating mosque. They were recruited to the trial at the mosque or through a home visit. We
 129 purposively selected household leads for interview to include men and women, smokers and non-smokers,
 130 with different descriptions of smoking in the home at 3-month follow-up (see Table 2). All imams who
 131 delivered the SFH intervention in six randomly selected mosques (3 from each intervention arm) were
 132 interviewed once intervention delivery was complete.

133

134 Two-thirds of household leads were men (n=20), and a similar proportion was aged <45 years (n=21). Over
 135 two-thirds (n=24) had no/only primary (1-5 years) education. At baseline, all men self-reported as smokers;
 136 no women were smokers. About two-thirds of participants (n=19) described their homes as smoke-free by
 137 3-month follow-up, defined as not permitting residents or visitors to smoke inside the home. The rest
 138 (n=11) described some/lots of smoking still occurring at home.

139

140 **Table 2:** Demographic characteristics and smoking/SFH status of interview participants

Characteristic	SFH (n=14)	SFH+IAQ (n=16)	All (n=30)
----------------	---------------	-------------------	---------------

		Men (n=10)	Women (n=4)	Men (n=10)	Women (n=6)	Men (n=20)	Women (n=10)
Age, years	18-25	2	0	0	0	2	0
	26-35	5	1	5	0	10	1
	36-45	2	2	1	3	3	5
	> 45	1	1	4	3	5	4
Education, total years	No education (0)	1	2	3	2	4	4
	Primary (1-5)	4	1	4	3	8	4
	Secondary (6-10)	2	1	3	1	5	2
	Higher secondary (10-12)	2	0	0	0	2	0
	University (>12)	1	0	0	0	1	0
Self-reported smoking status (at baseline)	Smoker	10	0	10	0	20	0
	Non-smoker	0	4	0	6	0	10
Description of smoking in the home (3-month follow-up) ^{a,b}	Nobody smoking	7	3	6	3	13	6
	Still some smoking	3	1	3	2	6	3
	Lots of smoking	0	0	1	1	1	1

141 ^aAll described smoking in the home at baseline. ^bThese descriptions may differ from the objective air quality
142 data collected in the trial.
143

144 All six imams were non-smokers (a pre-requisite of their mosque's inclusion in the trial). They had been an
145 imam for between 6 and 35 years, and 2 to 22 years in their current mosque. The size of their congregation
146 during Jum'ah prayers (a spiritually significant prayer offered during midday on Friday attended by men)
147 varied from 800 to 4500 men.

148

149 *Data collection*

150 Interviews were conducted in Bengali face-to-face in the household lead's home or at the imam's mosque.

151 All participants provided written informed consent before the interview commenced. Interviews with

152 household leads explored interaction with the SFH intervention/IAQ feedback, views about the

153 intervention(s), impact on SHS behaviours as well as individual or context barriers and drivers to creating a

154 SFH (Figure 1). These lasted 8-27 minutes. Interviews with imams explored acceptability of the SFH

155 intervention, and experiences of delivery including individual or context barriers and drivers. These lasted

156 25-53 minutes. All interviews were digitally audio-recorded.

157

158

Figure 1 in here

159

160

161 *Data analysis*

162 Interviews were transcribed verbatim, translated into English and checked by the interviewers. The data
163 were subjected to Framework analysis [29] by two researchers (ZAA, CJ). Excel 365 facilitated data
164 management.

165

166 An English language thematic framework was developed for each dataset based on the three process
167 evaluation functions (implementation, mechanisms of impact, context) and their components (e.g.
168 acceptability, social context barriers to SHS behaviour change). A sample of randomly selected interview
169 transcripts (seven – household lead, two – imam) were used to further refine the framework, e.g. identify
170 examples of social context barriers. The frameworks were piloted with more transcripts (three -household
171 lead, one - imam) before finalising. The data were then charted into the relevant frameworks. Summaries
172 of participant responses and verbatim quotes were entered. Both sets of charted data were then reviewed
173 and interrogated to compare views, seek patterns, connections, and explanations within the data.
174 Descriptive findings documents were written, organised by the components of the three process evaluation
175 functions.

176

177 **Questionnaire**

178 *Participants and data collection*

179 Household leads in the two intervention arms (SFH: 387 men, 33 women; SFH+IAQ: 461 men, 19 women;
180 75% response both arms) completed a short process evaluation questionnaire, administered face-to-face
181 by a researcher at 3-month follow up). It asked questions on which components of the SFH
182 intervention/IAQ feedback participants had received and perceived intervention usefulness.

183

184 *Data analysis*

185 Yes/no/don't know responses were used for the intervention receipt questions. Perceived intervention
186 usefulness was scored on a 7-point Likert scale from 1 (not at all useful) to 7 (extremely useful). Scores of 5
187 and above were classified as useful. Data were analysed using frequencies and proportions.

188

189 **Fidelity assessment**

190 *Data collection*

191 Delivery of the SFH intervention was observed in six (20%) randomly selected mosques. Trained researchers
192 conducted these checks and completed a fidelity index. Imams had previously received training on
193 delivering the linked Ayahs (verses from the Qur'an) and health messages. They were unaware that they
194 were being observed. In three mosques, delivery of Ayahs-messages scheduled for odd numbered weeks
195 (1,3,5 etc.) were checked. In the other three mosques, Ayahs-messages scheduled for even numbered
196 weeks (2,4,6 etc.) were checked. Each item in the index corresponded with the 12 weeks of Ayahs-
197 messages targeting five key barriers/drivers to SHS behaviours (see Figure 1 and Additional file 1). Delivery
198 of each Ayah-message was scored 0–not implemented, 1–Ayah recited with no message, 2–Ayah recited
199 with partial explanation of message, 3–Ayah recited with more than partial explanation but not full
200 explanation of message, and 4–fully implemented. Definitions were provided for each Ayah-message
201 (available from authors on request).

202

203 *Data analysis*

204 For each mosque, a total fidelity score was computed by summing the scores for Ayahs-messages from 0
205 (did not implement any Ayahs-messages) to 24 (all assessed Ayahs-messages were fully implemented). For
206 each target barrier/driver (Figure 1), we counted the number of times the Ayah-message was
207 fully/partially/not implemented and divided this by the total number of opportunities for full
208 implementation, for example, for "attitude" total number is 12 (3 mosques x 4 Ayahs-messages).

209

210 **Research team records**

211 *Data collection*

212 Records were collected from mosques on their self-reported delivery of the SFH intervention. Field
213 investigators self-recorded delivery of the IAQ feedback and a signature from the recipient was collected.

214

215 *Data analysis*

216 Counts and percentages were calculated for both delivery items.

217

218 **Triangulating findings**

219 To triangulate the findings from the different datasets, the key findings for each intervention (SFI, IAQ
220 feedback) from each dataset were displayed in a triangulation matrix (Additional files 2, 3) organised by the
221 three meta-themes [28]: implementation, mechanisms of impact and context [25]. For each meta-theme,
222 one or more datasets provided findings. Where there was more than one, these were compared to
223 consider if they were convergent (in agreement), complementary (partial agreement), contradictory
224 (disagreement) or silent (findings do not occur in a dataset but may have been expected to do so) [28].

225

226 **FINDINGS**

227 **IMPLEMENTATION**

228 **Frequency of SFH intervention delivery was judged moderate to good. There were mixed levels of**
229 **intervention fidelity and poor reach. Ayahs-messages targeting attitudes were most often fully**
230 **implemented and had greatest reach (along with those targeting social norms).**

231 Records showed that 29 of the 30 mosques (97%) reported delivering all 12 weeks of the SFH intervention.

232 The other mosque delivered 10 weeks. Imams typically reported they had delivered “almost all” of the SFH
233 intervention as instructed, during Jum’ah prayer, before Khutbah (formal sermon preached by the imam in
234 Arabic before the prayer) usually for 5-10 minutes. Two admitted to not delivering all 12 weeks. All
235 described using other opportunities to share the Ayahs-messages in the mosque including in the Madrasas
236 (educational institutions teaching Islamic subjects) and Maghrib (evening) prayers.

237

238 Whilst these convergent record and interview data indicated moderate-to-good frequency of intervention
239 delivery, the questionnaire data revealed poor intervention reach. Only half of men in both intervention
240 arms reported receiving the SFH intervention (SFH 49.4%; SFH+IAQ 55.5%). Women typically do not attend
241 Friday prayers, so were asked if any family members had heard the Ayahs-messages. Once again, only half
242 reported yes (SFH 51.5%; SFH+IAQ 52.6%). The interview data were more positive. All but three men
243 reported having received the SFH intervention and only one woman was unaware of family members
244 receiving it. For those men whom the intervention did reach, this was during Friday Jum'ah prayers (SFH
245 99.5%; SFH+IAQ 99.6%), with all women mentioning this for family members. Less than 3% of men reported
246 receiving the SFH intervention in other mosque sessions. This reach via Friday prayers was confirmed in the
247 interview data, thus both data sets supported the imams' delivery accounts.

248

249 *The imam said directly, "Never smoke at home." When he was delivering Khutbah, that time*
250 *he talked about it.*

251 [Man, SFH intervention, nobody smokes in home at 3-month follow-up]

252

253 *Yes. I have come to know about it from my younger son. He goes to Jumu'ah always. I need*
254 *not send him, he goes for his prayers by himself. Hujur (prayer leader at the mosque) tells*
255 *many Hadith (silent approvals of the prophet Muhammad) and gives speeches on smoking.*

256 [Woman, SFH intervention, nobody smokes in home at 3-month follow-up]

257

258 Regarding the detail of what was delivered by the imams, the mean fidelity score across six mosques was
259 19.6 (SD 2.51, range 16-22 of maximum 24). Ayahs-messages best delivered targeted attitudes and were
260 75.0% fully implemented. Ayahs-messages targeting self-efficacy and coping planning were 66.67% fully
261 implemented. Ayahs-messages targeting social norms and intention formation-action planning were only
262 50.0% fully implemented (see Table 3).

263

264 **Table 3:** Fidelity to delivery of SFH intervention

Target barrier/driver, n (%)	Level of implementation (%)					No data ^a
	Full	Partial – level 3	Partial – level 2	Partial – level 1	Not implemented	
Attitude, n=12	9 (75.0)	1 (8.3)	0 (0.0)	1 (8.3)	1 (8.3)	0 (0.0)
Self-efficacy, n=6	4 (66.7)	1 (16.7)	1 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)
Coping planning, n=6	4 (66.7)	1 (16.7)	1 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)
Social norms, n=6	3 (50.0)	0 (0.0)	1 (16.7)	0 (0.0)	1 (16.7)	1 (16.7)
Intention formation – action planning, n=6	3 (50.0)	3 (50.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

265 Note. Delivery of each Ayah-message was scored 0–not implemented, 1–Ayah recited with no message, 2– Ayah
266 recited with partial explanation of message, 3- Ayah recited with more than partial explanation but not full
267 explanation of message, 4-fully implemented. Ayahs-messages linked to attitudes were scheduled for delivery in four
268 weeks. The other four target barriers/drivers were scheduled for two weeks each. ^aNo assessment as this was
269 scheduled during the Eid festival.

270

271 Interview and questionnaire data partially confirmed this. Imams described focusing particularly on the
272 Ayahs-messages about risks of SHS to children, pregnant women, and others (targeting attitudes and social
273 norms). This preference was unrelated to the ease/difficulty of delivery (they were confident with all 12).
274 Instead, they believed their congregation were interested in learning about this, given that it is not usually
275 spoken about in the mosques.


276

277 These were also the Ayahs-messages that men most recalled hearing (79.1% to 94.8%, see Figure 2). All but
278 three men interviewed mentioned hearing Ayahs-messages about the risks of SHS, citing the dangers of
279 polluting their home and damaging the health of their family, particularly their children. Most also
280 remembered the clear direction from the imam within these Ayahs-messages to stop smoking near other
281 people.

282

283 *If I smoke, people who are around me are also harmed. Cause when I breathe out the*
284 *smoke, the people around inhale the oxygen or the air, they are also harmed. They are*
285 *harmed more than me. Then it is seen, when a child is born or a woman is pregnant,*
286 *smoking harms her children.*

287 [Man, SFH intervention, nobody smokes in home at 3-month follow-up]



288

289 **Figure 2:** Percentage recall (reach) of SFH intervention Ayahs-messages by men who had received the SFH
 290 intervention

291 *Note.* Att=attitude, SN=social norm, SE=self-efficacy, CP=coping planning, IF=intention formation

292

293 Noticeably less well recalled by men were Ayahs-messages targeting self-efficacy, coping planning and
 294 intention formation (37.5% to 45.0%, see Figure 2). Just five men who were interviewed mentioned that
 295 the imam provided guidance on “how” to change their smoking behaviours, whilst a similar minority
 296 declared the imam provided no advice at all.

297

298 Finally, the intention was that 100 copies of a short SFH booklet would be distributed in each mosque, thus
 299 reaching 3,000 households in total. Imams were unanimous that the booklets were popular, copies were
 300 distributed quickly, and more were needed. Some had targeted smokers, elders, or people they considered
 301 to be educated who would most benefit.

302

303 *We can understand who smokes. We tried to give it to them. Besides them, there are many*
 304 *educated people who want to know about it. We distributed among those educated and*
 305 *smokers. [Imam 2]*

306

307 The interview data suggested that reach of the booklet was poor. No men interviewed reported receiving it
308 and some added they could not have read it anyway. Three women mentioned that their sons had brought
309 the booklet home, two of whom could not read.

310

311 *We have received it, but we could not understand what the booklet was about, so we have*
312 *thrown it away. We are women so we don't understand all these things.*

313 [Woman, SFH+IAQ, nobody smokes in home at 3-month follow-up]

314

315 **Frequency and reach of IAQ feedback were good. Fidelity was not assessed.**

316 Research team records that included a signature from households showed that IAQ feedback was delivered
317 to all 640 households (100%) in that trial arm indicating good intervention frequency. Good reach was also
318 achieved with 98.9% of household leads and 13 of 15 interview participants (men and women) reporting
319 having received the IAQ feedback. Half of interview participants (men and women) mentioned that another
320 family member had received the report. A few commented they could not read the IAQ report, relying on
321 others to do this.

322

323 *Nobody can read in the home. The youngest daughter read it us twice or three times. After her*
324 *departure, we were unable get information from it.*

325 [Man, SFH+IAQ, nobody smokes in home at 3-month follow-up]

326

327 **MECHANISMS OF IMPACT**

328 **SFH intervention acceptability was good. Drivers were new SFH knowledge with corresponding positive**
329 **attitudes, social norms and intentions. Barriers were a lack of self-efficacy and plans.**

330 The male household lead and imam interview data were convergent indicating good acceptability of the
331 SFH intervention. The consensus amongst the men was that listening to the messages in the mosque "*felt*
332 *good*", informed them and motivated them to change their smoking behaviours.

333

334 *I felt deeply pleased because the message of the imam melts everyone's heart. I felt like if I*
335 *could give up smoking from today.*

336 [Man, SFH intervention, nobody smokes in home at 3-month follow-up]

337

338 One exception was a man who was not interested in the intervention, suggesting that he already knew this
339 information anyway.

340

341 The imams were also very enthusiastic. Their perception was that the Ayahs-messages were well received
342 by their congregations, and the SFH intervention was useful and appropriate.

343

344 *I believe that this is a very useful intervention and it is praiseworthy. The objectives are*
345 *very helpful for our society and it is a responsibility for us all to ensure that the objectives*
346 *are properly enforced. From Islamic approach and societal approach, this intervention is*
347 *praiseworthy on both fronts. [Imam 4]*

348

349 They also observed that delivering the messages during Jum'ah prayer was the right thing to do as that is
350 when the mosque was most crowded, would reach large numbers of people and potentially have greatest
351 impact.

352

353 *The Jumu'ah prayer time is the most suitable time for it because what I have seen in my 22*
354 *years' experience as an imam is that approximately 90% of people of our society attends*
355 *Jumu'ah prayer even though they do not perform the rest of the prayers. The best time to*
356 *discuss it is the time before Khutbah as there is no chance to discuss these topics after the*
357 *Jumu'ah prayer. Not all the partakers are present when the Jumu'ah speech starts around*
358 *12.25 or 12.30 pm but they are before the Khutbah. [Imam 5]*

359

360 The proposed individual drivers of behaviour change were attitudes, self-efficacy, social norms, intention
361 formation and planning (see Figure 1). Men's interview accounts clearly illustrated a development in their
362 knowledge and a shift in their attitudes and social norms about SHS, from the messages delivered in the
363 mosque (further confirming the recall data above). In fact, SHS and the risks to others appeared to be new
364 information for most, eliciting beliefs about the social consequences of their smoking, especially the
365 potential harm they were doing to their children. Several participants, both men and women, mentioned
366 having fresh air to breath, healthier children, and no bad smell in the house.

367

368 *If I want to keep my children healthy and safe then it is best for me to quit smoking*
369 *completely. He also said to advise others who smoke to quit as well since it does harm*
370 *those around you, particularly the children. Smoking is harmful for oneself and their*
371 *families. [Man, SFH intervention, still some smoking in home at 3-month follow-up]*

372

373 *I think that if I quit it will benefit everyone, not just one person. The smoke and smell will*
374 *not affect anyone if there is no one smoking at all.*

375 [Man, SFH intervention, still some smoking in home at 3-month follow-up]

376

377 Amongst many men, there was evidence of an intention to act, prompted by the words of the imam and a
378 corresponding new awareness of SHS.

379

380 *It was mostly due to the hujur's speech that inspired me. He always speaks to us keeping*
381 *our best interest in mind. He refers to Hadith so that we know what is best for our Muslim*
382 *community. I liked his messages very much and realized that it is for the best that I should*
383 *try to stop smoking at home.*

384 [Man, SFH+IAQ, lots of smoking in home at 3-month follow-up]

385

386 Notably whilst these men appeared motivated to change, they did not speak of "how" to translate their
387 intention into action or their self-efficacy in doing so. Just one man explicitly spoke of his confidence in
388 creating a SFH, instilled by the imam. Conversely, three men who were not motivated by the imam to
389 change, all alluded to a lack of strategies and low self-efficacy mentioning addiction and stress. One stated
390 that he never listens to the imam because he felt unable to apply this "education" into his life.

391

392 *Look everything that the hujur tells is very educative. We all actually know it but how many*
393 *of us listen to it? If I cannot apply those in my own life, then there is no meaning of this*
394 *educative lines. I never pay attention to the hujur's speech.*

395 [Man, SFH intervention, nobody smokes in home at 3-month follow-up]

396

397 **IAQ feedback acceptability was good. Drivers were new SFH knowledge with corresponding positive**
398 **attitudes, social norms and intentions. Barriers were a lack of plans.**

399 The IAQ machine that measured the air quality in the home, the personalised air quality report and
400 subsequent conversation with the field investigator were well received. They were seen by household leads
401 (men and women) to be educative and prompting intentions to create a SFH.

402

403 *I like the way you provide us report. It's a systematic way. They made us understand very*
404 *clearly with the help of that report. It was shown how smoking is causing harm. That's why*
405 *I liked it most.* [Man, SFH+IAQ, nobody smokes in home at 3-month follow-up]

406

407 As with the SFH intervention, interview accounts illustrated a development in SHS knowledge and a shift in
408 beliefs, attitudes and social norms. Approximately half the men and women interviewed spoke of learning
409 that the air pollution was at levels that were dangerous to their family's health; and the importance of the
410 smoker going outside or away from other people to smoke.

411

412 *We learnt from your initiative and nice report. We realized that it actually harms our*
413 *health or the children's health. So, it is better not to smoke. Even if I have to smoke, I can*
414 *do it outside home. [Man, SFH+IAQ, nobody smokes in home at 3-month follow-up]*

415

416 This new understanding elicited strong beliefs about the importance of having of a SFH, particularly to
417 improve their children's health. A few admitted the personalised feedback had "scared" smokers into
418 action.

419

420 *After this machine was set here, we felt one kind of fear in us and in our children as well.*
421 *They are afraid of it thinking, "If we smoke then something bad might happen to us", so we*
422 *will not smoke. [Woman, SFH+IAQ, nobody smokes in home at 3-month follow-up]*

423

424 All male participants had positive intentions to create a SFH following their IAQ feedback.

425

426 *You made me understand the facts while visiting my home and when I saw the facts with*
427 *proof in my own eyes then I thought it's better to give up this habit.*

428 [Man, SFH+IAQ, nobody smokes in home at 3-month follow-up]

429

430 Consistent with the SFH intervention, there was no mention of specific strategies that the men
431 planned to use to avoid smoking in the home or negotiation strategies that family members could
432 use.

433

434 **Mixed views on usefulness of SFH intervention. Moderate usefulness of IAQ feedback.**

435 Amongst men who reported receiving the SFH intervention 38.2% (SFH) and 79.2% (SFH+IAQ) said it
436 was useful in helping their family achieve a SFH, whilst 60.1% of household leads (men and women)
437 found the IAQ feedback useful.

438 In describing different levels of smoking in their homes, some interview participants referred to the
439 interventions.

440

441 *I used to smoke inside. Now when I buy a cigarette from a tea stall, I smoke beside that place*
442 *instead. When hujur said this, we heard and forgot. But after getting the machine, I got*
443 *scared. [Man, SFH+IAQ, nobody smokes in home at 3-month follow-up]*

444

445 *Since the machine, I mostly smoke outside, in my shop or where I buy the cigarettes. I plan*
446 *that in three months my house will be 80% less smoking inside. I still smoke near my*
447 *children. [Man, SFH+IAQ, still some smoking in home at 3-month follow-up]*

448

449 *After listening to the hujur's messages, my son has reduced his smoking in the house. He*
450 *used to smoke ten times inside and now it's decreased to three.*

451 [Woman, SFH intervention, still some smoking in home at 3-month follow-up]

452

453 Finally, just a small minority of interview participants (men and women) mentioned that they now request
454 other visitors to their home not to smoke indoors.

455

456 *I told them that I don't smoke inside my house, so you are not allowed to smoke here. If*
457 *you want, you may do this outside of my house.*

458 [Man, SFH intervention, still some smoking in home at 3-month follow-up]

459

460 This had resulted in one woman's brother no longer coming to the house. However, one man continued to
461 permit "special guests" to smoke in his home.

462

463 **CONTEXT**

464 **Social context drivers to SFH intervention implementation were in place and important. No context**
465 **barriers to implementation were reported.**

466 The consensus amongst imams was that they had faced no barriers in delivering the SFH intervention.
467 Social context seemed important. Permission from the Islamic Foundation was acknowledged as crucial to
468 demonstrate acceptance of the intervention and a united approach across mosques. Within their own
469 mosques, imams had felt supported by their mosque committees in the form of approval. One valued
470 sharing intervention delivery with a khatib, and another would have liked to have ongoing collaboration
471 about delivery with imams from other mosques.

472
473 **Context barriers/drivers to IAQ feedback implementation were not assessed.**

474 IAQ frequency and reach data suggested that there were no context barriers to implementation.
475

476 **Social context drivers to SHS behaviour change were children's requests. Barriers were a reluctance to**
477 **request male family members and visitors to smoke outside. (Not) having somewhere to smoke outside**
478 **was a physical context (barrier) and driver.**

479 Social and physical context barriers and drivers to SHS behaviour change emerged predominantly from
480 male household lead interview data. The key social driver to men smoking outside was having children in
481 the home, with children's direct requests providing further influence.

482
483 *It is important when my daughter says, "Father, please do not smoke and even if you need to then*
484 *smoke outside the home. Do not smoke in front of me." Is it not an important thing when the*
485 *daughter calls her father? [Man, SFH intervention, nobody smokes in home at 3-month follow-up]*

486
487 Social context barriers were evident. Some women remained reluctant to request male family members to
488 smoke outside seeing this request as "inappropriate". A few men and women did not want to ask *all* guests
489 to smoke outside. Others were happy to do so, confirming the mixed self-reported behaviour change data
490 above.

491

492 *I usually tell them not to smoke inside the house, but if it's a special guest then they are allowed.*

493 [Man, SFH intervention, nobody smokes in home at 3-month follow-up]

494

495 An additional perspective on social context was offered by several imams. They advocated taking a broader
496 societal approach to enhance message exposure and impact by involving the media and the internet,
497 engaging other institutions such as schools and workplaces, and additional influential community leaders
498 like politicians and celebrities.

499

500 *I think that if you can include those who are in charge of making decisions in a society,*
501 *community leaders, as well as committee of the mosques, then this will be more effective.*
502 *Political leaders have a lot of influence over many in our society. If you can include them*
503 *somehow then I think your intervention will have better impact. [Imam 1]*

504

505 *If you can look for these celebrities and large gatherings where multiple speakers offer*
506 *their speech, there are minimum two to three spokesman in these gatherings, you can*
507 *reach a huge audience by building up relationship with them to briefly include this topic in*
508 *his speech. He will proceed the discussion according to his rules but if he includes some*
509 *important facts about smoking, it will be better according to me. [Imam 5]*

510

511 Finally, physical context was also a driver and barrier to SHS behaviour change for men. Most readily
512 identified other places they could smoke, mentioning the road, at work or outside the tobacco shop. There
513 were two exceptions. One man complained he had nowhere to smoke outside late at night because the
514 gates to his compound are locked. Another did not want the shame of being seen smoking by other people.

515

516 *When I work at night and stay up late, the gates are locked by 11 or 11,30. I don't go out then. I*
517 *smoke at home. [Man, SFH+IAQ, still some smoking in home at 3-month follow-up]*

518

519 *I do not smoke outside at all. If I smoke outside now, people would say, "Uncle, as you are an elderly*
520 *person, you should not smoke." It is a matter of shame, thus, I do not smoke at all outside.* [Man,
521 SFH+IAQ, nobody smokes in home at 3-month follow-up]

522

523 **DISCUSSION**

524 Our investigation into the implementation, mechanisms of impact and context [25] of the SFH intervention
525 and IAQ feedback uncovered several explanations for their lack of effectiveness in reducing exposure to
526 SHS in the home (when objectively measured). In short, evidence of implementation of the SFH
527 intervention in the mosques was mixed, and good for IAQ feedback. Both interventions had high
528 acceptability but mixed perceptions of usefulness. Household leads described new SFH knowledge with
529 corresponding positive attitudes, social norms and intentions, whilst self-efficacy and plans were lacking.
530 Context for behaviour change was both positive (e.g. children's requests to smoke outside, places to
531 smoke) and negative (e.g. women's reluctance to ask men to smoke outside, nowhere for men to smoke
532 outside).

533

534 **Strengths and limitations**

535 Our mixed method process evaluation comprised four data sets that were triangulated to elucidate three
536 key process evaluation functions. This approach is recommended as good practice [25,28], ensured a
537 comprehensive process evaluation, and afforded confidence in our conclusions.

538

539 There were some gaps. Context barriers/drivers and fidelity for IAQ delivery were not assessed. The 100%
540 frequency and 98.9% reach data suggest there were limited/no barriers to delivery, and whilst we do not
541 know the quality of the IAQ verbal feedback provided, the IAQ written report was standardised. We have
542 very little interview data from women on their context barriers/drivers to achieving a SFH. Also, our sample
543 of imams interviewed (n=6) and mosques where fidelity assessment was conducted (n=6, 20%) was small.
544 However, they were randomly selected, we captured diversity in their accounts and intervention delivery,

545 and household data were confirmatory. We have no reason to think that other imams accounts or delivery
546 would be markedly different.

547

548 **Why did the interventions not work?**

549 Features of success for both interventions were good acceptability, good frequency of IAQ feedback and
550 moderate to good SFH intervention delivery within Friday Jum'ah prayers. Moreover, imams reported no
551 context barriers to delivery and important drivers (permission from the Islamic Foundation, support from
552 the mosque) were in place. These positive findings are not unexpected. We engaged stakeholders in our
553 intervention adaptation and development which is accepted good practice [15,30]. The IAQ feedback was
554 based on a format previously used in Europe [17-22] and adapted for Bangladesh with household lead
555 input. With hindsight we should have considered more carefully how the report would be used by those
556 who cannot read. The SFH intervention was developed using an iterative and collaborative approach (with
557 the Islamic Foundation, imams and household leads) [26] to ensure that it was truly "a religiously inspired
558 approach" [9, p1176] with acceptability and feasibility. Also, key lessons about intervention content (e.g.,
559 ensuring that the imams were credible "non-smoking" SHS messengers [7]) and delivery (e.g. support from
560 mosque committees) were gathered from an earlier pilot trial [31]. These informed careful preparation
561 work with mosques and imams to ensure they were ready for intervention delivery, a "success factor" of
562 effective faith-based health promotion programmes [32].

563

564 Less positive were findings of poor reach of the SFH intervention and mixed quality of delivery. Only half of
565 household leads recalled receiving the SFH intervention (or their family members receiving it) and no men
566 interviewed had received the booklet. Although Friday prayers are traditionally attended by most Muslim
567 men, the Khutbah sessions delivered before prayers are not mandatory. Anecdotally, attendance may be as
568 low as 10% of the total attendance in Friday prayers which may explain the poor reach. With hindsight, we
569 should probably have been more prescriptive about dissemination to other congregations (including
570 distribution of the SFH booklet), to increase frequency and reach. As an example, a "potentially effective"
571 Korean church-based intervention targeting SHS was more widely embedded across church activities that

572 lasted up to 1.5 hours, with dissemination of multiple resources (SHS brochures, quit-smoking guides, SHS
573 stickers, reusable grocery bags, and insulated lunch bags) [33].

574

575 Ayahs-messages targeting SHS attitudes and social norms were the self-declared focus of imams, with those
576 targeting attitudes implemented most fully. These were also the Ayahs-messages recalled by male
577 household leads, resulting in new knowledge with a corresponding shift in their SHS attitudes, social norms
578 and intentions to change their SHS behaviours. The SHS health messages e.g. risks to children, were best
579 remembered rather than the corresponding religious text. Even if they had remembered the religious
580 connection, this will only have impacted on motivation [9]. Ayahs-messages that targeted self-efficacy
581 (employing instruction, verbal persuasion and self-talk techniques [34]) and planning (using “if-then” plans
582 [35,36]) were not remembered and were less well delivered. It seems that imams can confidently educate
583 but lack skills or motivation to deliver strategies to turn knowledge into behaviour. The same outcome was
584 evident for the IAQ feedback, with interview participants self-reporting learning about the risks of SHS at
585 home, changing their attitudes, social norms and being motivated to create a SFH, yet plans for how to do
586 this were absent.

587

588 Both interventions were based on well-evidenced behaviour change techniques including those targeting
589 self-efficacy [34] and planning [34-36], yet they were remembered by recipients as educational
590 interventions. It seems likely that men were ill-equipped with confidence, coping and planning skills to
591 overcome significant context barriers and translate positive intentions into behaviour. This hypothesis is
592 consistent with a scoping review of fathers’ experiences of creating a SFH [37] and European evaluation of
593 an SFH intervention [20]. Our interview data with women suggest they found it difficult to request male
594 family members to smoke outside. Other studies reporting women’s inability to negotiate SFHs also report
595 these gendered power interactions [38,39]. Men-inclusive community interventions (like ours) that aim to
596 change social norms around smoking rather than relying on women to set household boundaries offer
597 potential to improve gender equity as well as health [37,40]. However, they need to be supported by
598 “gender transformative tobacco control” [41, p796] where gender theory is embedded into public health

599 policy [41]. Overall, it is unsurprising that there was a lack of perceived “usefulness in creating a SFH” for
600 both interventions, and no effect on the SHS exposure in homes (measured by 24-h mean household
601 airborne fine particulate matter (<2.5 microns in diameter [PM_{2.5}]) concentration) both at 3- and 12-
602 months post-intervention [24].

603

604 Literature reviews [42-45] consistently cite promising evidence for faith-based health promotion
605 interventions whilst advocating more rigorous evaluation. Our SFH intervention comprised many “success
606 factors” for effective faith-based programmes [32]. There is also support for IAQ feedback interventions in
607 Europe [16-22]. Our IAQ feedback was an adapted version of these European feedback tools, although our
608 frequency was less than other programmes that incorporate repeat measurement, follow-up visits or
609 phone calls [16-22]. What was different for both interventions is that we did not include one-to-one
610 practical support for behaviour change (including boosting confidence, developing coping and planning
611 skills) which is evident in other faith-based programmes via motivational coaches [33], lay volunteers [32]
612 or faith nurses [42]. We also did not include a motivational interview component [16-22] with the IAQ
613 report. A 2018 review concluded that the effectiveness of educational interventions in reducing SHS
614 exposure was unclear [3]. Whereas combining SHS interventions with smoking cessation support may
615 reduce SHS exposure [18].

616

617 Alturki [9] proposes that civil society including Muslim authorities should supplement smoking cessation
618 programmes delivered by health professionals. Unfortunately, in Bangladesh, smoking cessation services
619 are lacking, reflecting poor implementation of the World Health Organization Framework Convention on
620 Tobacco Control (FCTC) [46] Article 14 across LMICs [47]. A further challenge is the weak implementation of
621 SHS legislation (WHO FCTC Article 8) in Bangladesh, again consistent with other LMICs [47, 48]. The WHO
622 [8] and other authors in this field [7,9] advocate a community-wide strategy where faith-based
623 programmes are ‘one part of a comprehensive overall approach to tobacco control’ [8] including cessation
624 services and good policy. Embedding our two interventions within this wider community approach would
625 seem sensible. One example would be to link with the established network of community health workers

626 who deliver primary care and behaviour change counselling services in Bangladesh, to achieve a “multiplier
627 effect” [49].

628

629 **Conclusions**

630 Despite detailed development and adaption work with relevant stakeholders, the SFH intervention and IAQ
631 feedback became educational and motivational but were insufficient to overcome significant context
632 barriers to SHS behaviour change. Future interventions should include practical support for SFH behaviour
633 change. Moreover embedding these into community wide strategies that include practical cessation
634 support and enforcement of SFH legislation is needed.

635

636 **List of abbreviations**

637 FCTC: Framework Convention on Tobacco Control

638 IAQ: indoor air quality

639 LMIC: Low and middle-income countries

640 MCLASS: Muslim Communities Learning About SHS

641 SFH: Smoke-free homes

642 SHS: Second-hand smoke

643 WHO: World Health Organisation

644

645 **Declarations**

646 **Ethics approval and consent to participate**

647 All methods were carried out in accordance with relevant guidelines and regulations (Declaration of
648 Helsinki). Ethics approval was obtained from the Bangladesh Medical Research Council’s National Research
649 Ethics Committee (BMBC/NREC/2016–2019/358) and the University of York’s Health Sciences Research
650 Governance Committee. Written informed consent was obtained from imams or khatibs for their and their
651 mosques’ participation, heads of household for participation of households, and adults in
652 respective households for their own data collection.

653

654 **Consent for publication**

655 Not applicable

656

657 **Availability of data and material**

658 The datasets used and/or analysed during the current study are available from the corresponding author on
659 reasonable request.

660 De-identified individual participant data will be made available from the point of, and up to 5 years after
661 the acceptance for publication of the main findings from the final dataset. These data can be requested
662 from the Principal Investigator (Prof Kamran Siddiqi; kamran.siddiqi@york.ac.uk) and will be shared after
663 the provision of a methodologically sound proposal, and only under a data-sharing agreement that provides
664 for commitment to: using the data only for research purposes and not to identify any individual participant;
665 securing the data using appropriate computer technology; and destroying or returning the data after
666 analyses are completed. The proposals will be assessed and approved by members of the Programme
667 Management Group.

668

669 **Competing Interests**

670 The authors declare that they have no competing interests.

671

672 **Funding**

673 This trial was funded by the Medical Research Council UK under the Global Alliance for Chronic Diseases
674 research programme (MR/P008941/1). The funder was not involved in the design of the study, collection,
675 analysis, and interpretation of data or in writing the manuscript.

676

677 **Author's contributions**

678 CJ co-conceived, co-designed, and led the process evaluation and led the data analysis, data interpretation,
679 and drafting of the manuscript. ZA co-designed the process evaluation, conducted data collection and data

680 analysis, drafted some sections of the manuscript. IK co-conceived and co-designed the process evaluation
681 and co-drafted the manuscript. NDM co-conceived the process evaluation and revised the manuscript. CF
682 conducted data analysis and revised the manuscript. TF conducted data collection and revised the
683 manuscript. CH conducted data analysis and revised the manuscript. RH co-conceived the process
684 evaluation, supervised data collection and revised the manuscript. AM, SS, AS and KS co-conceived the
685 process evaluation and revised the manuscript. All authors approved the manuscript.

686

687 **Acknowledgements**

688 We would like to thank our MCLASS II trial team colleagues: Steve Parrott, Abdullah Sonnet, Shilpi Swami,
689 Han-I Wang, Qi Wu and members of the independent trial steering committee: Dr Andrew Fogarty (Chair),
690 and Professor Jo Leonardi-Bee, University of Nottingham, Nottingham, UK; and Dr Saidur Rahman
691 Mashreky, Bangladesh University of Health Sciences, Dhaka, Bangladesh. Our gratitude also goes to
692 members of our national-level reference group: Joint Secretary of the Ministry of Religious Affairs, Director
693 of Research of the Islamic Foundation, Head of the Imam Training Academy, and a representative from ARK
694 Foundation. We are also grateful to the project working group comprising an Islamic Foundation
695 representative, an imam, a research fellow, a public health expert, and a member of the public. Finally
696 thank you to the trial field investigators, the Islamic Foundation, participating mosques, their mosque
697 committee leaders, imams, and khatibs; all participants; professionals and other researchers who have
698 contributed to this trial.

699

700 **References**

- 701 1. Öberg M, Jaakkola MS, Woodward A, Peruga A, Prüss-Ustün A. Worldwide burden of disease from
702 exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet*.
703 2011;377:139-46.
- 704 2. Ferdous T, Siddiqi K, Semple S, Fairhurst C, Dobson R, Mdege N, et al. Smoking behaviours and indoor
705 air quality: a comparative analysis of smoking-permitted versus smoke-free homes in Dhaka,
706 Bangladesh. *Tob. Control*. 2020;Dec 16;tobaccocontrol-2020-055969 [Epub ahead of print].

- 707 3. Behbod B, Sharma M, Baxi R, Roseby R, Webster P. Family and carer smoking control programmes for
708 reducing children’s exposure to environmental tobacco smoke. *Cochrane Database of Systematic*
709 *Reviews*. 2018;1:CD00174.
- 710 4. Dherani M, Zehra SN, Jackson C, Satyanaryana V, Huque R, Chandra P, et al. Behaviour change
711 interventions to reduce second-hand smoke exposure at home in pregnant women – a systematic
712 review and intervention appraisal. *BMC Pregnancy and Childbirth*. 2017;17:378.
- 713 5. Tong TV, Dietz PM, Rolle IV, Kennedy SM, Thomas W, England LJ. Clinical interventions to reduce
714 secondhand smoke exposure among pregnant women: a systematic review. *Tob Control*. 2015;24:217–
715 23.
- 716 6. Nunziata L, Toffolutti. “Thou Shalt not Smoke”: Religion and smoking in a natural experiment of history.
717 *SSM-Popula Health*. 2019;8:1004012.
- 718 7. Byron MJ, Cohen JE, Gittelsohn J, Frattaroli S, Nuryunawati R, Jernigan DH. Influence of religious
719 organisations’ statements on compliance with a smoke-free law in Bogor, Indonesia: a qualitative
720 study. *BMJ Open*. 2015;5:e008111.
- 721 8. World Health Organization. *Tobacco Use and Religion*. Cairo: WHO Regional Office for the Eastern
722 Mediterranean; 2014.
- 723 9. Alturki K, Hamza A, Walton P. Islam and Motivation to Quit Smoking: Public Health Policy Implications. *J*
724 *Relig Health*. 2020;59:1175–88.
- 725 10. Garrusi B, Nakhaee N. Religion and smoking: A review of recent literature. *Int J Psychiatry Med*.
726 2012;43:279–29.
- 727 11. Mustafa Y, Baker D, Puligari P, Melody T, Yeung J, Gao-Smith F. The role of imams and mosques in
728 health promotion in Western societies—a systematic review protocol. *BMC Syst Rev*. 2017;6:25.
- 729 12. Radwan GN, Israel E, El-Setouhy M, Abdel-Aziz F, Mikhail N, Mohamed MK. Impact of religious rulings
730 (fatwa) on smoking. *J Egypt Soc Parasitol* 2003;33:1087–101.
- 731 13. Cultural Atlas. Bangladeshi Culture. [https://culturalatlas.sbs.com.au/bangladeshi-culture/bangladeshi-](https://culturalatlas.sbs.com.au/bangladeshi-culture/bangladeshi-culture-religion)
732 [culture-religion](https://culturalatlas.sbs.com.au/bangladeshi-culture/bangladeshi-culture-religion). Accessed 3 October 2021.

- 733 14. Liu JJ, Davidson E, Bhopal RS, White M, Johnson MRD, Netto G, et al. Adapting health promotion
734 interventions to meet the needs of ethnic minority groups: mixed-methods evidence synthesis. *Health*
735 *Technol Assess.* 2012;16:1-469.
- 736 15. Liu JJ, Wabnitz C, Davidson E, Bhopal RS, White M, Johnson MR, et al. Smoking cessation interventions
737 for ethnic minority groups—A systematic review of adapted interventions. *Prev. Med.* 2013;57:765-75.
- 738 16. Harutyunyan A, Movsisyan N, Petrosyan V, Petrosyan D, Stillman. Reducing children’s exposure to
739 secondhand smoke at home: a randomized trial. *Pediatrics.* 2013;132:1071–80.
- 740 17. Wilson I, Semple S, Mills LM, Ritchie D, Shaw A, O’Donnell R et al. REFRESH-reducing families’ exposure
741 to secondhand smoke in the home: a feasibility study. *Tob Control* 2013;22:e8.
- 742 18. Ratschen E, Thorley R, Jones L, Opazo Breton M, Cook J, McNeill A. A randomised controlled trial of a
743 complex intervention to reduce children’s exposure to secondhand smoke in the home. *Tob Control.*
744 2018;27:155–62.
- 745 19. Hughes SC, Bellettiere J, Nguyen B, Liles S, Klepeis NE, Qunitana PJE, et al. Randomized Trial to Reduce
746 Air Particle Levels in Homes of Smokers and Children. *Am J Prev Med.* 2018;54:359–67.
- 747 20. Dobson R, O’Donnell R, Tigova O, Fu M, Enriquez M, Fernandez E et al. Measuring for change: A multi-
748 centre pre-post trial of an air quality feedback intervention to promote smoke-free homes. *Environ Int.*
749 2020;140:105738.
- 750 21. Dobson R, O’Donnell R, de Bruin M, Turner S, Semple S. Using air quality monitoring to reduce second-
751 hand smoke exposure in homes: the AFRESH feasibility study. *Tob Prev Cessat.* 2017;3:117.
- 752 22. Semple S, Turner S, O’Donnell R, Adams L, Henderson T, Mitchell S, et al. Using air-quality feedback to
753 encourage disadvantaged parents to create a smoke-free home: Results from a randomised controlled
754 trial. *Environ Int.* 2018;120:104–10.
- 755 23. Mdege ND, Fairhurst C, Ferdous T; Hewitt C, Huque R, Jackson C, et al. Muslim Communities Learning
756 About Second-hand Smoke in Bangladesh (MCLASS II): study protocol for a cluster randomised
757 controlled trial of a community-based smoke-free homes intervention, with or without Indoor Air
758 Quality feedback. *Trials.* 2019;20:11.

- 759 24. Mdege ND, Fairhurst C, Wang Han-I, Ferdous T, Marshall AM, Hewitt C, et al. Muslim Communities
760 Learning About Second-hand Smoke in Bangladesh (MCLASS II): a three-arm, cluster randomised
761 controlled trial of the effectiveness and cost-effectiveness of a community-based smoke-free homes
762 intervention, with or without indoor air quality feedback. *Lancet Glob Health*. 2021;9:e639–50.
- 763 25. Medical Research Council. *Process evaluation of complex evaluations*. London: Medical Research
764 Council; 2015.
- 765 26. Kellar I, Azdi ZA, Jackson C, Huque R, Mdege ND, et al. Muslim Communities Learning About Second-
766 hand Smoke in Bangladesh (MCLASSII): a combined evidence and theory-based plus partnership
767 intervention development approach. *BMC Pilot and Feasibility Studies* 2022;8:136.
- 768 .
- 769 27. World Health Organization. Exposure to household air pollution for 2016. Last modified 2018.
770 https://www.who.int/airpollution/data/HAP_exposure_results_final.pdf?ua=1. Accessed 3 October
771 2021.
- 772 28. O Cathain A. *A Practical Guide to Using Qualitative Research with Randomized Controlled Trials*. Oxford:
773 Oxford University Press; 2018. Ritchie J, Lewis J, McNaughton Nicholls C, Ormston R. *Qualitative*
774 *Research Practice*. London: SAGE; 2014.
- 775 29. Ritchie J, Lewis J, McNaughton Nicholls C, Ormston R. *Qualitative Research Practice*. London: SAGE;
776 2014.
- 777 30. Vu M, Muhammad H, Peek ME, Padela AI. Muslim women's perspectives on designing mosque-based
778 women's health interventions—an exploratory qualitative study. *Women Health*. 2018;58:334-46.
- 779 31. King R, Warsi AS, Amos S, Shah S, Mir G, Sheikh A et al. Involving mosques in health promotion
780 programmes: a qualitative exploration of the MCLASS intervention on smoking in the home. *Health*
781 *Educ Res*. 2017;32 293-305.
- 782 32. Sternberg Z, Munschauer FE, Carrow SS, Sternberg E. Faith-placed cardiovascular health promotion: a
783 framework for contextual and organizational factors underlying program success. *Health Educ Res*.
784 2007;22:619-29.

- 785 33. Hughes SC, Corcos I, Hovell M, Hofstetter CR. Feasibility Pilot of a Randomized Faith-Based Intervention
786 to Reduce Secondhand Smoke Exposure Among Korean Americans. *Prev Chronic Dis.* 2017;14:1-8.
- 787 34. Carey RN, Connell LE, Johnstone M, Rothman AJ, de Bruin M, Kelly MP et al. Behavior Change
788 Techniques and Their Mechanisms of Action: A Synthesis of Links Described in Published Intervention
789 Literature. *Ann Behav Med.* 2019;53:693-707.
- 790 35. Hagerman CJ, Hoffman RK, Vaylay S, Dodge T. Implementation Intentions to Reduce Smoking: A
791 Systematic Review of the Literature. *Nicotine Tob Research.* 2021;7:1085–93
- 792 36. McWilliams, Bellhouse S, Yorke J, Lloyd K, Armitage CJ. Beyond “planning”: A meta-analysis of
793 implementation intentions to support smoking cessation. *Health Psychol.* 2019;38:1059-68.
- 794 37. O’Donnell R, Angus K, McCulloch P, Amos A, Greaves L, Semple S. Fathers' Views and Experiences of
795 Creating a Smoke-Free Home: A Scoping Review. *Int J Environ Res Public Health.* 2019;16:5164.
- 796 38. Jackson C, Huque R, Satyanarayana V, Nasreen S, Kaur M, Barua D et al. “He Doesn’t Listen to My
797 Words at All, so I Don’t Tell Him Anything” – A Qualitative Investigation on Exposure to Second Hand
798 Smoke among Pregnant Women, Their Husbands and Family Members from Rural Bangladesh and
799 Urban India. *Int J. Environ Res* 2016;13:1098.
- 800 39. Passey ME, Longman JM, Robinson J, Wiggers J, Jones LL. Smoke-free homes: What are the barriers,
801 motivators and enablers? A qualitative systematic review and thematic synthesis. *BMJ Open* 2016;6:
802 e010260
- 803 40. Padmawati RS, Prabandari YS, Istiyani T, Nichter M, Nichter M. Establishing a community-based smoke-
804 free homes movement in Indonesia. *Tob. Prev. Cessation* 2018;4:36.
- 805 41. Greaves L. Can tobacco control be transformative? Reducing gender inequity and tobacco use among
806 vulnerable populations. *Int J Environ Res Public Health.* 2014;11:792-803.
- 807 42. Bopp M, Peterson JA, Webb BL. A comprehensive review of faith-based physical activity interventions.
808 *Am J Lifestyle Med.* 2012;6:460-78.
- 809 43. Parra MT, Porfírio GJM, Arredondo EM, Atallah ÁN. Physical Activity Interventions in Faith-Based
810 Organizations: A Systematic Review. *Am J Health Promot.* 2018;32:677-90.
- 811

- 812 44. DeHaven MJ, Hunter IB, Wilder L, Walton JW, Berry J. Health Programs in Faith-Based Organizations:
813 Are They Effective? *Am J Pub Health*. 2004;94:1030-36.
- 814 45. Lancaster KJ, Carter-Edwards L, Grilo S, Shen C, Schoenthaler AM. Obesity interventions in African
815 American faith-based organizations: a systematic review. *Obes Rev*. 2014;15(suppl 4):159–76.
- 816 46. World Health Organization. WHO Framework Convention of Tobacco Control. <https://fctc.who.int/>
817 Accessed 26 October 2021.
- 818 47. Nilan K, Raw M, McKeever TM, Murray RL, McNeill A. Progress in implementation of WHO FCTC Article
819 14 and its guidelines: a survey of tobacco dependence treatment provision in 142 countries. *Addiction*.
820 2017;112:2023-31.
- 821 48. Nazar GP. Smoke-free legislation and active smoking, second hand exposure and health outcomes in
822 low- and middle-income countries. PhD thesis, London School of Hygiene and Tropical Medicine.
823 <https://researchonline.lshtm.ac.uk/id/eprint/4433694/> Accessed 8 October 2021.
- 824 49. Ahmed S, Khan JAM. Disseminating public health messages about second-hand smoking through
825 mosque congregations in Bangladesh. *Lancet*. 2021;9:e657-58.

826

827 **MCLASS II trial team**

828 Cath Jackson^{1,2}, Zunayed Al Azdi³, Ian Kellar⁴, Noreen Dadirai Mdege¹, Caroline Fairhurst¹, Tarana Ferdous³,
829 Catherine Hewitt¹, Rumana Huque³, Anna-Marie Marshall¹, Sean Semple⁵, Aziz Sheikh⁶, Kamran Siddiqi¹,
830 Steve Parrott¹, Abdullah Sonnet³, Shilpi Swami¹, Han-I Wang¹, Qi Wu¹

831

832 1. Department of Health Sciences, University of York, York, UK

833 2. Valid Research Ltd, Wetherby, UK

834 3. ARK Foundation, Dhaka, Bangladesh


835 4. School of Psychology, University of Leeds, Leeds, UK

836 5. Institute for Social Marketing and Health, University of Stirling, Stirling, Scotland, UK

837 6. Usher Institute, University of Edinburgh, Edinburgh, Scotland, UK

838

Figure 1: Intervention Programme Theory



Note. SFH is smoke-free homes where smokers do not smoke inside, smokers request residents and visitors to smoke outside

840
841

Additional file 1. Linked Ayah-messages and target constructs

Cycle	Week	Ayah	Message	Construct
1	1	Surah Al-Maaida - 4 (5:4) They ask you, [O Muhammad], what has been made lawful for them. Say, "Lawful for you are [all] good foods."	Though sometimes people think that smoking helps in some ways, the evidence that smoking, and second-hand smoke cause harm in many ways is clear. Would Allah permit you something harmful? No! Tobacco is harmful, and hence it is not permissible to Allah. The sin of smoking causes you spiritual as well as physical harm.	Attitude
1	2	Sura An-Nisaa – 59 (4:59) Believers! Obey Allah and obey the Messenger, and those from among you who are invested with authority.	Allah, in his grace, has given us experts who he has been given authority to tell us the facts about what heals us and what harms us. The evidence from scientists tells us that second-hand smoke contains more than 7,000 chemicals. Hundreds are toxic and about 70 can cause cancer. Second-hand smoke also causes numerous health problems in infants and children. Will you not listen to the facts? Will you not hear what your Imam says to you?	Attitude
1	3	Sura Al-Ahzaab – 58 (33:58) And those who harm believing men and believing women for [something] other than what they have earned have certainly born upon themselves a slander and manifest sin.	The evidence that second-hand smoke harms other is clear. It can result heart attack, stroke and lung cancer among innocent adults who are exposed to it. And children exposed to second-hand smoke are more prone to have chest infection, sneezing and coughing. Moreover, they have 50% higher chance of having ear infection. Now do you really want to do that to your family members and your children? Allah also said that – causing harm to others is a manifest sin.	Social norms

842

1	4	<p>Sura At-Takaathur – 8 (102:8) Then, on that Day, you will be called to account for all the bounties you enjoyed.</p>	<p>These messages to you are part of Allah’s bounty to you. But you need to make a commitment to enjoy his bounty. This means committing to either quitting or smoking outside. If you are going to do this, you need to make a plan.</p> <p>For planning to stop smoking at home, commit that if you reach for a cigarette – then leave the house before you light it. And for planning to quit smoking completely, commit that if you feel like smoking, then pray 2 rakat salat instantly.</p>	Intention formation (and prompt action planning)
1	5	<p>Sura Ar-Ra’d – 11 (13:11) The fact is that Allah does not change a people's lot unless they themselves change their own characteristics</p>	<p>You can trust Allah to help you, but to receive that support, you must take a step by yourself in faith. Trust that Allah will give you everything you need.</p> <p>You can find it difficult to stop smoking at home. But if YOU cannot make this simple change of behaviour for the sake of your family members, how can you expect Allah will help them in other ways? So, you need to make your plan of smoking outside home. For example, if you feel like smoking when you are at home – then leave the house before you light it. You can plan to remove your last cigarette before you come home.</p>	Self-efficacy (prompt Action Planning)
1	6	<p>Surah Al-Maaida - 9 (5:9) Allah has promised those who believe and do righteous deeds [that] for them there is forgiveness and great reward.</p>	<p>Allah knows you, Allah knows everything. He knows that you will need his forgiveness. Be quick to come to him. Trust that he will be with you as you come back to the right path.</p> <p>So make a plan that if you lapse, then you will call on Allah for forgiveness and recommit yourself and rehearse your plans.</p>	Coping planning

2	7	<p>Sura Al Maaida – 90 (5:90) Believers! Intoxicants, games of chance, idolatrous sacrifices at altars, and divining arrows are all abominations, the handiwork of Satan. So turn wholly away from it that you may attain to true success.</p>	<p>Tobacco is toxic. Your body becomes reliant on nicotine. It doesn't relieve stress. It only relieves withdrawal syndrome from your addiction.</p> <p>Tobacco is the handiwork of Satan. Do you want true success? Turn away wholly from tobacco.</p>	Attitude
2	8	<p>Surah Al-Maaida - 100 (5:100) Say, "Not equal are the evil and the good, although the abundance of evil might impress you." So, fear Allah, O you of understanding, that you may be successful.</p>	<p>Some of you may believe that smoking is good because it helps keep you warm, or stops you getting fat, or manage your stress. But Allah, in his grace, has given us eye to see, ears to hear and a mind to enquire. What do the experts tell us? Experts tell us that it does nothing but harm you and those who are staying beside you when you smoke. The only relief you feel getting after smoking is the relief from withdrawal syndrome which we mistakenly think as stress relief.</p>	Attitude
2	9	<p>Sura At-Baqara – 195 (2:195) And do good; indeed, Allah loves the doers of good.</p>	<p>Globally 600 thousand people die every year due to exposure to second-hand smoke. Those who smoke around us are directly causing harm to us though they are often not aware of the harm they are causing. Hence, we need to be aware and careful about smoking inside home and in front of others.</p> <p>We need to talk to others about the harm of smoking and second-hand smoke. We need to save our families from this harm. Allah also loves those who does good things.</p>	Social norms
2	10	<p>Surah Ash-Shams – 7-10 (91:7-10) And [by] the soul and He who proportioned it. And inspired it [with discernment of] its wickedness and its righteousness, He has succeeded who purifies it, and he has failed who instils it [with corruption].</p>	<p>Allah has given you wisdom, but to remember it, you have to act on it. Only then you and others will be benefitted by that.</p> <p>If you are going to do something, you need to make a plan. For example, if you reach for a cigarette when you are at home – then leave the house before you light it. And for quitting smoking, you should plan like this - if you feel the urge to do smoke, pray 2 rikat salat instantly.</p>	Intention formation

2	11	<p>Surah At-Taghaabun - 16 (64:16) So, fear Allah as much as you are able and listen and obey and spend [in the way of Allah]; it is better for yourselves. And whoever is protected from the stinginess of his soul - it is those who will be the successful.</p>	<p>Those who smoke can find it difficult to quit smoking or they can find it hard to go outside home every time they want to smoke. But believe it, Allah will help you if you wish to listen to him. One can make simple plans to overcome such issues. Just commit to yourself and others (if you can) that whenever you feel the urge of smoking, go outside home to light it or pray 2 rakat salat instantly.</p>	<p>Self-efficacy (prompt Action Planning)</p>
2	12	<p>Surah Al-Hajj - 77 (22:77) Oh you who have believed, bow and prostrate and worship your Lord and do good - that you may succeed</p>	<p>Allah knows best about his creatures. He understands that we may do things that will harm us and others. That is why, he encouraged us to enjoy all that is good and forbid all that is evil and keep patience in times of affliction.</p> <p>We must remind ourselves these words of Allah again and again. We must try to make our habits safe for others. We must remember the possible harms of our behaviour to others like smoking at home and repetitively plan to keep us and our families safe from its harm.</p>	<p>Coping planning</p>

845
846

847 **Additional File 2:** Triangulation matrix for SFH intervention

Meta-theme	Household lead interviews (N=20 men, N=10 women)	Imam interviews (N=6)	Household lead questionnaire (N=848 men, N=52 women)	Fidelity (N=6 mosques)	Research team records	Level of congruence	Conclusion
	SFH intervention						
Implementation	Frequency		4/6 (66.7%) reported delivering all 12 weeks. All reported distributing the SFH booklet.			29/30 (96.7%) mosques reported delivering all 12 weeks.	Moderate to good frequency of intervention
	Fidelity		All had delivered the intervention during Friday Juma'ah prayers (as per guidance). Consensus that had shared "most of the Ayahs-messages". Most focused on telling congregation about risks (Ayahs-messages targeting attitudes and social norms).		Mean fidelity score 19.6 (SD 2.51, range 16-22 of maximum 24). Ayahs-messages-linked to attitudes 75.0% fully implemented, self-efficacy/action planning and coping planning 66.7% fully implemented, social norms and intention formation/action planning 50.0% fully implemented.		Mixed levels of fidelity. Ayahs-messages targeting attitudes were most often fully implemented.

848

	Reach	<p>Majority of men recalled hearing Ayahs-messages during Friday Jumu'ah prayers. Majority of women reported that their partners/sons had received the same.</p> <p>Men recalled Ayahs-messages on risks of SHS (attitudes, social norms). Less well recalled were Ayahs-messages targeting self-efficacy, coping planning, and intention formation.</p> <p>No men had received the SFH booklet. 3 women had sons who received the booklet but 2 could not read it.</p>		<p>49.4% (SFH) and 55.5% (SFH+IAQ) of men had received the SFH intervention. Of these, 99.5% (SFH) and 99.6% (SFH+IAQ) received this during Friday Jumu'ah prayers.</p> <p>SFH 51.5% (SFH) and 52.6% (SFH+IAQ) of women reported family members receiving the SFH intervention, during Friday Jumu'ah prayers (100%)</p> <p>80%+ of these men recalled Ayahs-messages targeting attitudes and social norms. Just 37.5-45% recalled those targeting self-efficacy, coping planning, and intention formation.</p>			Complementary	Poor intervention reach. For those who it did reach, Ayahs-messages targeting attitudes and social norms had the best reach.
--	-------	---	--	---	--	--	---------------	--

Mechanisms of impact	Acceptability of the intervention	Majority view amongst men that SFH messages were informative and motivational.	Consensus that SFH intervention acceptable, appropriate and well received by congregation.				Convergent	Good acceptability of intervention
	Barriers and drivers to change (related to the individual)	Majority view amongst men that knowledge about SHS was new and changed their SHS attitudes, social norms and intended to smoke outside. They did not mention plans. Minority that was not motivated attributed this to lack of self-efficacy.					N/A	Drivers to change were new SFH knowledge with corresponding positive attitudes, social norms and intentions. Barriers were a lack of self-efficacy and plans.
	Usefulness of the intervention ^a	In describing different levels of smoking in their homes, some interview participants referred to the intervention.		38.2% (SFH alone) and 79.2% (SFH+IAQ) of men reported that the SFH intervention was useful in achieving a SFH				Convergent

Context	Impact on SFH implementation		Consensus that felt supported by mosque committee. Permission from Islamic Foundation seen as important. No barriers to delivery.				N/A	Social context drivers were in place and important for implementation. No context barriers reported.
	Impact on SHS behaviour change ^b	Minority of men/women spoke of children motivating men to smoke outside. Conversely social norms prevented some asking visitors to smoke outside and women to request this of male family members. Majority of men could identify other places to smoke. A minority could not.	Some imams advocated a wider societal approach to achieve behaviour change.				Complementary	Social and physical context barriers and drivers to creating a SFH were evident.

851 *Note.* Convergent = in agreement, complementary = partial agreement, contradictory = disagreement, silent = findings do not occur in a dataset but may have
852 been expected to do so [29]. ^aSame usefulness of the intervention data (mechanisms of impact) and ^bimpact on outcomes data (context) are reported in Additional
853 Files 2 and 3.

854

855 **Additional File 3:** Triangulation matrix for IAQ feedback

Meta-theme		Household lead interviews (N=20 men, N=10 women)	Household lead questionnaire (N=848 men, N=52 women)	Research team records	Level of congruence	Conclusion
Implementation	Frequency			IAQ feedback delivered to all 640 households	N/A	Good frequency of intervention
	Fidelity				No data	No data
	Reach	Majority remembered receiving the IAQ feedback, half reported another household member receiving the report. Minority commented that they could not read it.	98.9% of households received the IAQ feedback		Convergent	Good reach of intervention
Mechanisms of impact	Acceptability of the intervention	Consensus that IAQ feedback was informative and motivational				Good acceptability of intervention
	Barriers and drivers to change (related to the individual)	Consensus that that knowledge about SHS was new and changed their SHS attitudes, social norms and intended to create a SFH. No one mentioned planning for this.				Drivers to change were new SFH knowledge with corresponding positive attitudes, social norms and intentions. Barriers were a lack of plans.

	Usefulness of the intervention ^a	In describing different levels of smoking in their homes, some interview participants referred to the intervention.	60.1% of households reported that the IAQ feedback was useful in achieving a SFH		Convergent	Moderate usefulness of IAQ intervention.
Context	Impact on implementation					No data
	Impact on SHS behaviour change ^b	Minority of men/women spoke of children motivating men to smoke outside, conversely social norms prevented some asking visitors to smoke outside. Majority of men could identify other places to smoke. A minority could not.			N/A	Social and physical context barriers and drivers to creating a SFH were evident.

856 *Note.* Convergent = in agreement, complementary = partial agreement, contradictory = disagreement, silent = findings do not occur in a dataset but may have
857 been expected to do so [29]. ^aSame usefulness of the intervention data (mechanisms of impact) and ^bimpact on outcomes (context) data are reported in Additional
858 Files 2 and 3.

859

