




BMJ Open Prevalence of medically unexplained symptoms in adults who are high users of healthcare services and magnitude of associated costs: a systematic review

Ferozkhan Jadhakhan ¹, Daniel Romeu,^{2,3} Oana Lindner,⁴ Amy Blakemore ⁵, Elspeth Guthrie ⁶

To cite: Jadhakhan F, Romeu D, Lindner O, *et al.* Prevalence of medically unexplained symptoms in adults who are high users of healthcare services and magnitude of associated costs: a systematic review. *BMJ Open* 2022;**12**:e059971. doi:10.1136/bmjopen-2021-059971

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-059971>).

Received 08 December 2021
Accepted 20 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Ferozkhan Jadhakhan;
F.Jadhakhan@bham.ac.uk

ABSTRACT

Introduction Medically unexplained symptoms (MUS) is a common clinical syndrome in primary and secondary healthcare service. Outcomes for patients with persistent MUS include increased disability, poorer quality of life and higher healthcare costs. The aim of this systematic review was to determine the prevalence of MUS in patients who are high users of healthcare or high-cost patients in comparison with routine users and the magnitude of associated costs.

Design A systematic review of the available literature.

Data sources and eligibility criteria The following electronic databases were systematically searched without language restriction from inception to June 2018 and updated on 22 October 2021: MEDLINE, PsycINFO, EMBASE, CINAHL and PROSPERO. Inclusion criteria included studies investigating adults aged ≥18 years, who were high healthcare users or accrued high healthcare costs, in which the prevalence and/or associated costs of MUS was quantified. Two reviewers independently extracted information on study characteristics, exposure and outcomes.

Results From 5622 identified publications, 25 studies from 9 countries involving 31 650 patients were selected for inclusion. Due to high risk of bias in many studies and heterogeneity between studies, results are described narratively. There were wide variations in prevalence estimates for MUS in high users of healthcare (2.9%–76%), but MUS was more prevalent in high use groups compared with low use groups in all but one of the 12 studies that included a comparator group. Only three studies investigated healthcare costs associated with MUS, and all three reported greater healthcare costs associated with MUS.

Conclusion MUS has been found to be more prevalent in high use healthcare populations than comparator groups, but the magnitude of difference is difficult to estimate due to considerable heterogeneity between studies and potential for bias. Future studies should prioritise a standardised approach to this research area, with agreed definitions of MUS and high healthcare use.

PROSPERO registration number CRD42018100388.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ To reduce bias, this review was not restricted to the English language or publication date.
- ⇒ A wide range of medical databases and study types was used to identify potential papers for inclusion.
- ⇒ A broad search strategy with a wide spectrum of search terms, including healthcare cost/utilisation, frequent attenders, MUS and healthcare settings, was used.
- ⇒ Two reviewers conducted study selection, data extraction and quality assessment independently.
- ⇒ A meta-analysis of the results was not possible due to the high risk of bias among studies and methodological heterogeneity between them, thus a narrative summary of the outcome of the selected studies was presented in the final review.

BACKGROUND

Medically unexplained symptoms (MUS) is a general term that refers to the presence of persistent bodily symptoms without an obvious cause, or that cannot be explained by recognised pathological mechanisms.¹ It covers a wide spectrum of complaints that can vary in nature, site, severity and chronicity. In some people, MUS presents as mild discomfort that does not significantly impact functioning; at the more severe end, individuals can experience clinically severe symptoms that cause disability and functional impairment, particularly if appropriate treatment is not sought.² The term ‘MUS’ has received criticism as it suggests a classification based on exclusion and a newer term persistent physical symptoms is preferred by patient groups.³ MUS is a clinical construct, whereby a clinician decides whether symptoms have an organic cause or not based on clinical history, examination findings and investigation results. There are difficulties translating this to research settings, and broad definitions have been needed historically to

capture the breadth of presentations seen in clinical practice. This requirement for an exclusion of organic disease has been removed from Diagnostic and Statistical Manual of Mental Disorders V (DSM-V)⁴ and in the ‘somatic symptom and related disorders’ section, there is a focus on a person’s reaction to physical symptoms rather than the nature of the symptoms themselves. However, as most of the studies included in this review use the term MUS, and predate the publication of DSM-V, we have retained use of the term for the sake of clarity. Within the use of the term MUS, we include all relevant diagnostic terms including the somatoform disorders.

In recent years, MUS has received more attention and it is now routinely referred to in the literature.⁵ MUS is highly prevalent across all healthcare settings and accounts for approximately 45% of all general practice consultations and 20% of new consultations in primary care.^{6 7} MUS is also common in secondary care,⁸ and accounts for 20%–25% of all frequent attenders in medical clinics.^{9 10}

MUS is associated with a significant economic burden for healthcare systems. Patients with MUS are routinely referred for multiple assessments and investigations to little benefit and have longer doctor visits compared with other patients.^{11 12} They incur more sick leave and have significantly higher rates of unemployment.^{13–15} MUS accounts for approximately 10% of the NHS annual expenditure in adults of working age in England. The annual cost attributable to MUS due to lost productivity and decreased quality of life is over £14 billion to the UK economy.¹⁶ However, there is no satisfactory review of the available literature to support such estimations.

The overall purpose of this systematic review is to determine the prevalence of MUS in patients who are high users of healthcare and/or who accrue high healthcare costs and the magnitude of healthcare or associated costs.

Aims

This systematic review aims to:

- ▶ Determine the prevalence of MUS in adults aged ≥18 years who are high users of healthcare or ‘high-cost’ patients (those who accrue high healthcare costs), in comparison with routine users of healthcare.
- ▶ Determine the magnitude of the cost of use of healthcare associated with the presence of MUS among adults who are high users of healthcare.

METHODS

This systematic review was conducted according to the Preferred Reporting Items for Systematic Review and Meta-Analyses statement guidelines (online supplemental file 1).¹⁷ The review protocol is registered in the PROSPERO database. Assuming heterogeneity between studies, we planned to conduct a random effect meta-analysis with and without low-quality studies. The review protocol has been published elsewhere.¹⁸

Patient and public involvement

Patients and the public were not involved in the design and conduct of the study. The research question of this review was informed by the lack of relevant literature examining the prevalence of patients with MUS who are high users of healthcare or who accrue high healthcare costs. Therefore, patients and the public were not involved in defining the research question or outcome measures.

Eligibility criteria

This review included studies where cases are adults aged 18 years or over, who are high users of healthcare services or have high general healthcare costs and MUS. We included studies which described ‘patients who accrue high healthcare costs’, ‘high users’, ‘distressed high users or users of care’, ‘frequent attenders in primary care and secondary care’ and ‘frequent attenders at the emergency department’. In all studies, a recognised measure of the presence of MUS was required. This included application of any of the following: a standardised research interview (the Structured Clinical Interview for Mental Disorders)^{19 20} to generate a diagnosis of a somatoform disorder according to DSM-III,²¹ DSM-IV,²² International Classification of Diseases (ICD-10 F45 diagnoses),²³ structured clinical interview (SCID) for DSM-IV and abridged criteria for somatoform disorder,²⁴ the Diagnostic Criteria for Psychosomatic Research (DCPR),²⁵ General Health Questionnaire-28 (GHQ-28) somatisation subscale,²⁶ Patient Health Questionnaire-15 (PHQ-15),²⁷ Symptom Checklist-90-Revised (SCL-90-R),²⁸ the schedules for clinical assessment in neuropsychiatry (SCAN)²⁹ and clinical assessment for MUS in secondary care. Studies focusing on mental health services, or specific medical subspecialties, for example, oncology or obstetrics, were excluded. Observational studies, including retrospective and prospective cohort studies, case-control and cross-sectional studies were considered for this review. Single case studies and randomised controlled trials were excluded.

Search strategy

A comprehensive search strategy was developed to retrieve articles relevant to the principal aims of this review. The following electronic databases were systematically searched without language restriction from inception to June 2018 and updated on 22 October 2021: MEDLINE, PsycINFO, EMBASE, CINAHL and PROSPERO. The Cochrane library was also included in view of the significant proportion of non-observational studies currently published in the database. Ongoing studies, scientific literature and abstract proceedings were identified by searching the Cochrane Database of Systematic Reviews, Royal College of Psychiatrists, American Psychiatrists Association and Zetoc. Grey literature databases such as Grey Literature Report, OpenGrey, PubliCat and ScienceDaily.com were also examined. Open access theses and dissertations were retrieved from the ProQuest

Table 1 Review eligibility criteria checklist

Study design	Cohort studies (retrospective and prospective) Case-control and nested case-control studies Cross-sectional studies
Study characteristics	Full articles Reference lists of any recent review article Eligible manuscript identified by database search
Participants	Adult aged ≥18 years High user of healthcare Accrue high healthcare costs Presence of MUS
Comparator	Non-high cost and non-high users of healthcare
Outcome	Prevalence of MUS Patient characteristics and context associated with high service usage/costs among patients with MUS Magnitude of cost or use of healthcare associated with the presence of MUS

MUS, medically unexplained symptoms.

Dissertation Thesis Database and thesis.com. The reference lists of any recent review articles and from any other eligible manuscript identified by the above search were hand-searched. The Science Citation Index was used to scan and track study titles. Search strategies for each database are shown in online supplemental file 2.

Study selection

All records retrieved in the database search were imported into the literature management software EndNote to facilitate the management of references. Two reviewers (FJ and OL) independently reviewed the studies identified by the search strategy in two phases. Retrieved titles and abstracts were initially reviewed to identify eligibility for full-text screening. The full texts were then read to determine suitability for inclusion in the review. This was achieved by referring to an inclusion criteria checklist designed a priori (table 1) based on study eligibility criteria. Any discrepancies or differences in opinion were resolved by consensus.

Data extraction

Prior to data extraction, a standardised data extraction form was developed (online supplemental file 3) based on the Hayden *et al*'s framework.³⁰ This was developed iteratively with a focus on population, comparator, outcome and study design, then pilot-tested on known papers independently by two reviewers (FJ and OL). Following initial familiarisation with the included studies, two reviewers (FJ and DR) independently extracted the following information using the form: study design, study details (author(s), publication year and country), recruitment setting (eg, primary care), sample size, diagnostic and screening method used to diagnose MUS, sample characteristics (age and gender), reported prevalence of MUS, magnitude of costs associated with MUS and service use (eg, frequency of attendance). Data extraction using the same method was then completed by two other reviewers (AB and EG) to minimise the likelihood of missed or misinterpreted information. Any discrepancies were resolved by discussion and revisiting the relevant

study. Descriptive data extracted from included papers were summarised in a Microsoft Excel spreadsheet.

Quality assessment

Two reviewers (FJ and DR) completed a quality assessment of each included article independently to reduce bias. The quality assessment focused on sampling strategy, methods used to establish exposure and outcome, and analytical method employed. All selected articles were assessed using a modified form adapted from the Ottawa-Newcastle Scale³¹ assessing the quality of cohort, case-control and cross-sectional studies. The stages and domains of this modified tool are shown in online supplemental file 4. Quality assessment using the same method was then completed by two other reviewers (AB and EG) to minimise the likelihood of personal judgements and subjectivity influencing reported study quality. Any difference in opinion was resolved by further discussion and/or by involving a third reviewer. Risk of bias was presented according to the Cochrane Collaboration recommendations.³² Risk of bias was not displayed as a composite score; instead, an outcome of 'high risk', 'low risk' or 'unclear' was provided for each domain of the tool. A sensitivity analysis may be conducted to assess the effect of including or excluding poor quality studies on the main findings.

Statistical analysis

Estimates of MUS prevalence were considered separately for age, gender, ethnicity and definition of MUS, where applicable. Prevalence estimates were either reported as frequency (%), mean (SD) or OR with 95% CI between groups. Level of heterogeneity between study data was explored. We planned to quantify heterogeneity using the Cochrane Q-test and the I²-statistical test with 95% CI if appropriate. The magnitude of healthcare utilisation and costs defined by high users or patients who accrue high healthcare costs were extracted for each study. Where reported, differences between the cost or use of healthcare associated with the presence of MUS were recorded. Standardised mean difference with accompanying




Figure 1 PRISMA flow diagram of the study selection process. MUS, medically unexplained symptoms; PRISMA, Preferred Reporting Items for Systematic Review and Meta-Analyses; RCT, randomised controlled trial.

95% CI and median OR of costs or healthcare utilisation were extracted. In cases of insufficient data, authors were contacted at least twice by email.

RESULTS

Literature search

The search was updated at the time of manuscript preparation to capture recent and relevant studies. In total, the search strategy yielded 5622 articles. Additional reference searching and grey literature found 13 studies. After excluding 207 duplicates, the titles and abstracts of 5428 articles were screened for relevance. Title and abstract screening resulted in the exclusion of 5290 articles, mainly because these articles did not provide prevalence or healthcare utilisation cost/utilisation or were not primarily focused on MUS. Of the 138 full-text articles assessed, 113 were excluded at full-text review. Twenty-five articles were included in the final analysis. A flow diagram of the study identification process is presented in [figure 1](#).

STUDY CHARACTERISTICS

Twenty-five studies^{33–57} involving 31 650 individuals (high users) from 9 countries were included in the final analysis ([figure 1](#)). A great deal of variation was detected between

studies regarding study design, including study setting, data source and collection, classification and measures used to define MUS and reported prevalence rates. Most studies were performed in the UK (n=8), followed by the USA (n=5), Germany (n=3) and Finland (n=3). Most studies were conducted in primary care or used primary care data (n=15), followed by secondary care (n=7) and a combination of primary and secondary care (n=3). Eight studies^{33 35 36 38 41–43 51} had a cross-sectional design; nine were cohort studies,^{46 48 49 52–57} seven^{34 39 40 44 45 47 50} were case-control studies and one study³⁷ used case note review. Eighteen studies^{33 35 36 39–45 47–50 52–54 56} employed a purposive sampling strategy to recruit patients and seven studies^{34 37 38 46 51 55 57} recruited consecutive patients. Participants were predominantly female, constituting 66% of high users and 60.6% of comparators. Patients were slightly older in the comparator group (mean age: 71.3 years) compared with the high user group (mean age: 69.1 years). An overview of study characteristics can be found in [table 2](#).

Study quality

None of the included studies had a low risk of bias in all criteria of the checklist adapted from the Ottawa-Newcastle Scale.³¹ Sixteen^{34 36 37 39–44 46 48–50 54 56 57} of the 25

Table 2 Study characteristics

Study	High users						Comparators					
	Year	Origin	Setting	Number of participants	Mean age (years) (SD)	Gender (% female)	Ethnicity (% white)	Number of participants	Mean age (years) (SD)	Gender (% female)	Ethnicity (% white)	
Burton <i>et al</i> ³³	2011	UK	Primary and secondary care	267	Not reported	Not reported	Not reported	451	Not reported	Not reported	Not reported	
Ferrari <i>et al</i> ³⁴	2008	Italy	Primary care	50	50.7 (12.9)	76	Not reported	50	38.8 (14.8)	56	Not reported	
Gill <i>et al</i> ³⁵	2011	Spain	Primary care	318	53.3 (13.9)	67.6	Not reported	203	46.7 (14.3)	62.1	Not reported	
Haas <i>et al</i> ³⁶	1999	USA	Primary and secondary care	69	65 (not reported)	64	93	Compared with normative data from another study				
Hansen <i>et al</i> ³⁷	2002	Denmark	Secondary care	294	Not reported	45.9	Not reported	Not reported	Not reported	Not reported	Not reported	
Jacob <i>et al</i> ³⁸	2016	UK	Secondary care	100	Not reported	Not reported	Not reported	No control group				
Jyväsjärvi <i>et al</i> ³⁹	2001	Finland	Primary care	112	53.2 (not reported)	73.2	Not reported	106	51.3 (not reported)	70.8	Not reported	
Jyväsjärvi <i>et al</i> ⁴⁰	1999	Finland	Primary care	113	52.4 (17.0)	72.6	Not reported	107	42.7 (20.6)	71.0	Not reported	
Karlsson <i>et al</i> ⁴¹	1999	Finland	Primary care	67	49.9 (not reported)	68.7	Not reported	No control group				
Katon <i>et al</i> ⁴²	1990	USA	Primary care	119	45.1 (12.6)	62.1	77.2	No control group				
Little <i>et al</i> ⁴³	2001	UK	Primary care	630	Not reported	68	98.1	1898	Not reported	57	98.7	
McGorm <i>et al</i> ⁴⁴	2010	UK	Primary care	193	49 (10.0)	146(76)	Not reported	314	45.5 (11.5)	93 (61)	90 (59)	
Miranda <i>et al</i> ⁴⁵	1991	USA	Secondary care	54	50.1 (11.7)	34 (63)	14 (26)	160	52.1 (12.1)	92 (58)	70 (44)	
Norton <i>et al</i> ⁴⁶	2012	France	Primary care	Data presented for overall population: 61.8% female, age: 42 (range: 18–93), Ethnicity not reported								
Patel <i>et al</i> ⁴⁷	2015	UK	Primary care	71	57 (19)	Not reported	Not reported	71	56 (18)	Not reported	Not reported	
Portegijs <i>et al</i> ⁴⁸	1996	Netherlands	Primary care	45	37	56	Not reported	29	37	58	Not reported	
Reid <i>et al</i> ⁴⁹	2002	UK	Secondary care	61	<46 39 (63.9) ≥46 22 (36.1)	41 (67.2)	51 (83.6) white 10 (16.4) non-white	219	<46. 99 (45.2) ≥46. 120 (54.8)	140 (63.9)	181 (82.6) white 38 (17.4) non-white	
Schmitz and Kruse ⁵⁰	2002	Germany	Secondary care	389	45.4 (13.1)	63.7	Not reported	3337	40.8 (13.1)	48.2	Not reported	
Schneider <i>et al</i> ⁵¹	2011	Germany	Primary care	562	52.9 (17.5)	57.1	Not reported	159	33.5 (12.3)	83 (52.2)	Not reported	
Smith <i>et al</i> ⁵²	1986	USA	Secondary care	41	44 (range: 21–73)	83.7	Not reported	No comparator				
Smith <i>et al</i> ⁵³	2002	USA	Primary care	104	41.3	83	Not reported	66	39.7	65	Not reported	
Smits <i>et al</i> ⁵⁴	2009	Netherlands	Primary care	1008	15–65+	Not reported	Not reported	1601	15–65+	Not reported	Not reported	
Taylor <i>et al</i> ⁵⁵	2012	UK	Primary care	410	41.6 (15.3)	71.2	White=75.1 Black=16.8 Asian=4.8 Other=3.3	No comparator				
van den Bussche ⁵⁶	2016	Germany	Primary and secondary care	23590	73 (6.4)	46.3	Not reported	99 634	71.7 (6.1)	41.4	Not reported	
Williams <i>et al</i> ⁵⁷	2001	UK	Secondary care	35	Not reported	Not reported	Not reported	182	Not reported	Not reported	Not reported	

**Table 3** Quality assessment of included studies

Reference	Selection of participants	Adequate description of study population	Validated method for ascertaining exposure	Validated method confirm outcome	Analysis and controls for confounders	Sample size calculation	Analytical methods appropriate
Burton <i>et al</i> ³³	Low risk	High risk	Unclear	Low risk	Unclear	Low risk	Low risk
Ferrari <i>et al</i> ³⁴	Low risk	Low risk	Low risk	Low risk	High risk	High risk	Low risk
Gilli <i>et al</i> ³⁵	High risk	High risk	Low risk	Low risk	High risk	Unclear	Low risk
Haas <i>et al</i> ³⁶	Low risk	Low risk	Low risk	Low risk	Unclear	High risk	Low risk
Hansen <i>et al</i> ³⁷	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Jacob <i>et al</i> ³⁸	Low risk	High risk	Low risk	High risk	Unclear	Unclear	Low risk
Jyväsjärvi <i>et al</i> ³⁹	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Jyväsjärvi <i>et al</i> ⁴⁰	High risk	Low risk	High risk	Low risk	High risk	Unclear	Low risk
Karlsson <i>et al</i> ⁴¹	Low risk	Low risk	Low risk	Low risk	Unclear	High risk	Low risk
Katon <i>et al</i> ⁴²	Low risk	Low risk	Low risk	Low risk	Unclear	High risk	Low risk
Little <i>et al</i> ⁴³	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
McGorm <i>et al</i> ⁴⁴	Low risk	Low risk	Low risk	Low risk	High risk	Unclear	Low risk
Miranda <i>et al</i> ⁴⁵	Low risk	High risk	Low risk	High risk	High risk	High risk	Low risk
Norton <i>et al</i> ⁴⁶	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk
Patel <i>et al</i> ⁴⁷	Low risk	Unclear	Low risk	Low risk	High risk	High risk	Low risk
Portegijs <i>et al</i> ⁴⁸	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Low risk
Reid <i>et al</i> ⁴⁹	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Low risk
Schmitz and Kruse ⁵⁰	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Schneider <i>et al</i> ⁵¹	Low risk	Unclear	Low risk	Low risk	Low risk	Unclear	Low risk
Smith <i>et al</i> ⁵²	Low risk	Unclear	Low risk	Low risk	High risk	Unclear	Low risk
Smith <i>et al</i> ⁵³	Low risk	Unclear	Low risk	Low risk	Unclear	High risk	Low risk
Smits <i>et al</i> ⁵⁴	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Low risk
Taylor <i>et al</i> ⁵⁵	Low risk	Unclear	Unclear	Low risk	High risk	High risk	Low risk
van den Bussche ⁵⁶	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk
Williams <i>et al</i> ⁵⁷	Low risk	Low risk	Low risk	Low risk	Unclear	High risk	Low risk

'Low risk' indicates low risk of bias in that domain, 'high risk' indicates high risk of bias and 'unclear' indicates where risk of bias was unclear.

studies adequately described the study population (low risk of bias), and most studies^{33 34 36–39 41–57} (n=23) clearly defined the selection of participants, with an adequate description of the target population and inclusion/exclusion criteria. Exposure and outcome were almost universally well described; 22 studies^{34–39 41–54 56 57} presented valid measures for ascertaining exposure (the presence of MUS) and 23^{33–37 39–44 46–57} used appropriate outcome measures (the prevalence of MUS in high users of healthcare and/or the magnitude of costs among high users of healthcare). The other two studies^{38 45} did define the outcome measures but included these in subgroup analysis or cluster of symptoms. For all studies, the analytical approach used was considered appropriate. However, there were only two studies^{33 43} where the risk of bias in sample size considered to be low, because of either the absence or poor quality of sample size calculation in all other studies. Table 3 displays the review quality scores per item based on the adapted Ottawa-Newcastle scale.³¹

Diagnostic tool/measures used

We noted diversity in the methods used to ascertain the presence of MUS. Online supplemental table 1 provides

detailed descriptions of the diagnostic criteria and definitions of MUS used in the included studies. Four of the studies used a two-stage screening process.^{34 35 37 41} Diagnostic instruments with structured interviews were used in 7 studies,^{34 35 37 41 45 47 51} symptom checklists in 10 studies^{36 39 40 42 46 48 50 52 54 56} and the somatic symptoms scale in 1 study.⁴³ In three studies, MUS was identified by data extraction from electronic patient records.^{33 38 44} Furthermore, two studies used a GP assessment and rating sheet to ascertain MUS.^{53 55}

There were a variety of different definitions and thresholds used to identify populations of high users of healthcare. Thresholds involving the number of contacts with healthcare providers were used in twelve studies.^{35 38–41 43 44 47 48 51 53 56} However, there was considerable diversity in the cut-off chosen, and this ranged from three or more referrals in a 5 year period⁴⁴ to thirty or more consultations in 2 years.⁴⁷ Another definition used in four studies^{37 49 50 54} involved a top percentage of healthcare users, ranging from the top 20% to the top 5%. One study defined high users as those who exceeded the mean number of visits within a year, or the top 50% of

healthcare users.³⁶ Ferrari and colleagues generated a list of individuals with the highest number of primary care contacts and worked down this list until fifty participants were recruited.³⁴ The definition of high users was unclear in two studies.^{45 55}

Prevalence estimates

Online supplemental table 1 summarises prevalence rates and ORs of MUS in the included studies. Estimates of the prevalence of MUS ranged from 2.9%⁵⁷ to 76%⁴⁴ in high users of healthcare compared with between 1.1%³⁵ and 61%⁴⁴ in non-high user comparator groups. Only one of the 25 studies provided prevalence data by age group⁴⁹; prevalence rates for those aged <46 years was 63.9% and 22% for those aged over 46 years in high users of healthcare compared with 45.2% for those aged <46 years and 54.8% for those aged ≥46 years in the non-high user group. Two studies provided prevalence rates of MUS by gender.^{44 50} Prevalence rates ranged from 4.6% to 24% in males and from 8.3% to 76% in females in the high user group compared with from 2.9% to 39% in males and from 6.7% to 61% in females in the comparator group. Twelve studies^{34 35 39 43 44 47–51 54 57} included a comparator group, of those 11 studies^{34 35 39 43 44 47–51 54} reported a higher prevalence of MUS in high users of healthcare or 'high-cost' patients.

Seven studies^{34 35 37 39 43 44 47} reported ORs instead of or in addition to prevalence estimates, to summarise the difference between MUS groups and comparators. The ratios reported in the included studies ranged from 1.45 to 17, and of these, none were less than 1 (ie, in favour of comparators vs MUS). All but one of the reported ORs indicated statistical significance, supporting the claim that MUS is more prevalent among high users of healthcare compared with those who are not. The only exception was that reported by Little and colleagues⁴³ when a cut-off score of 1–2 for the Somatic Symptom Inventory was used to define MUS (95% CI 0.81 to 1.62).

Two studies^{48 56} used relative risk (RR) of being a high user of healthcare. Patients with MUS were twice as likely (RR: 2.33, 95% CI 2.20 to 2.50) to contact more than 10 different general practices and/or contact ≥3 different medical specialties.⁵⁶ Another study found that high healthcare utilisation was twice as high (RR: 2.0, 95% CI 1.10 to 3.60) in patients with a DSM-III-R diagnosis of somatisation.⁴⁸

For the reader's interest, we calculated unadjusted ORs and 95% CIs for studies where these were not reported, using the number of events in the high user and non-high user groups where possible. These calculations were conducted for six studies,^{35 49–51 54 57} and the results are summarised in online supplemental table 1, indicated by an asterisk. Generally, these ORs were greater than 1, suggesting that MUS is more prevalent among high healthcare users. There were two exceptions to this trend. The first was reported by Reid and colleagues, where participants aged ≥46 years who were in the top 5% of all outpatient appointments over a 3-year period had a

lower prevalence of MUS compared with those who were not high healthcare users (22% vs 54.8%; unadjusted OR: 0.46, 95% CI 0.25 to 0.87).⁴⁹ The other exception was found by Williams and colleagues, where routine attenders to an emergency department were associated with higher rates of somatoform disorder compared with frequent attenders (3.8% vs 2.9%; unadjusted OR: 0.73, 95% CI 0.01 to 6.04).⁵⁷ These unadjusted ORs varied considerably, ranging from 0.46⁴⁹ to 6.75,⁵¹ and CIs varied from narrow to as broad as 0.34–19.7,³⁵ suggesting that these anomalous results could be a product of the methodological heterogeneity between the studies.

Prevalence estimates according to diagnostic criteria

Numerous different criteria were used in the included studies to ascertain the diagnosis of MUS. The diagnostic criteria only used once included the DCPR,³⁴ the GHQ-28 somatisation subscale,³⁵ the Whitley index for somatisation,³⁷ the Somatic Symptom Inventory,⁴⁴ the International Classification of Primary Care code,⁵⁴ and diagnosis using the ICD-10 code for somatisation disorder.⁵⁶ Below, we summarise the prevalence estimates by grouping them according to diagnostic criteria used to define MUS.

Three studies^{38 44 49} reported prevalence estimates of MUS derived from patients' medical records; the rates ranged from 22%⁴⁹ to 76%⁴⁴ in high users of healthcare compared with between 39% and 61%⁴⁴ in non-high user comparator groups. DSM-IV criteria for somatoform disorder were used to define MUS in four studies,^{46 47 50 51} and the method used to collect the relevant information included PHQ-9,⁴⁶ PHQ-15⁵¹ and diagnostic interviews.^{47 50} The rates of MUS ranged from 4.6%⁵⁰ to 55.9%⁵¹ among high users of healthcare, compared with 2.9%⁵⁰ to 15.8%⁵¹ in non-high user comparator groups. Five studies^{41 42 45 48 52} used interviews of differing formats to diagnose somatisation according to DSM-III criteria. The prevalence of somatisation among high healthcare users ranged from 20.2%⁴² to 58%,⁴⁸ and of these 5 studies only Portegijs and colleagues reported prevalence of MUS among the comparator group (29%).⁴⁸ Two studies^{36 39} reported prevalence rates of somatoform disorder defined by SCL-36 and SCL-90; the rates of somatoform disorder were 28.6% in the high user group compared with 16% in the non-high user group³⁹ and 56.5% in the high user group.³⁶ Further two studies reported prevalence rates using SCAN to diagnose MUS,^{35 57} and prevalence among the high healthcare users ranged from 2.9%⁵⁷ to 17.3%,³⁵ and 3.4%³⁵ to 3.8%⁵⁷ among comparators. Finally, three studies used clinical assessments to diagnose MUS,^{33 53 55} although only Smith and colleagues reported prevalence rates. These were only presented for the high healthcare user group as 61.2% in 1995, 17.8% in 1996 and 13.1% in 1997.⁵³

Prevalence estimates according to different clinical settings

There were wide variations in the prevalence of MUS among high users in both primary care and secondary care settings. In primary care, prevalence estimates for

MUS ranged from 1.7%³⁵ to 76%⁴⁴ and in the secondary care setting from 4.6%⁵⁰ to 63.9%.⁴⁹ Given the large clinical heterogeneity between the studies included in this review, we determined not to proceed with a meta-analysis. The decision was also informed by the high risk of bias among the included studies due to insufficient attention to power.

Magnitude of healthcare costs

Only three studies included in the review investigated the magnitude of healthcare costs associated with the presence of MUS who are high users of healthcare.^{33 49 52} The comparator group was different across these three studies, precluding pooling of healthcare cost estimates. Therefore, we have summarised the main findings descriptively. Burton and colleagues³³ compared individuals referred at least 3 times from primary to secondary care in the previous 5 years with MUS (repeatedly referred with MUS, RRMUS) with those infrequently referred (IRS) and those frequently referred with medically explained symptoms (RRMES). The RRMUS group was associated with significantly greater costs per patient over a 5-year period than the IRS group, with a difference of £3539 (95% CI 1458 to 5261) in inpatient costs, £778 (95% CI 705 to 852) in outpatient costs, £99 (95% CI 74 to 123) in emergency department costs, £260 (95% CI 224 to 296) in investigation costs and £4416 (95% CI 2315 to 6517) in total costs. The RRMUS group incurred greater investigation costs than the RRMES group, with an average difference of £102 per patient over 5 years (95% CI 56 to 149). However, there was no significant difference in the average inpatient, outpatient, emergency department or total costs between the RRMUS and RRMES groups (difference (95% CIs) = £491 (-1737 to 2718), £25 (-78 to 127), £22 (-7 to 52) and £537 (-1723 to 2798), respectively). Reid and colleagues⁴⁹ investigated frequent attenders of secondary care services and identified patients with MUS and compared their healthcare use and costs with patients without MUS. Patients with MUS were associated with greater mean costs of investigations (£244 vs £124, mean difference = £120, 95% CI 68 to 172) and mean total costs in secondary care (£955 vs £882, mean difference = £73, 95% CI 39 to 185). However, average costs per consultation episode in secondary care were lower for those with MUS than those without (£226 vs £230, mean difference = £104, 95% CI 72 to 136). Smith and colleagues⁵² investigated healthcare utilisation of 41 patients meeting the essential features of somatisation disorder. Healthcare utilisation data were compiled in quarterly intervals. The mean inpatient charges averaged \$599 (SD: ±\$219), while outpatient charges averaged \$215 (SD: ±\$32). The mean combined charges (inpatient and outpatient) averaged \$814 (SD: ±299).

DISCUSSION

The purpose of this review was to systematically investigate the existing literature to determine the prevalence

of MUS in patients who are high users of healthcare and/or who accrue high healthcare costs and the magnitude of healthcare or associated costs. Although there is a vast body of literature estimating the prevalence of MUS and its associated costs, to the best of our knowledge, no previous study has focused on high users or high-cost patients. Our findings showed that there was great variation among studies on several different methodological parameters, including design; definitions and methods of identifying 'high use' or 'high-cost' populations; definitions and methods used to identify people with MUS within high use/high-cost populations; comparator groups; country and type of health service where the study was undertaken; and clinical setting (primary, secondary or ED).

Most studies adequately described the study population and inclusion/exclusion criteria. Exposure and outcome were almost universally well described and most studies used valid measures for ascertaining exposure (the presence of MUS) and outcome measures (the prevalence of MUS in high users of healthcare and/or the magnitude of costs among high users of healthcare). However, there was a 'high' risk of bias in most studies due to a lack of adequate consideration of power. Another frequently observed limitation was the lack of consecutive sampling in many studies, which could be explained by practical difficulties in reaching the target population. The degree of variation across the studies, combined with our quality findings that most studies were at high risk of bias, meant that we did not think it was appropriate to pool the results in a meta-analysis.

SUMMARY OF EVIDENCE

Prevalence estimates

The estimated prevalence of MUS was reported to be greater among high healthcare users compared with non-high user comparators for all but one of the 12 studies that included a comparator group^{34 35 39 43 44 47-51 54 57}; however, these estimates varied considerably between studies. This is not surprising given the variability in methodology across the studies. Prevalence estimates by age and gender were poorly recorded. Only one study reported an overall higher prevalence of MUS among the non-high user comparator compared with the high user group.⁵⁷ This could partly be explained by a disparity in sample size between the groups, with fewer in the high user group (n=77) than the comparator group (n=182). The authors also suggest that this unexpected finding could be attributed to suboptimal sensitivity of the SCAN tool to identify somatoform autonomic disorder (F45.3). This study importantly highlights that not all patients with a somatoform disorder are high users of healthcare.

Only two studies reported MUS prevalence among high users according to gender,^{43 49} and both found higher rates in females. One study provided prevalence estimates by age group,⁴⁸ and suggested high users were more likely to have MUS if they were aged under 46 years compared

with those who were older. A recent systematic review of the general characteristics of high-cost patients found costs were higher in older groups, but that mental health 'high-cost' patients tended to be younger.⁵⁸ Another study found that young adults (aged 18–24 years) with somatic symptoms and related disorders frequently used the healthcare system with substantial healthcare costs before and after diagnosis.⁵⁹ Despite the wide prevalence of MUS spectrum identified in this review, we believe MUS is a useful construct as it is consistently associated with increased morbidity and healthcare expenditure. Further research and interventions are required, incorporating a uniform definition and diagnostic approach.

Magnitude of cost

Only three studies^{33 49 52} investigated the magnitude of healthcare costs associated with the presence of MUS who are high users of healthcare. Although they provided estimates of the magnitude of costs associated with MUS among adults who are high users of healthcare, the comparability of these studies was limited by heterogeneity in terms of study design, follow-up period, outcome measures and definitions of comparator group. Two of these studies^{33 49} compared costs between those with MUS and those without, and unsurprisingly both reported greater investigation and total costs associated with MUS. The other study provided descriptions of quarterly inpatient and outpatient costs associated with somatisation disorder.⁵² Our findings provide preliminary evidence to suggest that MUS is associated with greater healthcare costs, and interventions aimed at identifying and treating MUS early could help to reduce these costs in addition to improving patient outcomes. Healthcare costs per patients repeatedly referred with MUS over a 5-year period were considerably higher compared with those who were infrequently referred. Our results concur with those of a previous study showing costs of hospitalised patients with MUS to different wards across several hospitals between 2008 and 2018 in Northern Italy, in which the overall estimated costs of hospitalisation was €475 410 with a mean annual cost per patient of approximately €48 000.⁶⁰ In both general practices and outpatient clinics of a regional community mental health service in greater Rotterdam (the Netherlands), the mean direct (use of healthcare) and indirect costs (absenteeism and presenteeism) were estimated at €6815 per patient per year.⁶¹ A recent systematic review investigating cost-of-illness studies and economic evaluations of MUS found that direct excess treatment costs (healthcare utilisation) per patient ranged from \$432 to \$5353 per year. There are also indirect costs (eg, presenteeism and sickness absence), which are estimated to be approximately seven times greater than the direct costs.⁶²

Strengths and limitations

Several recent studies have found that mental health problems are common in high use or high-cost populations.^{63–65} It is important, however, to begin to understand

the nature of these mental health problems to plan effective interventions. This is the first systematic review to identify and present an in-depth synthesis of the best available evidence describing the prevalence of MUS in patients who are high users of healthcare and/or who accrue high healthcare costs. Strengths of this systematic review include the rigorous methodological approach employed using an established methodological framework.^{17 30 31} Two independent reviewers were involved in study selection, data extraction and quality assessment, and a third reviewer was included to ensure overall methodological consistency and to resolve any disagreements. To ensure an exhaustive review of the available literature, a comprehensive search strategy was implemented with broad inclusion criteria. Additionally, the search was not restricted to the English language and grey literature sources were considered, to minimise the effects of language and publication bias, respectively. The search was repeated at the time of manuscript preparation to capture recent and relevant studies.

There are some limitations to the present study. First, the quality of these studies was variable and many did not report essential data, such as outcome measures, statistical power, reliability of measures and information about effect measures between intervention and control group. Second, given the limitations of the reported data, the high risk of bias among the included studies and the wide heterogeneity between them, we were unable to combine data in a meta-analysis, and instead results were reported as a narrative summary. Third, due to limited data on gender and age, we were unable to adequately measure the effect of these variables, although this represents an important area for future research. We also planned to assess publication bias but were unable to do so owing to the wide heterogeneity between the included studies. The generalisability of these findings may be uncertain, although each setting is inevitably unique and healthcare professionals may use different assessment criteria to ascertain MUS and definitions to identify high or costly healthcare users.

Implications of results

The findings suggest that people with MUS are over-represented in populations of high users of healthcare and high-cost patients, accounting for a disproportionate amount of healthcare use in both primary and secondary care settings. Given the use of healthcare resources by this population and the associated costs, interventions to identify those with MUS and to deliver targeted psychological interventions may reduce healthcare costs, optimise pharmacological interventions and improve integration of primary and secondary care while improving overall patient outcomes. van den Bussche and colleagues⁵⁶ argue that frequent attendance appears to involve various aspects of the healthcare system, including healthcare providers, patients and the disparaging healthcare system, contributing to high utilisation. Strategies to reduce



healthcare costs should, therefore, carefully consider these systemic issues.

In terms of future research, our findings demonstrate a clear need for a standardised approach to understanding MUS and high users of healthcare. There was a great deal of variety of methods used to ascertain the presence of MUS and to identify those who are high healthcare users. An agreed definition of MUS is required to allow comparison and synthesis of findings in the academic literature.⁶⁶ Similarly, a universal definition of high healthcare users would be helpful to integrate the estimates of sociodemographic and clinical characteristics of this high-need group. Hayes and colleagues found that there was significant variation in healthcare use and costs among those with high needs, defined as those with three or more chronic diseases associated with a functional impairment.⁶⁷

CONCLUSION

MUS is common among adults who are high users of healthcare and/or who accrue high healthcare costs. The present review quantifies the prevalence of MUS among high healthcare users and describes estimates of costs associated with this population. Significant heterogeneity was found between the included studies, particularly pertaining to methods of ascertaining MUS and definitions of high healthcare users, in addition to high risk of bias among the studies. These factors precluded meta-analysis. Nonetheless, this review indicates that this group of patients incurs a disproportionate level of healthcare resources compared with the general population, which should be considered by policymakers, clinicians and researchers. It also indicates that this group of individuals pursues specific form of health-seeking behaviour that should be adequately understood and addressed. Future studies should consider approaches to high users associated with MUS by carefully and consistently defining frequent attendance, measures used to define MUS and the study setting.

Dissemination

Any data generated from this systematic review will be made available from the corresponding author on reasonable request.

Author affiliations

¹Centre of Precision Rehabilitation for Spinal Pain School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, UK

²Leeds and York Partnership NHS Foundation Trust, Leeds, UK

³Division of Psychological and Social Medicine, Leeds Institute of Health Sciences, School of Medicine, University of Leeds, Leeds, UK

⁴Patient-Centred Outcomes Research Group, Leeds Institute of Medical Research at St. James's University Hospital, School of Medicine, University of Leeds, Leeds, UK

⁵Division of Nursing, Social Work and Midwifery, School of Health Sciences, Faculty of Biology, Medicine and Health, Manchester Academic Health Science Centre, The University of Manchester, Manchester, UK

⁶School of Medicine, Leeds Institute of Health Sciences, University of Leeds, Room 10.39, Worsley Building, Clarendon Way, Leeds, UK

Twitter Daniel Romeu @DanielJRomeu and Amy Blakemore @DrAmyBlakemore

Contributors All authors contributed to the focus of this systematic review topic and approved the final version for publication. FJ drafted the initial manuscript with guidance and feedback from DR, EG and AB at all stages. FJ drafted the final version, performed the search and is a guarantor. The search strategy was developed by FJ and iteration discussed with OL, EG and AB. FJ and OL performed the initial screening and study selection. FJ and DR collected data from the included studies, conducted quality assessment, performed data analysis/synthesis, drafted and critically reviewed the manuscript. EG ensured data extraction consistency.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This review does not require ethical approval as only existing published data available in scientific databases will be used. Findings of this systematic review will be presented for peer review in an appropriate journal.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data sharing not applicable as no datasets generated and/or analysed for this study.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Ferozkhan Jadhakhan <http://orcid.org/0000-0002-4545-3703>

Amy Blakemore <http://orcid.org/0000-0003-0972-100X>

Elspeth Guthrie <http://orcid.org/0000-0002-5834-6616>

REFERENCES

- Deary V, Chalder T, Sharpe M. The cognitive behavioural model of medically unexplained symptoms: a theoretical and empirical review. *Clin Psychol Rev* 2007;27:781–97.
- Kroenke K, Price RK. Symptoms in the community. prevalence, classification, and psychiatric comorbidity. *Arch Intern Med* 1993;153:2474–80.
- Marks EM, Hunter MS. Medically unexplained symptoms: an acceptable term? *Br J Pain* 2015;9:109–14.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. In: . 5Th. Washington DC: American Psychiatric Association, 2013.
- Nettleton S. 'I just want permission to be ill': towards a sociology of medically unexplained symptoms. *Soc Sci Med* 2006;62:1167–78.
- Haller H, Cramer H, Lauche R. Somatoform disorders and medically unexplained symptoms in primary care: a systematic review and meta-analysis of prevalence. *Dtsch Arztebl Int* 2015;112:279–87.
- Knapp M, McDaid D, Parsonage M. *Mental health promotion and mental illness prevention: the economic case*. London school of economics and political sciences: personal social services research unit, 2011.
- Nimnuan C, Hotopf M, Wessely S. Medically unexplained symptoms: an epidemiological study in seven Specialities. *J Psychosom Res* 2001;51:361–7.
- Fink P. The use of hospitalizations by persistent somatizing patients. *Psychol Med* 1992;22:173–80.

- 10 Reid S, Wessely S, Crayford T, *et al.* Medically unexplained symptoms in frequent attenders of secondary health care: retrospective cohort study. *BMJ* 2001;322:767–9.
- 11 Smith GR. The course of somatization and its effects on utilization of health care resources. *Psychosomatics* 1994;35:263–7.
- 12 Barsky AJ, Ettner SL, Horsky J, *et al.* Resource utilization of patients with hypochondriacal health anxiety and somatization. *Med Care* 2001;39:705–15.
- 13 Martin A, Rauh E, Fichter M, *et al.* A one-session treatment for patients suffering from medically unexplained symptoms in primary care: a randomized clinical trial. *Psychosomatics* 2007;48:294–303.
- 14 Hiller W, Fichter MM, Rief W. A controlled treatment study of somatoform disorders including analysis of healthcare utilization and cost-effectiveness. *J Psychosom Res* 2003;54:369–80.
- 15 Swartz M, Landerman R, George L. Somatization disorder. In: Regier DA, ed. *Robins In.* New York: Psychiatric Disorders in AmericaFree Press, 1991: 220–57.
- 16 Birmingham SL, Cohen A, Hague J, *et al.* The cost of somatisation among the working-age population in England for the year 2008–2009. *Ment Health Fam Med* 2010;7:71–84.
- 17 Page MJ, McKenzie JE, Bossuyt PM, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- 18 Jadhakhan F, Lindner OC, Blakemore A, *et al.* Prevalence of medically unexplained symptoms in adults who are high users of health care services: a systematic review and meta-analysis protocol. *BMJ Open* 2019;9:e027922.
- 19 First MB, Spitzer RL, Gibbon M. *Structured clinical interview for DSM-IV-TR axis disorders, research version, patient edition SCID/II-P.* New York: Biometrics Research, New York State Psychiatric Institute, 2002.
- 20 Spitzer RL, Williams JBW, Gibbon M. *User's guide for the structured clinical interview for DSM-III-R: SCID.* Arlington, VA, USA: American Psychiatric Association, 1990.
- 21 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* In: . 3rd. Washington DC: American Psychiatric Association, 1980.
- 22 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders.* In: . 4Th. Washington DC: American Psychiatric Association, 2000.
- 23 ICD. *Manual of the International classification of diseases, injuries, and causes of death. tenth revision, ICD-10.* GenevaWorld Health Organization, 1993.
- 24 First MB, Gibbon M, Spitzer RL. *Structured clinical interview for DSM-IV axis II personality disorders (SCID-II).* Washington DC: American Psychiatric Association, 1997.
- 25 Mangelli L, Rafanelli C, Porcelli P. Psychological factors affecting medical conditions. A new classification for DSM-V. Basel. In: Porcelli P, Sonino N, eds. *Interview for the diagnostic criteria for psychosomatic research.* CH: Karger, 2007: 174–81.
- 26 Goldberg DP, Hillier VF. A scaled version of the general health questionnaire. *Psychol Med* 1979;9:139–45.
- 27 Kroenke K, Spitzer RL, Williams JBW. The PHQ-15: validity of a new measure for evaluating the severity of somatic symptoms. *Psychosom Med* 2002;64:258–66.
- 28 Derogatis LR. *SCL-90-R: administration, scoring of procedures Manual-II for the R (revised version and other instruments of the psychopathology rating scale series.* Clinical Psychometric Research Incorporated, 1992.
- 29 Wing JKet *et al.* "SCAN. Schedules for Clinical Assessment in Neuropsychiatry". *Arch.Gen.Psychiatry* 1990;47:589–93.
- 30 Hayden JA, Côté P, Bombardier C. Evaluation of the quality of prognosis studies in systematic reviews. *Ann Intern Med* 2006;144:427–37.
- 31 Hartling L, Milne A, Hamm MP, *et al.* Testing the Newcastle Ottawa scale showed low reliability between individual reviewers. *J Clin Epidemiol* 2013;66:982–93.
- 32 Higgins JPT, Altman DG, Gotzsche PC, *et al.* The Cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928–9.
- 33 Burton C, McGorm K, Richardson G, *et al.* Healthcare costs incurred by patients repeatedly referred to secondary medical care with medically unexplained symptoms: a cost of illness study. *J Psychosom Res* 2012;72:242–7.
- 34 Ferrari S, Galeazzi GM, Mackinnon A, *et al.* Frequent attenders in primary care: impact of medical, psychiatric and psychosomatic diagnoses. *Psychother Psychosom* 2008;77:306–14.
- 35 Gill M, Luciano JV, Serrano MJ, *et al.* Mental disorders among frequent attenders in primary care: a comparison with routine attenders. *J Nerv Ment Dis* 2011;199:744–9.
- 36 Haas LJ, Spendlove DC, Silver MP, *et al.* Psychopathology and emotional distress among older high-utilizing health maintenance organization patients. *J Gerontol A Biol Sci Med Sci* 1999;54:M577–82.
- 37 Hansen MS, Fink P, Frydenberg M, *et al.* Use of health services, mental illness, and self-rated disability and health in medical inpatients. *Psychosom Med* 2002;64:668–75.
- 38 Jacob R, Wong ML, Hayhurst C, *et al.* Designing services for frequent attenders to the emergency department: a characterisation of this population to inform service design. *Clin Med* 2016;16:325–9.
- 39 Jyväskylä S, Joukamaa M, Väisänen E, *et al.* Somatizing frequent attenders in primary health care. *J Psychosom Res* 2001;50:185–92.
- 40 Jyväskylä S, Joukamaa M, Väisänen E, *et al.* Alexithymia, hypochondriacal beliefs, and psychological distress among frequent attenders in primary health care. *Compr Psychiatry* 1999;40:292–8.
- 41 Karlsson H, Joukamaa M, Lahti I, *et al.* Frequent attender profiles: different clinical subgroups among frequent attender patients in primary care. *J Psychosom Res* 1997;42:157–66.
- 42 Katon W, Von Korff M, Lin E, *et al.* Distressed high utilizers of medical care. DSM-III-R diagnoses and treatment needs. *Gen Hosp Psychiatry* 1990;12:355–62.
- 43 Little P, Somerville J, Williamson I, *et al.* Psychosocial, lifestyle, and health status variables in predicting high attendance among adults. *Br J Gen Pract* 2001;51:987–84.
- 44 McGorm K, Burton C, Weller D, *et al.* Patients repeatedly referred to secondary care with symptoms unexplained by organic disease: prevalence, characteristics and referral pattern. *Fam Pract* 2010;27:479–86.
- 45 Miranda J, Pérez-Stable EJ, Muñoz RF, *et al.* Somatization, psychiatric disorder, and stress in utilization of ambulatory medical services. *Health Psychol* 1991;10:46–51.
- 46 Norton J, David M, de Roquefeuil G, *et al.* Frequent attendance in family practice and common mental disorders in an open access health care system. *J Psychosom Res* 2012;72:413–8.
- 47 Patel S, Kai J, Atha C, *et al.* Clinical characteristics of persistent frequent attenders in primary care: case-control study. *Fam Pract* 2015;32:cmv076–30.
- 48 Portegijs PJ, van der Horst FG, Proot IM, *et al.* Somatization in frequent attenders of general practice. *Soc Psychiatry Psychiatr Epidemiol* 1996;31:29–37.
- 49 Reid S, Wessely S, Crayford T, *et al.* Frequent attenders with medically unexplained symptoms: service use and costs in secondary care. *Br J Psychiatry* 2002;180:248–53.
- 50 Schmitz N, Kruse J. The relationship between mental disorders and medical service utilization in a representative community sample. *Soc Psychiatry Psychiatr Epidemiol* 2002;37:380–6.
- 51 Schneider A, Hörlein E, Wartner E, *et al.* Unlimited access to health care—impact of psychosomatic co-morbidity on utilisation in German general practices. *BMC Fam Pract* 2011;12:51.
- 52 Smith GR, Monson RA, Ray DC. Patients with multiple unexplained symptoms. their characteristics, functional health, and health care utilization. *Arch Intern Med* 1986;146:69–72.
- 53 Smith RC, Gardiner JC, Lyles JS, *et al.* Minor acute illness: a preliminary research report on the "worried well". *J Fam Pract* 2002;51:24–9.
- 54 Smits FTM, Brouwer HJ, ter Riet G, *et al.* Epidemiology of frequent attenders: a 3-year historic cohort study comparing attendance, morbidity and prescriptions of one-year and persistent frequent attenders. *BMC Public Health* 2009;9:36.
- 55 Taylor RE, Marshall T, Mann A, *et al.* Insecure attachment and frequent attendance in primary care: a longitudinal cohort study of medically unexplained symptom presentations in ten UK general practices. *Psychol Med* 2012;42:855–64.
- 56 van den Bussche H, Kaduszkiewicz H, Schäfer I, *et al.* Overutilization of ambulatory medical care in the elderly German population? – an empirical study based on national insurance claims data and a review of foreign studies. *BMC Health Serv Res* 2016;16:1–16.
- 57 Williams ER, Guthrie E, Mackway-Jones K, *et al.* Psychiatric status, somatisation, and health care utilization of frequent attenders at the emergency department: a comparison with routine attenders. *J Psychosom Res* 2001;50:161–7.
- 58 Wammes JGG, van der Wees PJ, Tanke MAC, *et al.* Systematic review of high-cost patients' characteristics and healthcare utilisation. *BMJ Open* 2018;8:e023113.
- 59 Saunders NR, Gandhi S, Chen S. Healthcare use and costs of children, adolescents and young adults with somatic symptoms and related disorders. *JAMA* 2020;3:e2011295.
- 60 Poloni N, Caselli I, Ielmini M, *et al.* Hospitalized patients with medically unexplained physical symptoms: clinical context



- and economic costs of healthcare management. *Behav Sci* 2019;9:E80.
- 61 Zonneveld LNL, Sprangers MAG, Kooiman CG, *et al.* Patients with unexplained physical symptoms have poorer quality of life and higher costs than other patient groups: a cross-sectional study on burden. *BMC Health Serv Res* 2013;13:520.
- 62 Konnopka A, Schaefer R, Heinrich S, *et al.* Economics of medically unexplained symptoms: a systematic review of the literature. *Psychother Psychosom* 2012;81:265–75.
- 63 Ng SH-X, Rahman N, Ang IYH, *et al.* Characterization of high healthcare utilizer groups using administrative data from an electronic medical record database. *BMC Health Serv Res* 2019;19:452.
- 64 Johnson TL, Rinehart DJ, Durfee J, *et al.* For many patients who use large amounts of health care services, the need is intense yet temporary. *Health Aff* 2015;34:1312–9.
- 65 Young HW, Martin ET, Kwiatkowski E, *et al.* The association between emergency department Super-Utilizer status and willingness to participate in research. *Emerg Med Int* 2020;2020:1–6.
- 66 Burton C, Fink P, Henningsen P, *et al.* Functional somatic disorders: discussion paper for a new common classification for research and clinical use. *BMC Med* 2020;18:34.
- 67 Hayes SL, Salzberg CA, McCarthy D, *et al.* High-Need, high-cost patients: who are they and how do they use health care? a population-based comparison of demographics, health care use, and expenditures. *Issue Brief* 2016;26:1–14.