



UNIVERSITY OF LEEDS

This is a repository copy of *Genetic and Geographical Associations With Six Dimensions of Psychotic Experiences in Adolescence*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/193222/>

Version: Accepted Version

Article:

Maxwell, J, Ronald, A, Cardno, AG orcid.org/0000-0002-6136-5965 et al. (3 more authors) (2023) Genetic and Geographical Associations With Six Dimensions of Psychotic Experiences in Adolescence. *Schizophrenia Bulletin: The Journal of Psychoses and Related Disorders*, 49 (2). sbac149. pp. 319-328. ISSN 0586-7614

<https://doi.org/10.1093/schbul/sbac149>

© The Author(s) 2022. This is an author produced version of an article published in *Schizophrenia Bulletin: The Journal of Psychoses and Related Disorders*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 **GENETIC AND GEOGRAPHICAL ASSOCIATIONS WITH SIX DIMENSIONS OF PSYCHOTIC**
2 **EXPERIENCES IN ADOLESCENCE**

3
4 *Jessye Maxwell*^{*,1,2}, *Angelica Ronald*³, *Alastair G. Cardno*⁴, *Gerome Breen*^{1,2}, *Kaili Rimfeld*^{1,5},
5 *Evangelos Vassos*^{1,2}

6
7 ¹Social Genetic and Developmental Psychiatric Centre, Institute of Psychiatry, Psychology and
8 Neuroscience, King's College London, London, UK; ²NIHR Maudsley Biomedical Research
9 Centre, South London and Maudsley NHS Trust, London, UK; ³Department of Psychological
10 Sciences, Birkbeck, University of London, London, UK; ⁴Division of Psychological and Social
11 Medicine, University of Leeds, Leeds, UK; ⁵Department of Psychology, Royal Holloway,
12 University of London, Surrey, UK.
13
14
15

16 *To whom correspondence should be addresses; [Social Genetic and Developmental](#)
17 [Psychiatry Centre, Memory Lane, London, SE5 8AF](#); tel: [+44 \(0\)20 7848 0873](#); e-mail:
18 jessye.maxwell@kcl.ac.uk.
19

20 **Abstract word count:** 242

21 **Manuscript word count:** 3974

22

23 Abstract

24

25 **Background and hypothesis:** Large-scale epidemiological and genetic research have shown
26 that psychotic experiences in the community are risk factors for adverse physical and
27 psychiatric outcomes. We investigated the associations of 6 types of specific psychotic
28 experiences and negative symptoms assessed in mid-adolescence with well-established
29 environmental and genetic risk factors for psychosis.

30

31 **Study design:** Fourteen polygenic risk scores (PRS) and nine geographical environmental
32 variables from 3,590 participants of the Twins Early Development Study (mean age 16) were
33 associated with paranoia, hallucinations, cognitive disorganisation, grandiosity, anhedonia,
34 and negative symptoms scales. The predictors were modelled using LASSO regularisation
35 separately (Genetic and Environmental models) and jointly (GE model).

36

37 **Study results:** In joint GE models, we found significant genetic associations of negative
38 symptoms with educational attainment PRS ($\beta = -0.07$; 95%CI = -0.12 to -0.04); cognitive
39 disorganisation with neuroticism PRS ($\beta = 0.05$; 95%CI = 0.03 to 0.08); paranoia with MDD (β
40 = 0.07; 95%CI = 0.04 to 0.1), BMI ($\beta = 0.05$; 95%CI = 0.02 to 0.08), and neuroticism PRS ($\beta =$
41 0.05; 95%CI = 0.02 to 0.08). From the environmental measures only family SES ($\beta = -0.07$,
42 95%CI = -0.10 to -0.03) and regional education levels ($\beta = -0.06$; 95%CI = -0.09 to -0.02) were
43 associated with negative symptoms.

44

45 **Conclusions:** Our findings advance understanding of how genetic propensity for psychiatric,
46 cognitive, and anthropometric traits, as well as environmental factors, together play a role
47 in creating vulnerability for specific psychotic experiences and negative symptoms in mid-
48 adolescence.

49

50 **Key words:** environmental risk, geographical variables, polygenic risk scores, psychosis
51 scales, schizophrenia, prediction models

52

53

55 Introduction

56

57 It has been hypothesized that there is a psychosis continuum in the general population, with
58 clinical psychotic disorders such as schizophrenia and bipolar disorder at the extreme end of
59 continuously distributed phenotypes.^{1,2} Symptoms of psychotic disorders include positive
60 (delusions, hallucinations and disorganised symptoms), and negative (lack of volition, reduced
61 speech output, and flattening of affect) domains.³⁻⁷ 'Psychotic experiences' refers to
62 subclinical psychotic-like features measured in the general population, representing the full
63 range of severity.⁸⁻¹⁰ Consistent with clinical psychotic disorders, psychotic experiences have
64 been shown to have a multidimensional structure, although there is variation in the number
65 and definition of the dimensions across studies.^{8,11-15}

66

67 Large-scale epidemiological research has shown that psychotic experiences in the community
68 are risk factors for a range of later physical and mental health disorders and adverse
69 outcomes.¹⁶⁻¹⁸ Consequently, psychotic experiences may be an important target for providing
70 insights into the causes of psychosis and for preventative strategies.^{19,20} Studies have
71 demonstrated a shared aetiology between psychotic experiences and clinical psychotic
72 disorders, including both genetic and environmental risk factors.²¹⁻²³ Environmental
73 measures in common with clinical psychotic disorders include urbanicity, migrant status and
74 socio-economic status.^{3,21,24} Many studies also suggest that psychotic experiences in
75 childhood may be associated with a broader set of psychiatric disorders and behaviours with
76 onset in early adulthood, such as affective, anxiety, and substance use disorders and
77 suicidality.^{16,25-30}

78

79 Twin studies have estimated the heritability of psychotic experiences around 30-50%^{15,31-36}
80 and molecular genetic studies have attributed 3-17% of the variance in psychotic experiences
81 to common genetic variation.^{22,23,37} Schizophrenia polygenic risk score (PRS) has been
82 associated with psychotic experiences in adolescence,^{23,38} however, there is some
83 inconsistency across studies,^{39,40} which may be attributable to differences in sample
84 characteristics or definitions of psychotic experiences.^{22,41} Psychotic experiences are also
85 known to be associated with genetic predisposition to depression and neurodevelopmental
86 disorders.^{23,42,43} As such, in the current study, we employ a multi-PRS approach,⁴⁴ which
87 allows us to investigate the association of a variety of psychiatric, cognitive and
88 anthropometric trait PRS with psychotic experiences.

89

90 Psychotic experiences have been associated with urban upbringing.^{21,34,45-47} Studies aimed at
91 identifying what underlies these findings have discovered that psychotic experiences are
92 associated with certain characteristics of the urban environment including neighbourhood
93 adversity and deprivation^{48,49} or air pollution.⁵⁰ However, uncertainty over the interpretation
94 of these associations comes from the recent discovery that aspects of the environment,
95 including urban living, are themselves partially heritable, and the associations between
96 urbanicity and psychotic experiences may not be independent of a genetic predisposition to
97 psychiatric disorders.^{51,52} In the sample used in our study, psychotic experiences have been
98 shown to share some genetic influences with stressful life events and bullying,^{53,54} but more
99 distal environmental factors have not so far been investigated.

100
101
102
103
104
105
106
107
108
109
110

111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145

In light of the evidence that psychotic experiences show a range of transdiagnostic associations in mental health as well as associations with other types of adverse outcomes, we investigated the association of genetic predisposition to a range of psychiatric, cognitive, and anthropometric traits with a six-dimensional representation of psychotic experiences (i.e., paranoia, hallucinations, cognitive disorganization, grandiosity, anhedonia, and negative symptoms).⁸ Furthermore, we investigated the association of psychotic experiences with factors associated with urban living, a well-established environmental risk factor for psychotic disorders. Lastly, we modelled genetic and environmental measures in one model to assess whether the two predictors are independent of each other.⁵⁵ The study was preregistered at <https://osf.io/pts7m/>.

Methods

Sample

Participants included in the current study are part of the Longitudinal Experiences and Perceptions (LEAP) study, which is drawn from the Twins Early Development Study (TEDS).⁵⁶ TEDS is a community sample, which constitutes around 10,000 twin pairs who were born in England and Wales between 1994 and 1996. The recruitment of these participants was designed to obtain a sample of families that are representative of the population in England and Wales.⁵⁷ Of the 10,874 TEDS families that were contacted for inclusion in the LEAP study, 5,059 (47%) twin pairs provided psychotic experiences data (*mean age* = 16.32 years; *s.d.* = 0.68). Exclusions leading to the removal of 316 families included individuals who did not provide consent, had a severe medical disorder, perinatal complication, or had unknown zygosity. Further details on the LEAP study are found elsewhere.⁸ All twins provided written consent to participate in the study at age 16.

Genotyping

Genotyping was done either on the Affymetrix GeneChip 6.0 or Illumina HumanOmniExpressExome (61% of genotyped sample) DNA microarrays.⁵⁷ Genotypes from the two platforms were separately phased using EAGLE2,⁵⁸ and imputed into the Haplotype Reference Consortium (release 1.1). Further details are found elsewhere.⁵⁹ After merging, there were 7,363,646 genotyped or well-imputed SNPs (information score [INFO] > 0.75) available for analysis. After randomly selecting one twin from each pair, we obtained a sample of 4040 (56% female) individuals with genotype data at age 16.

The first 10 ancestry informed principal components were calculated using 39,353 autosomal SNPs with minor allele frequency > 5% and imputation INFO score of 1, selected after pruning to remove SNPs in linkage disequilibrium (r^2) > 0.1 and excluding regions with known high linkage disequilibrium.⁵⁹

After the exclusion of individuals without relevant genotype and phenotype data at age 16, we were left with a sample size of 3590 (complete cases only).

Polygenic risk scores

146
147 We calculated polygenic risk scores for 14 psychiatric, cognitive and anthropomorphic traits
148 (chosen based on their selection in a previous study⁵⁵) to estimate their association with
149 psychotic experience and negative symptoms at age 16. GWAS summary statistics were
150 downloaded for attention deficit hyperactivity disorder (ADHD),⁶⁰ anorexia nervosa,⁶¹
151 anxiety,⁶² autism spectrum disorders (ASD),⁶³ bipolar disorder,⁶⁴ major depressive disorder
152 (MDD),⁶⁵ education years,⁶⁶ extraversion,⁶⁷ intelligence (excluding the 3,414 TEDS
153 participants used in the reported GWAS),⁶⁸ schizophrenia,⁶⁹ subjective well-being (excluding
154 the 2,148 TEDS participants used in the reported GWAS),⁷⁰ neuroticism,⁷⁰ height⁷² and BMI⁷²
155 (**Supplementary Table 1**). We used PRSCs software with default parameter settings to
156 calculate posterior SNP effect sizes under continuous shrinkage priors for each of the GWAS
157 listed above.⁷³ Overlapping SNPs between the selected GWAS summary statistics and
158 7,363,646 genotyped or imputed SNPs available for TEDS data were used to generate the PRS.
159 PRS were then calculated using plink 1.9 as the sum of risk alleles weighted by SNP effect sizes
160 and standardized.

161 162 **Environmental measures**

163
164 Geographical variables from pollution, census, and landcover data were cross-referenced
165 with the participants' postcodes provided in 2008 (mean age of participants = 13, s.d. = 0.58).
166 We obtained 2008 pollution data for routinely collected particles (PM₁₀ and PM_{2.5}), nitrogen
167 dioxide (NO₂) and oxides of nitrogen (NO_x) from resources on annual pollution statistics based
168 on 1x1 km grid squares in the UK (<https://uk-air.defra.gov.uk/data/pcm-data>). These specific
169 pollutants were selected based on relevant literature.^{50,74-79}

170
171 We included measures of population density, urban/rural binary classification, Townsend
172 deprivation index, regional education levels, and regional levels of low social class based on
173 the 2011 census. Census statistics for output areas (OA, i.e., geographical regions created
174 specifically for collecting census data) from the 2011 census were downloaded from
175 Nomisweb (<https://www.nomisweb.co.uk/>; **Supplementary Methods 1**).

176
177 Measures of greenspace were based on land cover maps from the Centre of Ecology and
178 Hydrology (CEH). The percentage land cover types are derived from satellite images and
179 digital cartography. There are 23 classes of land cover that encompasses the entire range of
180 UK habitats. The greenspace variable was created by combining percentages for all 21 of the
181 rural land cover classes. Land cover maps used in the current study were generated in 2007
182 and based on 1x1 km grid squares (<https://www.ceh.ac.uk/services/land-cover-map-2007>).

183
184 Further details on the geographical variables included in the current study are shown in **Table**
185 **1**, including descriptive statistics for the TEDs sample and the means and standard deviations
186 of the age 16 population in England and Wales. T-tests for significant differences between
187 TEDs sample means and age 16 English and Welsh population means were all found to be
188 highly significant with P-values < 0.001 (details in **Supplementary Methods 2**).

189
190 We also included a family socio-economic status (SES) measure, which were collected at first
191 contact (mean age of participants = 18 months). This was calculated based on the mother's

192 and father’s qualification levels and employment status and the mother’s age at birth of the
 193 first child. A higher score corresponds to higher SES.

194
 195 Continuous measures were transformed using the Yeo-Johnson power transformation
 196 method (**Supplementary Methods 3 & Supplementary Figure 1**). Standardisation was applied
 197 after transformation of the data by subtracting the mean and dividing by the standard
 198 deviation. Details on all predictor variables included analysed are found in **Supplementary**
 199 **Table 2**. Correlations between variables are shown in **Supplementary Figure 2**. We removed
 200 variables if they had a correlation coefficient of > 0.8, i.e., PM₁₀ and NO₂.

201
 202 *Table 1: Descriptive statistics for geographical environmental variables in TEDs, compared*
 203 *with the age 16 population in England and Wales.*

Geographical variable	Sample mean [SD] or %	Population mean [SD] or %
PM _{2.5}	10 [1.8]	11 [2]
Oxides of nitrogen (NO _x)	24 [10.5]	28 [14]
Population density (people per hectare)	37 [32]	54 [62]
Urban classification	69%	81%
Townsend deprivation index (TDI)	-1.8 [2.3]	-0.07 [3.5]
Regional % persons aged 16+ with level-4 qualifications (university degree)	30 [13]	26 [13]
Regional % persons ages 16-74 in the lowest social class	19 [12]	26 [15]
Greenspace (% natural land cover)	48 [31]	41 [32]

204
 205 **Outcome measures**

206
 207 Dependent measures included six quantitative subscales of the Specific Psychotic Experiences
 208 Questionnaire (SPEQ) measures at mean age 16.3 (standard deviation = 0.68). These
 209 comprised paranoia, hallucinations, cognitive disorganisation, grandiosity, anhedonia, all self-
 210 reported, and parent-rated negative symptoms. Questionnaire items for each of the
 211 subscales are found in **Supplementary Table 3** Further details on how these scales were
 212 derived and validated are found elsewhere.⁸ Distributions of the psychotic experiences and
 213 negative symptom scales are shown in **Supplementary Figure 3** and correlations between
 214 these six scales in **Supplementary Figure 4**. Each of the psychotic experiences and parent-
 215 rated negative symptom scales were standardized by subtracting the mean and dividing by
 216 the standard deviation.

217
 218 **Statistical analyses**

219
 220 First, we performed partial correlations between each of the six psychotic experiences and
 221 negative symptom scales and each of the 23 genetic and environmental factors adjusted for

222 age and sex using the “ppcor” package in R. PRS included in partial correlations and in the
223 subsequent analyses were adjusted for genetic covariates by regressing each PRS on 10
224 ancestry informed principal components and genotyping chip and using the residuals in
225 analyses.

226

227 Next, we built three models for each of the six psychotic experiences and negative symptom
228 scales. These included separately and jointly modelled genetic and environmental models,
229 one with all the PRS (G), one with all the environmental factors (E) and a joint model of both
230 genetic and environmental factors (GE). Modelling was performed in R using “glmnet” and
231 “caret” packages.

232

233 We used LASSO (least absolute shrinkage and selection operator) regression,⁸⁰ as it reduces
234 overfitting and the sensitivity of the regression coefficients to multi-collinearity. LASSO
235 includes a penalty function that eliminates correlated coefficients, which improves the model
236 in case of collinearity but also conducts automated feature selection. This is especially
237 warranted when modelling multiple geographical measures due to their correlation. To
238 ensure that the age and sex covariates were not removed during the model selection
239 procedure, each of the psychotic experience scales was regressed on age and sex and the
240 residuals were used in the analysis.

241

242 We compared the variance explained R-squared (R^2) of the three models to assess whether
243 environmental measures had independent effects on psychotic experience and negative
244 symptoms scales when adjusting for the genetic effects, and vice versa.⁵⁵ We used a nested
245 cross-validation procedure with an inner and outer loop for model selection and measuring
246 model performance, respectively. The performance of each model was assessed by
247 computing the average R^2 in the hold-out set for each fold of the outer loop (details in
248 **Supplementary Methods 4**). Comparisons between the models were made using the
249 William’s test to calculate the significant difference in the correlation between the observed
250 and predicted values for each of these models (“paired.r” function from the “psych” package
251 in R).^{81,82} Predicted values were the average predictive values across each fold of the outer
252 loop in our nested cross-validation. A Bonferroni adjusted P-value threshold of 0.002 was
253 used (0.05 P-value adjusted for 20 tests, 5 model comparisons for each of the six outcomes).

254

255 For inspection of the model coefficients, we built final models using the whole dataset and
256 model selection was performed using the inner loop cross-validation procedure. Model
257 coefficients were estimated using post-selection inference methods to adjust for the variable
258 selection procedure.⁸³ Variables were reported as significant if they had p-values less than a
259 Bonferroni adjusted threshold of 0.008 (0.05 threshold adjusted for each model for the six
260 psychotic experiences scales). Further details in **Supplementary Methods 4**.

261

262 Results

263

264 Partial correlations between each of the six psychotic experiences and negative symptom
265 scales and the predictor variables, adjusted for age and sex, are shown in **Figure 1**. Cognitive
266 disorganisation and paranoia were both significantly positively correlated with neuroticism
267 (partial correlations of 0.08 and 0.06, respectively) and MDD PRS (0.07 and 0.09, respectively).

268 Cognitive disorganisation and parent-rated negative symptoms were both significantly
269 correlated with genetic and environmental factors relating to education and socio-economic
270 status (**Supplementary Table 5**).

271
272 Parent-rated negative symptoms achieved the highest overall variance explained in the
273 nested cross-validation. The joint modelling of genetic and environmental factors (GE model)
274 explained 2.3% of the variance (**Figure 2; Supplementary Table 6**). Nested comparisons
275 between the GE model and the more parsimonious models showed that the E but not the G
276 model contributed significantly to the variance explained in parent-rated negative symptoms
277 (William's test $p = 0.002$ and 0.05 , respectively; **Supplementary Table 7**). In post-selection
278 inference on the best performing GE model, parent-rated negative symptoms were
279 significantly negatively associated with both a genetic predisposition to education ($\beta = -0.07$;
280 $95\%CI = -0.12$ to -0.04 ; $p = 3 \times 10^{-4}$; **Figure 3; Supplementary Table 8**) and regional educational
281 attainment ($\beta = -0.06$; $95\%CI = -0.09$ to -0.02 ; $p = 0.004$; **Figure 3; Supplementary Table 8**).
282 Additionally, parent-rated negative symptoms were associated with lower SES ($\beta = -0.07$,
283 $95\%CI = -0.10$ to -0.03 ; $p = 0.002$; **Figure 3; Supplementary Table 8**).

284
285 The joint modelling of genetic and environmental factors (GE model) explained 1.3% of the
286 variance in cognitive disorganisation (**Figure 2; Supplementary Table 6**). However, only the G
287 model contributed significantly to the variance explained (William's test $p = 0.006$). In the
288 best performing GE model, the only significant association for cognitive disorganisation was
289 the neuroticism PRS ($\beta = 0.05$; $95\%CI = 0.03$ to 0.08 ; $p = 0.002$; **Figure 3; Supplementary**
290 **Tables 9**).

291
292 For the paranoia scale, it was found that the best performing E model shrunk all the
293 coefficients to zero, meaning no linear combination of any subset of the included
294 environmental factors were useful for predicting paranoia. The GE model explained 1 % of
295 the variance in paranoia (**Figure 2; Supplementary Table 6**). The paranoia scale was
296 associated with higher PRS for MDD ($\beta = 0.07$; $95\%CI = 0.04$ to 0.1 ; $p = 1 \times 10^{-4}$; **Figure 3;**
297 **Supplementary Table 10**, BMI ($\beta = 0.05$; $95\%CI = 0.02$ to 0.08 ; $p = 0.003$; **Figure 3;**
298 **Supplementary Table 10**), and neuroticism ($\beta = 0.05$; $95\%CI = 0.02$ to 0.08 ; $p = 0.008$; **Figure**
299 **3; Supplementary Table 10**).

300
301 For the hallucinations scale, the GE model achieved the highest variance explained (median
302 $R^2 = 0.4\%$; **Figure 2; Supplementary Table 6**). Model comparisons indicated that neither the
303 E nor G model contributed significantly to the variance explained (William's test $p = 0.09$ and
304 0.05 , respectively; **Supplementary Table 7**). Furthermore, we found no specific significant
305 associations with the hallucinations scale (**Figure 3; Supplementary Table 11**). The grandiosity
306 and anhedonia scales had no significant partial correlations for any of the included genetic
307 and environmental factors. Furthermore, in the LASSO models it was found that the best
308 performing model shrunk all the coefficients to zero, meaning no linear combination of any
309 subset of the included predictors were useful for predicting these two outcomes.

310

311 Discussion

312

313 In this study, we investigated the association of sets of polygenic scores and geographical
314 environmental risk factors with six psychotic experiences and negative symptom scales in
315 adolescents. Our findings strengthen previous literature demonstrating psychotic
316 experiences' transdiagnostic associations in mental health through highlighting differential
317 associations between PRS and the psychotic experience subscales, i.e., parent-rated negative
318 symptoms with educational attainment and subjective well-being PRS; cognitive
319 disorganisation with neuroticism PRS; paranoia with BMI, MDD and neuroticism PRS. We also
320 show that family SES and environmental measures of regional educational attainment are
321 associated with parent-rated negative symptoms. These results emphasize the value of
322 studying psychotic experience subscales separately. Furthermore, the results are from
323 models including both genetic and environmental factors, demonstrating that the effects
324 remain significant when adjusting for potential gene-environment correlation.

325
326 The highest variance explained was seen for parent-rated negative symptom scale, which was
327 associated with lower regional educational attainment and lower family socio-economic (SES)
328 measures. This is supported by previous research showing an association of negative
329 symptoms with lower SES and educational attainment.⁸⁶ The PRS for educational attainment
330 was also significantly associated with parent-rated negative symptoms,⁶⁶ suggesting that the
331 PRS influences negative symptoms independently of the educational attainment phenotype.
332 However, nested model comparisons demonstrated that only environmental factors had a
333 significant impact on the model's variance explained.

334
335 The second highest variance explained was seen for cognitive disorganisation, with only
336 genetic factors impacting on the prediction model. Despite the obvious difference of this scale
337 with the disorganisation dimension derived from factor analysis of cases with psychosis,
338 previous evidence suggests that disorganisation has the highest heritability and includes
339 influences that are independent of psychosis liability.⁸⁴ The predominant association for
340 cognitive disorganisation with neuroticism PRS is supported by a recent study demonstrating
341 a shared genetic aetiology between cognitive disorganisation and childhood emotional and
342 behavioural problems.⁸⁵

343
344 Comparing the hallucinations and paranoia scales, within the context of the current study, we
345 notice that hallucinations appear to be affected by both the genetic and environmental
346 factors, while paranoia only by the genetic factors examined. This is consistent with previous
347 findings in the same sample showing that paranoia has the highest heritability.⁸⁷
348 Furthermore, in the same study, shared environment only had a significant influence on
349 hallucinations and negative symptom scales,⁸⁷ which is supported by the association of these
350 scales with regional education levels and SES seen here.

351
352 Contrary to the hypothesis that psychotic experiences are part of a 'psychosis continuum',¹⁰
353 schizophrenia PRS was not associated significantly with any of the psychotic experiences or
354 negative symptom scales, even though it is the most powerful mental health PRS in predicting
355 case-control status. This adds to previous studies on this topic which provided mixed evidence
356 to date on the degree of association between schizophrenia PRS and psychotic experiences
357 or showed stronger association with PRS for other psychiatric disorders.^{23,42,88} However, most
358 of these studies examined older individuals, which may have different psychotic experience
359 profile. Using multiple PRS and the LASSO shrinkage method, our study provides evidence

360 that if psychotic experiences in adolescence are associated with the schizophrenia PRS, the
361 strength of the association is weaker than other disorder-trait genetic associations.⁸⁹

362
363 Here, we did not replicate associations between psychotic experiences and air pollution that
364 have previously been reported.⁵⁰ However, in the previous study, more fine-scale pollution
365 measures were used for multiple frequently visited addresses, which may account for the
366 differences in the reported associations. Another potential explanation for this disparity is the
367 inconsistent definition of psychotic experiences across studies, which include single
368 composite scores and, three- to six- factor dimensions.^{11–15,90} It has also been shown that
369 some co-occurring psychotic experiences, such as cognitive disorganisation and negative
370 symptoms, are associated with more schizophrenia-relevant variables.⁸⁶ Prevalence of
371 psychotic experiences in adolescence, depending on its definition, range from 7% to 95%,^{14,91}
372 which is an indicator of the challenge in defining these outcomes and may be the reason for
373 the inconsistency in study findings.⁴¹ A systematised definition of psychotic experiences in
374 non-clinical populations is necessary to achieve replicability and generalizability of research
375 findings.

376
377 There are several limitations to this study. Geographical variables were linked to the
378 participants' addresses in 2008, around three years prior to the collection of the psychotic
379 experience questionnaires, so did not include the effect of early life or cumulative exposure
380 to these factors. Furthermore, air pollution data was linked to the participants' home
381 addresses, so there was no information on exposures at school or other frequently visited
382 locations, which may lead to exposure misclassification. We also did not include any
383 information on indoor air pollution or smoking status or, more importantly, cannabis use, a
384 known risk factor for psychosis.⁹² Generally, our results are limited by the included genetic
385 and environmental variables as many other unmeasured variables may influence psychotic
386 experiences over and above those that were included. The predictive ability of the PRS
387 included are limited by the power of the training GWAS and we may see different results
388 when PRS explain more of the total genetic variation. Furthermore, the psychotic experience
389 and negative symptom questionnaire measures current symptoms rather than lifetime
390 symptoms, which may limit power to detect effects. TEDS is a twin cohort and results may
391 not generalize, although it has been shown that there is a similar prevalence of psychotic
392 experiences in twins vs non-twins cohorts.⁸⁷ Selection bias may also be present as our cohort
393 had better education, higher SES, and lower exposure to pollutants than the average
394 population of the same age. Finally, with regards to the inherent assumptions of the linear
395 regression method used in the current study, some may be violated, including 1) regression
396 coefficients reflect unconditional relationships only (i.e., no interaction effects between
397 predictors) and 2) normality and homoscedasticity of the residuals was not directly tested.
398 Furthermore, the models developed in the current study were not externally validated, thus
399 it is unknown how well the model coefficients and R^2 will generalize to other samples.

400
401 In conclusion, we find that both genetic risk and geographical environmental factors
402 contribute significantly to the reporting of psychotic experiences in adolescence. The
403 differential predictive ability of specific genetic and environmental risk factors with the six
404 specific psychotic experiences highlights the value of studying these domains separately. We
405 can conclude that cognitive disorganisation and negative symptoms during mid-adolescence
406 were most predicted from the genetic and environmental risk factors examined in our study.

407 Adolescence is known to be a critically vulnerable stage for mental health when most mental
408 health conditions begin. Identifying the genetic and environmental risk factors associated
409 with red flags for poor mental health in adolescence will help to identify suitable targets for
410 early intervention programmes.

411 Conflict of interest

412
413 The Authors have declared that there are no conflicts of interest in relation to the subject of
414 this study.

415 Acknowledgements

416
417 TEDS is supported by a programme grant to Professor Robert Plomin from the UK Medical
418 Research Council (MR/V012878/1 and previously MR/M021475/1), with additional support
419 from the US National Institutes of Health (AG046938). We gratefully acknowledge the
420 ongoing contribution of the participants in the Twins Early Development Study (TEDS) and
421 their families. The research leading to these results has also received funding from the
422 European Research Council under the European Union's Seventh Framework Programme
423 (FP7/2007-2013)/ grant agreement n° 602768. K.R. is supported by a Sir Henry Wellcome
424 Postdoctoral Fellowship. This work contains National Statistics data. Crown copyright and
425 database right (2021). Source: Office for National Statistics licensed under the Open
426 Government Licence v.3.0. LCM2007 and database right NERC 2011; all rights reserved. This
427 work contains data from Defra, UK. This report represents independent research partially
428 funded by the National Institute for Health Research (NIHR) Maudsley Biomedical Research
429 Centre at South London and Maudsley NHS Foundation Trust and King's College London. The
430 views expressed are those of the authors and not necessarily those of the NHS, the NIHR or
431 the Department of Health and Social Care.
432

433 References

- 434
- 435 1. Van Os, J., Hanssen, M., Bijl, R. V. & Ravelli, A. Strauss (1969) revisited: a psychosis
436 continuum in the general population? *Schizophr. Res.* **45**, 11–20 (2000).
 - 437 2. Van Os, J. & Linscott, R. J. Introduction: The Extended Psychosis Phenotype—
438 Relationship With Schizophrenia and With Ultrahigh Risk Status for Psychosis.
439 *Schizophr. Bull.* **38**, 227 (2012).
 - 440 3. Jauhar, S., Johnstone, M. & McKenna, P. J. Schizophrenia. *Lancet* **399**, 473–486
441 (2022).
 - 442 4. Liddle, P. F. The symptoms of chronic schizophrenia. A re-examination of the positive-
443 negative dichotomy. *Br. J. Psychiatry* **151**, 145–151 (1987).
 - 444 5. Andreasen, N. C., Arndt, S., Alliger, R., Miller, D. & Flaum, M. Symptoms of
445 schizophrenia. Methods, meanings, and mechanisms. *Arch. Gen. Psychiatry* **52**, 341–
446 351 (1995).
 - 447 6. Grube, B. S., Bilder, R. M. & Goldman, R. S. Meta-analysis of symptom factors in
448 schizophrenia. *Schizophr. Res.* **31**, 113–120 (1998).
 - 449 7. Shafer, A. & Dazzi, F. Meta-analysis of the positive and Negative Syndrome Scale
450 (PANSS) factor structure. *J. Psychiatr. Res.* **115**, 113–120 (2019).

- 451 8. Ronald, A. *et al.* Characterization of psychotic experiences in adolescence using the
452 specific psychotic experiences questionnaire: Findings from a study of 5000 16-Year-
453 Old Twins. *Schizophr. Bull.* **40**, 868–877 (2014).
- 454 9. Wigman, J. T. W. *et al.* Replication of the five-dimensional structure of positive
455 psychotic experiences in young adulthood. *Psychiatry Res.* **197**, 353–355 (2012).
- 456 10. Van Os, J., Linscott, R. J., Myin-Germeys, I., Delespaul, P. & Krabbendam, L. A
457 systematic review and meta-analysis of the psychosis continuum: Evidence for a
458 psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol.*
459 *Med.* **39**, 179–195 (2009).
- 460 11. Vollema, M. G. & Hoijtink, H. The multidimensionality of self-report schizotypy in a
461 psychiatric population: an analysis using multidimensional Rasch models. *Schizophr.*
462 *Bull.* **26**, 565–575 (2000).
- 463 12. Cyhlarova, E. & Claridge, G. Development of a version of the Schizotypy Traits
464 Questionnaire (STA) for screening children. *Schizophr. Res.* **80**, 253–261 (2005).
- 465 13. Fonseca-Pedrero, E., Paíno-Piñeiro, M., Lemos-Giráldez, S., Villazón-García, Ú. &
466 Muñiz, J. Validation of the Schizotypal Personality Questionnaire-Brief Form in
467 adolescents. *Schizophr. Res.* **111**, 53–60 (2009).
- 468 14. Wigman, J. T. W. *et al.* The Structure of The Extended Psychosis Phenotype in Early
469 Adolescence—A Cross-sample Replication. *Schizophr. Bull.* **37**, 850 (2011).
- 470 15. Ericson, M., Tuvblad, C., Raine, A., Young-Wolff, K. & Baker, L. A. Heritability and
471 longitudinal stability of schizotypal traits during adolescence. *Behav. Genet.* **41**, 499–
472 511 (2011).
- 473 16. McGrath, J. J. *et al.* The bidirectional associations between psychotic experiences and
474 DSM-IV mental disorders. *Am. J. Psychiatry* **173**, 997–1006 (2016).
- 475 17. Bromet, E. J. *et al.* Association Between Psychotic Experiences and Subsequent
476 Suicidal Thoughts and Behaviors: A Cross-National Analysis From the World Health
477 Organization World Mental Health Surveys. *JAMA psychiatry* **74**, 1136–1144 (2017).
- 478 18. Navarro-Mateu, F. *et al.* The association between psychotic experiences and
479 disability: results from the WHO World Mental Health Surveys. *Acta Psychiatr. Scand.*
480 **136**, 74–84 (2017).
- 481 19. Kelleher, I. & Cannon, M. Psychotic-like experiences in the general population:
482 characterizing a high-risk group for psychosis. *Psychol. Med.* **41**, 1–6 (2011).
- 483 20. Morrison, A. P. *et al.* Cognitive therapy for the prevention of psychosis in people at
484 ultra-high risk: Randomised controlled trial. *Br. J. Psychiatry* **185**, 291–297 (2004).
- 485 21. Linscott, R. J. & Van Os, J. An updated and conservative systematic review and meta-
486 analysis of epidemiological evidence on psychotic experiences in children and adults:
487 On the pathway from proneness to persistence to dimensional expression across
488 mental disorders. *Psychological Medicine* **43**, 1133–1149 (2013).
- 489 22. Ronald, A. & Pain, O. A systematic review of genome-wide research on psychotic
490 experiences and negative symptom traits: new revelations and implications for
491 psychiatry. *Hum. Mol. Genet.* **27**, R136–R152 (2018).
- 492 23. Pain, O. *et al.* Genome-wide analysis of adolescent psychotic-like experiences shows
493 genetic overlap with psychiatric disorders. *Am. J. Med. Genet. Part B Neuropsychiatr.*
494 *Genet.* **177**, 416–425 (2018).
- 495 24. McGrath, J., Saha, S., Chant, D. & Welham, J. Schizophrenia: a concise overview of
496 incidence, prevalence, and mortality. *Epidemiol. Rev.* **30**, 67–76 (2008).
- 497 25. Cederlöf, M. *et al.* A longitudinal study of adolescent psychotic experiences and later

- 498 development of substance use disorder and suicidal behavior. *Schizophr. Res.* **181**,
499 13–16 (2017).
- 500 26. Fisher, H. L. *et al.* Specificity of childhood psychotic symptoms for predicting
501 schizophrenia by 38 years of age: A birth cohort study. *Psychol. Med.* **43**, 2077–2086
502 (2013).
- 503 27. Kelleher, I., Cederlöf, M. & Lichtenstein, P. Psychotic experiences as a predictor of the
504 natural course of suicidal ideation: A Swedish cohort study. *World Psychiatry* **13**, 184–
505 188 (2014).
- 506 28. Kelleher, I. *et al.* Psychotic symptoms in adolescence index risk for suicidal behavior:
507 Findings from 2 population-based case-control clinical interview studies. *Arch. Gen.*
508 *Psychiatry* **69**, 1277–1283 (2012).
- 509 29. Werbeloff, N. *et al.* Self-reported attenuated psychotic symptoms as forerunners of
510 severe mental disorders later in life. *Arch. Gen. Psychiatry* **69**, 467–475 (2012).
- 511 30. Healy, C. *et al.* Childhood and adolescent psychotic experiences and risk of mental
512 disorder: a systematic review and meta-analysis. *Psychol. Med.* **49**, 1589–1599
513 (2019).
- 514 31. Hay, D. A. *et al.* Phenotypic and Genetic Analyses of a Short Measure of Psychosis-
515 proneness in a Large-scale Australian Twin Study. *Twin Res.* **4**, 30–40 (2003).
- 516 32. Hur, Y. M., Cherny, S. S. & Sham, P. C. Heritability of hallucinations in adolescent
517 twins. *Psychiatry Res.* **199**, 98–101 (2012).
- 518 33. Linney, Y. M. *et al.* A quantitative genetic analysis of schizotypal personality traits.
519 *Psychol. Med.* **33**, 803–816 (2003).
- 520 34. Polanczyk, G. *et al.* Etiological and clinical features of childhood psychotic symptoms
521 results from a birth cohort. *Arch. Gen. Psychiatry* **67**, 328–338 (2010).
- 522 35. Wigman, J. T. W. *et al.* A twin study of genetic and environmental determinants of
523 abnormal persistence of psychotic experiences in young adulthood. *Am. J. Med.*
524 *Genet. Part B Neuropsychiatr. Genet.* **156**, 546–552 (2011).
- 525 36. Zavos, H. M. S. *et al.* Consistent etiology of severe, frequent psychotic experiences
526 and milder, less frequent manifestations: A twin study of specific psychotic
527 experiences in adolescence. *JAMA Psychiatry* **71**, 1049–1057 (2014).
- 528 37. Sieradzka, D. *et al.* Heritability of Individual Psychotic Experiences Captured by
529 Common Genetic Variants in a Community Sample of Adolescents. *Behav. Genet.* **45**,
530 493–502 (2015).
- 531 38. Taylor, M. J. *et al.* Association of Genetic Risk Factors for Psychiatric Disorders and
532 Traits of These Disorders in a Swedish Population Twin Sample. *JAMA Psychiatry* **76**,
533 280–289 (2019).
- 534 39. Jones, H. J. *et al.* Phenotypic manifestation of genetic risk for schizophrenia during
535 adolescence in the general population. *JAMA Psychiatry* **73**, 221–228 (2016).
- 536 40. Sieradzka, D. *et al.* Are Genetic Risk Factors for Psychosis Also Associated with
537 Dimension-Specific Psychotic Experiences in Adolescence? *PLoS One* **9**, e94398
538 (2014).
- 539 41. Hinterbuchinger, B. & Mossaheb, N. Psychotic-Like Experiences: A Challenge in
540 Definition and Assessment. *Front. Psychiatry* **12**, 270 (2021).
- 541 42. Legge, S. E. *et al.* Association of Genetic Liability to Psychotic Experiences with
542 Neuropsychotic Disorders and Traits. *JAMA Psychiatry* **76**, 1256–1265 (2019).
- 543 43. Barkhuizen, W., Pain, O., Dudbridge, F. & Ronald, A. Genetic overlap between
544 psychotic experiences in the community across age and with psychiatric disorders.

- 545 *Transl. Psychiatry* **10**, (2020).
- 546 44. Krapohl, E. *et al.* Multi-polygenic score approach to trait prediction. *Mol. Psychiatry*
547 **23**, 1368–1374 (2018).
- 548 45. Vassos, E., Pedersen, C. B., Murray, R. M., Collier, D. A. & Lewis, C. M. Meta-Analysis
549 of the Association of Urbanicity With Schizophrenia. *Schizophr. Bull.* **38**, 1118–1123
550 (2012).
- 551 46. Newbury, J. *et al.* Why Are Children in Urban Neighborhoods at Increased Risk for
552 Psychotic Symptoms? Findings from a UK Longitudinal Cohort Study. *Schizophr. Bull.*
553 **42**, 1372–1383 (2016).
- 554 47. Singh, S. P., Winsper, C., Wolke, D. & Bryson, A. School mobility and prospective
555 pathways to psychotic-like symptoms in early adolescence: A prospective birth cohort
556 study. *J. Am. Acad. Child Adolesc. Psychiatry* **53**, 518–527.e1 (2014).
- 557 48. Newbury, J. B. *et al.* In the eye of the beholder: Perceptions of neighborhood
558 adversity and psychotic experiences in adolescence. *Dev. Psychopathol.* **29**, 1823–
559 1837 (2017).
- 560 49. Newbury, J. *et al.* Cumulative Effects of Neighborhood Social Adversity and Personal
561 Crime Victimization on Adolescent Psychotic Experiences. *Schizophr. Bull.* **44**, 348–
562 358 (2018).
- 563 50. Newbury, J. B. *et al.* Association of air pollution exposure with psychotic experiences
564 during adolescence. *JAMA Psychiatry* **76**, 614–623 (2019).
- 565 51. Colodro-Conde, L. *et al.* Association Between Population Density and Genetic Risk for
566 Schizophrenia. *JAMA Psychiatry* **75**, 901 (2018).
- 567 52. Maxwell, J. M., Coleman, J. R. I., Breen, G. & Vassos, E. Association Between Genetic
568 Risk for Psychiatric Disorders and the Probability of Living in Urban Settings. *JAMA*
569 *Psychiatry* (2021). doi:10.1001/JAMAPSYCHIATRY.2021.2983
- 570 53. Shakoor, S. *et al.* Association between stressful life events and psychotic experiences
571 in adolescence: evidence for gene-environment correlations. *Br. J. Psychiatry* **208**,
572 532–538 (2016).
- 573 54. Shakoor, S. *et al.* A shared genetic propensity underlies experiences of bullying
574 victimization in late childhood and self-rated paranoid thinking in adolescence.
575 *Schizophr. Bull.* **41**, 754–763 (2015).
- 576 55. Allegrini, A. G. *et al.* Multivariable G-E interplay in the prediction of educational
577 achievement. *PLOS Genet.* **16**, e1009153 (2020).
- 578 56. Haworth, C. M. A., Davis, O. S. P. & Plomin, R. Twins Early Development Study (TEDS):
579 a genetically sensitive investigation of cognitive and behavioral development from
580 childhood to young adulthood. *Twin Res. Hum. Genet.* **16**, 117–125 (2013).
- 581 57. Rimfeld, K. *et al.* Twins Early Development Study: A Genetically Sensitive Investigation
582 into Behavioral and Cognitive Development from Infancy to Emerging Adulthood.
583 *Twin Res. Hum. Genet.* **22**, 508–513 (2019).
- 584 58. Loh, P. R. *et al.* Reference-based phasing using the Haplotype Reference Consortium
585 panel. *Nat. Genet.* **48**, 1443–1448 (2016).
- 586 59. Selzam, S. *et al.* Evidence for gene-environment correlation in child feeding: Links
587 between common genetic variation for BMI in children and parental feeding
588 practices. *PLoS Genet.* **14**, e1007757 (2018).
- 589 60. Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for
590 attention deficit/hyperactivity disorder. *Nat. Genet.* **51**, 63–75 (2019).
- 591 61. Watson, H. J. *et al.* Genome-wide association study identifies eight risk loci and

- 592 implicates metabo-psychiatric origins for anorexia nervosa. *Nat. Genet.* **51**, 1207–
593 1214 (2019).
- 594 62. Purves, K. L. *et al.* A major role for common genetic variation in anxiety disorders.
595 *Mol. Psychiatry* (2019). doi:10.1038/s41380-019-0559-1
- 596 63. Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum
597 disorder. *Nat. Genet.* **51**, 431–444 (2019).
- 598 64. Stahl, E. A. *et al.* Genome-wide association study identifies 30 loci associated with
599 bipolar disorder. *Nat. Genet.* **51**, 793–803 (2019).
- 600 65. Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and
601 refine the genetic architecture of major depression. *Nat. Genet.* **50**, 668–681 (2018).
- 602 66. Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide
603 association study of educational attainment in 1.1 million individuals. *Nat. Genet.* **50**,
604 1112–1121 (2018).
- 605 67. van den Berg, S. M. *et al.* Meta-analysis of Genome-Wide Association Studies for
606 Extraversion: Findings from the Genetics of Personality Consortium. *Behav. Genet.* **46**,
607 170–182 (2016).
- 608 68. Savage, J. E. *et al.* Genome-wide association meta-analysis in 269,867 individuals
609 identifies new genetic and functional links to intelligence. *Nat. Genet.* **50**, 912–919
610 (2018).
- 611 69. Trubetsky, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in
612 schizophrenia. *Nat.* 2022 6047906 **604**, 502–508 (2022).
- 613 70. Okbay, A. *et al.* Genetic variants associated with subjective well-being, depressive
614 symptoms, and neuroticism identified through genome-wide analyses. *Nat. Genet.*
615 **48**, 624–633 (2016).
- 616 71. Jansen, P. R. *et al.* Genome-wide analysis of insomnia in 1,331,010 individuals
617 identifies new risk loci and functional pathways. *Nat. Genet.* **51**, 394–403 (2019).
- 618 72. Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body
619 mass index in ~700 000 individuals of European ancestry. *Hum. Mol. Genet.* **27**, 3641–
620 3649 (2018).
- 621 73. Ge, T., Chen, C. Y., Ni, Y., Feng, Y. C. A. & Smoller, J. W. Polygenic prediction via
622 Bayesian regression and continuous shrinkage priors. *Nat. Commun.* **10**, 1–10 (2019).
- 623 74. Pun, V. C., Manjourides, J. & Suh, H. Association of Ambient Air Pollution with
624 Depressive and Anxiety Symptoms in Older Adults: Results from the NSHAP Study.
625 *Environ. Health Perspect.* **125**, 342–348 (2017).
- 626 75. Szyzkowicz, M., Kousha, T., Kingsbury, M. & Colman, I. Air Pollution and Emergency
627 Department Visits for Depression: A Multicity Case-Crossover Study. *Environ. Health*
628 *Insights* **10**, (2016).
- 629 76. Klompmaker, J. O. *et al.* Associations of combined exposures to surrounding green,
630 air pollution and traffic noise on mental health. *Environ. Int.* **129**, 525–537 (2019).
- 631 77. Antonsen, S. *et al.* Exposure to air pollution during childhood and risk of developing
632 schizophrenia: a national cohort study. *Lancet Planet. Heal.* **4**, e64–e73 (2020).
- 633 78. Latham, R. M. *et al.* Childhood exposure to ambient air pollution and predicting
634 individual risk of depression onset in UK adolescents. *J. Psychiatr. Res.* **138**, 60–67
635 (2021).
- 636 79. Tzivian, L. *et al.* Effect of long-term outdoor air pollution and noise on cognitive and
637 psychological functions in adults. *International Journal of Hygiene and Environmental*
638 *Health* **218**, 1–11 (2015).

- 639 80. Friedman, J., Hastie, T. & Tibshirani, R. Regularization paths for generalized linear
640 models via coordinate descent. *J. Stat. Softw.* **33**, 1–22 (2010).
- 641 81. Williams, E. J. The Comparison of Regression Variables. *J. R. Stat. Soc. Ser. B* **21**, 396–
642 399 (1959).
- 643 82. Steiger, J. H. Tests for comparing elements of a correlation matrix. *Psychol. Bull.* **87**,
644 245–251 (1980).
- 645 83. Liu, K., Markovic, J. & Tibshirani, R. More powerful post-selection inference, with
646 application to the Lasso. *arXiv* (2018).
- 647 84. Rijdsdijk, F. V., Gottesman, I. I., McGuffin, P. & Cardno, A. G. Heritability estimates for
648 psychotic symptom dimensions in twins with psychotic disorders. *Am. J. Med. Genet.*
649 *B. Neuropsychiatr. Genet.* **156B**, 89–98 (2011).
- 650 85. Shakoor, S., McGuire, P., Cardno, A. G., Freeman, D. & Ronald, A. A twin study
651 exploring the association between childhood emotional and behaviour problems and
652 specific psychotic experiences in a community sample of adolescents. *J. Child Psychol.*
653 *Psychiatry.* **59**, 565 (2018).
- 654 86. Cardno, A. G., Selzam, S., Freeman, D. & Ronald, A. Psychotic-Like Experiences in
655 Adolescence Occurring in Combination or Isolation: Associations with Schizophrenia
656 Risk Factors. *Psychiatr. Res. Clin. Pract.* **3**, 67–75 (2021).
- 657 87. Zavos, H. M. S. *et al.* Consistent etiology of severe, frequent psychotic experiences
658 and milder, less frequent manifestations: A twin study of specific psychotic
659 experiences in adolescence. *JAMA Psychiatry* **71**, 1049–1057 (2014).
- 660 88. García-González, J. *et al.* The effects of polygenic risk for psychiatric disorders and
661 smoking behaviour on psychotic experiences in UK Biobank. *Transl. Psychiatry* **10**, 1–
662 10 (2020).
- 663 89. Ronald, A. Common (Genetic) Links Between Clinics and the Community: New
664 Evidence From a Tourette Syndrome Polygenic Score. *Biol. Psychiatry* **85**, 281–282
665 (2019).
- 666 90. Ronald, A. *et al.* Characterization of psychotic experiences in adolescence using the
667 specific psychotic experiences questionnaire: Findings from a study of 5000 16-Year-
668 Old Twins. *Schizophr. Bull.* **40**, 868–877 (2014).
- 669 91. Kelleher, I. *et al.* Clinicopathological significance of psychotic experiences in non-
670 psychotic young people: Evidence from four population-based studies. *Br. J.*
671 *Psychiatry* **201**, 26–32 (2012).
- 672 92. Marconi, A., Di Forti, M., Lewis, C. M., Murray, R. M. & Vassos, E. Meta-analysis of the
673 Association Between the Level of Cannabis Use and Risk of Psychosis. *Schizophr. Bull.*
674 **42**, 1262–1269 (2016).
- 675
- 676

677 Figure legends

678

679 **Figure 1:** Partial correlations estimated using Pearson's correlation and marked as statistically significant with
680 an asterisk (P -value < 0.0004).

681

682 **Figure 2:** Distribution of hold-out set variance explained (R^2) by the genetic (G), environmental (E), and joint
683 genetic and environmental (GE) models estimated in a nested cross-validation procedure. The median and
684 interquartile range of the R^2 from the 500-fold outer loop of the nested cross-validation are plotted in boxplots
685 with the top and bottom whiskers set at the 97.5th and 2.5th percentile.

686

687 **Figure 3:** Coefficient estimation for genetic and environmental associations with six psychotic experience and
688 negative symptom scales. Best GE model selected via 10-fold cross validation repeated 100 times. Coefficients
689 were estimated using post-selection inference analysis.

690