



This is a repository copy of *Reduced muscle strength (dynapenia) in women with obesity confers a greater risk of falls and fractures in the UK Biobank.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/194547/>

Version: Published Version

Article:

Dowling, L., Cuthbertson, D.J. and Walsh, J.S. (2023) Reduced muscle strength (dynapenia) in women with obesity confers a greater risk of falls and fractures in the UK Biobank. *Obesity*, 31 (2). pp. 496-505. ISSN 1930-7381

<https://doi.org/10.1002/oby.23609>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>




Medfyle Conference Coverage from ObesityWeek® 2022

Discover a new way to catch up with the latest advances in obesity research and care presented at ObesityWeek® 2022.

- Medfyle summaries
- Expert interviews
- Expert presentations
- Posters

[CLICK TO ACCESS NOW](#)



This activity is supported by an educational grant from Lilly

Reduced muscle strength (dynapenia) in women with obesity confers a greater risk of falls and fractures in the UK Biobank

Lisa Dowling¹  | Daniel J. Cuthbertson^{2,3}  | Jennifer S. Walsh¹

¹Oncology and Metabolism, Medical School, The University of Sheffield, Sheffield, UK

²Department of Cardiovascular and Metabolic Medicine, Institute of Life Course and Medical Sciences, The University of Liverpool, Liverpool, UK

³Liverpool University Hospitals NHS Foundation Trust, Liverpool, UK

Correspondence

Lisa Dowling, Oncology and Metabolism, Medical School, University of Sheffield, Beech Hill Road, Sheffield, S10 2RX, UK.
Email: imdowling1@sheffield.ac.uk

Funding information

Medical Research Council, Grant/Award Number: MR/R502182/1; Versus Arthritis

Abstract

Objective: This study aimed to determine the independent effects of obesity and dynapenia on falls risk, areal bone mineral density, and fracture risk (lower extremity or all other fractures).

Methods: A total of 16,147 women (aged 60–82 years) from the UK Biobank were categorized by handgrip strength (HGS; dynapenia status: HGS \leq 21 kg) and body weight (BMI: normal weight, overweight, or obesity). Multiple logistic regression models examined the association among dynapenia and obesity and self-reported falls (previous 12 months), lower extremity fractures, and all other fractures (previous 5 years).

Results: A total of 3793/16,147 women fell, and 1413/15,570 (9.1%) eligible women experienced fall-related fractures. Obesity (odds ratio [OR] 1.25; 95% CI: 1.12–1.38) and dynapenia (OR 0.87; 95% CI: 0.77–0.98) were both independently associated with greater lower extremity fracture risk, independently of areal bone mineral density. However, considering all other fracture sites, obesity conferred protection (OR 0.77; 95% CI: 0.61–0.96), except in those with low HGS, who had an equivalent fracture risk to those with normal weight (OR 1.06; 95% CI: 0.82–1.38).

Conclusions: Dynapenia further increases the increased risk of leg and ankle fractures in obesity and counteracts the protective effects of obesity on fracture risk at all other sites (wrist, arm, hip, spine, other bones).

INTRODUCTION

Obesity is associated with a greater risk of fracture at the ankle, lower leg, and proximal humerus [1–3] but a lower fracture risk at the wrist, hip, and spine [2,4,5]. A more injurious fall type may explain this site-specific fracture risk given the knowledge that obesity is associated with greater areal bone mineral density (BMD) and stronger bone structure, such as with denser cortices and increased trabecular number and thickness [5–7]. A greater understanding of fracture risk in a population with a growing prevalence of obesity could reduce the economic burden of fracture through targeted prevention strategies;

the medical and social cost of fragility fractures alone is estimated at £4.4 billion per year in the UK [8].

The measurement of lean mass in obesity is problematic owing to scaling with body size [9], and it is also becoming increasingly apparent that lean mass may not be the best surrogate of muscle mass [10]. Dynapenia, or low muscle strength, has been recommended in recent reports and definitions of sarcopenia [11,12]. As muscle strength reduces at a greater rate than muscle mass with age, and as this reduction is only partially explained by muscle mass, the concept of dynapenia has been identified as a distinct condition [13]. Dynapenic abdominal obesity refers to the combination of low muscle strength

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Obesity* published by Wiley Periodicals LLC on behalf of The Obesity Society.

and obesity, which is estimated to affect 3.6% to 23.4% of older adults [14–18]. Prevalence is higher in smaller studies [17,18] and is approaching ~10% in large and population-based studies [14–16]. There is a bidirectional relationship between obesity and dynapenia: obesity may exacerbate dynapenia (e.g., through secretion of proinflammatory cytokines), whereas dynapenia may exacerbate weight gain with an impaired ability to undertake physical activity [19].

Independently, dynapenia and obesity are associated with a greater risk of falls [20–22], with reports suggesting a synergistic effect on falls risk [17,18,23]. Dynapenia is associated with a greater risk of all fractures [24], whereas obesity is associated with site-specific fracture risk [1–5] and greater BMD [5,6]. Limited studies have explored the independent effects of dynapenia and obesity on BMD or fracture risk. A recent systematic review [25] reported no difference in lumbar spine BMD (eight studies included; total $n = 9014$) among groups with obesity and sarcopenic obesity; the clinical significance of a statistically cl neck BMD was unclear (six studies included; total $n = 5608$). It should be noted that heterogeneous definitions of sarcopenic obesity (low lean mass, low muscle strength, and a combination of both) were included in this review. In addition, the risk of nonvertebral fracture was similar in people with sarcopenic or dynapenic obesity compared with people with or without obesity, suggesting no cumulative effect. However, these two studies and respective subgroups of people with sarcopenic obesity ($n = 100$ – 128) were small [15,26], and nonvertebral fractures have heterogeneous risk factors and mechanisms, especially in obesity (e.g., obesity increases ankle fracture risk but decreases hip fracture risk).

The aim of this study was to determine whether the fracture pattern observed in women living with obesity is modulated by dynapenia. In order to address this aim, we examined the independent effects of obesity and dynapenia on falls, BMD, and fracture risk. Fractures were divided into lower extremity fractures (obesity-prone) and all other fractures (largely obesity-protective). We hypothesized that dynapenia increased the risk of lower extremity fractures and counteracted the protective effects of obesity on all other fractures, and we speculated that this may relate to differences in either falls risk or BMD.

METHODS

Participants

A description of the UK Biobank has been published elsewhere [27]. Briefly, between 2006 and 2010, the UK Biobank recruited ~500,000 participants (5.5% response rate) aged 40 to 69 years from the general UK population [27]. Participants attended 1 of 22 assessment centers across the UK where they completed a touch-screen questionnaire and an interview, underwent assessment of physical measures, and provided biological samples [27]. The results presented here are from the imaging visit, which began in 2014 and is ongoing. Recent comparison with population-based studies suggests that risk factor associations in the UK Biobank are generalizable [28]. Participants who were male (owing to the low incidence of fractures; <3%); aged <60 years; had body mass

Study Importance

What is already known?

- People with obesity have a greater risk of fractures at the ankle, lower leg, and proximal humerus but a lower risk of other fractures; however, the mechanisms are unclear.
- Low muscle strength (dynapenia) is associated with a greater risk of fracture.
- A recent systematic review found no cumulative effect of obesity and dynapenia on nonvertebral fractures; however, this group of fractures has heterogeneous risk factors and mechanisms, especially in obesity.

What does this study add?

- Dynapenia, irrespective of bone mineral density, further increases the risk of lower extremity (ankle and leg) fractures in women with obesity.
- Dynapenia counteracts the lower risk of all other fractures in women with obesity.

How might these results change the direction of research or the focus of clinical practice?

- These findings highlight the independent risks of these phenotypes (obesity and dynapenia) and contribute to the understanding of the site-specific fracture risk in this group.
- Clinically, these results highlight the importance of increasing physical activity and exercise in weight management programs.

index (BMI) < 18.5 kg/m² (owing to the association between low BMI and fracture risk and the low number of participants within this group); or had missing data for BMI, waist circumference, handgrip strength (HGS), or the question on prior falls were excluded. The UK Biobank was approved by the North West Multicenter Research Ethics Committee, UK. Written informed consent was obtained prior to study entry. A Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) checklist is included (Supporting Information Table S1) [29].

Measures

Falls and fractures

Participants were asked whether they had any falls in the last year. Participants could respond that they had no falls, only one fall, or more than one fall. Participants with one or more falls were classified as “fallers,” and those with no falls or those who could not recall were classified as “non-fallers.” Participants were asked whether they had

fractured or broken any bones in the last 5 years. Participants were subsequently asked whether the fracture had resulted from a fall (i.e., from standing height). Participants who responded that the fracture had resulted from a fall were classified as “injurers.” Participants who responded that the fracture had not resulted from a fall or that they did not know were excluded from the fracture analysis. Participants were asked to report the fractured bone site or sites (i.e., spine, hip, wrist, leg, ankle, arm, or other bones). Based on previous literature [1–5], fracture types were classified as follows: 1) lower extremity (ankle, leg) fractures, sites that are “obesity-prone;” and 2) other fractures (all other sites) or “obesity-protective.” Participants with ankle or leg fractures were preferentially categorized into the lower extremity fracture group irrespective of whether other sites were fractured.

Covariates

Sociodemographic factors included age, self-reported diabetes, alcohol status, and smoking status (prefer not to answer, never, previous, current). Individuals who preferred not to answer or did not know were assumed to never have smoked ($n = 75$), to never have had alcohol ($n = 4$), or to not have diabetes ($n = 49$). Weight was measured using a Tanita BC-418 MA body composition analyzer (Tanita Europe, Amsterdam, the Netherlands) without shoes and heavy outer clothing. Height was measured using a Seca 202 height measure (Seca, Hamburg, Germany). BMI was calculated as weight (kilograms) divided by height (meters squared). Waist circumference was measured at the level of the umbilicus using a Wessex non-stretchable sprung tape. Maximal HGS was measured once on both right and left hands using a Jamar J00105 hydraulic hand dynamometer (Lafayette Instrument Co., Lafayette, Indiana); the maximal measure from either hand was used in this analysis. The UK Biobank protocol for data entry included computer-generated warnings for implausible measurements. Dual-energy x-ray absorptiometry was performed using a GE-Lunar iDXA (GE Healthcare, Madison, Wisconsin). Ethnicity was identified from the available baseline data of the UK Biobank because only 22% had this information at the imaging visit. Owing to large numbers, ethnicity was grouped as White (British, Irish, any other White background) or other ethnic group (prefer not to answer; Asian or Asian British; Chinese; other ethnic group; White and Black Caribbean; White and Black African; White and Asian; any other mixed background; Indian; Pakistani; any other Asian background; Caribbean; African; and those with missing data).

Dynapenia and obesity

Dynapenia was defined as the lowest tertile of HGS (≤ 21 kg). A tertile approach was chosen, similar to others [15,16,30], owing to a lack of consensus and to allow for an exploratory approach. Normal weight, overweight, and obesity were classified according to consensus definitions based on BMI [31].

Statistical analysis

A χ^2 test was used for comparison among categorical variables. Comparison among more than two groups was conducted using either one-way ANOVA with Tukey post hoc test or Kruskal-Wallis test with Dunn post hoc test. Multiple logistic regression was used to examine the association among measures of obesity, dynapenia, or dynapenic obesity and self-reported falls in the past 12 months, lower extremity fractures, and all other fractures in the past 5 years. Regression models were adjusted for age, measurement site, smoking status, self-reported diabetes status, and alcohol status, with results expressed as odds ratios (ORs). Additional adjustment included right femoral neck BMD, HGS, or BMI. Physical activity (moderate activity minutes per day) was initially considered as a confounder for both falls and fractures, but adjustment did not alter results or conclusions, and it was deemed that the analysis was more impaired by missing or incomplete data for physical activity ($n = 2791$). Significance was accepted at $p < 0.05$. Analysis was undertaken using Stata version 16.1 (StataCorp LLC, College Station, Texas).

Individual variable models

Measures of obesity and dynapenia were explored separately as continuous variables. For continuous measures of dynapenia or obesity, z scores were calculated as an individual's result minus the population mean, divided by the population standard deviation.

Dynapenia by BMI category models

We included BMI categories divided by dynapenia status. There were six subgroups: 1) individuals with normal weight and no dynapenia; 2) individuals with dynapenia only; 3) individuals with overweight and without dynapenia; 4) individuals with overweight and dynapenia; 5) individuals with obesity only; and 6) individuals with dynapenic obesity. The group with normal weight without dynapenia was used as the reference group.

RESULTS

Baseline characteristics

Of the 17,175 women at the imaging visit aged 60 years or older, participants who did not answer the question about prior falls ($n = 152$); had a BMI < 18.5 kg/m² ($n = 206$; 2 of whom experienced fractures from falls); or had incomplete measures for HGS ($n = 617$), BMI ($n = 51$), or waist circumference ($n = 2$) were excluded (Supporting Information Figure S1). A total of 16,147 women (aged 60–82 years) were included and categorized according to dynapenia status (lowest tertile of HGS; < 21 kg) and BMI categories (normal weight, overweight, or obesity; Table 1). BMI

TABLE 1 Characteristics of participants according to BMI category and dynapenia status

	Normal weight (BMI = 18.5-24.9 kg/m ²)		Obesity (BMI ≥ 30 kg/m ²)		Overweight (BMI = 25-29.9 kg/m ²)	
	Normal strength, n = 5279 (32.7%)	Dynapenia, n = 2252 (14%)	Normal strength, n = 1812 (11.2%)	Dynapenia, n = 985 (6.1%)	Normal strength, n = 3941 (24.4%)	Dynapenia, n = 1878 (11.6%)
Age (y), median (IQR)	67 (7)	69 (8)	66 (7)	68 (8)	67 (8)	69 (8)
BMI (kg/m ²), median (IQR)	22.7 (2.5)	22.8 (2.5)	32.9 (4.2)	32.6 (4.1)	27.0 (2.3)	27.1 (2.5)
Waist circumference (cm), median (IQR)	75 (10)	75 (9)	99 (12)	99 (12)	86 (10)	86 (10)
HGS (kg), median (IQR)	26 (6)	18 (4)	26 (5)	18 (4)	26 (5)	18 (4)
Ethnicity, n (%)						
White	5191 (98)	2182 (97)	1751 (97)	961 (97)	3857 (98)	1823 (97)
Other ethnic group	88 (2)	70 (2)	61 (3)	24 (3)	84 (2)	56 (3)
BMD ^a , median (IQR)						
L1-L4 BMD (g/cm ²)	1.04 (0.21)	1.02 (0.21)	1.19 (0.25)	1.16 (0.25)	1.11 (0.23)	1.09 (0.23)
L1-L4 T score	-1.19 (1.76)	-1.31 (1.78)	0.04 (2.05)	-0.15 (2.10)	-0.56 (1.89)	-0.72 (1.93)
RFN BMD (g/cm ²)	0.83 (0.14)	0.81 (0.14)	0.93 (0.17)	0.90 (0.16)	0.88 (0.16)	0.86 (0.15)
RFN T score	-1.27 (1.19)	-1.42 (1.21)	-0.44 (1.43)	-0.71 (1.37)	-0.88 (1.31)	-1.03 (1.28)
Physical activity level, median (IQR)						
Mod. activity (min/d) ^b	60 (60)	60 (60)	40 (40)	45 (40)	60 (50)	60 (60)
Mod. activity (d/wk) ^c	5 (4)	5 (4)	4 (4)	3 (3)	4 (3)	4 (5)
Falls, n (%)	1051 (19.91)	576 (25.58)	465 (25.66)	294 (29.85)	887 (22.51)	520 (27.69)
More than one fall, n (%)	253 (4.79)	167 (7.41)	141 (7.78)	86 (8.73)	238 (6.04)	177 (9.42)
All fractures, n	470	291	152	114	361	184
Lower extremity, n	58	47	42	31	85	41
Other fractures, n	412	244	110	83	276	143
Diabetes, n (%)	77 (1.46)	48 (2.13)	185 (10.21)	119 (12.08)	150 (3.81)	111 (5.91)
Smokers, n (%)						
Never	3462 (65.58)	1523 (67.63)	1091 (60.21)	619 (62.84)	2484 (63.03)	1218 (64.86)
Previous	1707 (32.34)	669 (29.71)	683 (36.69)	344 (34.92)	1361 (34.53)	626 (33.33)
Current	110 (2.08)	60 (2.66)	38 (2.10)	22 (2.23)	96 (2.44)	34 (1.81)
Alcohol, n (%)						
Never	181 (3.43)	125 (5.55)	103 (5.68)	67 (6.80)	153 (3.88)	121 (6.44)
Previous	159 (3.01)	80 (3.55)	73 (4.03)	63 (6.40)	132 (3.35)	84 (4.47)
Current	4939 (93.56)	2047 (90.90)	1636 (90.29)	855 (86.80)	3656 (92.77)	1673 (89.08)

Abbreviations: BMD, bone mineral density; HGS, handgrip strength; Mod., moderate; RFN, right femoral neck.

^aMissing measurements for L1-L4 BMD: n = 3439; L1-L4 T score: n = 3445; RFN BMD: n = 3271; and RFN T score: n = 3278.

^bA total of 1334 participants did not know/answer this question, and 2357 participants had missing measurements.

^c578 participants did not know/answer this question.

ranged from 18.5 to 69.6 kg/m². Participants with dynapenia were older than their counterparts in the same BMI category ($p < 0.001$). BMI, waist circumference, and physical activity (minutes per day) were similar between participants with or without dynapenia in the same BMI category. Participants with dynapenic obesity ($p < 0.05$) and dynapenic overweight ($p < 0.01$) had lower BMD and T scores than counterparts in the same BMI category. Participants with normal weight and dynapenia had similar L1-L4 BMD ($p = 0.052$) and T scores ($p = 0.053$) but lower femoral neck BMD and T scores ($p < 0.001$). However, differences were not clinically relevant ($<5\%$ or $<0.5 T$ score) [32]. The proportion of fallers increased with both dynapenia and obesity status ($p < 0.001$).

Falls and fractures

A total of 3793 (23.5%) women fell, of whom 1062 reported more than one fall. Across all BMI categories, dynapenia appeared to




FIGURE 1 Association between dynapenia by BMI categories and falls risk. Adjusted for age, measurement center, smoking status, self-reported “diabetes” status and alcohol status. * $p > 0.05$. HGS, handgrip strength; OR, odds ratio

increase the risk of falling (Figure 1). In the past 5 years, 577 women reported a fracture from causes other than a fall and were excluded from the fracture analysis. Therefore, 1413 of 15,570 (9.1%) eligible women reported a fracture due to a fall. The proportion of injurers according to BMI and HGS tertile is presented in Table 2. Injurers were classified as lower extremity ($n = 295$; 1.9%) or other fractures ($n = 1118$; 7.2%). Of the 295 lower extremity injurers, 58 (19.6%) women also reported other fractures (spine, arm, wrist, other bones), and 9 (3%) reported both ankle and leg fractures. In total, 1572 fractures were recorded. The most frequently reported fractures were “other bones” (577; 37%), followed by wrist (461; 29%) and ankle (230; 15%).

Association between falls, fractures, and measures of muscle strength and obesity

The association between measures of obesity and dynapenia as continuous variables and falls, lower extremity fractures, and all other fracture risk are shown in Table 3. In a multivariable model, obesity, as determined either by greater BMI (OR 1.12; 95% confidence interval [CI]: 1.08-1.16) or waist circumference (OR 1.14; 95% CI: 1.09-1.18), and lower HGS (OR 0.86; 95% CI: 0.83-0.90) had independent effects on falls risk.

Obesity or dynapenia as continuous variables, irrespective of BMD, were associated with a greater risk of lower extremity fractures, with evidence of an independent effect of both obesity and dynapenia (Table 3). In contrast, obesity was protective of all other fractures, whereas low HGS appeared to be associated with a greater risk. These findings remained when analyzed according to BMI category, with an increased risk of lower extremity fracture and lower risk of other fractures in people with overweight or obesity (Tables S2 and S3). Mediation analyses showed that BMI and HGS as z scores partially mediate each other (direct and indirect effects $p < 0.05$) in their associations with both lower extremity or other fracture risk.

TABLE 2 Proportion of participants classified by both tertiles of HGS and BMI who were identified as being in the all other fractures or lower extremity fractures groups

HGS tertile	All other fractures		
	Normal weight, n (%)	Obesity, n (%)	Overweight, n (%)
High (≥ 26 kg)	191 (6.95)	45 (4.74)	142 (6.70)
Medium (21-25.9 kg)	178 (7.68)	55 (6.77)	106 (6.22)
Low (0-20.9 kg)	202 (9.41)	75 (7.81)	124 (6.84)
HGS tertile	Lower extremity fractures		
	Normal weight, n (%)	Obesity, n (%)	Overweight, n (%)
High (≥ 26 kg)	27 (0.98)	25 (2.63)	39 (1.84)
Medium (21-25.9 kg)	31 (1.13)	17 (2.09)	44 (2.58)
Low (0-20.9 kg)	41 (1.49)	30 (3.13)	41 (2.26)

Abbreviations: HGS, handgrip strength.

TABLE 3 Associations, expressed as ORs, among z scores of BMI, waist circumference, HGS, and falls (lower extremity fractures and all other fractures)

	Falls			Lower extremity fractures			All other fractures		
	OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>
Model 1									
BMI	1.12	1.08-1.16	0.000	1.25	1.13-1.39	0.000	0.91	0.85-0.97	0.005
Waist circumference	1.14	1.10-1.18	0.000	1.27	1.14-1.42	0.000	0.96	0.90-1.02	0.190
HGS	0.86	0.83-0.89	0.000	0.86	0.76-0.97	0.015	0.90	0.84-0.96	0.002
Model 1 + HGS z scores									
BMI	1.12	1.08-1.16	0.000	1.25	1.12-1.38	0.000	0.91	0.85-0.97	0.004
Waist circumference	1.14	1.09-1.18	0.000	1.27	1.13-1.41	0.000	0.96	0.90-1.02	0.168
HGS ^a	0.86	0.83-0.90	0.000	0.87	0.77-0.98	0.023	0.90	0.84-0.96	0.001
Model 1 + RFN BMD									
BMI	-	-	-	1.30	1.15-1.46	0.000	0.98	0.91-1.06	0.584
Waist circumference	-	-	-	1.33	1.17-1.52	0.000	1.02	0.95-1.10	0.601
HGS	-	-	-	0.84	0.74-0.97	0.014	0.92	0.86-0.99	0.033

Note: Z score for HGS = 5.71 kg; BMI = 4.56 kg/m²; and waist circumference = 11.57 cm. Model 1: adjusted for age, measurement center, diabetes status, smoking status, and alcohol status.

Abbreviations: BMD, bone mineral density; HGS, handgrip strength; OR, odds ratio; RFN, right femoral neck.

p < 0.05 are in bold.

^aHGS = Model 1 + BMI z scores.

Association between fractures and dynapenia according to BMI category

Compared with the group with normal weight, overweight and obesity were associated with a greater risk of lower extremity fractures (Figure 2A), with the greatest risk estimate seen in people with obesity (OR 2.08; 95% CI: 1.39-3.11) or with dynapenic obesity (OR 2.78; 95% CI: 1.77-4.37). The risk of lower extremity fracture among individuals with both normal weight and dynapenia was also greater than those with normal weight alone (OR 1.69; 95% CI: 1.12-2.54). These findings support our continuous analysis whereby obesity and dynapenia appear to have individual and independent negative associations with lower extremity fracture.

Findings from the categorical analysis of BMI and dynapenia (Figure 2B) confirm our continuous analysis (Table 3) that BMI within the obesity category was protective of all other fractures (OR 0.77; 95% CI: 0.61-0.96), but this effect was negated by the presence of low HGS in which fracture risk was similar to that of the group with normal weight (OR 1.06; 95% CI: 0.82-1.38). Participants who had overweight with (OR 0.90; 95% CI: 0.73-1.12) or without dynapenia (OR 0.88; 95% CI: 0.74-1.04) had a similar risk of all other fractures to women with normal weight. However, individuals with normal BMI and dynapenia had a greater risk of all other fractures (OR 1.29; 95% CI: 1.07-1.55) than those who had normal weight with normal strength. Owing to the large BMI range within this study, additional categorical analysis excluding participants with BMI ≥ 40 kg/m² (*n* = 188) found similar findings for both lower extremity and other fractures (Supporting Information Figure S2).

DISCUSSION

In this large, cross-sectional, retrospective study of women (aged 60-82 years), we have demonstrated the independent effects of obesity and dynapenia on risk of falls. We have shown that dynapenia confers an increased risk of all types of fractures. In contrast, the effects of obesity on fracture risk are site-specific. Lower extremity fracture risk is increased with both obesity (whether measured using BMI or waist circumference) and dynapenia, irrespective of BMD. For all other fractures (wrist, arm, spine, hip, other bones), obesity is associated with a reduced risk of fracture, except when accompanied by dynapenia.

A recent systematic review has shown that individuals with “sarcopenic obesity” (obesity with reduced muscle mass or strength) have a similar risk of nonvertebral fractures to individuals with obesity alone [25], thereby suggesting that there is no cumulative effect of sarcopenia on fracture risk. However, the available studies were relatively small, containing small subgroups [15,26] and failing to consider the site-specific nature of fractures in people with obesity. Our findings are particularly novel because we have used a large cohort and recognized the site-specific fracture risk in people living with obesity. Our results reinforce the evidence regarding the site-specific nature of fractures in people with obesity, with their greater risk of sustaining ankle or leg fractures [1-3] but a reduced risk of other fractures [2,4,5]. However, our findings of the interaction between obesity and dynapenia on fracture risk are novel, namely that obesity increases the risk of lower extremity fractures (ankle, leg) and that dynapenia negates the positive effect of obesity on other fracture risk (wrist, arm, hip, spine, other bones). Consideration of both body weight and



(A) Dynapenia by BMI categories - Lower Extremity Fractures**(B) Dynapenia by BMI categories - All Other Fractures**

FIGURE 2 Association among (A) lower extremity or (B) all other fractures by dynapenia and obesity status. Adjusted for age, measurement center, smoking status, self-reported “diabetes” status, and alcohol status. * $p > 0.05$. HGS, handgrip strength; OR, odds ratio

muscle strength are clearly relevant for determining and stratifying fall and fracture risk.

It is unclear the mechanism by which dynapenia may negate the protective effect of obesity on risk of other fractures. BMD is a well-acknowledged risk factor for fracture [32]; however, measurement of BMD is problematic in people with obesity because of confounding by body mass and bone size [33]. However, other quantitative computed tomography methods also support higher BMD, denser cortices, and increased trabecular thickness and number in people with obesity [7]. Comparison was made between groups with obesity and dynapenic obesity who would both be subject to the same measurement artifact, and, similar to previous studies [34,35], BMD was reduced in the group with dynapenic obesity, although the difference was small and unlikely to translate into a clinically significant

difference in fracture risk [32]. The pathophysiology of dynapenia is multifactorial and can result from malnutrition, reduced physical activity, or comorbid disease [19]; therefore, it is possible that these factors may also contribute to the negating effect of dynapenia, although we were unable to consider these in our analysis. Our findings tentatively suggest that the greater risk of fracture may relate to the cumulative falls risk observed, bearing in mind the differing recall period of both outcomes in the UK Biobank.

The majority of fractures result from falls from a standing height [36]. Separately, obesity and dynapenia are associated with a greater risk of falls [20–22], with growing evidence of a cumulative effect on falls risk, greater than from either phenotype alone [17,18,23]. Our findings support the notion of a greater impact or more injurious fall type in people with obesity [37]. Biomechanically, abdominal obesity

is associated with an anterior shift in the center of mass requiring greater torque for stabilization [38]. Moreover, reduced soft tissue padding at the lower extremities, in combination with a high impact fall, may render these sites more liable to fracture in obesity [1]. Our findings are similar to those of Nielson et al. [39] who demonstrated that obesity was associated with greater risk of lower extremity fracture in 5995 older men (mean follow-up of 7 years), but, in contrast with our results, the effect of obesity was attenuated by reduced physical performance and prior fracture history. Objective measures of lower limb performance were not available in the UK Biobank, and, as such, it requires further consideration as to whether other physical performance measures, especially of the lower extremities, may better relate to lower extremity fractures in women.

Our findings have important clinical implications suggesting that lifestyle interventions that target low muscle strength (either preventing or improving dynapenia), either alone or with measures to reduce excess body weight, could reduce both the risk of falls and fractures in women. Although evidence in younger adults suggests that modest weight reduction achieves greater benefits in balance [40,41], findings in older adults with obesity have shown that weight loss alone exacerbates dynapenia [42]. Furthermore, weight loss without concomitant resistance or weight-bearing exercise can result in loss of BMD, particularly around the hip [43,44], thereby potentially counteracting the reduction in fracture risk derived from a lesser falls risk. Notwithstanding, achieving a healthy weight is a public health priority considering the association of obesity with multiple chronic conditions/comorbidities (e.g., type 2 diabetes, cardiovascular disease, cancer) [31] in addition to falls and fractures risk. Indeed, the greatest gains in physical performance were derived from combined dietary and physical activity interventions [45], further highlighting the relevance of increasing physical activity and functional measurements, particularly of muscle strength, in weight management programs.

The limitations of this study must be acknowledged. First, we were unable to validate the occurrence of fractures because International Classification of Diseases codes were only available for those who were admitted to hospital. Therefore, it is possible that participants may have misclassified or underreported (e.g., spine) their (self-reported) fracture sites. Moreover, individuals with lower extremity fractures also had other fractures; therefore, this may have affected associations. Second, this was a retrospective study, and the inferences drawn should be viewed with caution because dynapenia or obesity may have occurred secondary to a fall or a fracture rather than being a driving pathophysiological factor. Longitudinal analysis was not feasible owing to either an inadequate sample size at both visits or variability in baseline measurement collection time (5-14 years) prior to the outcome (previous 5 years). Third, HGS is a surrogate measure of lower limb strength [13] but can be affected by other factors such as nutrition. Next, we could not replicate this present analysis in male individuals because of the low prevalence of fractures in this group; therefore, further work is required to confirm these findings in male individuals. Finally, the limitations of the UK Biobank in relation to generalizability must be considered, which include its low response rate, limited ethnic diversity, lower overall

prevalence of overweight and obesity than the UK population, and potential healthy volunteer bias (e.g., lower rates of diabetes, narrow socioeconomic backgrounds). However, strengths of the UK Biobank include its significant sample size and the detailed phenotyping available, with evidence supporting the notion that risk factor associations may be generalizable [28].

We have demonstrated that obesity and dynapenia have independent effects on falls risk, but the relationship among obesity, dynapenia, and fracture risk is anatomically site-specific. Dynapenia negates the protective effects of obesity on fractures of the wrist, arm, hip, spine, and other bones, suggesting that greater falls risk or other risk factors, rather than differences in BMD, may explain the negative effect of dynapenia. In contrast, both obesity and dynapenia are associated with a greater risk of lower extremity fractures, independently of BMD, suggesting fall type or fall force may be a mediating factor. These findings require validation in prospective analysis and in men but they may have important clinical implications relating to risk identification and prevention of falls and fractures in a growing population living with overweight and obesity. **O**

ACKNOWLEDGMENTS

This research has been conducted using the UK Biobank Resource under application number 65223.

FUNDING INFORMATION

This work was supported by a studentship from the Medical Research Council (MRC) and Versus Arthritis as part of the MRC Versus Arthritis Centre for Integrated Research into Musculoskeletal Ageing (CIMA; MR/R502182/1). The MRC Versus Arthritis Centre for Integrated Research into Musculoskeletal is a collaboration among the universities of Liverpool, Sheffield, and Newcastle.

CONFLICT OF INTEREST

The authors declared no conflict of interest.

ORCID

Lisa Dowling  <https://orcid.org/0000-0002-8332-358X>

Daniel J. Cuthbertson  <https://orcid.org/0000-0002-6128-0822>

REFERENCES

1. Compston JE, Watts NB, Chapurlat R, et al. Obesity is not protective against fracture in postmenopausal women: GLOW. *Am J Med*. 2011; 124:1043-1050.
2. Prieto-Alhambra D, Premaor MO, Fina Avilés F, et al. The association between fracture and obesity is site-dependent: a population-based study in postmenopausal women. *J Bone Miner Res* 2012;27: 294-300.
3. Johansson H, Kanis JA, Odén A, et al. A meta-analysis of the association of fracture risk and body mass index in women. *J Bone Miner Res* 2014;29:223-233.
4. Premaor MO, Compston JE, Fina Avilés F, et al. The association between fracture site and obesity in men: a population-based cohort study. *J Bone Miner Res* 2013;28:1771-1777.
5. Premaor MO, Pilbrow L, Tonkin C, Parker RA, Compston J. Obesity and fractures in postmenopausal women. *J Bone Miner Res*. 2010;25: 292-297.

6. Beck TJ, Petit MA, Wu G, LeBoff MS, Cauley JA, Chen Z. Does obesity really make the femur stronger? BMD, geometry, and fracture incidence in the women's health initiative-observational study. *J Bone Miner Res.* 2009;24:1369-1379.
7. Evans AL, Paggiosi MA, Eastell R, Walsh JS. Bone density, microstructure and strength in obese and normal weight men and women in younger and older adulthood. *J Bone Miner Res.* 2015;30:920-928.
8. Public Health England. *Falls and Fracture Consensus Statement: Supporting Commissioning for Prevention.* Public Health England; 2017.
9. Linge J, Heymsfield SB, Dahlqvist Leinhard O. On the definition of sarcopenia in the presence of aging and obesity-initial results from UK Biobank. *J Gerontol A Biol Sci Med Sci.* 2020;75:1309-131309-16.
10. Cawthon PM, Orwoll ES, Peters KE, et al. Strong relation between muscle mass determined by D3-creatine dilution, physical performance, and incidence of falls and mobility limitations in a prospective cohort of older men. *J Gerontol A Biol Sci Med Sci.* 2019;74:844-852.
11. Bhasin S, Travison TG, Manini TM, et al. Sarcopenia definition: the position statements of the sarcopenia definition and outcomes consortium. *J Am Geriatr Soc.* 2020;68:1410-1418.
12. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019; 48:16-31.
13. Manini TM, Clark BC. Dynapenia and aging: an update. *J Gerontol A Biol Sci Med Sci.* 2012;67:28-40.
14. Alexandre TDS, Scholes S, Ferreira Santos JL, Duarte YAO, de Oliveira C. The combination of dynapenia and abdominal obesity as a risk factor for worse trajectories of IADL disability among older adults. *Clin Nutr.* 2018;37:2045-2053.
15. Scott D, Chandrasekara SD, Laslett LL, Cicuttini F, Ebeling PR, Jones G. Associations of sarcopenic obesity and dynapenic obesity with bone mineral density and incident fractures over 5-10 years in community-dwelling older adults. *Calcif Tissue Int.* 2016;99:30-42.
16. Rossi AP, Bianchi L, Volpato S, et al. Dynapenic abdominal obesity as a predictor of worsening disability, hospitalization, and mortality in older adults: results from the InCHIANTI study. *J Gerontol A Biol Sci Med Sci* 2017;72:1098-1104.
17. Pereira JC, de Moraes EJ, Neri SGR, Gadelha AB, Lemos RR, Lima RM. Dynapenic abdominal obesity as a risk factor for falls in older women. *Top Geriatr Rehabil.* 2019;35:149-155.
18. Gadelha AB, Neri SGR, Vainshelboim B, Ferreira AP, Lima RM. Dynapenic abdominal obesity and the incidence of falls in older women: a prospective study. *Ageing Clin Exp Res.* 2020;32:1263-1270.
19. Batsis JA, Villareal DT. Sarcopenic obesity in older adults: aetiology, epidemiology and treatment strategies. *Nat Rev Endocrinol.* 2018;14:513-5-513-37.
20. Moreland JD, Richardson JA, Goldsmith CH, Clase CM. Muscle weakness and falls in older adults: a systematic review and meta-analysis. *J Am Geriatr Soc.* 2004;52:1121-1129.
21. Karlsson MK, Ribom E, Nilsson JÅ, et al. Inferior physical performance tests in 10,998 men in the MrOS study is associated with recurrent falls. *Age Ageing.* 2012;41:740-74-740-76.
22. Himes CL, Reynolds SL. Effect of obesity on falls, injury, and disability. *J Am Geriatr Soc.* 2012;60:124-129.
23. de Oliveira Máximo R, Santos JLF, Perracini MR, et al. Abdominal obesity, dynapenia and dynapenic-abdominal obesity as factors associated with falls. *Braz J Phys Ther.* 2019;23:497-505.
24. Harvey NC, Odén A, Orwoll E, et al. Measures of physical performance and muscle strength as predictors of fracture risk independent of FRAX, falls, and aBMD: a meta-analysis of the osteoporotic fractures in men (MrOS) study. *J Bone Miner Res.* 2018;33:2150-215-2150-217.
25. Gandham A, Mesinovic J, Jansons P, et al. Falls, fractures, and areal bone mineral density in older adults with sarcopenic obesity: a systematic review and meta-analysis. *Obes Rev* 2021;22:e13187. doi:10.1111/obr.13187
26. Scott D, Seibel M, Cumming R, et al. Sarcopenic obesity and its temporal associations with changes in bone mineral density, incident falls, and fractures in older men: the Concord Health and Ageing in Men Project. *J Bone Miner Res.* 2017;32:575-583.
27. Sudlow C, Gallacher J, Allen N, et al. UK Biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* 2015;12:e1001779. doi:10.1371/journal.pmed.1001779
28. Batty GD, Gale CR, Kivimäki M, Deary IJ, Bell S. Comparison of risk factor associations in UK Biobank against representative, general population based studies with conventional response rates: prospective cohort study and individual participant meta-analysis. *BMJ* 2020; 368:m131 doi:10.1136/bmj.m131.
29. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Bull World Health Organ.* 2007;85:867-872.
30. Cuthbertson DJ, Bell JA, Ng SY, Kemp GJ, Kivimaki M, Hamer M. Dynapenic obesity and the risk of incident type 2 diabetes: the English Longitudinal Study of Ageing. *Diabet Med.* 2016;33:1052-1059.
31. National Institute for Health and Care Excellence. Obesity: identification, assessment and management. NICE Guideline CG189. Published November 27 2014. Accessed December 2020. <https://www.nice.org.uk/guidance/cg189/resources/obesity-identification-assessment-and-management-pdf-35109821097925>
32. Johnell O, Kanis JA, Oden A, et al. Predictive value of BMD for hip and other fractures. *J Bone Miner Res.* 2005;20:1185-1194.
33. Reid IR. Fat and bone. *Arch Biochem Biophys.* 2010;503:20-27.
34. Buehring B, Hansen KE, Lewis BL, et al. Dymobility syndrome independently increases fracture risk in the osteoporotic fractures in men (MrOS) prospective cohort study. *J Bone Miner Res.* 2018;33:1622-1629.
35. Cawthon PM, Blackwell TL, Cauley J, et al. Evaluation of the usefulness of consensus definitions of sarcopenia in older men: results from the observational osteoporotic fractures in men cohort study. *J Am Geriatr Soc.* 2015;63:2247-2259.
36. Morrison A, Fan T, Sen SS, Weisenfluh L. Epidemiology of falls and osteoporotic fractures: a systematic review. *Clinicoecon Outcomes Res.* 2013;5:9.
37. Boussein ML, Szulc P, Munoz F, Thrall E, Sornay-Rendu E, Delmas PD. Contribution of trochanteric soft tissues to fall force estimates, the factor of risk, and prediction of hip fracture risk. *J Bone Miner Res.* 2007;22:825-831.
38. Corbeil P, Simoneau M, Rancourt D, Tremblay A, Teasdale N. Increased risk for falling associated with obesity: mathematical modeling of postural control. *IEEE Trans Neural Syst Rehabil Eng.* 2001;9:126-136.
39. Nielson CM, Marshall LM, Adams AL, et al. BMI and fracture risk in older men: the osteoporotic fractures in men study (MrOS). *J Bone Miner Res.* 2011;26:496-502.
40. Matrangola SL, Madigan ML. Relative effects of weight loss and strength training on balance recovery. *Med Sci Sports Exerc.* 2009;41:1488-1493.
41. Handrigan G, Hue O, Simoneau M, et al. Weight loss and muscular strength affect static balance control. *Int J Obes (Lond).* 2010;34:936-942.
42. Santanasto AJ, Glynn NW, Newman MA, et al. Impact of weight loss on physical function with changes in strength, muscle mass, and muscle fat infiltration in overweight to moderately obese older adults: a randomized clinical trial. *J Obes* 2011;2011:516576 doi:10.1155/2011/516576.

43. Ensrud KE, Ewing SK, Stone KL, et al. Intentional and unintentional weight loss increase bone loss and hip fracture risk in older women. *J Am Geriatr Soc* 2003;51:1740-1747.
44. Armamento-Villareal R, Aguirre L, Waters DL, Napoli N, Qualls C, Villareal DT. Effect of aerobic or resistance exercise, or both, on bone mineral density and bone metabolism in obese older adults while dieting: a randomized controlled trial. *J Bone Miner Res*. 2020; 35:430-439.
45. Villareal DT, Chode S, Parimi N, et al. Weight loss, exercise, or both and physical function in obese older adults. *N Engl J Med*. 2011;364: 1218-1229.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Dowling L, Cuthbertson DJ, Walsh JS. Reduced muscle strength (dynapenia) in women with obesity confers a greater risk of falls and fractures in the UK Biobank. *Obesity (Silver Spring)*. 2022;1-10. doi:[10.1002/oby.23609](https://doi.org/10.1002/oby.23609)