

This is a repository copy of *Tracking the British agricultural revolution through the isotopic analysis of dated parchment*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/195148/>

Version: Published Version

Article:

Doherty, Sean, Alexander, Michelle Marie orcid.org/0000-0001-8000-3639, Henderson, Stuart et al. (3 more authors) (2023) Tracking the British agricultural revolution through the isotopic analysis of dated parchment. *Scientific Reports*. 61. ISSN 2045-2322

<https://doi.org/10.1038/s41598-022-26013-4>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



OPEN Tracking the British agricultural revolution through the isotopic analysis of dated parchment

Sean P. Doherty^{1,2}✉, Michelle M. Alexander², Stuart Henderson³, Jason Newton⁴, Jonathan Finch² & Matthew J. Collins^{3,5}

Between the sixteenth and nineteenth century, British agriculture underwent a 'revolutionary' transformation. Yet despite over a century of research and the recognised centrality of agricultural developments to industrialisation and population growth, the character or chronology of any 'revolution' during this period remains contentious. Enquiry has been hampered by the fragmented and locally specific nature of historic accounts and the broad dating of early-modern zooarchaeological assemblages. To address this, we conducted stable isotope analysis on 658 legal documents written on sheepskin parchment; a unique biological resource that records the day, month and year of use (AD 1499 to 1969). We find these provide a high temporal resolution analysis of changing agricultural practices and episodes of disease. Most significantly, they suggest that if an 'Agricultural Revolution' occurred in livestock management, it did so from the mid-nineteenth century, in the aftermath of the Napoleonic Wars.

Historians agree the increase in British agricultural productivity between 1550 and 1880 was the result of major structural and technological innovations, notably: enclosure of open fields and commons, the adoption of new field rotation systems, the greater use of soil conditioners and fertilisers, and the improvement of livestock through selective breeding^{1–4}. But this is where the consensus ends. Enduring disagreement as to when and how rapidly these developments occurred persists due to the lack of representative farming data, particularly for livestock^{5–7}, prior to the start of annual agricultural returns in the 1860s. Consequently at least five periods of agricultural 'revolution' have been proposed⁵.

While the analysis of animal bones has provided valuable insights into the pastoral economy^{8–10}, early-modern zooarchaeological assemblages are rarely of a chronological resolution that permits examination on a scale below that of a century. In contrast, historic legal documents written on sheepskin parchment (Fig. 1) record the day, month and year the agreement was signed; a date likely only a few months after the death of the animal from which the parchment was produced¹¹. These documents provide an exceptional resource for high-resolution investigations of animal and land management strategies through stable isotope analysis, a tool used for reconstructing diet¹², discriminating the use of organic and inorganic fertilisers¹³, exploring stocking densities¹⁴ and identifying transhumance¹⁵. Recent analysis demonstrates that parchment is a viable analyte for isotope analysis, recording dietary and physiological signals from the weeks and months before death^{16,17}. Therefore to provide new insight on the timing, extent and drivers of agricultural change, we undertook the isotopic analysis ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of 658 historic legal documents written on sheepskin parchment (Table 1).

Results and discussion

Full $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and elemental composition results are presented in Supplementary Dataset 1, with summary statistics provided in Table 2, and plotted chronologically in Fig. 2. Of the 658 samples analysed, 23 failed to meet collagen quality criteria and were excluded from the analysis.

Procurement of skins. Parchment produced $\delta^{13}\text{C}$ values from -24.3‰ to -15.9‰ (mean: -22.4‰). Except for seven skins from the late 19th ($n=1$) and twentieth century ($n=6$) with values higher than -20‰ , all are consistent with sheep raised in the British Isles grazing on C_3 grasses in agreement with the documents' provenance¹⁸. Those with values above -20‰ indicate the consumption of C_4 plants¹⁹, although with the global

¹Department of Archaeology and History, University of Exeter, Exeter, UK. ²Department of Archaeology, University of York, York, UK. ³Department of Archaeology, University of Cambridge, Cambridge, UK. ⁴NERC National Environmental Isotope Facility, Scottish Universities Environmental Research Centre, East Kilbride, UK. ⁵Globe Institute, University of Copenhagen, Copenhagen, Denmark. ✉email: sean@palaeome.org




Figure 1. Legal deeds. (a) Title deed concerning the ownership of land in Enfield, Middlesex, signed and sealed 15th January 1499 (sample DL035); (b) title deed concerning property in Hanbury Woodend, Staffordshire, signed and sealed 11th August 1728 (sample DL110).

Collection	n		Date range (AD)	Collection information
	Total	Acceptable collagen quality		
Cheshire Records Office	15	15	1786–1813	Artificial collection of deeds concerning property in Cheshire
Campana et al. ⁵⁴	13	9	1730–1830	Artificial collection of various legal documents
Doherty Collection	8	8	1913–1940	Artificial collection of deeds concerning property across England and Wales
Hull History Centre	38	33	1596–1969	Artificial collection of deeds concerning property in the East Riding of Yorkshire
Lee Collection	254	245	1499–1907	Artificial collection of deeds concerning property in England, Wales and Scotland
Lincoln Records Office	9	9	1742–1907	Artificial collection of deeds concerning property in Lincolnshire
Lord Collection	50	48	1582–1893	Title deeds concerning Lower Winskill Farm, Settle, North Yorkshire
Tye Collection	254	253	1650–1904	Artificial collection of deeds concerning property in the City of London. Documents were discarded from the Sun Fire Office, London, company archives
Westminster City Archives	1	1	1707	Title deed from the City of Westminster
Wills Collection	16	14	1652–1790	Title deeds concerning property in Somerset
Total	658	635		

Table 1. Collection information of parchment documents analysed in the study (*n* = number of samples). Collagen quality criteria outlined in the methods section. The artificial collections contain documents of different provenance, while the others have grown organically around a single property.

Period	n	δ ¹³ C			δ ¹⁵ N		
		Mean (‰)	Range (‰)	SD	Mean (‰)	Range (‰)	SD
1499–1599	4	−22.4	0.8	0.4	8.4	3.1	1.3
1600–1624	7	−22.8	1.0	0.4	8.3	4.0	1.4
1625–1649	9	−22.9	1.9	0.5	9.0	5.7	2.1
1650–1674	22	−22.8	1.9	0.5	9.3	5.4	1.5
1675–1699	27	−22.8	2.1	0.4	9.2	7.9	1.6
1700–1724	28	−22.8	1.3	0.3	8.8	4.9	1.4
1725–1749	37	−22.8	1.5	0.4	9.4	6.1	1.4
1750–1774	65	−22.8	2.1	0.4	9.2	7.1	1.3
1775–1799	47	−22.8	1.7	0.4	9.0	5.5	1.2
1800–1824	109	−22.5	2.7	0.6	8.8	5.2	1.0
1825–1849	86	−22.3	3.0	0.5	9.3	7.8	1.2
1850–1874	102	−21.9	3.7	0.7	9.5	5.1	1.2
1875–1899	71	−21.9	4.1	0.8	9.3	5.9	1.0
1900–1924	15	−21.7	4.5	1.2	9.3	4.5	1.3
1925–1969	6	−18.1	4.2	1.5	7.8	0.7	0.3

Table 2. Summary of δ¹³C and δ¹⁵N values by 25-year periods (*n* = number of samples). Due to the small sample sizes, parchment dated between AD 1499–1599 and 1925–1969 are presented as single groups.




Figure 2. Isotope values from historic sheepskin parchment. **(a)** $\delta^{13}\text{C}$ (top) and $\delta^{15}\text{N}$ (bottom) of individual skins plotted against the year the document was signed; **(b)** 5 and 10-year averages of $\delta^{13}\text{C}$ (top) and $\delta^{15}\text{N}$ (bottom) from AD 1650 to AD 1900.

trade in both livestock and fodder in these later decades, it is difficult to say if it is the animal or their feed which are of a non-domestic origin.

Parchment produced $\delta^{15}\text{N}$ values from 5.6‰ to 14.2‰ (mean: 9.1‰), with many higher than those typically observed in terrestrial herbivores. Sheep bone collagen from the same period has yielded $\delta^{15}\text{N}$ values from 3.6 to 10.2‰ (mean: 6‰)^{20,21}, therefore the elevated values in parchment can be attributed in part to inter-tissue isotopic discrimination (mean $\delta^{15}\text{N}_{(\text{skin-bone})} + 1.1\text{‰}$ ¹⁷) and the impact of parchment production (mean $\delta^{15}\text{N}_{(\text{skin-parchment})} + 0.3\text{‰}$ ¹⁶). Additionally, these values likely reflect the finishing of sheep on well manured crops and pasture in an effort to increase their weight before slaughter. Across the English downlands where the parchment making industry was principally located¹¹, crop cultivation was supported by large flocks that were folded on fallow arable fields to enrich the thin soil with their dung²². To provide a concentrated covering, sheep were kept in moveable pens in high densities^{23,24}; a practice with the potential to substantially elevate $\delta^{15}\text{N}$ values in the crops and the consumer tissues^{25,26}. This supplementary feeding over the last 6–8 weeks²⁷ would not be detectable isotopically in bone collagen but would be in skin collagen due to its faster turnover rate^{28,29}. More detailed assessment of regional patterns is not possible as the location the legal agreement concerns, or the stationer through whom the parchment was sold, typically bear no relation to the location the animal was raised¹¹.

It is unlikely these high values reflect the use of young lambskins exhibiting a trophic enrichment from suckling, which elevates $\delta^{15}\text{N}$ in the offspring's tissues around 2‰ above that of the mother³⁰. Lambs were typically weaned around 16–18 weeks old during this period³¹; too young to yield a skin of the $\sim 70 \times 50$ cm typical of legal deeds. Legislation prohibiting the importation of leather gloves ensured lambskins were reserved for these and other expensive goods, while cheaper adult skins were used for parchment—as specified in contemporary manufacturing guides^{32–34}.

Episodes of famine and disease. The unusually high $\delta^{15}\text{N}$ of some skins may indicate they come from sheep which had been in poor health before death. Acute physiological stress due to starvation or illness typically results in an elevation of $\delta^{15}\text{N}$ due to catabolism of the body's own protein resulting in a fractionation characteristic of trophic level increases^{35,36}. While most nutritional stress studies have focussed on inert tissues, recent analysis demonstrates that the rapid turnover of skin collagen enables the recording of short-term stresses isotopically¹⁷.

A series of elevated nitrogen values coincide with an outbreak of 'sheep-rot' during the nineteenth century. Sheep-rot (bacterial pododermatitis) is a highly contagious and painful condition characterised by necrosis of the interdigital skin causing severe lameness, a reluctance to graze, emaciation, and ultimately death if untreated³⁷. Epidemics appeared throughout the early-modern period, but a particularly virulent form spread across Britain between 1828 and 1831, killing an estimated 8 million sheep; a quarter of the national flock³⁸. The twenty-one deeds covering these years have a mean $\delta^{15}\text{N}$ of 9.6‰, comparable with that of all samples, but five of these (five separate documents across three separate collections) have $\delta^{15}\text{N}$ values $> 10.5\text{‰}$, including one with an extraordinary high $\delta^{15}\text{N}$ value of 14.2‰, the highest observed in all samples.

Period	$\delta^{13}\text{C}$		$\delta^{15}\text{N}$	
	U	p	U	p
1650–1674 (22) vs. 1675–1699 (27)	273.5	0.64	265.5	0.53
1675–1699 (27) vs. 1700–1724 (28)	342.5	0.55	346.0	0.59
1700–1724 (28) vs. 1725–1749 (37)	495.0	0.76	422.5	0.21
1725–1749 (37) vs. 1750–1774 (65)	1153.0	0.73	1109.0	0.52
1750–1774 (37) vs. 1775–1799 (47)	1437.0	0.59	1446.5	0.63
1775–1799 (47) vs. 1800–1824 (109)	1815.5	0.004**	2334.5	0.38
1800–1824 (109) vs. 1825–1849 (86)	3695.0	0.01**	3669.0	0.009**
1825–1849 (86) vs. 1850–1874 (102)	2920.0	<0.001***	3875.5	0.17
1850–1874 (102) vs. 1875–1899 (71)	3463.0	0.63	3213.5	0.21
1875–1899 (71) vs. 1900–1924 (15)	519.0	0.88	504.0	0.75
1900–1924 (15) vs. 1925–1969 (6)	2.0	<0.001***	13.0	0.01**

Table 3. Significance of differences between 25-year periods in isotope values (Mann–Whitney *U* test). Number of samples in parentheses. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

It is tempting to interpret these five as victims of the rot which experienced acute inappetence and physiological stress prior to death. The disease can take months to reach an advanced stage and persist in chronic form for a duration more than sufficient to be recorded isotopically in skin³⁹. Contemporary accounts indicate sheep suffering from rot were sent to market as farmers sought to recoup their losses⁴⁰, providing the opportunity for their skins to be sent to a parchment maker. Yet the skins showed no visible sign of malnutrition, such as an impression of the ribs due to excessive thinness³⁴. Dearth and disease were perennial challenges for farmers prior to the advent of pesticides and antibiotics and one must be careful not to simply find historical accounts that support these anomalies. However this and other correlations invite future investigations of parchment for paleopathology studies using additional biomarkers of stress and nutritional deficiency.

Agricultural change after the Napoleonic Wars. $\delta^{13}\text{C}$ values between the early sixteenth century and early nineteenth century display remarkable uniformity, with almost all falling between -24 and -22‰ . When grouped into 25-year periods, no statistically significant difference was observed between quarter centuries (Table 3), with just 0.1‰ variation in mean values between 1600–1624 and 1800–1824 (Table 2). In contrast, $\delta^{15}\text{N}$ values exhibit a high degree of variability, although again with no statistically significant variation in mean values between 25-year periods.

The absence of statistically significant variation across the sixteenth to eighteenth century is surprising considering contemporary agricultural innovations, such as new field rotation systems which replaced fallows with grass leys and nitrogen-fixing root vegetables and legumes⁵. Biometrical data indicates an increase in the size of sheep across these centuries⁸, and as the dimensions of the appendicular skeleton are heavily influenced by diet⁴¹ it has been suggested that this size increase may reflect nutritional improvement⁴². Our results, as well as those of Fisher and Thomas⁴³ for cattle, suggest that early size increases were not accompanied by isotopically detectable changes in environment or diet, and were more likely to have been the result of genetic modifications through the introduction of new stock and selective breeding.

A significant change in isotopic values occurred from the early-nineteenth century, with mean values for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ increasing across the nineteenth century (Table 3). There is considerable ambiguity in interpreting complex data of this nature, with multiple possible readings of this increase. We suggest this reflects the shift in animal and land management in the wake of the Napoleonic Wars (1803–1815). The war created boom conditions for arable farmers, stimulating an extension of the land under cultivation beyond even the limits of the Second World War's reclamation campaign⁴⁴. As hostilities ceased and the continental blockade lifted, the era of high prices was replaced by decades of deflation. With the fresh supply of cultivable land almost exhausted and innovations introduced during the seventeenth and eighteenth centuries now running into diminishing returns⁷, farmers sought new ways to enhance their profits and productivity. Many met the fall in prices with intensive mixed farming, known as 'high farming' or 'high feeding', which achieved high outputs by maintaining large numbers of livestock on imported feeds, producing more manure, which in turn increased soil fertility and ultimately grain yields.

Previous improvements had essentially involved the recycling of materials produced on the farm itself. The essence of what F.M.L. Thompson termed the 'Second Agricultural Revolution' (1815–1880) was that "it broke the closed-circuit system and made the operations of the farmer much more like those of the factory owner"⁴⁴, relying on inputs imported from outside the farm and indeed the country, particularly oil-cake fodder and bonemeal fertiliser^{44,45}. Oil-cakes were a by-product of rape, linseed and cottonseed oil extraction. Virtually all were derived from imported materials, at first Prussia, then Russia from the 1820s, India from the 1850s, and Egypt from the 1860s⁴. American maize was added to oil cakes from the 1840s⁴⁵. Their consumption increased substantially during the nineteenth century, growing from 35,000 tons in the 1820s to 740,000 in the 1880s. Crops grown in continental Europe are characterised by higher $\delta^{13}\text{C}$ values than those grown in the British Isles, a difference which is recorded in the isotopic composition of the animal tissues^{18,46,47}. The consumption of

oil-cakes made from imported crops, including C₄ maize, is likely the key driver in the increasingly higher $\delta^{13}\text{C}$ values seen from 1815 onwards.

The growing use of organic fertilisers such as bonemeal in the nineteenth century is reflected in the significant elevation in mean $\delta^{15}\text{N}$ between 1800–1824 and 1825–1849 (Table 3). Bones were either crushed into half-inch pieces, ground into bonemeal or dissolved in sulphuric acid to be converted into superphosphate⁷. The quantity of domestic and imported bones is estimated to have increased from 30,000 tonnes in the 1810s to 115,000 by the 1880s⁴. Even the bones of those who fell at Waterloo were ground for fertiliser, performing, as one farmer regarded, “the less glorious, but more useful purpose of producing wheat for their brothers at home”⁴⁸. By the mid-eighteenth century, bones were supplemented with the nitrogenous Peruvian seabird guano and Chilean nitrates⁷ as increasingly distant sources of nutrients were used to replenish exhausted British fields. Despite the elevated $\delta^{15}\text{N}$ values of marine bird guano ($> +26\%$)⁴⁹ there is no clear evidence of the 1840–1880s ‘guano craze’ in sheepskins from this period, though contemporary surveys suggest it was infrequently used on the chalk downlands where the parchment making industry was principally located⁵⁰.

Conclusions

The isotopic compositions of sheepskin parchment indicate a substantial change in animal and land management from the second quarter of the nineteenth century. This transformation was driven by renewed and unfettered access to continental and American markets for manures and animal feed, which developed into ‘High Farming’ as agricultural production adapted to more intensive patterns of feeding and manuring. This quantitative data is a new and significant addition to the theory that there were staged episodes within a long ‘agricultural revolution’ much of which was contemporaneous with full industrialisation⁵¹. More broadly, the findings demonstrate that parchment is an extraordinary high-resolution biomolecular archive through which centuries of environmental history, agricultural history and animal health can be explored.

Materials and methods

Sampling. Samples were obtained from 645 individual skins from a total of 477 deeds. Of the deeds with multiple pages, each was of a size ($> 70 \times 50$ cm) to indicate they came from a single animal. Samples (approx. 1×5 mm) were removed from the edge of each skin from an area devoid of any ink, stamp, glue, seal or surface marking to avoid contamination.

Species identification via peptide mass fingerprinting, presented in Doherty et al.⁵², identified 622 (96.4%) as sheep (*Ovis aries*), whilst the remaining 23 (3.6%) could be classified as sheep or goat (*Capra aegagrus hircus*) as separation between the species was not possible due to a lack of diagnostic peptides. Sheepskin was preferentially used over goat or calfskin (*Bos taurus*) for the production of legal documents from the twelfth century onwards in England, Wales and Ireland due to its susceptibility to delaminate when scrapped serving to deter fraudulent textual erasure and modification⁵². No British legal deed has been identified as goatskin through biomolecular analysis^{52,53}. It is therefore highly likely that these 23 skins are from sheep and as such were included in the analysis.

Isotopic data from thirteen eighteenth and nineteenth century British legal deeds reported by Campana et al.⁵⁴ have been included in all statistical analyses, bringing the total to 658. While the original DNA analysis could not determine the species, subsequent identification via peptide mass fingerprinting has identified these as sheep (pers. comm. M.J Collins). Samples were prepared following the same methodology as this study.

Stable isotope analysis. Samples were prepared for stable isotope analysis at BioArCh facilities, Department of Archaeology, University of York, following the methodology outlined in Doherty et al.¹⁶. Lipids were removed via solvent extraction, dichloromethane/methanol (2:1 v/v), by ultrasonication for 1 h, with the supernatant removed and solvent replaced every 15 min. The samples were subsequently demineralised in 0.6 M HCl at 4 °C for 6 h to remove residual calcium carbonate/hydroxide, rinsed with distilled water, and gelatinised in 0.001 M pH 3 HCl at 80 °C for 48 h. The supernatant containing the collagen was filtered (60 μm Ezeze-Filter™, Elkay Laboratories, UK), frozen and freeze-dried.

Prepared collagen (0.9–1.1 mg) was weighed out in duplicate in 5×3.5 mm tin capsules (Elemental Microanalysis, Okehampton, UK) and analysed at the Natural Environment Research Council Life Sciences Mass Spectrometry Facility (NERC LSMSF) in East Kilbride, UK, where isotope ratio determinations were carried out on a ThermoElectron DeltaPlusXP (Thermo Fisher Scientific, Bremen, DE) with an Elementar PyroCUBE elemental analyser (Elementar UK Ltd). Sample data were reported in standard delta per mil notation (δ ‰) relative to V-PDB ($\delta^{13}\text{C}$) and AIR ($\delta^{15}\text{N}$) international standards. Three laboratory reference materials were interspersed within the measurement run to correct for linearity and instrument drift. Each of the laboratory reference materials is checked regularly against international standards USGS40 and USGS41. Following the calculations outlined in Szpak et al.⁵⁵, the total analytical uncertainty was estimated to be $\pm 0.18\%$ for $\delta^{13}\text{C}$ and $\pm 0.20\%$ for $\delta^{15}\text{N}$.

To account for the changing ratio of atmospheric $^{13}\text{C}\text{CO}_2$ to $^{12}\text{C}\text{CO}_2$ due to increased anthropogenic fossil fuel emissions, parchment $\delta^{13}\text{C}$ values were corrected following Dombrosky’s⁵⁶ Suess model to the average atmospheric $\delta^{13}\text{C}$ of AD 1760 (-6.4%), prior to the Industrial Revolution (Supplementary Dataset 1). This atmospheric correction increased the amount of ^{13}C in samples after AD 1807 by 0.1–0.9%.

Collagen quality indicators. Parchment produced an average collagen yield of 69.4% (range 32.4–98.3%), and %C ranged from 37.1 to 47.3% and %N from 12.3 to 16.9%; all consistent with modern parchment and skin¹⁶. Samples produced C:N ratios ranging from 3.1 to 3.5. Skin collagen has a theoretical C:N ratio of 3.11¹⁶ and the elevated ratios in parchment are likely due to collagen hydrolysis during the liming process. Twenty three samples produced C:N ratios > 3.5 and were excluded from the analysis.

Statistical analysis. Statistical testing was carried out using the IBM SPSS Statistics 27 software package. Shapiro-Wilks test for normality indicated $\delta^{13}\text{C}$ data did not conform to a normal distribution ($P < 0.05$), thus the resultant statistical tests were non-parametric in nature. Significance of difference in isotope values between period 25-year period groupings were evaluated with a Mann–Whitney U test.

Ethical compliance. No experiments were performed on live animals, or animal tissue deriving from previous experiments. The authors were not involved in the life or death of animals from which the parchment was made.

Data availability

All data generated in this study are presented in the article and Supplementary Information file.

Received: 25 February 2022; Accepted: 7 December 2022

Published online: 09 January 2023

References

- Jones, E. L. Agriculture and economic growth in England, 1660–1750: Agricultural change. *J. Econ. Hist.* **25**, 1–18 (1965).
- Chambers, J. D. & Mingay, G. E. *The Agricultural Revolution: 1750–1880* (Batsford, 1966).
- Kerridge, E. *The Agricultural Revolution* (Allen & Unwin, Paris, 1967).
- Thompson, F. M. L. The second agricultural revolution, 1815–1880. *Econ. Hist. Rev.* **21**, 62–77 (1968).
- Overton, M. *Agricultural Revolution in England: The Transformation of the Agrarian Economy 1500–1850* (Cambridge University Press, 1996).
- Turner, M. E., Beckett, J. V. & Afton, B. *Farm Production in England 1700–1914* (Oxford University Press, 2001).
- Williamson, T. *The Transformation of Rural England: Farming and the Landscape, 1700–1870* (University of Exeter Press, 2002).
- Davis, J. M. & Beckett, J. V. Animal husbandry and agricultural improvement: The archaeological evidence from animal bones and teeth. *Rural Hist.* **10**, 1–17 (1999).
- Thomas, R. Zooarchaeology, improvement and the British agricultural revolution. *Int. J. Hist. Archaeol.* **9**, 71–88 (2005).
- Sologestoa, I. G. & Albarella, U. (eds) *The Rural World in the Sixteenth Century: Exploring the Archaeology of Innovation in Europe* (Brepols, 2002).
- Doherty, S. P. & Henderson, S. Production of parchment legal deeds in England, 1690–1830. *Hist. Res.* **95**, 575–585 (2022).
- DeNiro, M. J. & Epstein, S. Influence of diet on the distribution of carbon isotopes in animals. *Geochim. Cosmochim. Acta* **42**, 495–506 (1978).
- Bateman, A. S. & Kelly, S. D. Fertilizer nitrogen isotope signatures. *Isotopes Environ. Health Stud.* **43**, 237–247 (2007).
- Szpak, P. Complexities of nitrogen isotope biogeochemistry in plant–soil systems: Implications for the study of ancient agricultural and animal management practices. *Front. Plant Sci.* **5**, 288 (2014).
- Trentacoste, A. *et al.* Heading for the hills? A multi-isotope study of sheep management in first-millennium BC Italy. *J. Archaeol. Sci. Rep.* **29**, 102036 (2020).
- Doherty, S., Alexander, M. M., Vnouček, J., Newton, J. & Collins, M. J. Measuring the impact of parchment production on skin collagen stable isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) values. *STAR Sci. Technol. Archaeol. Res.* **7**, 1–12 (2021).
- Doherty, S. P. *et al.* A modern baseline for the paired isotopic analysis of skin and bone in terrestrial mammals. *R. Soc. Open Sci.* **9**, 211587 (2022).
- Camin, F. *et al.* Multi-element (H, C, N, S) stable isotope characteristics of lamb meat from different European regions. *Anal. Bioanal. Chem.* **389**, 309–320 (2007).
- Kohn, M. J. Carbon isotope compositions of terrestrial C3 plants as indicators of (paleo)ecology and (paleo)climate. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 19691–19695 (2010).
- Clarkson, L. A. The manufacture of leather. In *The Agrarian History of England and Wales, Vol VI, 1750–1820* (ed. Mingay, G. E.) 466–485 (Cambridge University Press, 1989).
- Millard, A. R., Dodd, L. & Nowell, G. *Palace Green Library excavations 2013 (PGL13): Isotopic Studies Project Report* (2015).
- Bleasdale, M. *et al.* Multidisciplinary investigations of the diets of two post-medieval populations from London using stable isotopes and microdebris analysis. *Archaeol. Anthropol. Sci.* **11**, 6161–6181 (2019).
- Thirsk, J. *The English Rural Landscape* (Oxford University Press, 2000).
- Home, T. H. *The Complete Grazier* 5th edn. (Baldwin & Cradock, 1830).
- Ellman, J. On folding sheep. *The Farmers Magazine* 110 (1831).
- Bogaard, A., Heaton, T. H. E., Poulton, P. & Merbach, I. The impact of manuring on nitrogen isotope ratios in cereals: Archaeological implications for reconstruction of diet and crop management practices. *J. Archaeol. Sci.* **34**, 335–343 (2007).
- Schwertl, M., Auerswald, K., Schäufole, R. & Schnyder, H. Carbon and nitrogen stable isotope composition of cattle hair: Ecological fingerprints of production systems?. *Agric. Ecosyst. Environ.* **109**, 153–165 (2005).
- Trow-Smith, R. *A History of British Livestock Husbandry, 1700–1900* (Keegan & Paul, 1959).
- Babraj, J. A. *et al.* Collagen synthesis in human musculoskeletal tissues and skin. *Am. J. Physiol. Endocrinol. Metab.* **289**, E864–E869 (2005).
- El-Harake, W. A. *et al.* Measurement of dermal collagen synthesis rate in vivo in humans. *Am. J. Physiol.* **274**, E586–E591 (1998).
- Fuller, B. T., Fuller, J. L., Harris, D. A. & Hedges, R. E. M. Detection of breastfeeding and weaning in modern human infants with carbon and nitrogen stable isotope ratios. *Am. J. Phys. Anthropol.* **129**, 279–293 (2006).
- Houghton, J. Friday 7th 1694. In *Husbandry and Trade Improvd* (ed. Bradley, R.) 323–330 (Woodman and Lyon, 1728).
- de La Lande, J. & McCauley, G. The art of making parchment. *Art Transl.* **13**, 326–386 (2021).
- Reed, R. *Ancient Skins, Parchments and Leather* (Seminar Press, 1973).
- Mekota, A.-M., Grupe, G., Ufer, S. & Cuntz, U. Serial analysis of stable nitrogen and carbon isotopes in hair: Monitoring starvation and recovery phases of patients suffering from anorexia nervosa. *Rapid Commun. Mass Spectrom.* **20**, 1604–1610 (2006).
- Neuberger, F. M., Jopp, E., Graw, M., Püschel, K. & Grupe, G. Signs of malnutrition and starvation–reconstruction of nutritional life histories by serial isotopic analyses of hair. *Forensic Sci. Int.* **226**, 22–32 (2013).
- Hargis, A. M. & Myers, S. The intergument. In *Pathological Basis of Veterinary Disease* 6th edn (ed. Zachary, J. F.) 1009–1146 (Elsevier, 2017).
- Hansard's Parliamentary Debates, Volume 28, Third Series, comprising the period from 22nd May to 26th June 1835.* (T.C. Hansard, 1835).
- US Department of Agriculture. *Foot Rot of Sheep, FB2206* (USDA, 1972).
- The House of Commons. *Reports from Committees, Volume 8, Part 1 179–288* (Select Committee on Agricultural Distress, 1836).

41. Pálsson, H. & Vergés, J. B. Effects of the plane of nutrition on growth and the development of carcass quality in lambs Part I. The effects of High and Low planes of nutrition at different ages. *J. Agric. Sci.* **42**, 1–92 (1952).
42. Grau-Sologestoa, I. & Albarella, U. The 'long' sixteenth century: A key period of animal husbandry change in England. *Archaeol. Anthropol. Sci.* **11**, 2781–2803 (2019).
43. Fisher, A. & Thomas, R. Isotopic and zooarchaeological investigation of later medieval and post-medieval cattle husbandry at Dudley Castle, West Midlands. *Environ. Archaeol.* **17**, 151–167 (2012).
44. Jones, E. L. *The Development of English Agriculture, 1815–1873* (Palgrave, 1968).
45. Perren, R. *Agriculture in Depression 1870–1940* (Cambridge University Press, 1995).
46. Osorio, M. T., Moloney, A. P., Schmidt, O. & Monahan, F. J. Beef authentication and retrospective dietary verification using stable isotope ratio analysis of bovine muscle and tail hair. *J. Agric. Food Chem.* **59**, 3295–3305 (2011).
47. Zazzo, A. *et al.* Isotopic composition of sheep wool records seasonality of climate and diet. *Rapid Commun. Mass Spectrom.* **29**, 1357–1369 (2015).
48. Anonymous. Supplementary Chapter to 'An Essay on Calcareous Manures' in *The Farmer's Register. Vol. I.* 76–79 (Printed for Edmund Ruffin, 1834).
49. Szpak, P., Longstaffe, F. J., Millaire, J.-F. & White, C. D. Stable isotope biogeochemistry of seabird guano fertilization: Results from growth chamber studies with maize (*Zea mays*). *PLoS One* **7**, e33741 (2012).
50. Caird, S. J. *English Agriculture in 1850–51* (Longman, Brown, Green, and Longmans, 1852).
51. Prothero, R. E. *English Farming, Past and Present* (Longmans, Green, 1912).
52. Doherty, S. P., Henderson, S., Fiddymont, S., Finch, J. & Collins, M. J. Scratching the surface: The use of sheepskin parchment to deter textual erasure in early modern legal deeds. *Herit. Sci.* **9**, 29 (2021).
53. Fiddymont, S. *et al.* Animal origin of 13th-century uterine vellum revealed using noninvasive peptide fingerprinting. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 15066–15071 (2015).
54. Campana, M. G. *et al.* A flock of sheep, goats and cattle: Ancient DNA analysis reveals complexities of historical parchment manufacture. *J. Archaeol. Sci.* **37**, 1317–1325 (2010).
55. Szpak, P., Metcalfe, J. Z. & Macdonald, R. A. Best practices for calibrating and reporting stable isotope measurements in archaeology. *J. Archaeol. Sci. Rep.* **13**, 609–616 (2017).
56. Dombrosky, J. A. ~1000-year ^{13}C Suess correction model for the study of past ecosystems. *Holocene* **30**, 474–478 (2020).

Acknowledgements

We thank Tom Lord, Dave Lee, Ray Tye, Mr and Mrs Wills, Cheshire Records Office, Hull History Centre, Lincoln Records Office and Westminster City Archives for the generous donation of parchment for analysis.

Author contributions

S.P.D. and M.J.C. designed research; S.P.D. and J.N. performed research; S.P.D., M.M.A., S.H., J.F. and M.J.C. analysed data; S.P.D. wrote the paper. M.M.A., S.H., J.N., J.F. and M.J.C. reviewed the manuscript. For the purpose of open access, MJC has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Funding

This research was funded by NERC Life Sciences Mass Spectrometry Facility grant EK259–14/15 to S.P.D. and M.J.C., and ERC Investigator grant no. 295729-CodeX to M.J.C. S.P.D. was supported by the AHRC White Rose College of Arts and Humanities Doctoral Training Partnership (Award No. 1489527). M.J.C. is supported by the Danish National Research Foundation-DNRF128.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-26013-4>.

Correspondence and requests for materials should be addressed to S.P.D.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023