

This is a repository copy of *Oral hygiene and infective endocarditis: a case control study*.

White Rose Research Online URL for this paper: https://eprints.whiterose.ac.uk/196888/

Version: Supplemental Material

Article:

Lockhart, P., Chu, V., Zhao, J. et al. (12 more authors) (2023) Oral hygiene and infective endocarditis: a case control study. Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology, 136 (3). pp. 333-342. ISSN 2212-4403

https://doi.org/10.1016/j.oooo.2023.02.020

Article available under the terms of the CC-BY-NC-ND licence (https://creativecommons.org/licenses/by-nc-nd/4.0/).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Table 4. Blood Culture Results from IE Cases

Documented and Common Oral Species Group *	Number Reported
Streptococcus mitis†	6
Streptococcus sanguis (sanguinis)	4
viridans group streptococci	3
Streptococcus anginosus	2
Streptococcus oralis†	2
Streptococcus mutans	2
Streptococcus salivarius	2
Streptococcus gordonii	1
Haemophilus parainfluenzae	1
Granulicatella adiacens	1
Cardiobacterium spp.	1
Lactobacillus spp.	1
Rothia dentocariosa	1
	Subtotal 27 (44%)
Indeterminate [‡]	
Streptococcus not otherwise specified (NOS)	2
	Subtotal 2(3%)
Unlikely Oral Species [§]	
Staphylococcus aureus	11
Enterococcus faecalis	10
Group B Streptococcus/Streptococcus agalactiae	5
Enterococcus species	1
Staphylococcus saprophyticus	1
Staphylococcus epidermidis	1

Corynebacterium diphteriae	1
Corynebacterium	1
Group A beta hemolytic streptococcus	1
Streptococcus pneumoniae	1
	Subtotal 33 (53%)
	Total 62 (100%)

^{*} Documented oral species indicates that these are common to the mouth, and some are particularly common in dental plaque, or a gingival or periodontal pocket.³²

[†]Three cases had both *S. mitis* and *S. oralis* in their blood culture.

[‡] Indeterminate indicates that the bacteria could not be categorized without further identification.

[§] Unlikely oral species indicates that these bacteria are generally recognized as being from non-oral anatomical regions. However, some of these species have been cultured from specific regions of the mouth (*e.g.*, S. *epidermidis*^{32,33} and *Enterococcus faecalis*).³⁵