



This is a repository copy of *Climatic drivers of silicon accumulation in a model grass operate in low- but not high-silicon soils.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/197305/>

Version: Published Version

---

**Article:**

Johnson, S.N. [orcid.org/0000-0002-8388-8345](https://orcid.org/0000-0002-8388-8345), Vandegeer, R.K., Borevitz, J.O. et al. (3 more authors) (2023) Climatic drivers of silicon accumulation in a model grass operate in low- but not high-silicon soils. *Plants*, 12 (5). 995. ISSN 2223-7747

<https://doi.org/10.3390/plants12050995>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

Communication

# Climatic Drivers of Silicon Accumulation in a Model Grass Operate in Low- but Not High-Silicon Soils

Scott N. Johnson<sup>1,\*</sup>, Rebecca K. Vandeger<sup>1</sup>, Justin O. Borevitz<sup>2</sup>, Susan E. Hartley<sup>3</sup>, David T. Tissue<sup>1</sup>  
and Casey R. Hall<sup>1</sup><sup>1</sup> Hawkesbury Institute for the Environment, Western Sydney University, Penrith, NSW 2751, Australia<sup>2</sup> Research School of Biology, Australian National University, Canberra, ACT 2601, Australia<sup>3</sup> School of Biosciences, University of Sheffield, Sheffield S10 2TN, UK

\* Correspondence: scott.johnson@westernsydney.edu.au; Tel.: +61-024-570-1374

**Abstract:** Grasses are hyper-accumulators of silicon (Si), which is known to alleviate diverse environmental stresses, prompting speculation that Si accumulation evolved in response to unfavourable climatic conditions, including seasonally arid environments. We conducted a common garden experiment using 57 accessions of the model grass *Brachypodium distachyon*, sourced from different Mediterranean locations, to test relationships between Si accumulation and 19 bioclimatic variables. Plants were grown in soil with either low or high (Si supplemented) levels of bioavailable Si. Si accumulation was negatively correlated with temperature variables (annual mean diurnal temperature range, temperature seasonality, annual temperature range) and precipitation seasonality. Si accumulation was positively correlated with precipitation variables (annual precipitation, precipitation of the driest month and quarter, and precipitation of the warmest quarter). These relationships, however, were only observed in low-Si soils and not in Si-supplemented soils. Our hypothesis that accessions of *B. distachyon* from seasonally arid conditions have higher Si accumulation was not supported. On the contrary, higher temperatures and lower precipitation regimes were associated with lower Si accumulation. These relationships were decoupled in high-Si soils. These exploratory results suggest that geographical origin and prevailing climatic conditions may play a role in predicting patterns of Si accumulation in grasses.



**Citation:** Johnson, S.N.; Vandeger, R.K.; Borevitz, J.O.; Hartley, S.E.; Tissue, D.T.; Hall, C.R. Climatic Drivers of Silicon Accumulation in a Model Grass Operate in Low- but Not High-Silicon Soils. *Plants* **2023**, *12*, 995. <https://doi.org/10.3390/plants12050995>

Academic Editor: Ivana Puglisi

Received: 17 January 2023

Revised: 14 February 2023

Accepted: 15 February 2023

Published: 22 February 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Keywords:** false brome; phytoliths; precipitation; rainfall; seasonality; silica

## 1. Introduction

Most grasses take up and accumulate silicon (Si) more than any other inorganic constituent [1]. Si accumulation is increasingly recognised as playing an important functional role in plant ecology [2], particularly in terms of its role in relieving the adverse effects of environmental stress [3,4]. These include drought, heat, and salinity as well as biotic agents of plant stress (e.g., pathogens and herbivorous insects) [5]. The mechanisms for stress alleviation vary and are often incompletely understood, but usually involve silicification of tissues, which can have direct impacts (e.g., phytoliths that inhibit herbivory) [6] or indirect impacts on plant chemical and physiological processes (e.g., stomatal closure reducing transpiration losses) [7].

Given that Si has been reported to play an important role in the alleviation of drought stress in plants [8], a pervasive idea is that Si accumulation may be highly beneficial and therefore common in seasonally arid environments [9]. While Si accumulation may alleviate the impacts of drought, water limitation may in itself reduce the ability of plants to take up Si from the soil since passive uptake is highly dependent on the transpiration stream and stomatal conductance [10]. Water limitation has mostly been reported to limit Si accumulation in experimental situations, although some plant species appear to be able to maintain Si accumulation under drought/osmotic stress [11–13]. Maintaining, or even increasing, Si accumulation under these conditions may be achievable via

increased evapotranspiration under hot and dry conditions [14]. Si accumulation is also attained via active, energy-dependent transport of Si [15–18], which may be less affected by water limitation.

Brightly, et al. [14] sampled 36 grass species grown under different temperature and irrigation regimes and generally found positive effects of high temperatures and water limitation on Si accumulation, although they concluded the relationships with water limitation were weak. In natural systems, Johnston, et al. [19] did not detect any patterns between precipitation and Si concentrations of grasses from Northern America. Using rain exclusion shelters, drought conditions increased Si concentrations in some grass species while decreasing them in others [20]. Sampling across a rainfall gradient, Katz, et al. [21] reported that Si (phytoliths) concentrations were positively correlated with water availability in a grass (*Avena sterilis*), but this was not seen in Asteraceae species. Sustained drought reduced Si accumulation in a barley (*Hordeum vulgare*) landrace and cultivar, whereas intermittent drought had little impact [22]. In short, consistent patterns between climatic variables and grass silicification have proved elusive. Moreover, climatic impacts on Si accumulation may be affected by the availability of bioavailable Si in the soil [23].

The objective of this study was to investigate potential relationships between geographic patterns of temperature and precipitation and Si accumulation in the model grass *Brachypodium distachyon*. Using a common garden experiment, we compared the Si accumulation capacity of 57 accessions with 19 climatic variables (Table S1) [24] for the Mediterranean region of origin (Figure 1). We hypothesised that accessions from seasonally arid regions have a greater capacity for Si uptake, which would be reflected in positive correlations with relevant temperature and precipitation climatic variables (e.g., precipitation seasonality). We predicted that these relationships may be altered by bioavailable Si in the soil.




**Figure 1.** Map showing the location of the 57 *Brachypodium distachyon* accessions used in this research.

## 2. Results


When plants were grown under low-Si conditions, leaf Si concentrations were negatively correlated with four climatic variables (Table 1): annual mean diurnal temperature range (Figure 2a), temperature seasonality (Figure 2b), annual temperature range (Figure 2c), and precipitation seasonality (Figure 2d). In contrast, leaf Si concentrations were positively correlated with precipitation variables (Table 1): annual precipitation (Figure 3a), precipitation of the driest month (Figure 3b), precipitation of the driest quarter (Figure 3c), and precipitation of the warmest quarter (Figure 3d). When plants were grown under high-Si soil conditions, none of the climatic variables were correlated with leaf Si concentrations (Table S2).

**Table 1.** Correlation test results for leaf Si concentrations and climatic variables for plants grown under low-Si soil conditions. *p* values corrected to account for multiple testing using the Benjamini and Hochberg False Discovery method [25].

Statistical Significance	Correlation	Climatic Variable	Figure	$r_s$	<i>p</i>
Significant	Negative	Bio 2—Annual Mean Diurnal Temperature Range	2a	−0.262	< 0.001
		Bio 4—Temperature Seasonality (Standard Deviation)	2b	−0.188	0.010
		Bio 7—Annual Temperature Range	2c	−0.210	0.010
		Bio 15—Precipitation Seasonality	2d	−0.156	0.033
	Positive	Bio 12—Annual Precipitation	3a	0.186	0.019
		Bio 14—Precipitation of Driest Month	3b	0.187	0.014
		Bio 17—Precipitation of Driest Quarter	3c	0.184	0.011
		Bio 18—Precipitation of Warmest Quarter	3d	0.192	0.011
Non-significant	No correlation	Bio 1—Annual Mean Temperature		0.073	0.319
		Bio 3—Isothermality		0.068	0.338
		Bio 5—Max Temperature of Warmest Month		−0.116	0.119
		Bio 6—Min Temperature of Coldest Month		0.139	0.061
		Bio 8—Mean Temperature of Wettest Quarter		0.048	0.506
		Bio 9—Mean Temperature of Driest Quarter		−0.078	0.303
		Bio 10—Mean Temperature of Warmest Quarter		−0.001	0.999
		Bio 11—Mean Temperature of Coldest Quarter		0.125	0.095
		Bio 13—Precipitation of Wettest Month		0.102	0.174
		Bio 16—Precipitation of Wettest Quarter		0.080	0.310
		Bio 19—Precipitation of Coldest Quarter		0.039	0.575




**Figure 2.** Significant negative correlations between leaf Si concentrations and (a) annual mean diurnal temperature range, (b) temperature seasonality, (c) annual temperature range, and (d) precipitation seasonality. *N* = 247. Regression line (solid line) fitted with 95% confidence displayed as dashed lines. See Table 1 for statistical analysis.




**Figure 3.** Significant positive correlations between leaf Si concentrations and (a) annual precipitation, (b) precipitation of the driest month, (c) precipitation of the driest quarter, and (d) precipitation of the warmest month. N = 247. Regression line (solid line) fitted with 95% confidence displayed as dashed lines. See Table 1 for statistical analysis.

Plants grown in high-Si soil had significantly higher concentrations of leaf Si compared with those grown in low-Si ( $2.41\% \pm 0.04$  and  $1.32\% \pm 0.02$ , respectively, mean  $\pm$  standard error given) ( $H_1 = 320.2, p < 0.001$ ). There was, however, a significant negative correlation between this increase in Si concentrations, either in terms of absolute (Figure 4a) or percentage (Figure 4b) increase, and the Si concentrations observed when growing in low- (untreated) Si soil. In other words, accessions achieving relatively greater Si accumulation under low-Si conditions tended to show comparatively smaller increases in Si when the soil was supplemented with Si.



**Figure 4.** Cont.



**Figure 4.** The relationship between shoot Si concentrations when growing in low-Si soil and the (a) absolute and (b) relative (%) increase in Si accumulation when accessions were grown in high (Si-supplemented) soil. Accessions indicated (1) Gaz2, (2) Adi6, (3) Adi15, (4) Bd23–01, (5) BdTR3a, (6) Mig3, (7) BdTR11a, (8) BdTR9a, (9) Bd3–1 RR 2009, (10) BdTR1k, (11) BdTR1d, (12) BdTR12a, (13) Adi1, (14) BdTR13c, (15) Bd3–1 Bd3–2, (16) BdTR11g, (17) Bd21–3 ANU, (18) Adi18, (19) Bd21, (20) Adi8, (21) BdTR10f, (22) BdTR1j, (23) Bd18–1, (24) Kah6, (25) Koz2, (26) Koz3, (27) Gaz6, (28) Kah4, (29) Kah5, (30) BdTR3s, (31) BdTR10i, (32) Bdis05–07, (33) BdTR2a, (34) BdTR10c, (35) Bd21–3 INRA, (36) Bdis28–08, (37) Adi16, (38) Gal1, (39) Cas2, (40) BdTR9f, (41) BdTR5m, (42) Bdis05–09, (43) BdTR3b, (44) BdTR2j, (45) Gaz3, (46) BdTR2p, (47) Bdis25–04, (48) Bdis22–06, (49) Bd25–01, (50) Bdis31–02, (51) Bd21 RR 2009, (52) Bdis31–01, (53) Bdis22–01, (54) Bdis32–02, (55) Bdis25–10, (56) BdTR1c, and (57) Pal2032.

### 3. Discussion

Accessions of *B. distachyon* with the greatest capacity for Si accumulation under low-Si conditions tended to originate in regions with lower temperatures and higher precipitation patterns. It seems possible that water limitation hinders Si accumulation, which is potentially exacerbated by warmer temperatures. This refutes our hypothesis that accessions from seasonally arid regions would have the highest concentrations of Si. On the contrary, a climatic variable associated with seasonal aridity (precipitation seasonality) was negatively correlated with leaf Si concentrations. This relationship was decoupled, however, when we increased soil Si availability.

#### 3.1. Interpreting the Bioclimatic Variables

Many of the bioclimatic variables used in this study will correlate with one another given their similar nature; hence, we must be cautious about overinterpreting statistically significant correlations. What can we infer if we consider the strongest negative and positive correlations (*annual mean diurnal temperature range* and *precipitation of the warmest month*, respectively)? The first correlation reflects the range between minimum and maximum temperatures with larger values reflecting more extreme temperatures; the second correlation reflects the level of precipitation in the warmest three months (Table S1). Greater fluctuations in temperature might hinder Si accumulation because higher temperatures may exacerbate the effects of water limitation via evaporation from the soil surface [26], whereas colder temperatures may slow down Si absorption [27]. In contrast, higher amounts of precipitation during the active growing season (i.e., the warmest quarter) are likely to increase plant growth and nutrient uptake in general. This may explain why *B. distachyon* accessions from regions with low temperature fluctuations and ample precipitation during

growth periods tended to accumulate more Si than accessions from regions with highly fluctuating temperatures and dry summers. Overall, increasing the availability of soil Si seems to mask this underlying capacity.

### 3.2. Linkages between Temperature, Precipitation, and Bioavailable Si in the Soil

Bioavailable Si derives from weathering and desilication of primary silicate minerals (lithogenic Si) or the deposition of silicified plant tissues (e.g., phytoliths) into the soil via plant litter (biogenic or phytogenic Si) [28]. In grassland and forest systems, the predominant source of silicic acid for plants is biogenic Si. In essence, Si is cycled from plant litter through the biogenic pool of soil Si and back to plants [29]. While higher temperatures generally increase Si release from phytoliths in the soil [30], this is inhibited in dry soils because soil water is required for dissolution [31]. Blecker, et al. [32] showed that soil biogenic Si (e.g., phytoliths) decreased with precipitation, proposing that faster soil phytolith turnover rates under higher rainfall regimes underpinned this relationship.

Quigley, et al. [23] also found that amorphous Si (pre-weathered silicates) in the soil was negatively correlated with precipitation. Moreover, they reported that dissolved Si (bioavailable silicic acid;  $H_4SiO_4$ ) was also negatively correlated with precipitation, namely, concentrations were lowest under wetter conditions. If this was due to increased Si uptake by plants (i.e., depletion of dissolved soil Si), one would assume a negative relationship between dissolved soil Si and leaf Si concentrations, but, in fact, they reported a positive correlation between the two [23]. Considering multiple plant taxa, Cooke and Leishman [33] determined that soil Si availability does not generally correlate with Si accumulation, suggesting that other factors may influence these relationships.

### 3.3. Accessions Differ in Their Capacity for Si Uptake

We observed that accessions achieving relatively higher Si accumulation under low-Si conditions attained smaller increases in Si, when supplemented with Si, compared with accessions with lower Si concentrations that showed stronger patterns of Si uptake when it was made more available. A possible interpretation of this result is that some *B. distachyon* accessions are well adapted to Si accumulation under low bioavailable Si conditions, whereas other accessions show greater plasticity to Si availability in the soil. When compost was supplemented with Si (without drought exposure), Thorne, et al. [13] observed that ‘high-Si accumulator’ wheat (*Triticum aestivum*) landraces achieved similar increases in Si accumulation to ‘low-Si accumulator’ landraces. In contrast, when grown hydroponically (without osmotic stress) the ‘high-Si accumulators’ showed disproportionately larger increases in shoot Si concentrations compared with ‘low-Si accumulators’ [13]. This suggests that the soil itself places some constraints on the plasticity of Si accumulation, which is also affected by factors such as root exudation [34].

### 3.4. Do Trade-Offs Operate in Water Limited Environments?

Many studies show that nutrient limitation promotes Si accumulation [35,36]. Quigley, et al. [35] also observed a leaf-level trade-off between Si and carbon, which was stronger at arid sites than mesic sites. They concluded that plants used Si rather than invest in C-based leaf construction and C fixation, which is relatively costly when water is limited. Si accumulation may not necessarily be higher in plants growing in arid conditions because of physiological constraints; therefore, it remains possible that Si could play a relatively more important role in arid regions. Si-reinforced cell walls may prevent cell collapse during water shortages, which could be beneficial in seasonally arid environments [9].

## 4. Materials and Methods

### 4.1. Common Garden Experiment

Fifty-seven *Brachypodium distachyon* accessions (Table S3) were used. These were mainly obtained from the Australian National University (Canberra, ACT, Australia) with additional accessions from Western Sydney University (Richmond, NSW, Australia).

Germplasm was originally collected from its native Mediterranean region [37] (Figure 1). The accessions were grown from seed in soil recovered from the Hawkesbury Campus of Western Sydney University (33.6138° S, 150.7500° E). Seeds were initially stratified for five days at 4 °C before planting. Plants were grown in a 50:50 composite of sandy loam and loam soil with low bioavailable Si levels of 16.00 mg kg<sup>-1</sup> (extracted with CaCl<sub>2</sub> [38]); see Table S4 for further soil chemistry. The experiment involved growing half of the plants in 'low-Si soil' (–Si) and half in 'high-Si soil' (+Si). To obtain high-Si soil, half of the pots were watered with a solution consisting of potassium silicate (Agsil32, PQ Australia, SA, Australia) at a concentration of 2 mM (SiO<sub>2</sub> equivalent) and adjusted to pH 7 using HCl. The low-Si soil pots were watered with a control solution containing 1.6 mM KCl to balance additional K<sup>+</sup> and Cl<sup>–</sup> in the Si-supplemented soils; see [39] for details.

Plants were grown in a growth cabinet (Climatron, Thermoline Scientific) maintained at 22/16 °C on an 8:16 h light:dark cycle. The humidity was kept at ca. 60% and light levels were ca. 150 μmol m<sup>-2</sup> s<sup>-1</sup>. Plants were watered with 25 mL of either –Si or +Si solutions (low- and high-Si soil, respectively) every 10 days for the first 10 weeks of the experiment and then every 7 days for the remaining four weeks of the experiment for a total of 14 weeks. Plants were harvested, oven dried, and analysed for leaf Si concentrations at this point. There was a median of five replicates for each accession under low- and high-Si soil conditions at the end of the experiment. We obtained 19 climatic variables (temperature and precipitation; Table S1) for each accession from the WorldClim database [24] to examine any relationships between leaf Si accumulation and climatic variables (see Table S1).

#### 4.2. Leaf Si Accumulation

We determined leaf Si concentrations using ca. 100 mg of ground shoot tissue placed into small receptacles (Malvern PANalytical, Malvern UK), which were then analysed with an X-ray fluorescence spectrometer (Epsilon 3×, PANalytical) using the procedure and certified reference material described in Hiltbold et al. [40]. This method was based on the methodology developed by Reidinger, et al. [41].

#### 4.3. Statistical Analysis

Relationships between leaf concentrations of Si and the 19 climatic variables (Table S1) were examined for all plants grown under low- and high-Si soil conditions separately using Spearman's correlation tests. To account for repeated tests, which can increase false-positive (significance) rates, we adjusted *p* values using the Benjamini and Hochberg False Discovery Method [25]. Differences in leaf Si concentrations between plants grown under low- and high-Si soil conditions were examined with a Kruskal–Wallis test. Relationships between Si concentrations in plants grown in low-Si soil and corresponding increases in Si concentrations when growing in high-Si soil were conducted using mean values for each accession. All analyses were conducted in Genstat version 21 (VSN International Ltd., Hemel Hempstead, UK).

### 5. Conclusions

This exploratory study suggests that *B. distachyon* accessions from regions with higher precipitation during the warmest months have a greater capacity for Si accumulation than those from drier regions. Warmer temperatures and adequate rainfall are prerequisites for the dissolution of phytoliths [28,29], which may go some way to explaining this result. Given the important functional role of Si accumulation in many grasses, understanding how climatic factors affect patterns of Si uptake and accumulation may become increasingly important as the planet experiences environmental change at an unprecedented rate.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/plants12050995/s1>, Table S1: The 19 climatic variables used in the study. Definitions and interpretations summarises descriptions outlined in O'Donnell and Ignizio (2012); Table S2: Correlation test results for foliar Si concentrations and climatic variables for plants grown in high Si soils (+Si); Table S3: Collection locations for the *Brachypodium distachyon*

accessions; Table S4: Characteristics of the homogenised soil used in the common garden experiment. References [42–44] are cited in the supplementary file.

**Author Contributions:** Conceptualisation, S.N.J., S.E.H. and D.T.T.; methodology, S.N.J., C.R.H., R.K.V. and J.O.B.; formal analysis, S.N.J.; investigation, C.R.H. and R.K.V.; data curation, C.R.H.; writing—original draft preparation, S.N.J.; writing—review and editing, S.N.J., C.R.H., S.E.H., R.K.V., D.T.T. and J.O.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by Australian Research Council, grant numbers FT170100342 and DP170102278.

**Data Availability Statement:** The dataset associated with this research will be made available via the FigShare repository [details to follow pending review].

**Acknowledgments:** We thank members of the J.O.B. group for discussions and assistance collecting the plant germplasm, Jeff Powell and Rohan Riley for providing extra germplasm, and Andrew Gherlenda for assistance with the growth cabinets.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Epstein, E. Silicon. *Annu. Rev. Plant Physiol. Plant Molec. Biol.* **1999**, *50*, 641–664. [[CrossRef](#)]
- Cooke, J.; Leishman, M.R. Is plant ecology more siliceous than we realise? *Trends Plant Sci.* **2011**, *16*, 61–68. [[CrossRef](#)] [[PubMed](#)]
- Cooke, J.; Leishman, M.R. Consistent alleviation of abiotic stress with silicon addition: A meta-analysis. *Funct. Ecol.* **2016**, *30*, 1340–1357. [[CrossRef](#)]
- Coskun, D.; Deshmukh, R.; Sonah, H.; Menzies, J.G.; Reynolds, O.L.; Ma, J.F.; Kronzucker, H.J.; Bélanger, R.R. The controversies of silicon’s role in plant biology. *New Phytol.* **2019**, *221*, 67–85. [[CrossRef](#)]
- Debona, D.; Rodrigues, F.A.; Datnoff, L.E. Silicon’s role in abiotic and biotic plant stresses. *Annu. Rev. Phytopathol.* **2017**, *55*, 85–107. [[CrossRef](#)]
- Hall, C.R.; Dagg, V.; Waterman, J.M.; Johnson, S.N. Silicon alters leaf surface morphology and suppresses insect herbivory in a model grass species. *Plants* **2020**, *9*, 643. [[CrossRef](#)]
- Vandeger, R.K.; Zhao, C.; Cibils-Stewart, X.; Wuhner, R.; Hall, C.R.; Hartley, S.E.; Tissue, D.T.; Johnson, S.N. Silicon deposition on guard cells increases stomatal sensitivity as mediated by K<sup>+</sup> efflux and consequently reduces stomatal conductance. *Physiol. Plant.* **2021**, *171*, 358–370. [[CrossRef](#)]
- Thorne, S.J.; Hartley, S.E.; Maathuis, F.J.M. Is silicon a panacea for alleviating drought and salt stress in crops? *Front. Plant Sci.* **2020**, *11*, 1221. [[CrossRef](#)] [[PubMed](#)]
- Coughenour, M.B. Graminoid responses to grazing by large herbivores—Adaptations, exaptations, and interacting processes. *Ann. Missouri Bot. Gard.* **1985**, *72*, 852–863. [[CrossRef](#)]
- McLarnon, E.; McQueen-Mason, S.; Lenk, I.; Hartley, S.E. Evidence for active uptake and deposition of Si-based defenses in tall fescue. *Front. Plant Sci.* **2017**, *8*, 1199. [[CrossRef](#)]
- Quigley, K.M.; Anderson, T.M. Leaf silica concentration in Serengeti grasses increases with watering but not clipping: Insights from a common garden study and literature review. *Front. Plant Sci.* **2014**, *5*, 568. [[CrossRef](#)]
- Ryalls, J.M.W.; Moore, B.D.; Johnson, S.N. Silicon uptake by a pasture grass experiencing simulated grazing is greatest under elevated precipitation. *BMC Ecol.* **2018**, *18*, 53. [[CrossRef](#)] [[PubMed](#)]
- Thorne, S.J.; Hartley, S.E.; Maathuis, F.J.M. The effect of silicon on osmotic and drought stress tolerance in wheat landraces. *Plants* **2021**, *10*, 814. [[CrossRef](#)] [[PubMed](#)]
- Brightly, W.H.; Hartley, S.E.; Osborne, C.P.; Simpson, K.J.; Strömberg, C.A.E. High silicon concentrations in grasses are linked to environmental conditions and not associated with C<sub>4</sub> photosynthesis. *Glob. Change Biol.* **2020**, *26*, 7128–7143. [[CrossRef](#)] [[PubMed](#)]
- Ma, J.F.; Yamaji, N. Silicon uptake and accumulation in higher plants. *Trends Plant Sci.* **2006**, *11*, 392–397. [[CrossRef](#)] [[PubMed](#)]
- Ma, J.F.; Tamai, K.; Yamaji, N.; Mitani, N.; Konishi, S.; Katsuhara, M.; Ishiguro, M.; Murata, Y.; Yano, M. A silicon transporter in rice. *Nature* **2006**, *440*, 688–691. [[CrossRef](#)]
- Deshmukh, R.; Bélanger, R.R. Molecular evolution of aquaporins and silicon influx in plants. *Funct. Ecol.* **2016**, *30*, 1277–1285. [[CrossRef](#)]
- Ma, J.F.; Yamaji, N. A cooperative system of silicon transport in plants. *Trends Plant Sci.* **2015**, *20*, 435–442. [[CrossRef](#)]
- Johnston, A.; Bezeau, L.M.; Smoliak, S. Variation in silica content of range grasses. *Can. J. Plant Sci.* **1967**, *47*, 65–71. [[CrossRef](#)]
- Power, S.A.; Barnett, K.L.; Ochoa-Huesco, R.; Facey, S.L.; Gibson-Forty, E.V.-J.; Hartley, S.E.; Nielsen, U.N.; Tissue, D.T.; Johnson, S.N. DRI-Grass: A new experimental platform for addressing grassland ecosystem responses to future precipitation scenarios in south-east Australia. *Front. Plant Sci.* **2016**, *7*, 1373. [[CrossRef](#)]
- Katz, O.; Lev-Yadun, S.; Bar, P. Plasticity and variability in the patterns of phytolith formation in Asteraceae species along a large rainfall gradient in Israel. *Flora* **2013**, *208*, 438–444. [[CrossRef](#)]

22. Wade, R.N.; Donaldson, S.M.; Karley, A.J.; Johnson, S.N.; Hartley, S.E. Uptake of silicon in barley under contrasting drought regimes. *Plant Soil* **2022**, *477*, 69–81. [[CrossRef](#)]
23. Quigley, K.M.; Donati, G.L.; Anderson, T.M. Variation in the soil ‘silicon landscape’ explains plant silica accumulation across environmental gradients in Serengeti. *Plant Soil* **2017**, *410*, 217–229. [[CrossRef](#)]
24. Hijmans, R.J.; Cameron, S.E.; Parra, J.L.; Jones, P.G.; Jarvis, A. Very high resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* **2005**, *25*, 1965–1978. [[CrossRef](#)]
25. Benjamini, Y.; Hochberg, Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **1995**, *57*, 289–300. [[CrossRef](#)]
26. An, N.; Tang, C.-S.; Xu, S.-K.; Gong, X.-P.; Shi, B.; Inyang, H. Effects of soil characteristics on moisture evaporation. *Eng. Geol.* **2018**, *239*, 126–135. [[CrossRef](#)]
27. Barber, D.A.; Shone, M.G.T. The absorption of silica from aqueous solutions by plants. *J. Exp. Bot.* **1966**, *17*, 569–578. [[CrossRef](#)]
28. Haynes, R.J. A contemporary overview of silicon availability in agricultural soils. *J. Plant Nutr. Soil Sci.* **2014**, *177*, 831–844. [[CrossRef](#)]
29. Haynes, R.J. The nature of biogenic Si and its potential role in Si supply in agricultural soils. *Agric. Ecosyst. Environ.* **2017**, *245*, 100–111. [[CrossRef](#)]
30. Frayssé, F.; Pokrovsky, O.S.; Schott, J.; Meunier, J.-D. Surface properties, solubility and dissolution kinetics of bamboo phytoliths. *Geochim. Cosmochim. Acta* **2006**, *70*, 1939–1951. [[CrossRef](#)]
31. Cabanes, D.; Shahaack-Gross, R. Understanding fossil phytolith preservation: The role of partial dissolution in paleoecology and archaeology. *PLoS ONE* **2015**, *10*, e0125532. [[CrossRef](#)] [[PubMed](#)]
32. Blecker, S.W.; McCulley, R.L.; Chadwick, O.A.; Kelly, E.F. Biologic cycling of silica across a grassland bioclimate sequence. *Global Biogeochem. Cycles* **2006**, *20*, GB3023. [[CrossRef](#)]
33. Cooke, J.; Leishman, M.R. Tradeoffs between foliar silicon and carbon-based defences: Evidence from vegetation communities of contrasting soil types. *Oikos* **2012**, *121*, 2052–2060. [[CrossRef](#)]
34. de Tombeur, F.; Cornelis, J.T.; Lambers, H. Silicon mobilisation by root-released carboxylates. *Trends Plant Sci.* **2021**, *26*, 1116–1125. [[CrossRef](#)] [[PubMed](#)]
35. Quigley, K.M.; Griffith, D.M.; Donati, G.L.; Anderson, T.M. Soil nutrients and precipitation are major drivers of global patterns of grass leaf silicification. *Ecology* **2020**, *101*, e03006. [[CrossRef](#)]
36. Johnson, S.N.; Waterman, J.M.; Wuhrer, R.; Rowe, R.C.; Hall, C.R.; Cibils-Stewart, X. Siliceous and non-nutritious: Nitrogen limitation increases anti-herbivore silicon defences in a model grass. *J. Ecol.* **2021**, *109*, 3767–3778. [[CrossRef](#)]
37. Catalán, P.; Müller, J.; Hasterok, R.; Jenkins, G.; Mur, L.A.J.; Langdon, T.; Betekhtin, A.; Siwinska, D.; Pimentel, M.; López-Alvarez, D. Evolution and taxonomic split of the model grass *Brachypodium distachyon*. *Ann. Bot.* **2012**, *109*, 385–405. [[CrossRef](#)] [[PubMed](#)]
38. Sauer, D.; Saccone, L.; Conley, D.J.; Herrmann, L.; Sommer, M. Review of methodologies for extracting plant-available and amorphous Si from soils and aquatic sediments. *Biogeochemistry* **2006**, *80*, 89–108. [[CrossRef](#)]
39. Hall, C.R.; Mikhael, M.; Hartley, S.E.; Johnson, S.N. Elevated atmospheric CO<sub>2</sub> suppresses jasmonate and silicon-based defences without affecting herbivores. *Funct. Ecol.* **2020**, *34*, 993–1002. [[CrossRef](#)]
40. Hiltbold, I.; Demarta, L.; Johnson, S.N.; Moore, B.D.; Power, S.A.; Mitchell, C. Silicon and other essential element composition in roots using X-ray fluorescence spectroscopy: A high throughput approach. In Proceedings of the Invertebrate Ecology of Australasian Grasslands, Hawkesbury, NSW, Australia, 4–7 April 2016; pp. 191–196.
41. Reidinger, S.; Ramsey, M.H.; Hartley, S.E. Rapid and accurate analyses of silicon and phosphorus in plants using a portable X-ray fluorescence spectrometer. *New Phytol.* **2012**, *195*, 699–706. [[CrossRef](#)]
42. Morgan, M.F. Chemical soil diagnosis by the universal soil testing system. *Conn. Agric. Exp. Stn. Circ.* **1941**, *450*, 579–628.
43. O’Donnell, M.S.; Ignizio, D.A. *Bioclimatic Predictors for Supporting Ecological Applications in the Conterminous United States*; U.S. Geological Survey Data Series 691; U.S. Geological Survey: Reston, VA, USA, 2012.
44. Rayment, G.E.; Lyons, D.J. *Soil Chemical Methods—Australasia*; CSIRO Publishing: Collingwood, VIC, Australia, 2011.

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.