



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/197742/>

Version: Published Version

Article:

Adhikari, G., Carlin, N., Choi, J. et al. (2023) An induced annual modulation signature in COSINE-100 data by DAMA/LIBRA's analysis method. *Scientific Reports*, 13. 4676. ISSN: 2045-2322

<https://doi.org/10.1038/s41598-023-31688-4>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



OPEN

An induced annual modulation signature in COSINE-100 data by DAMA/LIBRA's analysis method

Govinda Adhikari¹, Nelson Carlin², JaeJin Choi^{3,4}, Seonho Choi³, Anthony Ezeribe⁵, Luis Eduardo França², Chang Hyon Ha⁶, In Sik Hahn^{7,8,9}, Sophia J. Hollick¹, Eunju Jeon⁴, Jay Hyun Jo¹, Han Wool Joo³, Woon Gu Kang⁴, Matthew Kauer¹⁰, Bongho Kim⁴, Hongjoo Kim¹¹, Jinyoung Kim⁶, Kyungwon Kim⁴, SungHyun Kim⁴, Sun Kee Kim³, Won Kyung Kim^{4,9}, Yeongduk Kim^{4,9,12}, Yong-Hamb Kim^{4,9,13}, Young Ju Ko⁴, Doo Hyok Lee¹¹, Eun Kyung Lee⁴, Hyunseok Lee^{4,9}, Hyun Su Lee^{4,9}✉, Hye Young Lee⁴, In Soo Lee⁴, Jaison Lee⁴, Jooyoung Lee¹¹, Moo Hyun Lee^{4,9}, Seo Hyun Lee^{4,9}, Seung Mok Lee³, Yu Jin Lee⁶, Douglas Leonard⁴, Bruno B. Manzato², Reina H. Maruyama¹, Robert J. Neal⁵, James A. Nikkel¹, Stephen L. Olsen⁴, Byung Ju Park^{4,9}, Hyang Kyu Park¹⁴, Hyeonseo Park¹³, Kangsoon Park⁴, Se Dong Park¹¹, Ricardo L. C. Pitta², Hafizh Prihtyadi⁴✉, Sejin Ra⁴, Carsten Rott^{15,16}, Keon Ah Shin⁴, Andrew Scarff⁵, Neil J. C. Spooner⁵, William G. Thompson¹, Liang Yang¹⁷ & Gyun Ho Yu^{4,15}

The DAMA/LIBRA collaboration has reported the observation of an annual modulation in the event rate that has been attributed to dark matter interactions over the last two decades. However, even though tremendous efforts to detect similar dark matter interactions were pursued, no definitive evidence has been observed to corroborate the DAMA/LIBRA signal. Many studies assuming various dark matter models have attempted to reconcile DAMA/LIBRA's modulation signals and null results from other experiments, however no clear conclusion can be drawn. Apart from the dark matter hypothesis, several studies have examined the possibility that the modulation is induced by variations in detector's environment or their specific analysis methods. In particular, a recent study presents a possible cause of the annual modulation from an analysis method adopted by the DAMA/LIBRA experiment in which the observed annual modulation could be reproduced by a slowly varying time-dependent background. Here, we study the COSINE-100 data using an analysis method similar to the one adopted by the DAMA/LIBRA experiment and observe a significant annual modulation, however the modulation phase is almost opposite to that of the DAMA/LIBRA data. Assuming the same background composition for COSINE-100 and DAMA/LIBRA, simulated experiments for the DAMA/LIBRA without dark matter signals also provide significant annual modulation with an amplitude similar to DAMA/LIBRA with opposite phase. Even though this observation does not directly explain

¹Department of Physics and Wright Laboratory, Yale University, New Haven, CT 06520, USA. ²Physics Institute, University of São Paulo, São Paulo 05508-090, Brazil. ³Department of Physics and Astronomy, Seoul National University, Seoul 08826, Republic of Korea. ⁴Center for Underground Physics, Institute for Basic Science (IBS), Daejeon 34126, Republic of Korea. ⁵Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, UK. ⁶Department of Physics, Chung-Ang University, Seoul 06973, Republic of Korea. ⁷Department of Science Education, Ewha Womans University, Seoul 03760, Republic of Korea. ⁸Center for Exotic Nuclear Studies, Institute for Basic Science (IBS), Daejeon 34126, Republic of Korea. ⁹IBS School, University of Science and Technology (UST), Daejeon 34113, Republic of Korea. ¹⁰Department of Physics and Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin-Madison, Madison, WI 53706, USA. ¹¹Department of Physics, Kyungpook National University, Daegu 41566, Republic of Korea. ¹²Department of Physics, Sejong University, Seoul 05006, Republic of Korea. ¹³Korea Research Institute of Standards and Science, Daejeon 34113, Republic of Korea. ¹⁴Department of Accelerator Science, Korea University, Sejong 30019, Republic of Korea. ¹⁵Department of Physics, Sungkyunkwan University, Suwon 16419, Republic of Korea. ¹⁶Department of Physics and Astronomy, University of Utah, Salt Lake City, UT 84112, USA. ¹⁷Department of Physics, University of California, San Diego, La Jolla, CA 92093, USA. ✉email: hyunsulee@ibs.re.kr; hafizhp@ibs.re.kr

the DAMA/LIBRA results directly, this interesting phenomenon motivates more profound studies of the time-dependent DAMA/LIBRA background data.

Several groups have attempted to develop experiments aiming at reproducing or refuting DAMA/LIBRA's results using the same NaI(Tl) target material^{1–3}. The COSINE-100 experiment^{4,5} is one of these that is currently operating with 106 kg of low-background NaI(Tl) crystals at the Yangyang underground laboratory (Y2L). Dark matter interpretations of the COSINE-100 data from the background spectra presented null observations^{6,7} that were inconsistent with an explanation of the DAMA/LIBRA signals as a spin-independent interaction between weakly interacting massive particles (WIMPs), the stringent candidate of the dark matter particle, and sodium or iodine nuclei in the specific context of the standard halo model. Model independent searches of the annual modulation were reported^{15,8} but with insufficient statistics to corroborate the DAMA/LIBRA observation yet. Another experiment currently in operation, ANAIS-112, also reported the model-independent annual modulation search results that exhibited more than 2σ tension with DAMA/LIBRA's signals⁹ but has not yet made a conclusion. These model-independent searches considered time-dependent backgrounds based on the time-dependence of the rate induced by individual isotopes studied by cosmogenic activations^{10,11} and precise background modeling^{12–14}. However, the DAMA/LIBRA experiment used the residual rate by subtracting an average rate in every one-year cycle of data-taking roughly starting in September^{15,16}. If the background rate is not constant over time, this procedure can generate annually modulated event rates^{17,18}. Specifically, a slowly increasing event rate as a function of the time in the region of interest (ROI) can provide an annual modulation similar to the one observed by DAMA/LIBRA without dark matter signals as studied in literature¹⁷. To verify this phenomenon, it is interesting to apply DAMA/LIBRA's analysis technique to other experimental data. Here we analyze the COSINE-100 data in terms of the annual modulation, but applying the analysis methods adopted by DAMA/LIBRA.

Results

Experiment. The COSINE-100 experiment⁴ started physics operation in September 2016 at Y2L in South Korea with about 700 m of rock overburden. It utilizes eight low-background NaI(Tl) crystals arranged in a 4×2 array, with a total target mass of 106 kg. Each crystal is coupled to two photomultiplier tubes (PMTs) to measure the amount of energy deposited in the crystal. The NaI(Tl) detectors are immersed in a 2200 liter liquid scintillator, which allows for the identification and subsequent reduction of radioactive backgrounds observed by the crystal¹⁹. The liquid scintillator is surrounded by copper, lead, and plastic scintillators to reduce the background contribution from external radiation as well as tag cosmic-ray muons^{20,21}.

In contrast with the COSINE-100 detector, DAMA/LIBRA does not employ plastic- or liquid scintillator-based veto detectors. We, therefore, do not use information from those detectors in this analysis. In the ROI, PMT-induced noise events predominantly contribute to the single-hit physics data. They typically have fast decay times of less than 50 ns compared with typical NaI(Tl) scintillation of about 250 ns. The DAMA/LIBRA experiment developed a parameter to discriminate the PMT-induced noise by using a ratio of fast charge between 0 and 50 ns, X_1 , and slow charge between 100 and 600 ns, X_2 , each defined from relative to a time in the rising edge of PMT's waveform. The event selection (ES) parameter based on X_1 and X_2 provided good separation of the PMT-induced noise. DAMA/LIBRA claimed that they could efficiently remove the PMT-induced noise events by only using the ES parameter cut and achieve a 1 keV energy threshold with almost no noise contamination²². It should be noted that the 1 keV energy threshold from the COSINE-100 data was achieved by a multivariable machine learning technique (COSINE-100 nominal event selection) that used multiple parameters including, but not limited to X_1 and X_2 , mean decay time, charge asymmetry between two PMTs, and likelihood parameters for signal-like and noise-like templates^{7,23}.

We found slightly different selection efficiencies when we applied exactly the same criteria used by the DAMA/LIBRA for the ES parameter. Instead of the same value of the ES parameter cut, we choose values of the ES cut that result in a selection efficiency similar to the DAMA/LIBRA-phase2²². We evaluate the selection efficiency with a ⁶⁰Co calibration dataset, which has been used for the COSINE-100 data analyses^{5–8}, although DAMA/LIBRA used ²⁴¹Am calibration data. Our selection criteria developed with the ⁶⁰Co calibration data and applied to the physics data are presented in Fig. 1. The event selection efficiency compared with DAMA/LIBRA's efficiencies are shown in Fig. 2.

Figure 3 shows energy spectra of the single-hit and the multiple-hit events using the ES parameter cut. These spectra are compared with those of the COSINE-100 nominal event selections. DAMA/LIBRA's event selection introduces an excess below 2 keV compared with the nominal COSINE-100 event selection. These events were categorized as PMT-induced noise in the nominal COSINE-100 event selection^{7,23}. In the DAMA/LIBRA-phase2 energy spectrum, they also found a mild increase of the event rate below 2 keV¹⁶, although these excess events were claimed as possible dark matter interactions²². However, the possibility of remnants from PMT-induced noise for excess events of the DAMA/LIBRA data are not fully excluded.

Because DAMA/LIBRA's analysis always used residual spectra of the event rate by subtracting average backgrounds for the modulation fit, we first model the time-dependent background. Here we use two different approaches to account for the time-dependent backgrounds. The first model uses an exponential function to describe the time-dependent background used for the initial annual modulation studies in COSINE-100⁵ and ANAIS-112². The second model uses the yearly averaged rate in the DAMA/LIBRA experiment¹⁶.

After subtracting the background, the residual rates are fitted with a sinusoidal function,




Figure 1. The ES parameter vs energy applied to data from COSINE-100. The DAMA’s ES parameters for the PMT-induced noise rejection as a function of energy are presented for the ^{60}Co calibration data of the multiple-hit events (left) and three years of COSINE-100 physics data of the single-hit events (right). Solid red lines present selection criteria that provide selection efficiencies similar to the DAMA/LIBRA-phase2.




Figure 2. Event selection efficiency with the DAMA/LIBRA’s event selection applied to COSINE-100 data. Event selection efficiencies determined from the ^{60}Co calibration data using the multiple-hit events of the COSINE-100 crystal (crystal 6) (black points) are compared with those of DAMA/LIBRA-phase1 (red points)²⁴ and DAMA/LIBRA-phase2 (blue points)²⁵. Here selection criteria shown in Fig. 1 are determined to provide efficiencies similar to the DAMA/LIBRA-phase2.

$$R(t) = S_m \cos \frac{2\pi(t - t_0)}{T}, \tag{1}$$

where $R(t)$ is the residual event rate as a function of time, S_m is the modulation amplitude, t_0 is a phase, and T is a period. In the typical dark matter interaction assuming the standard halo model, t_0 and T are expected to be June 2nd, and 365.25 days (1 year), respectively.

Figure 4 shows 1–6 keV single-hit data and the time-dependent background model with the single-exponential model (a). The residual spectrum (c) is fitted with the sinusoidal function to obtain the modulation amplitude $S_m = 0.0048 \pm 0.0055$ counts/kg/keV/day. A similar procedure for the 2–6 keV single-hit events ((e) and (g)) obtains $S_m = 0.0041 \pm 0.0056$ counts/kg/keV/day. Even though the event selection only using the ES parameter contains significant PMT-induced noise events in the COSINE-100 data, the fitted results are consistent with no modulation when we model the time-dependent background using the single exponential function.

The same 1–6 keV single-hit data are modeled with the DAMA-like year average method (b). Considering the 13-year cycles of DAMA/LIBRA shown in Table 1, we divide the COSINE-100 data into three-year cycles




Figure 3. Low-energy spectra of a NaI(Tl) crystal in the COSINE-100 experiment. Energy spectra of one crystal (crystal 6) in COSINE-100 using nominal COSINE-100 event selection (black filled circles) and the DAMA/LIBRA's event selection (red open circles) are presented for the single-hit events (a) and multiple-hit events (b). Here, selection efficiencies are corrected for proper comparison. Because we do not use the muon and LS detectors for the DAMA/LIBRA's event selection, the COSINE-100 nominal analysis obtains significantly larger numbers of multiple-hit events. Due to the remnants of the PMT-induced events from the DAMA/LIBRA's event selection, a significant excess of the event rate below 2 keV in the single-hit events is observed.

presented in Table 2. The vertical lines in Fig. 4b represent the start and end of each year cycle. The residual rates from this model are shown in Fig. 4d with the annual modulation fit. Here we obtain a significant modulation amplitude $S_m = -0.044 \pm 0.006$ counts/kg/keV/day, which has about 7σ significance. Similarly, in the 2–6 keV energy region ((f) and (h)), $S_m = -0.046 \pm 0.006$ counts/kg/keV/day is obtained. The negative sign of S_m indicates an opposite phase compared to that of DAMA/LIBRA and to the predicted phase from the WIMP dark matter model. Because the COSINE-100 data have time-dependent backgrounds from cosmogenically activated nuclides and ^{210}Pb , event rates are clearly decreasing as a function of time. Simple yearly averages provide a bias on the modulation fit due to the mismodeling of the time-dependent background.

Multiple-hit events in both 1–6 keV and 2–6 keV regions do not show significant modulation behaviors even though we apply the DAMA-like method in Fig. 5). This is because the multiple-hit requirement removes the majority of radioisotopes that have high enough activities and short enough half-lives to affect the multi-hit event rate appreciably. Table 3 summarizes the fit results for the single-hit 1–6, 2–6 keV events and comparisons of results from COSINE-100 (nominal analysis)⁸, ANAIS-112⁹, and DAMA/LIBRA¹⁶.

Following up on the observation of the significant negative modulation of the COSINE-100 data using the DAMA-like method, we perform simulation studies about the DAMA/LIBRA's time-dependent background. Although the DAMA/LIBRA collaboration claimed no time-dependent background in their data²², a clear decrease of the event rate in the 2–6 keV range from DAMA/LIBRA-phase1²⁶ and DAMA/LIBRA-phase2¹⁶ is observed, approximately from 1.2 to 0.7 counts/kg/keV/day. The major update from DAMA/LIBRA-phase1 to DAMA/LIBRA-phase2 was the replacement of PMTs from ET (Electron tubes) Enterprises to Hamamatsu Photonics for the high quantum efficiency²⁵ but using the same NaI(Tl) crystals. The observed decrease in backgrounds between phases is likely not due to the use of different PMTs, as there is a 10 cm quartz light guide between PMTs and crystals. Background contribution from the PMT's radioisotopes in the COSINE-100, which used PMTs from the Hamamatsu photonics, was less than 0.05 counts/kg/keV/day without the quartz block¹⁴. We also noted that similar radiopurities of Electron tubes were reported and Hamamatsu PMTs²⁵.

Therefore, one can suspect that the background rate decreasing between phase1 and phase2 was indeed due to decays of short-lived radioisotopes such as ^3H and ^{210}Pb that were reported by COSINE-100^{10,14} and ANAIS-112¹¹. In this simulation study, we assume that DAMA/LIBRA's crystals have the same background composition as crystal 6 of COSINE-100, which has the lowest background among the COSINE-100 crystals. The total average rate in the COSINE-100 crystal data is 2.5 times higher than the total DAMA/LIBRA rate averaged over phase1 and phase2. We scale each COSINE-100 crystal 6 background component by factor 1/2 to simulate the DAMA/LIBRA data. After this scaling, the average rate of 2–6 keV single-hit events is 1.3 counts/kg/keV/day corresponds to the background rate at the beginning of the DAMA/LIBRA-phase1. Because the dominant backgrounds in this region from the COSINE-100 crystal are from ^3H and ^{210}Pb , a decreased background rate is obtained, as shown in Fig. 6. This background model plausibly describes the rate decrease from DAMA/LIBRA-phase1 to DAMA/LIBRA-phase2 (See Methods).

Simulated data are generated from the aforementioned time-dependent background without the dark matter signals. Figure 7a presents an example of the simulated data for 13-year cycles of the DAMA/LIBRA-like experiment. Vertical lines present the start and end of one year cycle used by DAMA/LIBRA (see Table 1). The residual spectrum subtracting the yearly average rate in the DAMA-like method is presented in Fig. 7b with the




Figure 4. Single-hit event rates in the unit of counts/keV/kg/day as a function of time. The top four panels present time-dependent event rates and the residual rates in the single-hit 1–6 keV regions with 15 days bin. Here, the event rates are averaged for the five crystals with weights from uncertainties in each 15-day bin. Purple solid lines present background modeling with the single exponential (a) and the yearly averaged DAMA-like method (b). Residual spectra for the single exponential model (c) and the DAMA-like model (d) are fitted with the sinusoidal function (red solid lines). Same for 2–6 keV in the bottom four panels. Strong annual modulations are observed using the DAMA-like method while the result using the single-exponential models are consistent with no observed modulation.

Cycle	Date period	Exposure (kg × day)
1	Sept. 9, 2003–July 21, 2004	51,405
2	July 21, 2004–Oct. 28, 2005	52,597
3	Oct. 28, 2005–July 18, 2006	39,445
4	July 19, 2006–July 17, 2007	49,377
5	July 17, 2007–Aug. 29, 2008	66,105
6	Nov. 12, 2008–Sept. 1, 2009	58,768
7	Sept. 1, 2009–Sept. 8, 2010	62,098
–	Dec. 23, 2010–Sept. 9, 2011	Commissioning
8	Nov. 2, 2011–Sept. 11, 2012	62,917
9	Oct. 8, 2012–Sept. 2, 2013	60,586
10	Sept. 8, 2013–Sept. 1, 2014	73,792
11	Sept. 1, 2014–Sept. 9, 2015	71,180
12	Sept. 10, 2015–Aug. 24, 2016	67,527
13	Sept. 7, 2016–Sept. 25, 2017	75,135

Table 1. DAMA/LIBRA annual cycles Thirteen annual cycles used by DAMA/LIBRA are obtained from Refs. ^{15,16,26}.

Cycle	Date period	Exposure (kg × day)
COSINE-1	Oct. 20, 2016–Sept. 16, 2017	20,323
COSINE-2	Sept. 17, 2017–Sept. 26, 2018	22,963
COSINE-3	Sept. 27, 2018–Sept. 20, 2019	22,042

Table 2. Three-year cycles of COSINE-100 data for the DAMA-like method^{15,16}.




Figure 5. Multiple-hit event rates in the unit of counts/keV/kg/day as a function of time. The top four panels present time-dependent events rates and the residual rates in the multiple-hit 1–6 keV regions with 15-day bin. Here, the event rates are averaged for the five crystals with weights from uncertainties in each 15 days bin size. Purple solid lines present background modeling with the single exponential (a) and with the yearly averaged DAMA-like method (b). Residual spectra for the single exponential model (c) and the DAMA-like model (d) are fitted with the sinusoidal function (red solid lines). The bottom four panels show the results for the 2–6 keV region using the same methods. In the multiple-hit events, no strong modulations from both methods are observed.

Counts/kg/keV/day	1–6 keV	2–6 keV
This work	-0.0441 ± 0.0057	-0.0456 ± 0.0056
DAMA/LIBRA	0.0105 ± 0.0011	0.0095 ± 0.0008
COSINE-100	0.0067 ± 0.0042	0.0050 ± 0.0047
ANAIS-112	-0.0034 ± 0.0042	0.0003 ± 0.0037

Table 3. Annual modulation amplitudes from various experiments. The amplitudes of the annual modulation fits using the DAMA-like method to the COSINE-100 3 years data (this work) are compared with results from DAMA/LIBRA^{15,16}, COSINE-100⁸, and ANAIS-112⁹ in both 1–6 keV and 2–6 keV regions.




Figure 6. Background model for DAMA/LIBRA using the COSINE-100 background compositions. Background compositions of the COSINE-100’s crystal (crystal 6) in the 2–6 keV single-hit regions are scaled to have 1.3 counts/keV/kg/day at the beginning of DAMA/LIBRA-phase1. Here dominant background contributions are ^{210}Pb (half-life of 8140 days) and ^3H (half-life of 4494 days). The total model (black solid line) is compared with the averaged rate in the 2–6 keV region from the initial 4 years of the DAMA/LIBRA-phase1 and 6 years of the DAMA/LIBRA-phase2¹⁶. Due to the decays of the time-dependent backgrounds, this model presents a decreasing rate as a function of time. This model describes the rate of decrease obtained from the DAMA/LIBRA-phase1 to DAMA/LIBRA-phase2 denoted in the plot.




Figure 7. An example of a simulation of the DAMA/LIBRA experiment in the single-hit 2–6 keV region. (a) One sample of the simulated experiment of the time-dependent event rates for the DAMA/LIBRA assuming COSINE-100’s background composition (points) is presented for 13-year cycles. Vertical lines represent the start and end of each cycle used by the DAMA/LIBRA experiment. (b) The residual spectrum applying the DAMA-like method (points) is fitted with the sinusoidal function (solid line). Although no dark matter signals are inserted in this simulated data, strong modulation is observed from the DAMA-like method.

modulation fit (solid line). Here we observe strong negative modulation of $S_m = -0.0098 \pm 0.0008$ counts/kg/keV/day corresponding to approximately 12σ significance.

We perform 1,000 simulated experiments and obtain the modulation amplitude distribution in Fig. 8a. Although negative signs of the modulation amplitudes are obtained, the magnitude of the modulation amplitude is consistent with that obtained by DAMA/LIBRA. We also perform the phase floated fits using the same simulated data. Figure 8b and c show the modulation amplitudes and phases, respectively, compared with those of the DAMA/LIBRA experiment. Once again, results of similar modulation amplitude and opposite phase were observed.

Conclusion

The direct detection of dark matter interactions is one of the main challenges of contemporary physics. Because of its significance, any hints of dark matter evidence need to be precisely examined. COSINE-100 continues to collect and analyze data to confirm or refute DAMA/LIBRA's hints at dark matter signals. Alternative explanations of the signals that do not involve dark matter, are investigated with the same NaI(Tl) target materials from the COSINE-100 experiment. We have observed PMT-induced noise contamination by adopting the selection criteria of DAMA/LIBRA. The time-dependent background model using a single exponential decay function provided no modulation from the COSINE-100 data, although significant PMT-induced noise in the low-energy signal region was observed. However, a DAMA-like yearly averaging method generates clear annual modulation signals because of the mismodeling of the time-dependent background. Furthermore, the DAMA-like method as applied to the simulated background-only data of DAMA/LIBRA, which are based on the time-dependent background in the COSINE-100 crystals, provides a consistent annual modulation signal of DAMA/LIBRA in the modulation amplitudes but a modulation phases that is almost opposite to that observed by DAMA/LIBRA. Although the observed phase is opposite, the consistent magnitude of the modulation amplitude may indicate an interesting phenomenon hidden in DAMA/LIBRA's background subtraction procedure.

Methods

Following DAMA/LIBRA's analysis method for the COSINE-100 data. The shield structure of the COSINE-100 experiment is similar to that of the DAMA/LIBRA experiment. However, the COSINE-100 detector has additional active veto detectors including the outermost plastic scintillators for the muon veto^{20,21} and the inner 2200 L of liquid scintillator for the external or internal radiation veto¹⁹. However, in this study we follow DAMA/LIBRA's event selection such that we do not use information from the plastic scintillators or the liquid scintillator, though they are still working as passive shields for external radioactivities. We defined two different categories of the events as the single-hit and the multiple-hit. The single-hit events correspond to any hit in a single NaI(Tl) crystal having no measurable energy in the other crystals. The multiple-hit events are defined in such a way that two or more NaI(Tl) crystals have measurable hits.

In the data acquisition of DAMA/LIBRA, PMT signals from the two ends of the crystal are digitized by a waveform analyzer for a time window of 2048 ns²⁴. In the data analysis, only a 600 ns time window integration of each pulse starting from the rising edge of each event was used to evaluate deposited energies.

Low energies corresponding up to 100 keV energy region were calibrated with external sources of ²⁴¹Am (59.5 keV) and ¹³³Ba (30.4 keV and 81.0 keV) and with internal X-rays or γ s (3.2 keV, 40.4 keV, 67.3 keV etc). A linear fits for those points was used for the energy calibration²⁴. However, in the DAMA/LIBRA-phase2 they




Figure 8. Results of 1000 simulated experiments for the DAMA/LIBRA experiment. We generate 1000 simulated experiments with a background derived from COSINE-100 with an exposure and analysis method derived from the DAMA/LIBRA experiment without the dark matter signals. (a) Results of the modulation amplitudes (S_m) using the phase-fixed fit are presented with DAMA/LIBRA's modulation amplitude. Although negative signs (opposite phase) are required, the simulated experiments obtain a modulation amplitude consistent with DAMA/LIBRA. Results for the phase-floated fits are shown for the modulation amplitude (b) and the modulation phase (c). Here we also observe consistent modulation amplitude but opposite phase by shifting the phase by π between the simulated experiments and DAMA/LIBRA.

observed about 0.2 keV shift for the tagged 3.2 keV line and applied an additional correction in the low energy region between the software energy threshold and 15 keV¹⁶.

The data acquisition system of the COSINE-100 experiment took waveforms of events for 8000 ns time windows^{4,27}. Nominal analysis used a 5,000 ns time window starting from the rising edge of each event for the deposited energy. We used only internal X-rays or γ lines for the energy calibration to avoid position dependencies from external sources. The nonproportional scintillation behavior of the NaI(Tl) crystals studied in Ref.²⁸ was applied for energy calibration in the low-energy region¹⁴. However, in this analysis, we follow the DAMA/LIBRA's method as closely as possible. Integrated charge in a 600 ns time window is used for the deposited energy. A linear fit relating charge to energy is determined using the 59.5 keV from external ²⁴¹Am source and 3.2 keV from internal ⁴⁰K, providing an energy scale for the low-energy events.

In the low-energy signal region below 10 keV, PMT-induced noise events predominantly contribute to the single-hit physics data. The dominant noise has a fast decay time of less than 50 ns compared with typical NaI(Tl) scintillation of about 250 ns. The DAMA/LIBRA collaboration developed a good parameter based on ratios of slow (X_1) and fast (X_2) charges defined as following^{24,25},

$$X_1 = \text{Charge (100 to 600 ns)}/\text{Charge (0 to 600 ns)}, \quad (2)$$

$$X_2 = \text{Charge (0 to 50 ns)}/\text{Charge (0 to 600 ns)}, \quad (3)$$

The typical PMT-induced noise deposited most of the pulses in X_2 while scintillation events of the NaI(Tl) crystals deposited about 70 % scintillation charge in the X_1 area. For an effective rejection of PMT-induced noise events, DAMA/LIBRA defined an event selection (ES) parameter as follows,

$$\text{ES} = \frac{1 - (X_2 - X_1)}{2}. \quad (4)$$

Typical signals have an ES parameter of around 0.8 while for the PMT-induced noise it is around 0.3, as one can see in Fig 1. DAMA/LIBRA collected pure scintillation events using 59.5 keV γ rays from an ²⁴¹Am source. DAMA/LIBRA determined the selection criteria as ES greater than 0.72 (0.85) in the energy region of 1–3 keV (3–6 keV)²⁴.

Initially, we tried to use identical selection criteria for the COSINE-100 data, but the resulting selection efficiency was seen to differ significantly from that reported by DAMA/LIBRA^{24,25}. This may be caused by slightly different scintillation characteristics of crystals due to different environmental conditions and different analysis methods used to determine the rising edge in the two experiments. Instead of the same ES parameter, we empirically develop cut criteria based on the selection efficiencies, making them similar between DAMA/LIBRA-phase2 and COSINE-100, as shown in Figs. 1 and 2.

Because DAMA/LIBRA does not have the muon veto detectors, muon-related events are not directly removed. Therefore, we tag high-energy events in each crystal requiring energies above 4 MeV. After the high energy event, we remove a 1 s period of events in each crystal. In addition, data is monitored in two-hour periods which are occasionally removed when any large variation in the environmental or detector parameters are recorded.

The low-energy spectra from the COSINE-100 data using the aforementioned calibration and event selection are presented in Fig. 3 and compared with the energy spectra from the nominal COSINE-100 data analysis¹⁴. Noticeably, we observe a significant increase of the event rate below 2 keV following DAMA/LIBRA's event selection method. Those excess events are categorized as PMT-induced noise events by the typical COSINE-100 data analysis.

Fitting procedure. We calculate the event rate for each NaI(Tl) detector binned in 15-day intervals. We then evaluate the detector livetime in each time bin and normalize the event rate based on its relative exposure. This process accounts for variations in exposure induced by both detector-off periods and data periods that are removed due to detector instability.

Although time-dependent event rates were never presented by DAMA/LIBRA, they claimed no time-dependent backgrounds in their data²². The time dependence is reported for the residuals. The published DAMA/LIBRA residuals were found by subtracting the time-averaged ROI event rate for each one-year cycle from the measured rate.

The DAMA/LIBRA cycles start every year around September, as summarized in Table 1^{15,16,26}. In addition to DAMA/LIBRA's method of the yearly averaged rate, we try to model the time-dependent background using a single exponential function to check the method bias.

The residual rates are fitted to the sinusoidal functions. The data from five detectors are fitted simultaneously. We perform the fit by minimizing the value of the computed χ^2 , defined as

$$\chi^2 = \sum_{ij} \frac{(n_{ij} - R_{ij})^2}{\sigma_{ij}^2}, \quad (5)$$

where n_{ij} is the residual rate for i th crystal and j th time bin, and σ_{ij} is the corresponding uncertainty of the data. R_{ij} is the expected modulation rate,

Isotopes	Half-lives (day)	Contributions (Counts/day/kg/keV)
^{238}U , ^{232}Th , ^{40}K	$> 10^{10}$	0.043
^{210}Pb	8140	0.687
^3H	4494	0.474
^{113}Sn	115.1	0.055
^{109}Cd	462	0.025
^{121m}Te	164.2	0.004
^{127m}Te	106.1	0.011

Table 4. Time-dependent background contributions in the NaI(Tl) crystals. The model of DAMA/LIBRA background model, assuming COSINE-100's background composition is summarized with half-lives and initial background contributions in the 2–6 keV single-hit events.

$$R_{ij} = S_m \cos \frac{2\pi(j - j_0)}{T}. \quad (6)$$

In the phase-fixed fit, j_0 is fixed to be 152 days with period $T=365.25$ days. A phase-floated fit with no constraints on j_0 is also performed.

Time dependent background in DAMA/LIBRA and COSINE-100. DAMA/LIBRA has not explicitly reported their background in detail, although a few individual radioisotope contaminations were studied²⁴. In a recent paper²², they presented the single-hit 1–100 keV energy spectrum for the first time (Fig. 20 in Ref.²²). They described the data with ^{129}I , ^{40}K , and ^{210}Pb decays, with continuum (constant) background due to high-energy γ/β , and signals (possibly dark matter interactions). Here, ^{210}Pb can contribute to the time-dependent background due to its half-life of 22.3 years. In their modeling, ^{210}Pb contribution in the ROI is negligible, so an assumption of no time-dependent background seems to be legitimate, assuming the validity of this particular background model.

In the ROI, dominant background contributions are caused by internal ^{40}K and the continuum background. The continuum background may be caused by internal contamination of ^{238}U , ^{232}Th or external PMT radioactivities such as ^{238}U , ^{232}Th , and ^{40}K . However, a precise modeling of the NaI(Tl) crystals performed by ANAIS-112 and COSINE-100 did not show a continuum background from such components^{13,14}. Those backgrounds have decreasing rates in the low-energy region. In this case, other backgrounds with the potential to increase rates at low energy may need to be considered, examples are ^3H and ^{210}Pb in the crystal surface^{14,29}. When considered, a time-dependent event rate that decreases in time is natural and similar to what is observed in ANAIS-112⁹ and COSINE-100⁸. It is worthwhile to mention that the background event rate in the 2–6 keV region in DAMA/LIBRA has decreased from DAMA/LIBRA-phase1 to DAMA/LIBRA-phase2 as indicated in Fig. 6.

For the simulated experiments of DAMA/LIBRA, we try to generate the time-dependent background describing DAMA/LIBRA's background behavior. Because there is no explicit description of time-dependent backgrounds from DAMA/LIBRA, we simply consider the same background components as the COSINE-100 crystals. The approximately 2.5 times lower background is scaled from the COSINE-100 background using the same fractional compositions of different background components. Short-lived radioisotopes with half-lives of less than 100 days were not included. Averaged rate of each time-dependent background component at the beginning of the experiment in the 2–6 keV region is summarized in Table 4.

Figure 6 shows the model of the time-dependent background in the 2–6 keV region derived from the COSINE-100 data. Interestingly, this decreasing rate agrees with the change in average event rates between DAMA/LIBRA-phase1 and phase2. With this model, we generate simulated data and performed the fit for the year-subtracted residual rate as shown in Fig 7.

Data availability

The datasets used and/or analyzed during the current study available from the corresponding author on reasonable request.

Received: 9 November 2022; Accepted: 15 March 2023

Published online: 22 March 2023

References

- de Barbosa Souza, E. *et al.* First search for a dark matter annual modulation signal with NaI(Tl) in the Southern Hemisphere by DM-Ice17. *Phys. Rev. D* **95**, 032006. <https://doi.org/10.1103/PhysRevD.95.032006> (2017).
- Amaré, J. *et al.* First results on dark matter annual modulation from ANAIS-112 experiment. *Phys. Rev. Lett.* **123**, 031301. <https://doi.org/10.1103/PhysRevLett.123.031301> (2019).
- Antonello, M. *et al.* The SABRE project and the SABRE Proof-of-Principle. *Eur. Phys. J. C* **79**, 363. <https://doi.org/10.1140/epjc/s10052-019-6860-y> (2019).
- Adhikari, G. *et al.* Initial performance of the COSINE-100 experiment. *Eur. Phys. J. C* **78**, 107. <https://doi.org/10.1140/epjc/s10052-018-5590-x> (2018).

5. Adhikari, G. *et al.* Search for a dark matter-induced annual modulation signal in NaI(Tl) with the COSINE-100 experiment. *Phys. Rev. Lett.* **123**, 031302. <https://doi.org/10.1103/PhysRevLett.123.031302> (2019).
6. Adhikari, G. *et al.* An experiment to search for dark-matter interactions using sodium iodide detectors. *Nature* **564**, 83. <https://doi.org/10.1038/s41586-018-0739-1> (2018).
7. Adhikari, G. *et al.* Strong constraints from COSINE-100 on the DAMA dark matter results using the same sodium iodide target. *Sci. Adv.* **7**, eabk2699. <https://doi.org/10.1126/sciadv.abk2699> (2021).
8. Adhikari, G. *et al.* Three-year annual modulation search with cosine-100. *Phys. Rev. D* **106**, 052005. <https://doi.org/10.1103/PhysRevD.106.052005> (2022).
9. Amaré, J. *et al.* Annual modulation results from three-year exposure of ANAIS-112. *Phys. Rev. D* **103**, 102005. <https://doi.org/10.1103/PhysRevD.103.102005> (2021).
10. Barbosa de Souza, E. *et al.* Study of cosmogenic radionuclides in the COSINE-100 NaI(Tl) detectors. *Astropart. Phys.* **115**, 102390. <https://doi.org/10.1016/j.astropartphys.2019.102390> (2020).
11. Villar, P. *et al.* Study of the cosmogenic activation in NaI(Tl) crystals within the ANAIS experiment. *Int. J. Mod. Phys. A* **33**, 1843006. <https://doi.org/10.1142/S0217751X18430066> (2018).
12. Adhikari, P. *et al.* Background model for the NaI(Tl) crystals in COSINE-100. *Eur. Phys. J. C* **78**, 490. <https://doi.org/10.1140/epjc/s10052-018-5970-2> (2018).
13. Amaré, J. *et al.* Analysis of backgrounds for the ANAIS-112 dark matter experiment. *Eur. Phys. J. C* **79**, 412. <https://doi.org/10.1140/epjc/s10052-019-6911-4> (2019).
14. Adhikari, G. *et al.* Background modeling for dark matter search with 1.7 years of COSINE-100 data. *Eur. Phys. J. C* **81**, 837 (2021).
15. Bernabei, R. *et al.* Final model independent result of DAMA/LIBRA-phase1. *Eur. Phys. J. C* **73**, 2648. <https://doi.org/10.1140/epjc/s10052-013-2648-7> (2013).
16. Bernabei, R. *et al.* First model independent results from DAMA/LIBRA-phase2. *Nucl. Phys. At. Energy* **19**, 307. <https://doi.org/10.15407/jnpae2018.04.307> (2018).
17. Buttazzo, D., Panci, P., Rossi, N. & Strumia, A. Annual modulations from secular variations: Relaxing DAMA?. *JHEP* **20**, 137. [https://doi.org/10.1007/JHEP04\(2020\)137](https://doi.org/10.1007/JHEP04(2020)137) (2020).
18. Messina, A., Nardecchia, M. & Piacentini, S. Annual modulations from secular variations: Not relaxing DAMA?. *JCAP* **20**, 037. <https://doi.org/10.1088/1475-7516/2020/04/037> (2020).
19. Adhikari, G. *et al.* The COSINE-100 liquid scintillator veto system. *Nucl. Instrum. Methods A* **1006**, 165431. <https://doi.org/10.1016/j.nima.2021.165431> (2021).
20. Prihtiadi, H. *et al.* Muon detector for the COSINE-100 experiment. *JINST* **13**, T02007. <https://doi.org/10.1088/1748-0221/13/02/T02007> (2018).
21. Prihtiadi, H. *et al.* Measurement of the cosmic muon annual and diurnal flux variation with the COSINE-100 detector. *JCAP* **02**, 013. <https://doi.org/10.1088/1475-7516/2021/02/013> (2021).
22. Bernabei, R. *et al.* The DAMA project: Achievements, implications and perspectives. *Prog. Part. Nucl. Phys.* **114**, 103810. <https://doi.org/10.1016/j.pnpnp.2020.103810> (2020).
23. Adhikari, G. *et al.* Lowering the energy threshold in COSINE-100 dark matter searches. *Astropart. Phys.* **130**, 102581. <https://doi.org/10.1016/j.astropartphys.2021.102581> (2021).
24. Bernabei, R. *et al.* The DAMA/LIBRA apparatus. *Nucl. Instrum. Meth. A* **592**, 297–315. <https://doi.org/10.1016/j.nima.2008.04.082> (2008).
25. Bernabei, R. *et al.* Performances of the new high quantum efficiency PMTs in DAMA/LIBRA. *JINST* **7**, P03009. <https://doi.org/10.1088/1748-0221/7/03/P03009> (2012).
26. Bernabei, R. *et al.* First results from DAMA/LIBRA and the combined results with DAMA/NaI. *Eur. Phys. J. C* **56**, 333–355. <https://doi.org/10.1140/epjc/s10052-008-0662-y> (2008).
27. Adhikari, G. *et al.* The COSINE-100 data acquisition system. *JINST* **13**, P09006. <https://doi.org/10.1088/1748-0221/13/09/P09006> (2018).
28. Swiderski, L. Response of doped alkali iodides measured with gamma-ray absorption and Compton electrons. *Nucl. Instrum. Methods A* **705**, 42 (2013).
29. Yu, G. H. *et al.* Depth profile study of ^{210}Pb in the surface of an NaI(Tl) crystal. *Astropart. Phys.* **126**, 102518. <https://doi.org/10.1016/j.astropartphys.2020.102518> (2021).

Acknowledgements

We thank the Korea Hydro and Nuclear Power (KHNP) Company for providing underground laboratory space at Yangyang. This work is supported by: the Institute for Basic Science (IBS) under project code IBS-R016-A1 and NRF-2016R1A2B3008343, Republic of Korea; NSF Grants No. PHY-1913742, DGE-1122492, WIPAC, the Wisconsin Alumni Research Foundation, United States; STFC Grant ST/N000277/1 and ST/K001337/1, United Kingdom; Grant No. 2021/06743-1 FAPESP, CAPES Finance Code 001, CNPq 131152/2020-3, Brazil.

Author contributions

The main analysis was led by H.P. supervised by H.S.L. The paper was written by H.P. and H.S.L. and edited by other members of the collaboration. All authors have participated in online shift and approved the manuscript. Authors are alphabetically listed by their last names.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to H.S.L. or H.P.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023