



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/198274/>

Version: Supplemental Material

Article:

Elder, FCT, O'Neill, AJ, Collins, LM et al. (2023) A framework to assess the terrestrial risk of antibiotic resistance from antibiotics in slurry or manure amended soils. *Environmental Science: Advances*, 2 (5). pp. 780-794. ISSN: 2754-7000

<https://doi.org/10.1039/D2VA00306F>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Supplementary Information

A framework to assess the terrestrial risk of antibiotic resistance from antibiotics in slurry or manure amended soils.

Felicity C T Elder^{1*}, Alex J O'Neill², Lisa M Collins³, Laura J Carter¹

¹ School of Geography, Faculty of Environment, University of Leeds, United Kingdom

² School of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds

³ School of Biology, Faculty of Biological Sciences, University of Leeds

Corresponding author: Felicity C T Elder, E-mail: f.elder@leeds.ac.uk

Table of Contents

SI Table 1 Predicted No Effect Concentrations

SI Table 2 Literature Concentrations of Antibiotics in Farmyard Manure

SI Table 3 Tetracyclines – Predicted Environmental Concentrations Pore Water

SI Table 5 Sulfonamides and Lincosamides– Predicted Environmental Concentrations Pore Water

SI Table 1 Predicted No Effect Concentrations (1)

Antibiotic	PNEC environment ($\mu\text{g.L}^{-1}$)	PNEC resistance ($\mu\text{g.L}^{-1}$)
Tetracyclines		
Tetracycline (TET)	3.2	1
Oxytetracycline (OXY)	18	0.5
Fluoroquinolones		
Ciprofloxacin (CIPRO)	0.45	0.06
Enrofloxacin (ENRO)	1.9	0.06
Sulfonamides		
Sulfamethoxazole (SMZ)	NA	NA
Lincosamides		
Lincomycin (LINCO)	1.8	2

SI Table 2 Literature Concentrations of Antibiotics in Farmyard Manure (FYM)

Antibiotic	Slurry Conc. (mg Kg ⁻¹)	FYM type	Reference
Tetracyclines			
Tetracycline (TET)	0.45	Calf slurry	Huygens et al, 2021(2)
	0.16	Pig slurry	Rasschaert et al, 2020 (3)
	12.10	Pig slurry	Qiao et al, 2012(4)
	16.00	Pig slurry	Gros et al, 2019(5)
	43.50	manure	Hu et al, 2010(6)
	300.0	Pig slurry	Widyasari-Mehta et al, 2016 (7)
	227.0	Pig slurry	Widyasari-Mehta et al, 2016 (7)
	5.90	Pig slurry	Widyasari-Mehta et al, 2016 (7)
Oxytetracycline (OXY)	40.78	Calf slurry	Huygens et al, 2021(2)
	4.819	Pig slurry	Rasschaert et al. 2020 (3)
	770.0	Pig slurry	Gros et al, 2019(5)
	75.29	Pig slurry	Gros et al, 2019(5)
	34.23	Pig slurry	Gros et al, 2019(5)
	183.5	manure	Hu et al, 2010(6)
	211	Pig slurry	Widyasari-Mehta et al, 2016 (7)
	58.27	Pig slurry	Huygens et al, 2021(5)
Fluoroquinolones			
Ciprofloxacin (CIPRO)	0.48	Calf slurry	Huygens et al, 2021(2)
	0.06	Pig slurry	Huygens et al, 2021(3)
	8.80	Pig slurry	Gros et al, 2019(5)
	34.00	Pig slurry	Gros et al, 2019(5)
	43.00	manure	Hu et al, 2010(6)
Enrofloxacin (ENRO)	0.31	Calf slurry	Huygens et al, 2021(2)
	0.14	Pig slurry	Rasschaert et al, 2020(3)
	60.21	Pig slurry	Gros et al, 2019(5)
	53.00	Pig slurry	Gros et al, 2019(5)
	4.70	Pig slurry	Widyasari-Mehta et al, 2016 (7)
Sulfonamides			
Sulfamethazine (SMZ)	0.03	Calf slurry	Huygens et al, 2021(2)
	0.458	Pig slurry	Gros et al, 2019(5)
	3.0	Pig slurry	Van den Meersche et al, 2016(8)
	4.9	Pig manure	Ji et al, 2012(9)
	7.3	Pig manure	Hu et al, 2010(6)
	4.6	Cattle manure	Ji et al, 2012(9)
	8.0	Chicken manure	Ho et al, 2014 (10)
5.8	Chicken manure		
Lincosamides			
Lincomycin (LINCOO)	0.36	Calf slurry	Huygens et al, 2021(2)
	1.767	Pig slurry	Rasschaert et al, 2020(3)
	227	Pig slurry	Gros et al, 2019(5)
	97.8	Pig slurry	Kuchta and Cessna, 2009(11)
	3.8	manure	Hu et al, 2010(6)

SI Table 3 Tetracyclines – Predicted Environmental Concentrations Pore Water

Equ.2 – Droge and Goss (2013)(12)

Equ.3 – Franco et al (2009)(13).

Soil pH	% Ionised	Model	K_d (L.Kg ⁻¹)	FYM conc (mg.kg-1)	Non-Ploughed			Ploughed		
					PEC soil (µg.kg-1)	PEC _{PW} (µg.L-1)	RQ	PEC soil (µg.kg-1)	PEC _{PW} (µg.L-1)	RQ
Tetracycline (TET)										
8.7	66.86	Equ.3	0.0007	0.45	0.1	21.54	21.54	0.02	4.31	4.31
5	37.75	Equ.2	5958.22			2.56E-05	2.57E-05		5.14E-06	5.14E-06
8.7	66.86	Equ.3	0.0007	0.16	0.036	7.71	7.71	0.007	1.54	1.54
5	37.75	Equ.2	5958.22			9.18E-06	9.19E-06		1.84E-06	1.84E-06
8.7	66.86	Equ.3	0.0007	12.1	2.74	579.08	597.08	0.55	115.82	115.82
5	37.75	Equ.2	5958.22			6.9E-04	6.90E-04		1.38E-04	1.38E-04
8.7	66.86	Equ.3	0.0007	16	3.62	765.73	765.73	0.73	153.15	153.15
5	37.75	Equ.2	5958.22			9.13E-04	9.13E-04		1.83E-04	1.83E-04
8.7	66.86	Equ.3	0.0007	43.5	9.86	2081.82	2001.82	1.97	416.36	416.36
5	37.75	Equ.2	5958.22			2.48E-03	2.48E-03		4.96E-04	4.96E-04
8.7	66.86	Equ.3	0.0007	300	68.0	14357.41	14357.41	13.6	2871.48	2871.48
5	37.75	Equ.2	5958.22			1.71E-02	1.71E-02		3.42E-03	3.42E-03
8.7	66.86	Equ.3	0.0007	227	51.45	10863.77	10863.77	10.29	2172.75	2172.75
5	37.75	Equ.2	5958.22			1.29E-05	1.30E-02		2.59E-03	2.59E-03
8.7	66.86	Equ.3	0.0007	5.9	1.34	282.36	282.36	0.27	56.47	56.47
5	37.75	Equ.2	5958.22			3.36E-04	3.37E-04		6.73E-05	6.73E-05
Oxytetracycline (OXY)										
8.7	74.67	Equ.3	0.0002	40.78	9.24	6770.30	13540.60	1.85	1354.06	2708.12
5	32.89	Equ.2	6172.00			0.0023	4.49E-03		4.49E-04	8.99E-04
8.7	74.67	Equ.3	0.0002	4.82	1.09	800.05	1600.10	0.22	160.01	320.02
5	32.89	Equ.2	6172.00			0.0003	5.31E-04		5.31E-05	1.06E-04
8.7	74.67	Equ.3	0.0002	770	174.53	127835.45	255670.89	34.91	25567.09	51134.18
5	32.89	Equ.2	6172.00			0.0424	0.085		8.48E-03	0.017
8.7	74.67	Equ.3	0.0002	75.29	17.07	12499.65	24999.30	3.41	2499.93	4.999.86
5	32.89	Equ.2	6172.00			0.0042	8.30E-03		8.30E-04	1.66E-03
8.7	74.67	Equ.3	0.0002	34.23	7.76	5682.87	11365.73	1.55	1136.57	2273.15
5	32.89	Equ.2	6172.00			0.0019	3.77E-03		3.77E-04	7.54E-04
8.7	74.67	Equ.3	0.0002	183.5	41.59	30464.68	60929.36	8.32	6092.94	12185.87
5	32.89	Equ.2	6172.00			0.0101	2.02E-02		2.02E03	4.04E-03
8.7	74.67	Equ.3	0.0002	211	47.82	35030.23	70060.47	9.57	7006.05	14012.09
5	32.89	Equ.2	6172.00			0.0116	2.33E-02		2.32E-03	4.65E-05
8.7	66.86	Equ.3	0.0002	58.27	13.21	9673.99	19347.98	2.64	1934.80	3869.60
5	37.75	Equ.2	6172.00			0.00321	6.42E-03		6.42E-04	1.28E-05

SI Table 4 Fluoroquinolones– Predicted Environmental Concentrations Pore Water

Equ.2 – Droge and Goss (2013)(12)

Equ.3 – Franco et al (2009)(13).

Soil pH	% Ionised	Model	K_d (L.Kg ⁻¹)	FYM conc (mg.kg ⁻¹)	Non-Ploughed			Ploughed		
					PEC soil (µg.Kg-1)	PEC _{PW} (µg.L-1)	RQ	PEC soil (µg.Kg-1)	PEC _{PW} (µg.L-1)	RQ
Ciprofloxacin (CIPRO)										
8.7	70.79	Equ.3	0.26	0.48	0.11	0.63	10.55	0.02	0.13	2.11
5	2.24	Equ.2	5038.76			3.24E-05	5.40E-04		6.48E-06	1.08E-04
8.7	70.79	Equ.3	0.26	0.06	0.003	0.08	1.38	0.003	0.02	0.28
5	2.24	Equ.2	5038.76			4.25E-06	7.09E-05		8.50E-07	1.42E-05
8.7	70.79	Equ.3	0.26	8.8	1.99	11.60	198.38	0.04	2.32	38.68
5	2.24	Equ.2	5038.76			5.94E-04	9.88E-03		1.19E-04	1.98E-03
8.7	70.79	Equ.3	0.26	34	7.71	44.83	747.16	1.54	8.97	149.43
5	2.24	Equ.2	5038.76			2.29E-03	8.83E-02		4.59E-04	7.65E03
8.7	70.79	Equ.3	0.041	90	20.4	118.67	1977.79	4.08	23.73	395.56
5	2.24	Equ.2	5038.76			6.07E-03	1.01E-01		1.21E-03	2.02E-02
Enrofloxacin (ENRO)										
8.7	92.9	Equ.3	5.73	0.31	0.07	0.02	0.31	0.01	3.68E-03	0.06
5	3.82	Equ.2	6434.84			1.63E-05	2.73E-04		2.28E-06	5.46E-05
8.7	92.9	Equ.3	5.73	0.142	0.03	0.01	0.14	6.44E-03	1.69E-03	0.03
5	3.82	Equ.2	6434.84			7.5E-06	1.25E-04		1.5E-06	2.50E-05
8.7	92.9	Equ.3	5.73	60.21	13.65	3.57	59.57	2.73	0.71	11.91
5	3.82	Equ.2	6434.84			3.18E-03	5.30E-02		6.36E-04	1.06E-02
8.7	92.9	Equ.3	5.73	53	12.01	3.15	52.43	2.4	0.63	10.49
5	3.82	Equ.2	6434.84			2.8E-03	4.67E-02		5.6E-04	9.34E-03
8.7	92.9	Equ.3	5.73	4.7	1.07	0.28	4.65	0.21	0.06	0.93
5	3.82	Equ.2	6434.84			2.48E-04	4.14E-03		4.97E-05	8.28E-04

SI Table 5 Sulfonamides and Lincosamides– Predicted Environmental Concentrations Pore Water

Equ.2 – Droge and Goss (2013)(12)

Equ.3 – Franco et al (2009)(13).

*RQ calculated using PNECr for sulfamethoxazole.

Soil pH	% Ionised	Model	K_d (L.Kg ⁻¹)	FYM conc (mg.kg ⁻¹)	Non-Ploughed			Ploughed		
					PEC soil (µg.Kg ⁻¹)	PEC _{PW} (µg.L ⁻¹)	RQ	PEC soil (µg.Kg ⁻¹)	PEC _{PW} (µg.L ⁻¹)	RQ
Sulfamethazine (SMZ)										
8.7 5	97.10 50	Equ.3 NA	8.57 NA	0.45	6.8E-03	0.001 NA	7.43E-05*	0.001	0.0002 NA	1.49E-05*
8.7 5	97.10 50	Equ.3 NA	8.57 NA	0.46	0.1	0.018 NA	1.14E-03*	0.021	0.004 NA	2.27E-04*
8.7 5	97.10 50	Equ.3 NA	8.57 NA	3.0	0.68	0.119 NA	7.43E-03*	0.136	0.024 NA	1/49E-03*
8.7 5	97.10 50	Equ.3 NA	8.57 NA	4.9	1.11	0.194 NA	1.21E-02*	0.222	0.039 NA	2.43E-03*
8.7 5	97.10 50	Equ.3 NA	8.57 NA	7.3	1.65	0.289 NA	1.81E-02*	0.331	0.058 NA	3.62E-03*
8.7 5	97.10 50	Equ.3 NA	8.57 NA	4.6	1.04	0.182 NA	1.14E-02*	0.209	0.036 NA	2.28E-03*
8.7 5	97.10 50	Equ.3 NA	8.57 NA	8.0	1.81	0.317 NA	1.98E-02*	0.363	0.063 NA	3.96E-03*
8.7 5	97.10 50	Equ.3 NA	8.57 NA	5.8	1.31	0.230 NA	1.44E-02*	0.263	0.046 NA	2.87E-03*
Lincomycin (LINCO)										
8.7 5	22.7 0.01	Equ.2 Equ.2	7818.34	0.36	0.08	1.57E-05	7.83E-06	0.016	3.13E-06	1.56E-06
8.7 5	22.7 0.01	Equ.2 Equ.2	7818.34	1.77	0.4	7.68E-05	3.84E-05	0.08	1.54E-05	7.68E-06
8.7 5	22.7 0.01	Equ.2 Equ.2	7818.34	227	51.45	9.87E-03	4.94E-03	10.291	1.97E-03	9.87E-04
8.7 5	22.7 0.01	Equ.2 Equ.2	7818.34	3.8	0.86	1.65E-04	8.26E-05	0.172	3.31E-05	1.65E-05

References

1. AMR Industry Alliance. AMR Industry Alliance Antibiotic Discharge Targets -List of Predicted No-Effect Concentrations (PNECs)-. 2018;(October 2000):1–5. Available from: https://www.amrindustryalliance.org/wp-content/uploads/2018/09/AMR_Industry_Alliance_List-of-Predicted-No-Effect-Concentrations-PNECs.pdf
2. Huygens J, Daeseleire E, Mahillon J, Van Elst D, Decrop J, Meirlaen J, et al. Presence of antibiotic residues and antibiotic resistant bacteria in cattle manure intended for fertilization of agricultural fields: A one health perspective. *Antibiotics*. 2021;10(4).
3. Rasschaert G, Van Elst D, Colson L, Herman L, Ferreira HC de C, Dewulf J, et al. Antibiotic residues and antibiotic-resistant bacteria in pig slurry used to fertilize agricultural fields. *Antibiotics*. 2020;9(1).
4. Qiao M, Chen W, Su J, Zhang B, Zhang C. Fate of tetracyclines in swine manure of three selected swine farms in China. *J Environ Sci (China)* [Internet]. 2012;24(6):1047–52. Available from: [http://dx.doi.org/10.1016/S1001-0742\(11\)60890-5](http://dx.doi.org/10.1016/S1001-0742(11)60890-5)
5. Gros M, Mas-Pla J, Boy-Roura M, Geli I, Domingo F, Petrović M. Veterinary pharmaceuticals and antibiotics in manure and slurry and their fate in amended agricultural soils: Findings from an experimental field site (Baix Empordà, NE Catalonia). *Sci Total Environ*. 2019;654:1337–49.
6. Hu X, Zhou Q, Luo Y. Occurrence and source analysis of typical veterinary antibiotics in manure, soil, vegetables and groundwater from organic vegetable bases, northern China. *Environ Pollut* [Internet]. 2010;158(9):2992–8. Available from: <http://dx.doi.org/10.1016/j.envpol.2010.05.023>
7. Widiasari-Mehta A, Hartung S, Kreuzig R. From the application of antibiotics to antibiotic residues in liquid manures and digestates: A screening study in one European center of conventional pig husbandry. *J Environ Manage* [Internet]. 2016;177:129–37. Available from: <http://dx.doi.org/10.1016/j.jenvman.2016.04.012>
8. Van den Meersche T, Pamel E Van, Poucke C Van, Herman L, Heyndrickx M, Rasschaert G, et al. Development, validation and application of an ultra high performance liquid chromatographic-tandem mass spectrometric method for the simultaneous detection and quantification of five different classes of veterinary antibiotics in swine manure. *J Chromatogr A* [Internet]. 2016;1429:248–57. Available from: <http://dx.doi.org/10.1016/j.chroma.2015.12.046>
9. Ji X, Shen Q, Liu F, Ma J, Xu G, Wang Y, et al. Antibiotic resistance gene abundances associated with antibiotics and heavy metals in animal manures and agricultural soils adjacent to feedlots in Shanghai; China. *J Hazard Mater* [Internet]. 2012;235–236:178–85. Available from: <http://dx.doi.org/10.1016/j.jhazmat.2012.07.040>
10. Ho Y Bin, Zakaria MP, Latif PA, Saari N. Occurrence of veterinary antibiotics and progesterone in broiler manure and agricultural soil in Malaysia. *Sci Total Environ* [Internet]. 2014;488–489(1):261–7. Available from: <http://dx.doi.org/10.1016/j.scitotenv.2014.04.109>
11. Kuchta SL, Cessna AJ. Lincomycin and spectinomycin concentrations in liquid swine manure

and their persistence during simulated manure storage. Arch Environ Contam Toxicol. 2009;57(1):1–10.

12. Droge STJ, Goss KU. Development and evaluation of a new sorption model for organic cations in soil: Contributions from organic matter and clay minerals. Environ Sci Technol. 2013;47(24):14233–41.
13. Franco A, Fu W, Trapp S. Influence of soil pH on the sorption of ionizable chemicals: Modeling advances. Environ Toxicol Chem. 2009;28(9):2018.