



This is a repository copy of *What influences GPs' use of pelvic examination? A qualitative investigation in primary care.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/199586/>

Version: Published Version

Article:

Williams, P., Murchie, P., Cruickshank, M.E. et al. (2 more authors) (2023) What influences GPs' use of pelvic examination? A qualitative investigation in primary care. *British Journal of General Practice*, 73 (732). e528-e536. ISSN 0960-1643

<https://doi.org/10.3399/bjgp.2022.0363>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

What influences GPs' use of pelvic examination?

A qualitative investigation in primary care

Abstract

Background

Omission of pelvic examination (PE) has been associated with diagnostic delay in women diagnosed with gynaecological cancer. However, PEs are often not carried out by GPs.

Aim

To determine the perceptions of GPs about the role of PEs, the barriers to and facilitators of PEs, and GPs' experience of PEs in practice.

Design and setting

Qualitative semi-structured interview study conducted in one health board in Scotland (mixed urban and rural) with an approximate population of 500 000.

Method

Interviews were conducted face-to-face or by telephone between March and June 2019. Framework analysis used the COM-B behaviour change model concepts of capability, opportunity, and motivation.

Results

Data was compatible with all three domains of the COM-B framework. Capability related to training in and maintenance of skills. These went beyond carrying out the examination to interpreting it reliably. Opportunity related to the clinical environment and the provision of chaperones for intimate examination. Interviewees described a range of motivations towards or against PEs that were unrelated to either capability or opportunity. These all related to providing high-quality care, but this was defined in different ways: 'doing what is best for the individual', 'doctors examine', and 'GPs as pragmatists'.

Conclusion

GPs' reasons for carrying out, or not carrying out, PEs in women with symptoms potentially indicating cancer are complex. The COM-B framework provides a way of understanding this complexity. Interventions to increase the use of PEs, and critics of its non-use, need to consider these multiple factors.

Keywords

clinical examination; gynaecological cancer; pelvic examination; primary health care; qualitative research; general practice.

INTRODUCTION

Clinical examination is traditionally viewed as an essential skill for a doctor.¹ Pelvic examination (PE) consists of inspection of the vulva, bimanual examination of the pelvic organs, and visualisation of the cervix by speculum examination. The intimate nature of a PE can make it a challenging examination for both clinicians and patients but there is evidence that lack of a PE is associated with diagnostic delay.²⁻⁴ PE can provoke feelings of awkwardness, discomfort, embarrassment around getting undressed, and loss of dignity,⁵⁻⁷ engendering a feeling of vulnerability, and some women avoid PEs altogether.⁸

GPs have a professional duty to manage the intimacy of PEs as well as to ensure the examination is carried out when clinically required. This responsibility has been acknowledged by the General Medical Council (GMC), who have produced relevant guidance.⁹ One study has shown that GPs would carry out a pre-referral PE in only 37% of women who present with a gynaecological problem requiring referral to a specialist.¹⁰

PE requires dexterity and sensitivity; however, if a clinician does not carry out the examination frequently, the skills, or confidence in those skills, may decline along

with the ability to distinguish normal from abnormal findings, and the willingness to undertake the procedure.^{11,12}

Sensitive topics such as clinical behaviour can be explored more fully using qualitative methods. This qualitative study answered the question: 'What are the perceptions and experiences of GPs on the role of primary-care-based PE, and the barriers to and facilitators of undertaking it?'

The theoretical framework used was the established COM-B behaviour change model. This model was used because it identifies what component of behaviour, capability, opportunity, or motivation needs to be modified for an intervention, in this instance, PE, to be successful (Figure 1).^{13,14} COM-B has been used previously to understand variation in practice by GPs where social awkwardness is an issue: for instance, in relation to implementing testing for sexually transmitted infections.¹⁵ It also links to the behavioural change wheel that aids intervention development.¹³

METHOD

Qualitative semi-structured telephone or face-to-face interviews were conducted with practising GPs in one health board in Scotland between March and June 2019. Fourteen NHS Grampian GP practices

P Williams (ORCID: 0000-0003-1575-5417), MBChB (Comm), BSc (Hons), PhD, MRCP, GP and NHS Research Scotland research fellow; **ME Cruickshank** (ORCID: 0000-0002-8893-8620), MBChB, MD, FRCOG, professor of gynaecology, Aberdeen Centre for Women's Health Research, Institute of Applied Health Sciences, University of Aberdeen, Aberdeen Maternity Hospital, Aberdeen. **P Murchie** (ORCID: 0000-0001-9968-5991), MBChB, BSc (MedSci), MSc, FRCGP, PhD, professor of primary care; **CM Bond** (ORCID: 0000-0003-0429-8205), BPharm (Hons), Med, PhD, FFPH, FRCPE, FRPS, FFRPS, FRCGP, FHEA, emeritus professor of primary care, Academic Primary Care, Institute of Applied Health Sciences, University of Aberdeen, Aberdeen. **CD Burton** (ORCID: 0000-0003-1159-1494), MBChB, MD,

FRCGP, professor of primary care, Academic Unit of Primary Care, University of Sheffield, Sheffield.

Address for correspondence

Christopher D. Burton, Academic Unit of Medical Education, Sam Fox House, Northern General Hospital, Herries Road, Sheffield S5 7AU, UK.

Email: chris.burton@sheffield.ac.uk

Submitted: 11 July 2022; **Editor's response:** 6 September 2022; **final acceptance:** 27 January 2023.

©The Authors

This is the full-length article (published online 18 Apr 2023) of an abridged version published in print. Cite this version as: **Br J Gen Pract 2023; DOI: <https://doi.org/10.3399/BJGP.2022.0363>**

How this fits in

Not carrying out a pelvic examination (PE) is associated with diagnostic delay in gynaecological cancers; however, PEs are often not carried out by GPs. This qualitative study reveals that GPs' reasons for carrying out, or not carrying out, PEs in women with symptoms potentially indicating cancer are complex. GPs need to have the skills to do, and interpret the findings of, a PE; have the opportunity to carry out the examination in terms of time and availability of chaperones; and believe that it will influence the referral process.

that had participated in a previous linked data analysis study involving patients diagnosed with a gynaecological cancer were approached (unpublished data, Pauline Williams, 2021). An introductory email containing a letter of invitation, study information sheet, and an expression of interest reply slip was sent to practice managers who cascaded it to the GPs. Reply slips were returned by post or email.


Semi-structured interviews were conducted at a mutually convenient time and place, and face-to-face or by telephone according to interviewee preference. Written consent was obtained

at face-to-face interviews, and recorded verbal consent was obtained from telephone participants. Participants could withdraw from the study at any stage either during or after the interview and any collected data was deleted. Interviews were conducted until data saturation was reached (three successive interviews with no new subthemes identified¹⁶).

Interviews followed a semi-structured topic guide (see Supplementary Appendix S1) based on the COM-B behaviour change model and developed iteratively as interviews progressed. All interviews were digitally recorded, transcribed verbatim, then imported into NVivo (version 12). A Framework approach¹⁷ was used. Analysis followed the steps of familiarisation, identifying a thematic framework, indexing, charting, and mapping/interpreting.

However, although COM-B was the framework, the generation of additional themes outside of the framework and overarching themes across elements of the framework were permitted. Indexing and charting were carried out on full transcripts by two authors. Mapping and interpretation were led by the first author, with contributions to this in meetings with all authors. Analysis was conducted iteratively alongside interviews.¹⁷

Figure 1. Based on the COM-B model.¹³ © 2011 Michie S, et al; licensee BioMed Central Ltd. Reproduced under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>).



Box 1. Summary of themes using the COM-B framework

COM-B component	Theme	Theme details
Capability	Learning	Early training Further training
	Continuing practice	Limited opportunities Loss of capability
	Different forms of capability	Procedural skill Confidence in interpretation
Opportunity	Chaperones	Patient preference Practical availability
	Clinical environment	Enhanced versus generic
Motivation	Doing the best for the individual	Prioritise patient over rules
	Doctors examine	Importance of checking Professional reputation
	GPs as pragmatists	Question added value of PE Referral as acceptable alternative

PE = pelvic examination.

RESULTS

Fifteen GPs (11 female) from nine practices were interviewed. Twelve worked in urban settings and three in rural. Five had advanced training in PE, as evidenced by holding a letter of competence in intrauterine techniques. Time since full GMC registration ranged from 9 to 35 years: two <10 years; three 11–20 years; eight 21–30 years; and two >30 years.

Themes in each of the COM-B components were identified (three within capability, two within opportunity, and three within motivation). In addition, two overarching themes of patient choice and practitioner gender were identified. In reporting the analysis, the authors recognise that some items could be described within multiple COM-B components. For instance, the delivery of much cervical screening and sexual and reproductive health by nurses in primary care reduces GPs opportunities to carry out PEs, and in turn leads to reduced capability through lack of continuing practice. Where this arises, factors have been described under one heading. Box 1 summarises the themes according to the COM-B framework.

Capability

Participants acknowledged that capability was dependent on training opportunities, experience, and exposure in current clinical practice. There were three themes related to capability: learning, continuing practice, and different forms of capability.

Learning. Most interviewees regarded opportunities for learning PE in their

undergraduate training as inadequate. Rotation through obstetrics and gynaecology as part of GP vocational training was considered a valuable component of learning although not all found it useful.

Other than one female interviewee, most did not recall any consolidation or further training of PE skills during GP registrar training; however, new approaches to GP registrar training with required competences listed in an e-portfolio were regarded as beneficial, likely to produce GPs with improved skills, as the following quote illustrates:

'So I think the new trainees coming out are much better than we were, because they've had to prove throughout their e-portfolio that they are doing intimate examinations [...] so they're usually examining and then the patient being re-examined by either a trainer or another colleague.' (GP4, female)

Interviewees were asked whether they thought continuing education in PE should be mandatory or voluntary and there was no consensus. In terms of what learning opportunities might be offered, suggestions included didactic lectures, attending gynaecological outpatient clinics, and use of pelvic models.

Interviewees recognised that capability required both training and continuing practice of PE.

Continuing practice. All GPs described the way that changes to primary care practice (particularly nurse-led cervical screening) had reduced their continuing practice of PEs:

'I think it might be very hard to get day to day experience with gynaecology and you could go weeks [...] where you wouldn't see anything.' (GP 15, female)

Male GPs were seen as less likely to have opportunities for continuing practice of PEs. Women were seen as more likely to choose a female GP for gynaecological problems and some felt that practice receptionists might triage women to female practitioners (GP or nurse) or those perceived as specialists. For example, practitioners trained to fit intrauterine devices were seen as more 'specialised' within the practice and may gain extra experience both by carrying out non-diagnostic PEs as part of device fitting and also by more general awareness of their expertise:

'You'll have one person in the practice who does coils, so that even decreases skill sets for other female GPs, never mind the male GPs.' (GP 14, male)

There was consensus that higher levels of exposure contributed to higher levels of confidence in examination skills and capability, and was determined partly by the sex of the GP, as male GPs were less likely to carry out PEs. The requirement for chaperones (see later section) was perceived to be more problematic for male GPs and patients were perceived to self-select female GPs when presenting with gynaecological symptoms, as in UK primary care, patients can select their clinician. Those certified to fit intrauterine devices felt more skilled and confident because of increased non-diagnostic exposure, but this had an impact on other GPs' exposure levels.

Different forms of capability. Perceptions of capability varied across the different elements of PE. Visualisation of the cervix was considered an essential part of the examination by all interviewees, although one male GP stated using a speculum would 'cause him alarm'. However, several interviewees were concerned that they had insufficient capability to accurately identify cervical abnormalities. Bimanual examination was considered more difficult and subjective in interpretation. Its diagnostic value was questioned with concern about both false-negative and false-positive examinations. Most felt that overt changes would be correctly identified, but subtle changes were more likely to be misinterpreted or missed. No one mentioned examination of the vulva, the third component of PE.

In summary, capability in PE was seen as requiring adequate training and learning, continuing practice, and the ability not just to carry out examination but to interpret it accurately.

Opportunity

Two themes related specifically to opportunity: chaperones and clinical environment.

Chaperones. Beliefs about the use of chaperones varied between GPs as did their actual use of chaperones; however, similar beliefs did not always result in the same behaviour.

All GPs felt the need for chaperones was an important obstacle to opportunity. All were aware of guidance from the Royal College of

General Practitioners and the GMC. Although most GPs routinely offered chaperones, some did not. Although the offer of a chaperone was made, some felt the guidance did not benefit patients and were happy to carry out the examination without one:

'It [using a chaperone] makes the whole thing less natural, and more awkward [...] it's embarrassing enough for them to have me looking at their down belows, without somebody else.' (GP7, female)

Some female participants did not always offer a chaperone. Many said they had never had a patient request a chaperone:

'I'd say, are you okay if I examine you, and if they say yes, and they jump up on the bed, then that's good enough for me.' (GP4, female)

Clear sex differences were observed regarding the belief that chaperones were essential, but one male GP indicated a situation where they would not always use a chaperone:

'I'll do it on a post-menopausal lady [...] or I'll say if you've had your kids, and you're comfortable with me doing it, and quite often they'll say yes [...] I'm exposing myself to some risk, but I think it's pretty low, but anybody my age or younger certainly not.' (GP13, male)

This behaviour was driven by assumptions the GP had about patients' beliefs:

'Elderly ladies [...] I could be wrong, but they say this is just a procedure that needs to be done, I'm sure younger women do, but I do it, to protect myself, rather than them.' (GP13, male)

This need to protect themselves from potential litigation by offering a chaperone was also expressed by female GPs, despite some feeling that offering a chaperone was a tick-box exercise:

'I think that it's one of these tick box sentences, that feels like, it feels so stupid to ask it [...] do you want someone come in with you, they [the patient] look at you, like, what, what?' (GP5, female)

Female GPs also acknowledged not using a chaperone contradicted the guidance, but it was normal practice for several participants. This decision was influenced by familiarity with or age of the patient. The majority of

those who offered a chaperone did not expect the patient to want one. Some GPs who routinely did not use a chaperone would suggest a chaperone if the patient made them feel uncomfortable, with GPs of both sexes acknowledging their vulnerability when carrying out PEs:

'I do offer the patient one, but if they don't wish one [...] I should get one anyway, for my own defence [...] because I often know the patients quite well, I, I chose to take that risk.' (GP8, female)

The guidance states chaperones should preferably be a health professional and familiar with PE procedures, however, participants commented that practice staff with these attributes are often engaged in their own clinical practice and not free to chaperone. One GP's practice had trained reception staff to be chaperones to increase availability. However, for some practices even larger obstacles were present:

'We have a, a satellite surgery in [location] where it's just us, you know, there's no one, no admin or anybody.' (GP15, female)

Only one GP explicitly described the value of chaperones for those patients who were especially nervous about the examination.

Clinical environment. Participants described a number of issues relating to the clinical environment that either facilitated or obstructed the conduct of PEs. These included time and equipment such as adjustable examination couches. Interviewees gave examples of where these were absent. Routine consultations of 10 min were seen as insufficient for an adequate history and examination when compounded by the need to find a chaperone with no notice and ensure sufficient dignity for the patient. In contrast, designated appointments with appropriate time and equipment, especially where the practitioner had been selected for their expertise (for instance, by referral within the practice between clinicians) were seen as facilitating PEs. The issue of in-practice referral, especially from a male GP to female was discussed by several participants and was generally felt to be appropriate.

Motivation

Interviewees described a range of motivations towards or against PEs that were independent of either capability or opportunity. Importantly all related to providing high-quality care, but this was defined in different ways. These are

categorised broadly as three themes: 'doing what is best for the individual', 'doctors examine', and 'GPs as pragmatists'.

Doing what is best for the individual. Some GPs made it clear that there were few hard and fast rules about when PEs should be carried out and that the highest priority in clinical decisions was the patient:

'We should be tailoring the investigation to the individual, with respect for their dignity, and asking the question, what will this investigation give me?' (GP3, female)

'It's an intimate examination, it should only be done if it's going to add to your, to your diagnosis, or management.' (GP8, female)

However, some GPs assumed that patients did not want to be examined because of embarrassment. They also assumed that patients would not wish to be examined by students, resulting in reduced learning opportunities.

Doctors should examine. Some participants emphasised the importance of clinical examination as something that doctors do. Thus, although the intimacy of PE should be acknowledged, this should not prevent it:

'I think it would be embarrassing to miss something for the want of, of their embarrassment or our discomfort at having to do it [...] because if we miss that then, you know, they're on a completely different path.' (GP6, female)

Some participants acknowledged that lack of examination could lead to misattribution of symptoms:

'I guess even post-menopausal bleeding for query endometrial [cancer], usually I would examine them, just to make sure if there was anything local.' (GP8, female)

For some, not carrying out examination before referral, even if it was unlikely to add clinical value, was seen as unprofessional:

'You wouldn't put [refer] someone to cardiology, if you hadn't listened to their chest, they would laugh at you.' (GP2, female)

GPs as pragmatists. Some GPs explicitly questioned the diagnostic value of PE by a non-specialist. For instance, several argued that imaging offered more clinical yield than

clinical examination and that they would refer for ultrasound scanning rather than use clinical examination for diagnosis, especially for a patient with post-menopausal bleeding.

For many women presenting with symptoms that might be indicative of gynaecological cancer, some GPs argued that a decision to refer could be made on history alone. Once 'red flag' features had been identified in the history, examination findings would not positively have an impact on triaging of the referral by the specialist, or the patient's diagnostic journey. One GP argued that although a rectal examination was clinically useful for a suspected rectal cancer, a PE was less useful for at least some types of suspected gynaecological cancer:

'In the past couple of months, two women with post-menopausal bleeding, I didn't even bother to examine them [...] they need [to be] referred [...] I'm not going to find anything of value [...] I can see the utility of the, the rectal exam, in you know, for rectal cancer, because if you feel the tumour [...] that's going to potentially change the initial management.' (GP14, male)

This argument was often seen in GPs who described low capability and opportunity (and thus, for whom the probability of a well conducted and interpreted examination was likely to be lower).

Although GPs recognised clinical guidelines for urgent referral of patients with suspected cancer they described their limitations, especially when advice changed:

'I just have a gut feeling there's something not right, they don't necessarily fit a guideline, but they need [to be] referred.' (GP2, female)

Pragmatic GPs also saw limitations to the value of providing extra information at the time of referral:

'My referral letters tend to be quite short and to the point, so maybe mine are read more, but I know I've signed letters for some people who just write screeds of stuff, and think what's the point, because who's going to read this at the end?' (GP7, female)

Integrating findings

While the analysis has been presented by deconstructing interviews into the COM-B framework, it is clear that a number of different factors influence GPs' approach to carrying out PEs. These are presented in Box 2, which summarises the findings in a set of conditional statements, each of which makes PEs more or less likely

in the GP consultation. This has value in two ways: increasing understanding and planning service innovation. In terms of understanding, it makes it apparent why factors (such as male sex of the GP) may have effects at multiple levels of this integrated model. In terms of service innovation, it suggests that action at a single level is unlikely to have a substantial impact without addressing others.

DISCUSSION

Summary

The reasons why GPs do, or do not, use PEs in assessing gynaecological symptoms that might indicate cancer are complex. The COM-B model provides a way to understand GPs behaviour that shows how contrasting views about the value and practice of examination depend on factors relating to all three components: capability, opportunity, and motivation. A framework that integrates these findings allows these factors to be considered together and could be used both in understanding individual decisions and in planning training or innovation in the way services are delivered.

Strengths and limitations

To the authors' knowledge, this is the first study of its kind that explores GPs' perceptions of PE using the COM-B behaviour change model as the analytical framework. Analysis was carried out by all researchers: three GPs, a pharmacist, and a consultant gynaecologist. Preconceptions of researchers can be a challenge, but the multidisciplinary approach helped to mitigate this.

There were, however, some limitations. Study participants knew the interviewer (the first author) was a GP, potentially influencing responses. As with all exploratory studies the results are not intended to be generalisable, but interviewee selection should aim to recruit a range of characteristics likely to affect views and experiences. Undertaking the study in one health board in Scotland and the self-selection of GPs agreeing to be interviewed may mean there are other views that have not been identified.

Comparison with existing literature

Existing literature is limited. Paluska and D'Amico presented data in 2000 confirming doctors are more comfortable examining patients of their own sex.¹⁸ They assumed this was because male doctors felt patients would be embarrassed; however, questioning patients has revealed that despite feeling nervous about PEs,

Box 2. Summary of conditions that make GP PE more or less likely

	Conditions	PE in the GP consultation more likely	PE in the GP consultation less likely
Capability	GPs who ...	are well trained in PE have a substantial PE caseload to maintain confidence in their skills	have not had detailed training in PE have a limited PE caseload and less confidence in their skills
Opportunity	GPs who practice in where chaperone needs and where patients with gynaecological symptoms ...	a clinical environment which has been enhanced to provide PE are easily met choose, or are directed, to see them because of their expertise	a generic clinical environment are at best hard to meet have little choice in who they see
Motivation	GPs who believe that by carrying out PE ...	they commonly add important value to the clinical process before or at the time of referral	they commonly add no value to a decision to refer either for imaging or specialist assessment
Behaviour	Outcome	GP is more likely to offer and conduct PE before referral	GP is less likely to offer or conduct PE, but, instead to either bypass PE (by direct referral) or to arrange for PE by a practice colleague

PE = pelvic examination.

patients regard it as necessary.¹⁹ Although patients report being exposed during the examination, they also describe how clinicians can influence this discomfort. Good communication during the examination, a trusted relationship with the clinician, and a comfortable environment can reduce patient distress.¹⁹⁻²¹ Clinician discomfort can increase patient's distress.²¹ Clinician discomfort can also influence their use of chaperones. Reflecting the findings in the current study, discordance between clinical practice and guideline recommendations has been observed previously and has been quantified; whereas 72% of clinician's thought patients should be offered a chaperone, only 27% of the study population used chaperones.²² Another study suggested that guidance on the use of chaperones should be flexible, taking into consideration the relationship between patients and their doctors while considering practice staffing and space;²³ this pragmatic approach was observed in this study.

Chaperone use was acknowledged as a barrier in terms of resource. But there was also recognition of clinical vulnerability and potential litigation while acknowledging that patients generally did not want chaperones. The use of chaperones

to alleviate feelings of vulnerability to false allegations or to sexualisation of the examination by the patient has been reported by Hine and Smith.²⁴

Implications for research and practice

The identification in the current study of important factors in all three elements of the COM-B framework indicates that interventions to change practice either through training or service delivery innovations need to address multiple components. As suggested in Box 2, changing one element without addressing others is likely to have limited effect. It is therefore important to consider all three components.

Increasing capability could involve more skills training with the use of pelvic mannequins; however, these are not anatomically representative of many women and do not allow the development of communication skills that are key in reducing patient concerns. An alternative could be gynaecological teaching associates (GTA). A GTA is a non-doctor woman who has been trained to teach PE using their own bodies.²⁵ They are able to develop a patient-centred approach while removing the homogeneous depictions of women in mannequins. Their value in improving medical student education is well established,²⁶ but it is not clear how effective they are for maintaining skills after initial training.

Several practices described ways of delivering care that addressed opportunity, for instance, supporting patient choice of, or explicitly directing patients to, GPs with greater experience. Further consideration is required to determine how patients access those confident GPs at the earliest possible opportunity. Ten-minute consultations and difficulties in locating appropriate chaperones clearly reduce opportunity. Increased length of consultation has also been highlighted by the Royal College of Obstetricians and Gynaecologists report, 'Better for Women', which proposes 15-min GP consultations.²⁷ Additionally, primary care has changed its consultation model because of the COVID-19 pandemic. The authors do not know in what way this has affected how clinicians carry out PEs.

Motivation factors about PE are clearly complex. GPs remain unsure about how valuable examination findings are to a good gynaecological referral. This could be addressed in further database studies, but as a starting point, gynaecologists should be asked whether they value PEs carried out in primary care. More research

is also needed to understand how much GPs' assumptions about what women prefer match to the reality.

The unheard voice in this research is of women themselves. While the authors have knowledge of patients' views on PE, the authors do not know how this influences their help-seeking behaviour. Delayed help seeking is associated with poorer outcomes.²⁸ Research is needed to

understand how PE influences patients' help-seeking behaviour.

In conclusion, GPs' reasons for using, or not using, PE in women with symptoms potentially indicating cancer are complex. The COM-B framework provides a way of understanding this complexity. Interventions to increase the use of PE, and critics of its non-use, need to consider these multiple factors.

Funding

Pauline Williams completed this work during a clinical academic fellowship funded by the Chief Scientist's Office of the Scottish Government (grant reference: RG13111-10).

Ethical approval

The study was approved by the North of Scotland Research Ethics Committee on the 26 September 2017 (reference: 17/NS/0094), and by NHS Research and Development on 18 October 2017 (reference: NRS17/213808).

Provenance

Freely submitted; externally peer reviewed.

Competing interests

The authors have declared no competing interests.

Acknowledgements

The authors would like to thank the Chief Scientist Office, Scotland, who funded Pauline Williams through a Clinical Academic Training Fellowship.

Open access

This article is Open Access: CC BY 4.0 licence (<http://creativecommons.org/licenses/by/4.0/>).

Discuss this article

Contribute and read comments about this article: bjgp.org/letters

REFERENCES

1. Russi EW. [Physical examination of the lung--value and limits]. [Article in German]. *Ther Umsch* 2006; **63(7)**: 471–474.
2. Yu CK, Chiu C, McCormack M, *et al*. Delayed diagnosis of cervical cancer in young women. *J Ob Gynae* 2005; **25(4)**: 356–370.
3. Macleod U, Mitchell ED, Burgess C, *et al*. Risk factors for delayed presentation and referral of symptomatic cancer: evidence for common cancers. *Br J Cancer* 2009; **101(suppl 2)**: S92–S101.
4. Vandborg MP, DePont Christensen R, Kragstrup J, *et al*. Reasons for diagnostic delay in gynecological malignancies. *Int J Gynecol Cancer* 2011; **21(6)**: 967–974.
5. Bernet R, Buddeberg C. Physician–patient relations in gynecology: expectations and experiences of patients. *Gynakol Geburtshilfliche Rundsch* 1994; **34(11)**: 43–49.
6. Domar AD. Psychological aspects of the pelvic exam: individual needs and physician involvement. *Women Health* 1985; **10(4)**: 75–90.
7. Yanikkerem E, Özdemir M, Bingol H, *et al*. Women's attitudes and expectations regarding gynaecological examination. *Midwifery* 2009; **25(5)**: 500–508.
8. Hilden M, Sidenius K, Langhoff-Roos J, *et al*. Women's experiences of the gynecologic examination: factors associated with discomfort. *Acta Obstet Gynecol Scand* 2003; **82(11)**: 1030–1036.
9. General Medical Council. *Intimate examinations and chaperones (summary)*. 2013. <https://www.gmc-uk.org/ethical-guidance/ethical-guidance-for-doctors/intimate-examinations-and-chaperones> (accessed 15 Mar 2023).
10. Hjörleifsson S, Bjorvatn B, Meland E, *et al*. The when and how of the gynaecological examination: a survey among Norwegian general practitioners. *Scand J Prim Health Care* 2019; **37(2)**: 264–270.
11. Ladouceur R. Recommendations for the routine screening pelvic examination: could they have a negative effect on physician competence? *Can Fam Physician* 2016; **62(6)**: 460.
12. Saleh N, Abu-Gariba M, Yehoshua I, *et al*. Barriers to implementation of a pelvic examination among family doctors in primary care clinics. *Postgrad Med* 2018; **130(3)**: 341–347.
13. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011; **6**: 42.
14. West R, Michie S. A brief introduction to the COM-B model of behaviour and the PRIME theory of motivation. *Qeios* 2020; DOI: 10.32388/WW04E6.
15. McDonagh LK, Saunders JM, Cassell J, *et al*. Application of the COM-B model to barriers and facilitators to chlamydia testing in general practice for young people and primary care practitioners: a systematic review. *Implement Sci* 2018; **13(1)**: 130.
16. Francis JJ, Johnston M, Robertson C, *et al*. What is an adequate sample size? Operationalising data saturation for theory-based interview studies. *Psychol Health* 2010; **25(10)**: 1229–1245.
17. Marks DF, Yardley L. Content and thematic analysis. In: Marks DF, Yardley L, eds. *Research Methods for Clinical and Health Psychology*. London: SAGE, 2004.
18. Paluska SA, D'Amico FJ. The comfort of family practice residents with health care of patients of the opposite gender. *Fam Med* 2000; **32(9)**: 612–617.
19. Larsen M, Oldeide CC, Malterud K. Not so bad after all... Women's experiences of pelvic examinations. *Fam Pract* 1997; **14(2)**: 148–152.
20. Cook C. 'About as comfortable as a stranger putting their finger up your nose': speculation about the (extra)ordinary in gynaecological examinations. *Cult Health Sex* 2011; **13(7)**: 767–780.
21. Oscarsson M, Benzein E. Women's experiences of pelvic examination: an interview study. *J Psychosom Obstet Gynecol* 2002; **23(1)**: 17–25.
22. Guidozzi Y, Gardner J, Dhali A. Professionalism in the intimate examination: how healthcare practitioners feel about having chaperones present during an intimate consultation and examination. *SAfr Med J* 2013; **108(1)**: 25–27.
23. Price DH, Shawn Tracy C, Upshur REG. Chaperone use during intimate examinations in primary care: postal survey of family physicians. *BMC Fam Pract* 2005; **6**: 52.
24. Hine P, Smith H. Attitudes of UK doctors to intimate examinations. *Cult Health Sex* 2014; **16(8)**: 944–959.
25. Beckmann CRB, Barzansky BM, Sharf BF, *et al*. Training gynaecological teaching associates. *Med Educ* 1988; **22(2)**: 124–131.
26. Duffy JMN, Chequer S, Braddy A, *et al*. Educational effectiveness of gynaecological teaching associates: a multi-centre randomised controlled trial. *BJOG* 2016; **123(6)**: 1005–1010.
27. Royal College of Obstetricians and Gynaecologists. *Better for women: improving the wellbeing of girls and women*. 2019. <https://www.rcog.org.uk/media/h3smwohw/better-for-women-full-report.pdf> (accessed 15 Mar 2023).
28. Allemani C, Matsuda T, di Carlo V, *et al*. Global surveillance of trends in cancer survival 2000–14 (CONCORD-3): analysis of individual records for 37 513 025 patients diagnosed with one of 18 cancers from 322 population-based registries in 71 countries. *Lancet* 2018; **391(10125)**: 1023–1075.