

This is a repository copy of *On Spin Calogero-Moser system at infinity*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/200163/>

Preprint:

Sklyanin, Evgeny orcid.org/0000-0003-2194-0643, Khoroshkin, Sergey and Matushko, M. G. (2016) On Spin Calogero-Moser system at infinity. [Preprint]

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

On Spin Calogero-Moser system at infinity

S.M. Khoroshkin^{*°}, M.G. Matushko[°] and E.K. Sklyanin^{*}

^{*}*Institute for Theoretical and Experimental Physics, Moscow 117259, Russia;*

[°]*National Research University Higher School of Economics, Myasnitskay Ulitsa, Moscow 101000, Russia;*

^{*}*Department of Mathematics, University of York, York YO10 5DD, United Kingdom*

Abstract

We present a construction of a new integrable model as an infinite limit of Calogero models of N particles with spin. It is implemented in the multicomponent Fock space. Explicit formulas for Dunkl operators, the Yangian generators in the multicomponent Fock space are presented. The classical limit of the system is examined.

1 Introduction

The Calogero-Moser-Sutherland (CMS) model of N bosonic particles with spins has two properties: integrability and the presence of an algebra of symmetries [6, 9]. The symmetries provide a representation of the Yangian $Y(\mathfrak{gl}_s)$. Both of these properties relate to the Dunkl operators: the integrals of motion may be expressed through its symmetric combinations, and the corresponding representation of the Yangian uses the same Dunkl operators and N copies of the algebra \mathfrak{gl}_s . It is natural to develop analogues of the CMS model for systems with an infinite number of particles. Such systems were defined and studied by I. Andric, A. Jevicki and H. Levine [1], H. Awata, Y. Matsuo, S. Odake, J. Shiraishi [3] and others in the framework of “collective field theory” in the spinless case and by H. Awata, Y. Matsuo and T. Yamamoto [5] for models with spins.

Effective and rigorous constructions of limits of quantum Calogero-Sutherland (CS) systems have attracted the attention of mathematicians for many years. Note first the fundamental research of D.Uglov [12], where he defined and studied a fermionic inductive limit of the CS system. His construction waited for more than 15 years for a further development until M. Nazarov and E.Sklyanin suggested a precise construction of higher Hamiltonians for an Uglov type limit of the scalar CS system using the Sekiguchi determinant and the machinery of symmetric functions [7]. In [10] A.Veselov and A.Sergeev suggested to define a bosonic limit of the CS system as a projective limit of finite models. Precise bosonic constructions of higher Hamiltonians in a Fock space were then presented by M.Nazarov and E.Sklyanin in [8] and by A.Veselov and A.Sergeev in [11]. It was also shown in [8] that the classical limit of the infinite system can be identified with the Benjamin-Ono hierarchy whose Lax presentation can

be derived from the suggested description of the quantum system. Moreover, the equivariant family of Heckman–Dunkl operators can be regarded as a quantum L -operator for the CS system.

This paper can be regarded as a extension of the latter ideas to spin CMS models. Our motivation to perform such an exercise was to reproduce an expression for the first Hamiltonian given in [5] and to find a regular way to obtain similar expressions for the Yangian generators and higher Hamiltonians of the systems. We first perform a translation of the finite spin CMS model to the language of polysymmetric functions and then realize the projective type limit of the spin CMS model in the multicomponent Fock space. Here we present explicit formulas for the Dunkl operators, and the L -operator, which are then used for construction of the Yangian generators and the commuting Hamiltonians. Note that almost the same formulas are valid in the finite case as well. In particular, we reproduce an expression for the first Hamiltonian from [5]. The answers are given now by integral operators with kernels composed of products of vertex operators.

We study the classical limit of the obtained system by sending the coupling constant to infinity. In the resulting model the Hamiltonians are constructed, and the equations of motion are presented. The classical system, which may be regarded as a multicomponent version of the Benjamin-Ono hierarchy, is integrable. It possesses an infinite number of commuting integrals of motion. Moreover, it is possible to express the equations of motion in Lax form: we present an explicit expression for the Lax pair.

2 CMS model and related algebraic structures

2.1 Calogero-Sutherland model

Consider the quantum Calogero-Sutherland spin model of N bosonic particles on the circle [6]. Its Hamiltonian is

$$H^{CS} = - \sum_{i=1}^N \left(\frac{\partial}{\partial q_i} \right)^2 + \sum_{i,j=1}^N \frac{\beta(\beta-1)}{\sin^2(q_i - q_j)}.$$

After conjugating by the function $|\prod_{i<j} \sin(q_i - q_j)|^\beta$ which represents the vacuum state, and passing to the exponential variables $x_i = e^{2\pi i q_i}$ we come to the Hamiltonian

$$H = \sum_{i=1}^N \left(x_i \frac{\partial}{\partial x_i} \right)^2 + \beta \sum_{i<j} \frac{x_i + x_j}{x_i - x_j} \left(x_i \frac{\partial}{\partial x_i} - x_j \frac{\partial}{\partial x_j} \right) - 2\beta \sum_{i<j} \frac{x_i x_j}{(x_i - x_j)^2} (1 - K_{ij}), \quad (1)$$

where the operator K_{ij} permutes the variables x_i and x_j . The wave function of the Hamiltonian depends on positions x_i of the particles and of their spins $\sigma(i)$, which take values in the set $\{1, 2, 3, \dots, s\}$. We will consider polynomial wave functions invariant under the exchanges

$$\psi \left(\dots, x_i^{(a)}, \dots, x_j^{(b)}, \dots \right) = \psi \left(\dots, x_j^{(b)}, \dots, x_i^{(a)}, \dots \right), \quad (2)$$

where the notation $x_i^{(a)}$ means that i -th particle has the position x_i and spin $a = 1, \dots, s$. The Dunkl operators

$$d_i = x_i \frac{\partial}{\partial x_i} + \beta \sum_{j < i} \frac{x_j}{x_i - x_j} (1 - K_{ij}) + \beta \sum_{i < j} \frac{x_i}{x_i - x_j} (1 - K_{ij}) + \beta (i - 1) \quad (3)$$

act on the space of wave functions and respect the condition (2), see Section 2.2 for more details. These operators commute

$$[d_i, d_j] = 0$$

and give the expression of the Hamiltonian

$$H = \sum_i (d_i^2 - \beta d_i),$$

which is regarded as a linear operator on the space of functions with property (2). The higher commuting Hamiltonians, which act on the same space have the form

$$H_n = \sum_i d_i^n. \quad (4)$$

2.2 Yangian and degenerate affine Hecke algebra

Denote by M_N the space of vector-valued polynomials

$$M_N = \underbrace{(\mathbb{C}^s \otimes \mathbb{C}[z]) \otimes \dots \otimes (\mathbb{C}^s \otimes \mathbb{C}[z]) \otimes (\mathbb{C}^s \otimes \mathbb{C}[z])}_N. \quad (5)$$

Then the space of the model can be identified with the space of S_N -invariants in M_N denoted by $M_N^{S_N}$, where the symmetric group exchanges both the variables and vector components in the tensor product. The spaces of spins and of coordinates are denoted by \mathbb{C}^s and $\mathbb{C}[z]$, respectively.

More precisely, we fix a basis $\{e^1, e^2, \dots, e^s\}$ in the spin space \mathbb{C}^s . The operators K_{ij} of permutation of the coordinates, P_{ij} of permutation of the spins, and $\sigma_{ij} = K_{ij}P_{ij}$ of the corresponding total action of the symmetric group S_n can be expressed by the following formulas

$$K_{ij} : (\dots \otimes (e^{a_i} \otimes x^{k_i}) \otimes \dots \otimes (e^{a_j} \otimes x^{k_j}) \otimes \dots) \rightarrow (\dots \otimes (e^{a_i} \otimes x^{k_j}) \otimes \dots \otimes (e^{a_j} \otimes x^{k_i}) \otimes \dots)$$

$$P_{ij} : (\dots \otimes (e^{a_i} \otimes x^{k_i}) \otimes \dots \otimes (e^{a_j} \otimes x^{k_j}) \otimes \dots) \rightarrow (\dots \otimes (e^{a_j} \otimes x^{k_i}) \otimes \dots \otimes (e^{a_i} \otimes x^{k_j}) \otimes \dots)$$

$$\sigma_{ij} : (\dots \otimes (e^{a_i} \otimes x^{k_i}) \otimes \dots \otimes (e^{a_j} \otimes x^{k_j}) \otimes \dots) \rightarrow (\dots \otimes (e^{a_j} \otimes x^{k_j}) \otimes \dots \otimes (e^{a_i} \otimes x^{k_i}) \otimes \dots)$$

Any element in $M_N^{S_N}$ is invariant under the action of the symmetric group $\sigma_{ij}\psi = \psi$ by definition. Hence,

$$K_{ij}\psi = P_{ij}\psi, \quad \psi \in M_N^{S_N}.$$

Next we describe the representation of the degenerate affine Hecke algebra and Lie algebra \mathfrak{gl}_s in the space M_N . The Heckman–Dunkl operators $D_i^{(N)} : M_N \rightarrow M_N$ are given by

$$D_i^{(N)} = x_i \frac{\partial}{\partial x_i} + \beta \sum_{j \neq i} \frac{x_i}{x_i - x_j} (1 - K_{ij}). \quad (6)$$

These operators do not change spins and satisfy the relations

$$K_{ij} D_i^{(N)} = D_j^{(N)} K_{ij}, \quad (7)$$

$$[D_i^{(N)}, D_j^{(N)}] = \beta (D_j^{(N)} - D_i^{(N)}) K_{ij}, \quad (8)$$

which coincide with the relations of the degenerate affine Hecke algebra after the renormalization $D_i^{(N)} = \beta \tilde{D}_i$. One can use another set of commuting elements d_i of the degenerate affine Hecke algebra:

$$d_i = D_i^{(N)} + \beta \sum_{j < i} K_{ij}, \quad (9)$$

so that

$$[d_i, d_j] = 0, \quad K_{i,i+1} d_i = d_{i+1} K_{i,i+1} - \beta.$$

The action of elements d_i in the space M_N is given by equation (3). The center of the degenerate affine Hecke algebra is represented by symmetric functions of d_i . Moreover, symmetric polynomials of d_i represent integrals of motion of the quantum Calogero Sutherland model (4).

The spin Calogero–Sutherland model admits Yangian symmetry, namely there is an action of the Yangian on the space $M_N^{S_N}$ which commutes with the Hamiltonians (4). Further we recall the definition of the Yangian and describe its action on space $M_N^{S_N}$. Let

$$R_{12}(u - v) = 1 - \frac{1}{u - v} \sum_{a,b} E^{ab} \otimes E^{ba} \in \text{End}(\mathbb{C}^s \otimes \mathbb{C}^s),$$

where $E^{ab} \in \text{End} \mathbb{C}^s$ is the matrix unit, $E^{ab}(e^c) = \delta_{bc} e^a$ for basic vectors $e^c \in \mathbb{C}^s$. By definition, the Yangian $Y(\mathfrak{gl}_s)$ is a unital associative algebra over \mathbb{C} with generators $t_{(i)}^{ab}$, $a, b = 1, \dots, s$, $i = 0, 1, \dots$ subject to the relations encoded in the Yang–Baxter equation

$$R_{12}(u - v) t_1(u) t_2(v) = t_2(v) t_1(u) R_{12}(u - v), \quad (10)$$

where $t_1(u) = t(u) \otimes \text{Id}$, $t_2(u) = \text{Id} \otimes t(u)$, and

$$t(u) = \sum_{a,b} t^{ab}(u) \otimes E^{ab} \in Y(\mathfrak{gl}_s) \otimes \text{End}(\mathbb{C}^s), \quad t^{ab}(u) = \delta^{ab} + \sum_{i=0}^{\infty} t_{(i)}^{ab} \frac{1}{u^{i+1}}.$$

Equivalently, the relations (10) are

$$[t^{ab}(u), t^{cd}(v)] = \frac{t^{cb}(u) t^{ad}(v) - t^{cb}(v) t^{ad}(u)}{u - v}.$$

We have an action of N copies of the Lie algebra \mathfrak{gl}_s on M_N :

$$E_i^{ab} \left(\cdots \otimes \underbrace{(e^c \otimes x^k)}_i \otimes \cdots \right) = \delta_{bc} \left(\cdots \otimes \underbrace{(e^a \otimes x^k)}_i \otimes \cdots \right).$$

The global action of \mathfrak{gl}_s on M_N is denoted by $\mathbb{E}^{ab} = \sum_i E_i^{ab}$.

We have a representation of the Yangian $Y(\mathfrak{gl}_s)$ on the space $M_N^{S_N}$. We use the additive version of the Drinfeld functor to define the Yangian action [2]:

$$t^{ab}(u) = \delta_{ab} + \beta \sum_i \frac{E_i^{ab}}{\beta u + D_i^{(N)}} \quad (11)$$

on the space $M_N^{S_N}$. The following result is well known [4]:

Theorem *The representation (11) of Yangian $Y(\mathfrak{gl}_s)$ commutes with the Hamiltonians of the CS system. The algebra of higher Hamiltonians coincides with the center of the Yangian, which is generated by the coefficients of the quantum determinant*

$$q \det t(u) = \sum_{\sigma \in S_m} (-1)^{\text{sgn}(\sigma)} t^{\sigma(1),1}(u) t^{\sigma(2),2}(u-1) \dots t^{\sigma(m),m}(u-m+1). \quad (12)$$

3 CS system in the limit of infinitely many particles

3.1 Review of the scalar system

Here we review recent results on the scalar CS system [8, 11] mainly following the approach of [8]. The main idea is to regard the equivariant Heckman-Dunkl operators as a quantum L-operator acting on the space of polynomial functions of one variable with coefficients being symmetric polynomials of the remaining $N-1$ variables.

In the scalar CS model the action of the Hamiltonian (1) on symmetric functions can be reduced to

$$H = \sum_{i=1}^N \left(x_i \frac{\partial}{\partial x_i} \right)^2 + \beta \sum_{i < j} \frac{x_i + x_j}{x_i - x_j} \left(x_i \frac{\partial}{\partial x_i} - x_j \frac{\partial}{\partial x_j} \right). \quad (13)$$

The corresponding Heckman–Dunkl operators $D_i^{(N)} : \mathbb{C}[x_1, \dots, x_N] \rightarrow \mathbb{C}[x_1, \dots, x_N]$ are defined by (6). Clearly, symmetric functions of $D_i^{(N)}$ preserve the ring of symmetric polynomials $\Lambda_N = \mathbb{C}[x_1, \dots, x_N]^{S_N}$. This algebra is generated by the Newton polynomials $p_k^N = x_1^k + \dots + x_N^k$, $k \in 0, 1, \dots, N$ (sometimes we omit the upper index N and simply write p_k). The Dunkl operator $D_i^{(N)}$ itself preserves the symmetry involving all variables other than x_i and therefore it acts on the space $\Lambda_{N,i}$ of functions symmetric in all variables except x_i :

$$F(x_i, \{p_n\}) \in \Lambda_{N,i} \simeq \mathbb{C}[x_i] \otimes \mathbb{C}[x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N]^{S_{N-1}}. \quad (14)$$

In the following we use the notation

$$\tilde{V}(z) = \exp \left(\sum_{n>0} z^n \frac{\partial}{\partial p_n} \right). \quad (15)$$

for the linear map, which changes each occurrence of a Newton sum p_k^N by $p_k^{N-1} + z^k$.

Let $F \in \Lambda_N$ be a symmetric function of $\{p_n\}$ depending on N variables.

Lemma 1 *The natural embedding $\iota_{N,i} : \Lambda_N \rightarrow \Lambda_{N,i}$ is given by*

$$\iota_{N,i}(F) = \tilde{V}(x_i)F. \quad (16)$$

Here $\tilde{V}(x_i)F$ is a function of x_i and $\{p_n\}$ depending on $(N-1)$ variables.

Proof. The embedding $\iota_{N,i}$ can be regarded as the presentation of a symmetric function F by a polynomial in x_i with coefficients being symmetric functions of the remaining variables:

$$\begin{aligned} F(\{p_n\}) &= F_0(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N) + F_1(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N)x_i + \\ &\quad F_2(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N)x_i^2 + \dots \end{aligned}$$

This expansion can be obtained by means of a substitution

$$p_n^N \rightarrow p_n^{N-1} + x_i^n \quad (17)$$

which in its turn can be obtained by applying the vertex operator (15) due to the Taylor formula

$$f(z+t) = \exp\left(t \frac{\partial}{\partial z}\right) f(z) = f(z) + f'(z)t + \frac{1}{2}f''(z)t^2 + \dots$$

which yields a finite sum for polynomials. Observe that the formula (16) is correct for any expression of the symmetric function in terms of Newton polynomials p_n irrespective of their dependencies. ■

Let $\phi^\pm(\xi)$ be the following power series in $\xi^{\pm 1}$:

$$\phi^+(\xi) = \sum_{n=1}^{\infty} \xi^n \frac{n \partial}{\partial p_n}, \quad \phi^-(\xi) = \sum_{n=0}^{\infty} \frac{p_n}{\xi^n}, \quad (18)$$

where $p_0 = N$. We also use the notation

$$\tilde{V}'(z) = \exp \left(\sum_{n>0} -z^n \frac{\partial}{\partial p_n} \right). \quad (19)$$

By definition the operator $\tilde{V}'(z)$ changes each occurrence of the formal variable p_k^{N-1} by the difference $p_k^N - z^k$. The operator $\tilde{V}'(x_i)$ maps the space $\Lambda_{N,i}$ to $\Lambda_N \otimes \mathbb{C}[x_i]$. Note that

$$\tilde{V}'(z)\tilde{V}(z)F = F \quad \forall F \in \Lambda_N. \quad (20)$$

Define the operator $\mathcal{E} : \Lambda_{N,i} \rightarrow \Lambda_N$ of summation over i . For any $F(x_1, \dots, x_{i-1}; z; x_{i+1}, \dots, x_N) \in \Lambda_{N,i}$:

$$\begin{aligned} \mathcal{E}F(x_1, \dots, x_N) = & F(x_i, \dots, x_{i-1}; x_1; x_{i+1}, \dots, x_N) + F(x_1, x_i, \dots, x_{i-1}; x_2; x_{i+1}, \dots, x_N) + \dots \\ & + F(x_1, \dots, x_{i-1}; x_N; x_{i+1}, \dots, x_i). \end{aligned}$$

Lemma 2 *Let $F(x_i; \{p_n\}) \in \Lambda_{N,i}$. Then*

$$\mathcal{E}F(\{p_n\}) = \oint \frac{d\xi}{\xi} \phi^-(\xi) \left(\tilde{V}'(\xi) F \right) (\xi; \{p_n\}).$$

Here on the RHS the function $F(\xi; \{p_n\})$ depends on symmetric functions of $(N-1)$ variables, while $F(\{p_n\})$ on the LHS and $\tilde{V}'(\xi)F$ on the RHS both depend on symmetric functions on N variables.

Lemma 3 *The action of the Dunkl operator $D_i^{(N)}$ on functions $F(x_i, \{p_n\}) \in \Lambda_{N,i}$ can be expressed by the following formula:*

$$D_i^{(N)}(F(x_i, \{p_n\})) = x_i \frac{\partial}{\partial x_i} F(x_i, \{p_n\}) + \beta x_i \oint \frac{d\xi}{\xi^2} \frac{\phi^-(\xi) - 1}{1 - \frac{x_i}{\xi}} \left(\tilde{V}'(\xi) \tilde{V}(x_i) F \right) (\xi, \{p_n\}), \quad (21)$$

The integral on the right hand side counts the residue at infinity:

$$\oint f(\xi) d\xi = f_{-1} \text{ for } f(\xi) = \sum_i f_i \xi^i.$$

Proof. In detail, the Dunkl operator $D_i^{(N)}$ transforms the space $\Lambda_{N,i}$ of functions with chosen variable x_i into itself:

$$\begin{aligned} D_i^{(N)}(F(x_i, \{p_n\})) &= x_i \frac{\partial}{\partial x_i} F(x_i, \{p_n\}) + \beta \sum_{j \neq i} \frac{x_i}{x_i - x_j} (1 - K_{ij}) F(x_i, \{p_n\}) = \\ &= x_i \frac{\partial}{\partial x_i} F(x_i, \{p_n\}) + \beta \sum_{j \neq i} \frac{x_i}{x_i - x_j} \left(\left(\tilde{V}(x_j) F \right) (x_i, \{p_n\}) - \left(\tilde{V}(x_i) F \right) (x_j, \{p_n\}) \right) = \end{aligned}$$

In each occurrence of $F(x_i; \{p_n\})$ we regard $\{p_n\}$ as symmetric functions of $(N-1)$ variables, while in $\left(\tilde{V}(x_j) F \right) (x_i, \{p_n\})$ $\{p_n\}$ depend on $(N-2)$ variables (all except x_i and x_j). Using the absence of singularities on the diagonals $x_i = x_j$ for Dunkl operators, we first present each fraction in the series as a function of x_i/x_j , then replace them by Cauchy integrals, to get:

$$D_i^{(N)}(F(x_i, \{p_n\})) = x_i \frac{\partial}{\partial x_i} F(x_i, \{p_n\}) -$$

$$\beta \sum_{j \neq i} \frac{\frac{x_i}{x_j}}{1 - \frac{x_i}{x_j}} \left(\left(\tilde{V}(x_j) F \right) (x_i, \{p_n\}) - \left(\tilde{V}(x_i) F \right) (x_j, \{p_n\}) \right) = x_i \frac{\partial}{\partial x_i} F(x_i, \{p_n\}) +$$

$$\beta x_i \oint \frac{d\xi}{\xi^2} \frac{\phi^-(\xi) - 1}{1 - \frac{x_i}{\xi}} \left(\tilde{V}'(\xi) \tilde{V}(x_i) F \right) (\xi, \{p_n\}) - \beta x_i \oint \frac{d\xi}{\xi^2} \frac{\phi^-(\xi) - 1}{1 - \frac{x_i}{\xi}} \left(\tilde{V}'(\xi) \tilde{V}(\xi) F \right) (x_i, \{p_n\}).$$

In the last summand the vertex operators cancel each other due to (20), and the corresponding integral vanishes since it contains ξ only in negative powers. We then obtain formula (21). ■

The ring Λ_∞ of symmetric functions with infinite number of variables is defined as the projective limit $\Lambda_\infty = \varprojlim \Lambda_N$ with respect to the projection $\Lambda_{N+1} \rightarrow \Lambda_N$:

$$f(x_1, x_2, \dots, x_N, x_{N+1}) \rightarrow f(x_1, x_2, \dots, x_N, 0).$$

An element of Λ_∞ can be represented by a sequence of symmetric functions:

$$f_1(x_1), f_2(x_1, x_2), \dots, f_N(x_1, x_2, \dots, x_N), \dots, \quad (22)$$

that stabilizes $f_{N+1}(x_1, x_2, \dots, x_N, 0) = f_N(x_1, x_2, \dots, x_N)$.

The ring Λ_N is generated by Newton power sums $p_k^{(N)}(x_1, x_2, \dots, x_N) = \sum_{i=1}^N x_i^k$ ($n \leq N$). The Newton polynomials satisfy the stability condition (22) and thus correctly define an element $p_k \in \Lambda_\infty$ that can be presented as a series $p_k = \sum_i x_i^k$. The elements p_k , $k = 0, 1, \dots$ freely generate the ring Λ_∞ . Note that we add to Λ_∞ the formal variable p_0 . The canonical projection $\Lambda_\infty \rightarrow \Lambda_N$ can be described by the relation:

$$\Lambda_\infty \rightarrow \Lambda_N : \quad p_k \rightarrow p_k^{(N)} = \sum_{i=1}^N x_i^k, \quad p_0 \rightarrow N. \quad (23)$$

As is well known, Λ_∞ is a realization of the bosonic Fock space, which is an irreducible representation of the Heisenberg algebra \mathcal{U} , generated by elements α_n , $n \in \mathbb{Z}$ subject to relations

$$[\alpha_n, \alpha_m] = n\delta_{n,-m},$$

$$\alpha_n \rightarrow n \frac{\partial}{\partial p_n}, \quad \alpha_{-n} \rightarrow p_n, \quad n > 0, \quad \alpha_0 \rightarrow p_0.$$

We identify Λ_∞ with the Fock space \mathcal{F} .

We use the operator $V(z) : \mathcal{F} \otimes \mathbb{C}[z] \rightarrow \mathcal{F} \otimes \mathbb{C}[z]$

$$V(z) = \exp \left(\sum_{n \geq 0} z^n \frac{\partial}{\partial p_n} \right). \quad (24)$$

Define an operator $D : \mathcal{F} \otimes \mathbb{C}[z] \rightarrow \mathcal{F} \otimes \mathbb{C}[z]$

$$D(F(z, \{p_n\})) = z \frac{\partial}{\partial z} F(z, \{p_n\}) + \beta z \oint \frac{d\xi}{\xi^2} \frac{\phi^-(\xi)}{1 - \frac{z}{\xi}} V^{-1}(\xi) V(x_i) F(\xi, \{p_n\}), \quad (25)$$

There is a map $\pi_i^{(N)} : \mathcal{F} \otimes \mathbb{C}[z] \rightarrow \Lambda_{N,i}$:

$$p_k \rightarrow \sum_{j \neq i}^N x_j^k, \quad z \rightarrow x_i.$$

By (21) and (25), we get the following commutative diagram:

$$\begin{array}{ccc} \mathcal{F} \otimes \mathbb{C}[z] & \xrightarrow{\pi_i^{(N)}} & \Lambda_{N,i} \\ D \downarrow & & \downarrow D_i^{(N)} \\ \mathcal{F} \otimes \mathbb{C}[z] & \xrightarrow{\pi_i^{(N)}} & \Lambda_{N,i} \end{array} \quad (26)$$

There is a difference between the definitions (15) and (24) for the vertex operators $\tilde{V}(x)$ and $V(z)$ in the zero mode $\frac{\partial}{\partial p_0}$. This difference is compensated by the term $\phi^-(\xi) - 1$ in (21). The presence of $\frac{\partial}{\partial p_0}$ in $V(z)$ is responsible for a decrease in the number of particles in the zero-mode p_0 . The commutativity of the diagram (26) implies

Proposition 1 *The operator D (25) is a projective limit of Dunkl operators $D_i^{(N)}$, see [8] and [11].*

Define the linear map $\mathcal{E} : \mathcal{F} \otimes \mathbb{C}[z] \rightarrow \mathcal{F}$ as

$$\mathcal{E}F(z, \{p_n\}) = \oint \frac{d\xi}{\xi} \phi^-(\xi) V^{-1}(\xi) F(\xi, \{p_n\}).$$

Then the operators $\mathcal{H}_k = \mathcal{E} D^k : \mathcal{F} \rightarrow \mathcal{F}$,

$$\mathcal{H}_k : \mathcal{F} \xrightarrow{\iota} \mathcal{F} \otimes \mathbb{C}[z] \xrightarrow{D^k} \mathcal{F} \otimes \mathbb{C}[z] \xrightarrow{\mathcal{E}} \mathcal{F},$$

generate a commutative family of Hamiltonians of the limiting system [8, 11].

As an example let us calculate the Hamiltonian \mathcal{H}_2 :

$$\begin{aligned} \mathcal{H}_2 F(p) &= \oint_{\xi} \frac{d\xi}{\xi} \phi^-(\xi) V^{-1}(\xi) \left(\xi \frac{\partial}{\partial \xi} \right)^2 V(\xi) F(p) + \\ &+ \beta \oint_{\eta < \xi} \frac{d\xi d\eta}{\xi^2} \phi^-(\eta) V^{-1}(\eta) \frac{\phi^-(\xi)}{1 - \frac{\eta}{\xi}} V(\eta) V^{-1}(\xi) \left(\xi \frac{\partial}{\partial \xi} \right) V(\xi) F(p) = \\ &= \oint_{\xi} \frac{d\xi}{\xi} \phi^-(\xi) \left(\sum_{k>0, n>0} k n \xi^{k+n} \frac{\partial}{\partial p_k} \frac{\partial}{\partial p_n} + \sum_{n>0} n^2 \xi^n \frac{\partial}{\partial p_n} \right) F(p) + \\ &+ \beta \oint \frac{d\xi d\eta}{\xi^2} \phi^-(\eta) \left(\frac{\phi^-(\xi)}{1 - \frac{\eta}{\xi}} - \frac{1}{(1 - \frac{\eta}{\xi})^2} \right) \sum_{n>0} \xi^n \frac{\partial}{\partial p_n} F(p). \end{aligned}$$

We have taken the second derivative in the first integral and used the commutator relations

$$[\phi^-(\xi), V(\eta)] = -\frac{V(\eta)}{(1 - \frac{\eta}{\xi})}$$

in the second. Thus

$$\mathcal{H}_2 = \sum_{k>0, n>0} knp_{k+n} \frac{\partial}{\partial p_k} \frac{\partial}{\partial p_n} + \sum_{n>0} n^2 p_n \frac{\partial}{\partial p_n} - \beta \sum_{n>0} n^2 p_n \frac{\partial}{\partial p_n} + \beta \sum_{k>0, n \geq 0} (k+n) p_k p_n \frac{\partial}{\partial p_{k+n}}.$$

3.2 The Spin CS system and multicomponent Fock space

In this section we present a generalization of the calculations in section 3.1 to the spin CS model. Let us split the difference part of the Dunkl operator into two sums

$$D_i = x_i \frac{\partial}{\partial x_i} + \beta \sum_{j \neq i, \sigma(i)=\sigma(j)} \frac{x_i}{x_i - x_j} (1 - K_{ij}) + \beta \sum_{j, \sigma(i) \neq \sigma(j)} \frac{x_i}{x_i - x_j} (1 - K_{ij}). \quad (27)$$

The Dunkl operators act on the space M_N and the symmetric functions of D_i preserve the subspace of global S_N -invariants $\bar{M}_N := M_N^{S_N}$, which we further identify with the direct sum of spaces of symmetric functions by groups of variables. The space M_N can be decomposed into the direct sum of weight spaces with respect to the diagonal action of the Cartan subalgebra of \mathfrak{gl}_s :

$$M_N = \bigoplus_{|\lambda|=N} M_\lambda,$$

where $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_s)$ is a collection of nonnegative integers, $\sum_{c=1}^s \lambda_c = N$. In other words, M_λ is the space of vector-valued polynomials generated by tensors of the form

$$\mathbf{m} = (e_1 \otimes z^{a_{11}} \dots e_1 \otimes z^{a_{1\lambda_1}}) \otimes (e_2 \otimes z^{a_{21}} \dots e_2 \otimes z^{a_{2\lambda_2}}) \otimes \dots \otimes (e_s \otimes z^{a_{s1}} \dots e_s \otimes z^{a_{s\lambda_s}}) \quad (28)$$

together with all their permutations. Denote by $\bar{M}_\lambda := M_\lambda^{S_N}$ the space of S_N -invariants in M_λ . Hence,

$$\bar{M}_N = \bigoplus_{|\lambda|=N} \bar{M}_\lambda.$$

Let A_λ be the ring of polynomials in N variables $(x_{11}, x_{12}, \dots, x_{1\lambda_1}, x_{21}, x_{22}, \dots, x_{2\lambda_2}, \dots, x_{s1}, x_{s2}, \dots, x_{s\lambda_s})$. Denote by $\Lambda_\lambda \in A_\lambda$ the ring of functions symmetric within each group, defined by the partition λ , of variables $(x_{11}, x_{12}, \dots, x_{1\lambda_1}), (x_{21}, x_{22}, \dots, x_{2\lambda_2}), \dots, (x_{s1}, x_{s2}, \dots, x_{s\lambda_s})$. Define a linear map $\gamma_\lambda : A_\lambda \rightarrow M_\lambda$. This map associates to each monomial

$$x = x_{11}^{a_{11}} \dots x_{1\lambda_1}^{a_{1\lambda_1}} \dots x_{s1}^{a_{s1}} \dots x_{s\lambda_s}^{a_{s\lambda_s}}$$

the total symmetrization of the corresponding tensor \mathbf{m} , see (28), divided by the product $\lambda_1! \dots \lambda_s!$:

$$\gamma_\lambda(x) = \frac{1}{\lambda_1! \dots \lambda_s!} \sum_{\sigma \in S_N} \sigma(\mathbf{m}). \quad (29)$$

The restriction of the map γ_λ to Λ_λ establishes an isomorphism of vector spaces

$$\gamma_\lambda : \Lambda_\lambda \simeq \bar{M}_\lambda.$$

Using this isomorphism we push forward the ring structure of Λ_λ to \bar{M}_λ . The ring Λ_λ is generated by the (not indepent) elements

$$p_{n,a}^\lambda = \sum_{i=1}^{\lambda_a} x_{ai}^n,$$

with $a = 1, \dots, s$, $n = 1, 2, \dots$. Further we present elements of the ring Λ_λ as polynomial functions $F(p)$ of the elements $p_{n,a}^\lambda$.

For each $i = 1, \dots, N$ denote by $\bar{M}_{\lambda,i}$ the subspace of M_λ which consists of tensors invariant with respect to the symmetric group (isomorphic to S_{N-1}) of permutations of all tensor factors except the i -th one. We have a natural inclusion (decomposition of a tensor with respect to its i -th factor):

$$\alpha_{\lambda,i} : \bar{M}_\lambda \rightarrow \bar{M}_{\lambda,i}.$$

Denote by $A_{\lambda,\cdot}$ and by $\Lambda_{\lambda,\cdot}$ the rings

$$A_{\lambda,\cdot} = \bigoplus_{c=1}^s (A_{\lambda-\epsilon_c} \otimes \mathbb{C}[z]), \quad \Lambda_{\lambda,\cdot} = \bigoplus_{c=1}^s (\Lambda_{\lambda-\epsilon_c} \otimes \mathbb{C}[z]).$$

Here $\epsilon_1 = (1, 0, \dots, 0)$, $\dots \epsilon_s = (0, \dots, 0, 1)$. We present elements of the ring $\Lambda_{\lambda,\cdot}$ as vector valued polynomial functions on elements $p_{n,a}^\lambda$ and z :

$$\bar{F}(p, z) = \begin{pmatrix} F_1(p, z) \\ \dots \\ F_s(p, z) \end{pmatrix}.$$

For each $i = 1, \dots, N$ we define a map $\gamma_{\lambda,i} : A_{\lambda,\cdot} \rightarrow \bar{M}_{\lambda,i}$. For the reasons of typographic convenience we first describe the map $\gamma_{\lambda,N}$. This map attaches to a monomial $x_c \in A_{\lambda-\epsilon_c} \otimes \mathbb{C}[z]$,

$$x_c = x_{11}^{a_{11}} \dots x_{1\lambda_1}^{a_{1\lambda_1}} \dots x_{c1}^{a_{c1}} \dots x_{c,\lambda_c-1}^{a_{c,\lambda_c-1}} \dots x_{s1}^{a_{s1}} \dots x_{s\lambda_s}^{a_{s\lambda_s}} \cdot z^d$$

the symmetrization over the first $N-1$ tensor components of the tensor

$$\mathbf{m}' = (e_1 \otimes z^{a_{11}} \dots e_1 \otimes z^{a_{1\lambda_1}}) \dots (e_c \otimes z^{a_{c1}} \dots e_c \otimes z^{a_{c,\lambda_c-1}}) \dots (e_s \otimes z^{a_{s1}} \dots e_s \otimes z^{a_{s\lambda_s}}) \otimes (e_c \otimes z^d),$$

divided by the product $\lambda_1! \dots (\lambda_c - 1)! \dots \lambda_s!$:

$$\gamma_{\lambda,N}(x_c) = \frac{1}{\lambda_1! \dots (\lambda_c - 1)! \dots \lambda_s!} \sum_{\sigma \in S_{N-1}} \sigma(\mathbf{m}').$$

Next we set

$$\gamma_{\lambda,i} = \sigma_{i,N} \cdot \gamma_{\lambda,N},$$

where $\sigma_{i,j}$ is the transposition of the i -th and j -th tensor factors. By construction, the restriction of the map $\gamma_{\lambda,i}$ to $\Lambda_{\lambda,\cdot}$ establishes an isomorphism of vector spaces

$$\gamma_{\lambda,i} : \Lambda_{\lambda,\cdot} \approx \bar{M}_{\lambda,i}.$$

Restrictions of $\gamma_{\lambda,i}$ to $\Lambda_{\lambda,\cdot}$ form an equivariant family:

$$\gamma_{\lambda,i} = \sigma_{i,j} \cdot \gamma_{\lambda,i}.$$

For each $a = 1, \dots, s$ we define the ‘vertex’ operators $\tilde{V}_a(z) : \Lambda_\lambda \rightarrow \Lambda_{\lambda-\epsilon_a} \otimes \mathbb{C}[z]$

$$\tilde{V}_a(z) = \exp \left(\sum_{n>0} z^n \frac{\partial}{\partial p_{n,a}^\lambda} \right).$$

Thus we have the linear map

$$\iota_\lambda : \Lambda_\lambda \rightarrow \Lambda_{\lambda,\cdot}, \quad \iota_\lambda : F(p) \rightarrow \tilde{V}_1(z, p)F(p) \oplus \dots \oplus \tilde{V}_s(z, p)F(p), \quad (30)$$

that we also represent using the column-vector notation:

$$\iota_\lambda F(p) = \begin{pmatrix} \tilde{V}_1(z, p)F(p) \\ \vdots \\ \tilde{V}_s(z, p)F(p) \end{pmatrix}.$$

Lemma 4 *We have the equality $\gamma_{\lambda,i} \iota_\lambda = \alpha_{\lambda,i} \gamma_\lambda$.*

The commutative diagram below illustrates the statement from Lemma 4.

$$\begin{array}{ccc} \Lambda_\lambda & \xrightarrow{\gamma_\lambda} & \bar{M}_\lambda \\ \iota_\lambda \downarrow & & \downarrow \alpha_{\lambda,i} \\ \Lambda_{\lambda,\cdot} & \xrightarrow{\gamma_{\lambda,i}} & \bar{M}_{\lambda,i} \end{array} \quad (31)$$

Next we translate the Dunkl operators to the language of polysymmetric functions. Let $\phi_a^\pm(\xi)$ be the power series in ξ and in ξ^{-1} :

$$\phi_a^+(\xi) = \sum_{n=1}^{\infty} \xi^n n \frac{\partial}{\partial p_{n,a}^\lambda}, \quad \phi_a^-(\xi) = \sum_{n=0}^{\infty} \frac{p_{n,a}^\lambda}{\xi^n}, \quad (32)$$

where we set $p_{0,a}^\lambda = \lambda_a$. Let $\tilde{V}'_a(z)$ be the following operator from the polynomial ring of formal variables $p_{n,a}^\lambda$, $n > 0$, $a = 1, \dots, s$ to $\Lambda_{\lambda+\epsilon_a} \otimes \mathbb{C}[z]$:

$$\tilde{V}'_a(z) = \exp \left(- \sum_{n>0} z^n \frac{\partial}{\partial p_{n,a}^\lambda} \right).$$

Introduce the operator $D : \Lambda_{\lambda,\cdot} \rightarrow \Lambda_{\lambda,\cdot}$. It acts on a vector function $F(z, p)$ as a diagonal matrix, such that

$$D(F_a(z, p)) = z \frac{\partial}{\partial z} F_a(z, p) + \beta \sum_b z \oint \frac{d\xi}{\xi^2} \frac{\phi_b^-(\xi) - \delta_{a,b}}{1 - \frac{z}{\xi}} \tilde{V}_b'(\xi) \tilde{V}_b(z) F_a(\xi, p). \quad (33)$$

Lemma 5 *For any $i = 1, \dots, N$ we have the equality*

$$\gamma_{\lambda,i} D = D_i \gamma_{\lambda,i}.$$

For $a, b = 1, \dots, s$ define the linear map $\mathbb{E}^{ab} : \Lambda_{\lambda,\cdot} \rightarrow \Lambda_{\lambda+\epsilon_a-\epsilon_b}$ by the relation

$$\mathcal{E}^{ab} F_c(z, p) = \delta_{bc} \oint_{\xi} \frac{d\xi}{\xi} \phi_a^-(\xi) \tilde{V}_a'(\xi) F_c(\xi, p). \quad (34)$$

Lemma 6 *For any $F \in \Lambda_{\lambda,\cdot}$ we have the relation*

$$\gamma_{\lambda} \mathcal{E}^{ab} F = \sum_{i=1}^N E_i^{ab} \gamma_{\lambda,i} F.$$

In other words, \mathcal{E}^{ab} is the operator of matrix averaging of a collection of equivariant tensors.

The proof of lemmas 3, 4, 5 is analogous to that of lemmas 1, 2, 3.

Set $\Lambda^{s,N} = \oplus_{|\mu|=N} \Lambda_{\mu}$. As a direct corollary of statements above we get the Yangian action on $\Lambda^{s,N}$ translated from that on \bar{M}_N .

Proposition 2 *The Yangian generator $t_{(k)}^{ab}$ maps the space Λ_{μ} to the space $\Lambda_{\mu+\epsilon_a-\epsilon_b}$ and is given by the relation*

$$t_{(k)}^{ab} F = \frac{(-1)^k}{\beta^k} \mathcal{E}^{ab} D^k \iota_{\mu} F \quad \text{for any } F \in \Lambda_{\mu}. \quad (35)$$

where the maps ι_{μ} , D and \mathcal{E}^{ab} are given by the relations (30), (33), (34).

In terms of generating functions this action looks as follows:

$$t^{ab}(u) = \delta_{ab} + \beta \mathcal{E}^{ab} (\beta u + D)^{-1} \iota_{\mu}. \quad (36)$$

Let \mathcal{U}^s be the tensor product of s copies of the Heisenberg algebra with generators $\alpha_{n,a}$ and $Q_a^{\pm 1}$, where $n \in \mathbb{Z}$, $a = 1, \dots, s$ and relations

$$[\alpha_{n,a}, \alpha_{m,b}] = n \delta_{a,b} \delta_{n,-m}, \quad Q_a \alpha_{n,b} Q_a^{-1} = \alpha_{n,b} + \delta_{n,0} \delta_{a,b}.$$

The multicomponent Fock space \mathcal{F}^s is an irreducible representation of \mathcal{U}^s in the space of polynomials $\mathbb{C}[p_{n,a}, q_b^{\pm 1} \equiv e^{\pm \frac{\partial}{\partial p_{0,b}}}]$, $n \neq 0$, $a, b = 1, \dots, s$:

$$\alpha_{n,a} \rightarrow n \frac{\partial}{\partial p_{n,a}}, \quad \alpha_{-n,a} \rightarrow p_{n,a}, \quad n > 0, \quad \alpha_{0,a} \rightarrow p_{0,a} - \lambda_a, \quad Q_a^{\pm 1} \mapsto q_a^{\pm 1}, \quad \alpha_{0,a} \cdot 1 = 0,$$

where $\lambda_1, \dots, \lambda_s \in \mathbb{C}$ are arbitrary constants. For any $\nu \in \mathbb{Z}^s$ denote by $\mathcal{F}_\nu^s \in \mathcal{F}^s$ the subspace $\mathbb{C}[p_{n,a}]q_1^{\nu_1} \cdots q_s^{\nu_s}$. We have the decomposition

$$\mathcal{F}^s = \bigoplus_{\nu \in \mathbb{Z}^s} \mathcal{F}_\nu^s.$$

The subspace \mathcal{F}_ν^s is invariant with respect to all operators $\alpha_{n,a}$ and $\alpha_{0,a}v = \nu_a v$ for any $v \in \mathcal{F}_\nu^s$, or, in terms of $p_{0,a}$,

$$p_{0,a}v = (\nu_a + \lambda_a)v \quad \text{for any } v \in \mathcal{F}_\nu^s. \quad (37)$$

The multicomponent Fock space may be regarded as a projective limit of the spaces of polysymmetric functions in the following sense. Fix a natural number N and a weight $\lambda = (\lambda_1, \dots, \lambda_s)$ consisting of nonnegative integers such that $\sum_a \lambda_a = N$. For any $\nu \in \mathbb{Z}^s$ define a map $\pi_{\lambda\nu} : \mathcal{F}_\nu^s \rightarrow \Lambda_{\lambda+\nu}$ by setting

$$p_{n,a} \mapsto p_{n,a}^{\lambda+\nu}, \quad n > 0, \quad \text{and} \quad p_{0,a} \mapsto \lambda_a + \nu_a.$$

By definition, this map is nonzero only when all numbers $\nu_a + \lambda_a$ are nonzero. It is compatible with zero evaluation maps $\tau_{\mu,a} : \Lambda_\mu \rightarrow \Lambda_{\mu-\epsilon_a}$,

$$\begin{aligned} f(x_{11}, \dots, x_{1\mu_1}, \dots, x_{a1}, \dots, x_{a\mu_a-1}, x_{a\mu_a}, \dots, x_{s1}, \dots, x_{s\mu_s}) &\mapsto \\ f(x_{11}, \dots, x_{1\mu_1}, \dots, x_{a1}, \dots, x_{a\mu_a-1}, 0, \dots, x_{s1}, \dots, x_{s\mu_s}). \end{aligned}$$

in the sense that

$$\pi_\lambda \cdot \tau_a = \pi_{\lambda-\epsilon_a},$$

where $\pi_\lambda = \bigoplus_{\nu \in \mathbb{Z}^s} \pi_{\lambda\nu}$, $\tau_a = \bigoplus_{\mu \in \mathbb{Z}_{\geq 0}^s} \tau_{\mu,a}$.

For any $a = 1, \dots, s$ denote by $V_a(z) : \mathcal{F}^s \rightarrow \mathcal{F}^s \otimes \mathbb{C}[z]$ the vertex operator

$$V_a(z) = \exp \left(\sum_{n \geq 0} z^n \frac{\partial}{\partial p_{n,a}} \right). \quad (38)$$

and use them for the embedding $\iota : \mathcal{F}^s \xrightarrow{\iota} \mathcal{F}^s \otimes (\mathbb{C}^s \otimes \mathbb{C}[z])$, where $\iota(F) = V_1(z)F \oplus \cdots \oplus V_s(z)F$, or, in vector notation

$$\iota F = \begin{pmatrix} V_1(z)F \\ \vdots \\ V_s(z)F \end{pmatrix}, \quad F = F(p, q) \in \mathcal{F}^s. \quad (39)$$

We also use the fields

$$\phi_a^+(\xi) = \sum_{n=1}^{\infty} \xi^n n \frac{\partial}{\partial p_{n,a}}, \quad \phi_a^-(\xi) = \sum_{n=0}^{\infty} \frac{p_{n,a}}{\xi^n}.$$

Let us introduce the operator D . It acts on a vector function $F(z) \in \mathcal{F}^s \otimes \mathbb{C}[z]$ as a diagonal matrix $D : \mathcal{F}^s \otimes (\mathbb{C}^s \otimes \mathbb{C}[z]) \rightarrow \mathcal{F}^s \otimes (\mathbb{C}^s \otimes \mathbb{C}[z])$ by

$$D(F_a(z)) = z \frac{\partial}{\partial z} F_a(z) + \beta \sum_b z \oint \frac{d\xi}{\xi^2} \frac{\phi_b^-(\xi)}{1 - \frac{z}{\xi}} V_b^{-1}(\xi) V_b(z) F_a(\xi). \quad (40)$$

The operator $\mathcal{E}^{ab} : \mathcal{F}^s \otimes (\mathbb{C}^s \otimes \mathbb{C}[z]) \rightarrow \mathcal{F}^s$ of matrix averaging is now given by the relation

$$\mathcal{E}^{ab} F_c(z) = \delta_{bc} \oint_{\xi} \frac{d\xi}{\xi} \phi_a^-(\xi) V_a^{-1}(\xi) F_c(\xi). \quad (41)$$

The maps ι, D and \mathcal{E}^{ab} are compatible with finite projections π_λ in the following sense. We formulate the compatibility conditions with the starting point at $\mathcal{F}_\nu^s \in \mathcal{F}^s$.

Proposition 3 *For any $v \in \mathcal{F}^s$ and $u \in \mathcal{F}^s \otimes (\mathbb{C}^s \otimes \mathbb{C}[z])$ we have the following equalities:*

$$\begin{aligned} (\pi_\lambda \otimes 1) D(u) &= D(\pi_\lambda \otimes 1)(u), & \iota_{\lambda+\nu} \pi_\lambda(v) &= (\pi_\lambda \otimes 1) \iota(v), \\ \mathcal{E}^{ab}(\pi_\lambda \otimes 1)(u) &= (\pi_\lambda \otimes 1) \mathcal{E}^{ab}(u). \end{aligned}$$

As a consequence of proposition 2 and proposition 3 we obtain the representation T_k^{ab} of the Yangian generators $t_{(k)}^{ab}$ in the multicomponent Fock space:

Proposition 4 *The assignment $t_{(k)}^{ab} \rightarrow T_k^{ab}$, where*

$$T_k^{ab} F = \frac{(-1)^k}{\beta^k} \mathcal{E}^{ab} D^k \iota F \quad \text{for any } F \in \mathcal{F}^s, \quad (42)$$

defines a representation of the Yangian $Y(\mathfrak{gl}_s)$ in the space \mathcal{F}^s .

Here the maps ι, D and \mathcal{E}^{ab} are given by the relations (39), (40), (41). In generating functions this action looks as follows:

$$T^{ab}(u) = \delta_{ab} + \beta \mathcal{E}^{ab} (\beta u + D)^{-1} \iota.$$

The Yangian action can be illustrated by the following scheme:

$$\beta^k T_k^{ab} : \mathcal{F}^s \xrightarrow{\iota} \mathcal{F}^s \otimes (\mathbb{C}^s \otimes \mathbb{C}[z]) \xrightarrow{(-D)^k} \mathcal{F}^s \otimes (\mathbb{C}^s \otimes \mathbb{C}[z]) \xrightarrow{\mathcal{E}^{ab}} \mathcal{F}^s. \quad (43)$$

As an example we present corresponding integral expression for the zero and the first generators of the Yangian $Y(\mathfrak{gl}_s)$:

$$T_0^{ab} = \oint \frac{d\xi}{\xi} \phi_a^-(\xi) V_a^{-1}(\xi) V_b(\xi), \quad (44)$$

$$\begin{aligned} T_1^{ab} &= -\frac{1}{\beta} \oint \frac{d\xi}{\xi} \phi_a^-(\xi) V_a^{-1}(\xi) \phi_b^+(\xi) V_b(\xi) \\ &\quad - \sum_c \oint \frac{d\xi d\eta}{\xi \eta} \frac{\frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_c^-(\xi) V_c(\eta) V_c^{-1}(\xi) V_b(\xi). \end{aligned} \quad (45)$$

3.3 Hamiltonians

In this section we provide explicit expressions for the first few Hamiltonians constructed by means of the procedure (43). The proof and calculation are presented in Appendix 4. The Hamiltonians $H_n = \sum_i d_i^n$ can be expressed in terms of D_i by means of (9):

$$H_1 = \sum_i D_i + \frac{\beta}{2} \sum_{a,b} \mathbb{E}^{ab} \mathbb{E}^{ba} - \frac{\beta}{2} s p_0, \quad (46)$$

$$\begin{aligned} H_2 = & \sum_i D_i^2 + \beta \sum_{i,j,a,b} E_i^{ab} E_j^{ba} D_j - \beta s \sum_i D_i + \\ & + \frac{\beta^2}{3} \sum_{a,b,c} \mathbb{E}^{ab} \mathbb{E}^{bc} \mathbb{E}^{ca} - \frac{2s\beta^2}{3} \sum_{a,b} \mathbb{E}^{ab} \mathbb{E}^{ba} + \beta^2 \frac{2s^2 + p_0 - 1}{6} p_0, \end{aligned} \quad (47)$$

where s and $p_0 = \sum_a p_{0,a}$ are the numbers of spins and particles, respectively. Using (4) we can rewrite these formulas in terms of the Yangian generators T_k^{ab} :

$$\frac{H_1}{\beta} = - \sum_a T_1^{aa} + \frac{1}{2} \sum_{a,b} T_0^{ab} T_0^{ba} - \frac{s}{2} \sum_a T_0^{aa}, \quad (48)$$

$$\begin{aligned} \frac{H_2}{\beta^2} = & \sum_a T_2^{aa} - \sum_{a,b} T_0^{ab} T_1^{ba} + s \sum_a T_1^{aa} + \frac{1}{3} \sum_{a,b,c} T_0^{ab} T_0^{bc} T_0^{ca} \\ & - \frac{2s}{3} \sum_{a,b} T_0^{ab} T_0^{ba} + \frac{1}{6} \sum_{a,b} T_0^{aa} T_0^{bb} + \frac{2s^2 - 1}{6} \sum_a T_0^{aa} \end{aligned} \quad (49)$$

On the other hand, the Hamiltonians can be obtained as elements of the q -determinant (Th1). Consider the representation of the q -determinant :

$$\text{q-det } T(u) = \sum_{\sigma \in S_m} (-1)^{\text{sgn}(\sigma)} T^{\sigma(1),1}(u) T^{\sigma(2),2}(u-1) \dots T^{\sigma(m),m}(u-m+1) = 1 + \frac{\Delta_0}{u} + \frac{\Delta_1}{u^2} + \frac{\Delta_2}{u^3} + \dots,$$

where

$$T^{ab}(u) = \delta^{ab} + \frac{T_0^{ab}}{u} + \frac{T_1^{ab}}{u^2} + \frac{T_2^{ab}}{u^3} + \dots$$

The elements Δ_i can be expressed in terms of T_k^{ab} . Explicit expressions for $\Delta_0, \Delta_1, \Delta_2$ are presented in Appendix 3. The Hamiltonians can be rewritten in the following form:

$$\frac{H_1}{\beta} = -\Delta_1 + \frac{1}{2} \Delta_0^2 - \frac{1}{2} \Delta_0, \quad (50)$$

$$\frac{H_2}{\beta^2} = \Delta_2 - \Delta_0 \Delta_1 + \Delta_1 + \frac{1}{3} \Delta_0^3 - \frac{1}{2} \Delta_0^2 + \frac{1}{6} \Delta_0. \quad (51)$$

Proposition 5 *The first Hamiltonians in the multicomponent Fock space have the following form*

$$\mathcal{H}_1 = \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) + \frac{\beta}{2} (p_0^2 - p_0), \quad (52)$$

$$\begin{aligned} \mathcal{H} = \mathcal{H}_2 - \beta \mathcal{H}_1 = & \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) (\phi_a^+(\xi))^2 + (1 - \beta) \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) (\phi_a^+(\xi))' + \\ & -\beta \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) + \beta \sum_{a,b} \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_b^-(\xi) \phi_a^+(\xi) + \\ & \beta \sum_{a>b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) \phi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\xi) V_a(\xi). \end{aligned} \quad (53)$$

4 Classical limit

In this section we investigate the classical limit of the Hamiltonian (53). In the spinless case it leads to the periodic Benjamin-Ono equation [8]. By multiplying (53) by β , we obtain:

$$\begin{aligned} \beta \mathcal{H} = & \beta \oint_{\xi} \frac{d\xi}{\xi} \phi_a^-(\xi) (\phi_a^+(\xi))^2 + \beta \oint_{\xi} \frac{d\xi}{\xi} \phi_a^-(\xi) (\phi_a^+(\xi))' + \beta^2 \sum_{a,b} \oint_{\eta} \frac{d\eta}{\eta} \phi_a^-(\eta) \phi_b^-(\eta) \phi_a^+(\eta) \\ & + \beta^2 \sum_{a>b} \oint_{\eta, \xi} \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) \phi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\xi) V_a(\xi), \end{aligned} \quad (54)$$

where for convenience we denote $V_a(x) V_b^{-1}(x)$ by $V_{ab}(x)$.

Introducing the classical variables $\alpha_{n,a}$, $n \in \mathbb{Z}$

$$\alpha_{n,a} = \begin{cases} \beta p_{-n,a}, & n \leq 0; \\ n \frac{\partial}{\partial p_{n,a}}, & n > 0, \end{cases}$$

with Poisson bracket relations

$$\{\alpha_{n,a}, \alpha_{m,b}\} = n \delta_{a,b} \delta_{n+m,0}, \quad (55)$$

combine them into generating functions

$$\varphi_a^-(\xi) = \sum_{n=0}^{\infty} \frac{\alpha_{-n,a}}{\xi^n}, \quad \varphi_a^+(\xi) = \sum_{n=1}^{\infty} \alpha_{n,a} \xi^n, \quad \text{and} \quad \varphi_a(x) = \varphi_a^+(x) + \varphi_a^-(x).$$

Then

$$\{\varphi_a^-(x), \varphi_b^+(y)\} = \delta_{a,b} \frac{y}{(1 - \frac{y}{x})^2}, \quad \{\varphi_a^{\pm}(x), \varphi_b^{\pm}(y)\} = 0.$$

or

$$\{\varphi_a(x), \varphi_b(y)\} = \delta'(x/y) \delta_{a,b}.$$

The classical counterparts $\mathcal{V}_a(\xi)$ of the vertex operators $V_a(\xi)$ satisfy the relations

$$\xi \frac{d \log \mathcal{V}_a(\xi)}{d\xi} = \varphi_a^+(\xi),$$

and

$$\{\varphi_a^-(x), \mathcal{V}_b(y)\} = -\delta_{a,b} \frac{\mathcal{V}_b(y)}{1 - \frac{y}{x}}, \quad \{\varphi_a^+(x), \mathcal{V}_b(y)\} = 0. \quad (56)$$

As before, we use the notation $\mathcal{V}_{ab}(\xi)$ for $\mathcal{V}_a(\xi) \mathcal{V}_b^{-1}(\xi)$. Set

$$\begin{aligned} \mathcal{H} = & \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) (\varphi_a^+(\xi))^2 + \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) (\varphi_a^+(\xi))' + \sum_{a,b} \oint_{\eta} \frac{d\eta}{\eta} \varphi_a^-(\eta) \varphi_b^-(\eta) \varphi_a^+(\eta) \\ & + \sum_{a>b} \oint_{\eta, \xi} \frac{d\xi d\eta}{\xi \eta} \varphi_a^-(\eta) \varphi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) \mathcal{V}_{ab}^{-1}(\eta) \mathcal{V}_{ab}(\xi). \end{aligned} \quad (57)$$

The operator \mathcal{H} is the classical limit of the Hamiltonian (54) ($\beta \rightarrow 0$). The rule between the quantum commutator and Poisson bracket is $\beta^{-1}[\ , \] \rightarrow \{ \ , \ }$.

Proposition 6¹ *The equations of motion determined by the Hamiltonian \mathcal{H} are:*

$$\begin{aligned} \{\varphi_a^+(x), \mathcal{H}\} = & x \frac{\partial}{\partial x} (\varphi_a^+(x))^2 + \left(x \frac{\partial}{\partial x} \right)^2 (\varphi_a^+(x)) + \sum_b x \frac{\partial}{\partial x} (\varphi_b^-(x) \varphi_a^+(x))^+ + \\ & + \sum_b x \frac{\partial}{\partial x} (\varphi_b^-(x) \varphi_b^+(x))^+ + \sum_{b \neq a} x \frac{\partial}{\partial x} \left(\mathcal{V}_{ab}^{-1}(x) x \frac{\partial}{\partial x} \left((\varphi_b^-(x) \mathcal{V}_{ab}(x))^+ - (\varphi_b^-(x) \mathcal{V}_{ab}(x))^- \right) \right)^+ = \\ & x \frac{\partial}{\partial x} (\varphi_a^+(x))^2 + \left(x \frac{\partial}{\partial x} \right)^2 (\varphi_a^+(x)) + 2 \sum_b x \frac{\partial}{\partial x} (\varphi_b^-(x) \varphi_b^+(x))^+ \\ & + 2 \sum_{b \neq a} x \frac{\partial}{\partial x} \left(\mathcal{V}_{ab}^{-1}(x) x \frac{\partial}{\partial x} (\varphi_b^-(x) \mathcal{V}_{ab}(x))^+ \right), \end{aligned} \quad (58)$$

$$\begin{aligned} \{\varphi_a^-(x), \mathcal{H}\} = & 2x \frac{\partial}{\partial x} (\varphi_a^-(x) \varphi_a^+(x))^- - \left(x \frac{\partial}{\partial x} \right)^2 (\varphi_a^-(x)) + \sum_b x \frac{\partial}{\partial x} (\varphi_b^-(x) \varphi_a^-(x)) \\ & + \sum_{b \neq a} \left(\varphi_a^-(x) \mathcal{V}_{ab}^{-1}(x) x \frac{\partial}{\partial x} \left((\varphi_b^-(x) \mathcal{V}_{ab}(x))^+ - (\varphi_b^-(x) \mathcal{V}_{ab}(x))^- \right) \right)^- \\ & - \sum_{b \neq a} \left(\varphi_b^-(x) \mathcal{V}_{ab}(x) x \frac{\partial}{\partial x} \left((\varphi_a^-(x) \mathcal{V}_{ab}^{-1}(x))^+ - (\varphi_a^-(x) \mathcal{V}_{ab}^{-1}(x))^- \right) \right)^- \end{aligned} \quad (59)$$

¹For a formal series $f(z) = \sum_{n \in \mathbb{Z}} f_n z^n$ we denote by $f^+(z)$ the series $f^+(z) = \sum_{n \geq 1} f_n z^n = \oint \frac{zf(\xi)d\xi}{(1-z/\xi)\xi}$ and by $f^-(z)$ the series $f^-(z) = \sum_{n \leq 0} f_n z^n = \oint \frac{f(\xi)d\xi}{(1-\xi/z)\xi}$.

Remark Unlike the Yangian generators (44), (45), the Hamiltonian (53) does not contain dual zero modes $\frac{\partial}{\partial p_{0a}}$. The same holds for the classical limit, where we can freely use the operators $\tilde{\mathcal{V}}_a(\xi) = \exp \sum_{n \geq 1} \frac{a_n}{n} \xi^n$ instead of $\mathcal{V}_a(\xi)$. The Hamiltonian and the equations of motion do not change, while the brackets (56) turn into

$$\{\varphi_a^-(x), \tilde{\mathcal{V}}_b(y)\} = -\delta_{a,b} \frac{y/x \tilde{\mathcal{V}}_b(y)}{1 - \frac{y}{x}}, \quad \{\varphi_a^+(x), \tilde{\mathcal{V}}_b(y)\} = 0.$$

The quantum system is integrable: it has an infinite number of integrals of motion that can be obtained from the q -determinant of the Yangian generator function $T^{ab}(u)$. It is natural to assume that the classical system is integrable as well. In particular, it should admit a Lax pair presentation. Consider the operators L and M :

$$\begin{aligned} Lf &= z \frac{\partial}{\partial z} f(z) + \sum_a \mathcal{V}_a(z) (\varphi_a^-(z) \mathcal{V}_a^{-1}(z) f(z))^+, \\ Mf &= \left(z \frac{\partial}{\partial z} \right)^2 f(z) + 2 \sum_b (\varphi_b^+(z) \varphi_b^-(z))^+ f(z) + 2 \sum_b \mathcal{V}_b(z) z \frac{\partial}{\partial z} (\varphi_b^-(z) \mathcal{V}_b^{-1}(z) f(z))^+. \end{aligned} \quad (60)$$

They act on the space of analytic functions

$$f(z) = f_0 + f_1 z + f_2 z^2 + \dots,$$

where coefficients f_i are polysymmetrical functions $f_i(p_{n,a})$.

Proposition 7 *The operators L and M (60) represent a Lax pair of the classical system (57):*

$$\frac{dL}{dt} = [M, L].$$

Appendix

A.1 Rational Calogero system

In this section we present the analogous formulas for the rational Calogero model in the limit of infinitely many particles. We use all notations as before. The Dunkl operators for the rational system are defined as:

$$D_i = \frac{\partial}{\partial x_i} + \beta \sum_{j \neq i, \sigma(i)=\sigma(j)} \frac{1}{x_i - x_j} (1 - K_{ij}) + \beta \sum_{j, \sigma(i) \neq \sigma(j)} \frac{1}{x_i - x_j} (1 - K_{ij}).$$

We use a similar construction for the rational case and here present the analog of formulas (40) and (41):

$$D(F_a(z)) = \frac{\partial}{\partial z} F_a(z, p) + \beta \sum_b \oint \frac{d\xi}{\xi^2} \frac{\phi_b^-(\xi)}{1 - \frac{z}{\xi}} V_b(z) V_b^{-1}(\xi) F_a(\xi), \quad (61)$$

$$\mathcal{E}^{ab}(F_c(z)) = \delta_{bc} \oint_{\xi} \frac{d\xi}{\xi} \phi_a^-(\xi) V_a^{-1}(\xi) F_c(\xi). \quad (62)$$

The answer for the second Hamiltonian is

$$\begin{aligned} \mathcal{H}_2 = & \sum_{k,n \geq 1, a} k n p_{n+k-2,a} \frac{\partial}{\partial p_{k,a}} \frac{\partial}{\partial p_{n,a}} + (1-\beta) \sum_{n \geq 1, a} n(n-1) p_{n-2,a} \frac{\partial}{\partial p_{n,a}} + \\ & \beta \sum_{a,b} \sum_{n,k \geq 0} (k+n+2) p_{k,a} p_{n,b} \frac{\partial}{\partial p_{k+n+2}} + \\ & + \beta \sum_{a > b} \oint \frac{d\xi d\eta}{\xi^2 \eta^2} \phi_a^-(\eta) \phi_b^-(\xi) \left(n \frac{\xi^n}{\eta^n} + n \frac{\eta^n}{\xi^n} \right) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) V_a^{-1}(\eta). \end{aligned}$$

A.2 Calculations of Hamiltonians

In this section the proofs for formulas (52),(53) are presented.

$$\begin{aligned} \mathcal{H}_1 = & \sum_i D_i + \frac{\beta}{2} \sum_{a,b} \mathbb{E}^{ab} \mathbb{E}^{ba} - \frac{\beta}{2} s p_0 = \\ = & \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) V_a^{-1}(\xi) \left(\xi \frac{\partial}{\partial \xi} \right) V_a(\xi) + \beta \sum_{a,b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \frac{\phi_b^-(\xi) \frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} V_b(\eta) V_b^{-1}(\xi) V_a(\xi) \\ & + \frac{\beta}{2} \sum_{a,b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) V_b(\eta) \phi_b^-(\xi) V_b^{-1}(\xi) V_a(\xi) - \frac{\beta}{2} s p_0 \end{aligned}$$

We split the second sum into two parts $a \neq b$ and $a = b$, swap $V_b(\eta)$ and $\phi_b^-(\xi)$ in the third item using commutator relations $[V_b(\eta), \phi_b^-(\xi)] = \frac{1}{1-\frac{\eta}{\xi}} V_b(\eta)$.

$$\begin{aligned} \mathcal{H}_1 = & \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) + \beta \sum_{a \neq b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) \phi_b^-(\xi) \frac{\frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) + \\ & + \beta \sum_a \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_a^-(\xi) \frac{\frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} V_a(\eta) \\ & + \frac{\beta}{2} \sum_{a,b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) \\ & + \frac{\beta}{2} \sum_{a,b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \frac{1}{1 - \frac{\eta}{\xi}} V_b(\eta) V_b^{-1}(\xi) V_a(\xi) - \frac{\beta}{2} s p_0 \end{aligned}$$

The third item vanishes as ξ enters only in negative powers. The last integral can be simplified

$$\oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \frac{1}{1 - \frac{\eta}{\xi}} V_b(\eta) V_b^{-1}(\xi) V_a(\xi) = \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\eta) V_a(\eta) = p_{0,a}$$

Thus it cancels with $-\frac{\beta}{2}sp_0$. We halve the second integral and change the indices of summing a, b and variables ξ, η in the second one. Thus we obtain

$$\begin{aligned}\mathcal{H}_1 = & \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) + \frac{\beta}{2} \sum_{a \neq b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) \phi_b^-(\xi) \frac{\frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) + \\ & + \frac{\beta}{2} \sum_{a \neq b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_b^-(\xi) \phi_a^-(\eta) \frac{\frac{\xi}{\eta}}{1 - \frac{\xi}{\eta}} V_b^{-1}(\xi) V_a(\xi) V_a^{-1}(\eta) V_b(\eta) \\ & + \frac{\beta}{2} \sum_{a \neq b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) + \frac{\beta}{2} \sum_a \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_a^-(\xi) V_a(\eta)\end{aligned}$$

The second, third and forth integrals are combined into an integral with delta-function $\delta\left(\frac{\xi}{\eta}\right) = \sum_{n \in \mathbb{Z}} \frac{\xi^n}{\eta^n}$, which is expressed by the following formula

$$\oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) \phi_b^-(\xi) \delta\left(\frac{\xi}{\eta}\right) V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) = \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) \phi_b^-(\xi) = p_{0,a} p_{0,b},$$

while the last integral equals $p_{0,a}^2 - p_{0,a}$. The final answer is the following

$$\mathcal{H}_1 = \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) + \frac{\beta}{2} (p_0^2 - p_0),$$

where $p_0 = \sum_a p_{0,a}$. Now we calculate the second Hamiltonian:

$$\begin{aligned}\mathcal{H}_2 = & \sum_i D_i^2 + \beta \sum_{i,j,a,b} E_i^{ab} E_j^{ba} D_j - \beta s \sum_i D_i + \frac{\beta^2}{3} \sum_{a,b,c} \mathbb{E}^{ab} \mathbb{E}^{bc} \mathbb{E}^{ca} \\ & - \frac{2s\beta^2}{3} \sum_{a,b} \mathbb{E}^{ab} \mathbb{E}^{ba} + \beta^2 \frac{2s^2 + p_0 - 1}{6} p_0,\end{aligned}\tag{63}$$

In this expression we separately calculate the operator coefficients at β^0, β, β^2 :

$$\mathcal{H} = \mathcal{H}_2 - \beta \mathcal{H}_1 = U_0 + \beta U_1 + \beta^2 U_2$$

$$U_0 = \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) V_a^{-1}(\xi) \left(\xi \frac{\partial}{\partial \xi} \right)^2 V_a(\xi) = \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) (\phi_a^+(\xi))^2 + \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) (\phi_a^+(\xi))'$$

The first three items in (63) contribute to U_1 :

$$\begin{aligned}U_1 = & \sum_{a,b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) \left(\frac{\phi_b^+(\eta) \frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} + \frac{\frac{\eta}{\xi}}{(1 - \frac{\eta}{\xi})^2} \right) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) + \\ & + \sum_{a,b} \oint \frac{d\xi d\eta}{\xi \eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) \frac{\frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} V_b(\eta) V_b^{-1}(\xi) \phi_a^+(\xi) V_a(\xi) +\end{aligned}$$

$$+ \sum_{a,b} \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) V_b(\eta) \phi_b^-(\xi) V_b^{-1}(\xi) \phi_a^+(\xi) V_a(\xi) - (s+1) \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) =$$

In the first item we can take the sum $a \neq b$ as in case $a = b$ it vanishes. In the third integral we use the commutator relations $[V_b(\eta), \phi_b^-(\xi)] = \frac{1}{1-\frac{\eta}{\xi}} V_b(\eta)$ and add it to the second item.

$$\begin{aligned} U_1 &= \sum_{a \neq b} \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) \left(\frac{\phi_b^+(\eta) \frac{\eta}{\xi}}{1 - \frac{\eta}{\xi}} + \frac{\frac{\eta}{\xi}}{(1 - \frac{\eta}{\xi})^2} \right) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) + \\ &\quad + \sum_{a,b} \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) \frac{1}{1 - \frac{\eta}{\xi}} V_b(\eta) V_b^{-1}(\xi) \phi_a^+(\xi) V_a(\xi) + \\ &\quad + \sum_{a,b} \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) \frac{1}{1 - \frac{\eta}{\xi}} V_b(\eta) V_b^{-1}(\xi) \phi_a^+(\xi) V_a(\xi) - (s+1) \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) = \end{aligned}$$

In the second item we change the indices of summing a, b and variables ξ, η and as in H_1 we combine it with the first integral and obtain the delta-function. The third integral can be simplified:

$$\oint \frac{d\eta}{\eta} \phi_a^-(\eta) V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\eta) \phi_a^+(\eta) V_a(\eta) = \oint \frac{d\eta}{\eta} \phi_a^-(\eta) \phi_a^+(\eta)$$

Thus we obtain

$$\begin{aligned} U_1 &= \sum_{a \neq b} \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) \delta\left(\frac{\xi}{\eta}\right) V_b(\eta) V_b^{-1}(\xi) \phi_a^+(\xi) V_a(\xi) \\ &\quad + \sum_{a \neq b} \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) \frac{\frac{\eta}{\xi}}{(1 - \frac{\eta}{\xi})^2} V_b(\eta) V_b^{-1}(\xi) V_a(\xi) - \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) \\ &\quad + \sum_a \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_a^-(\xi) \frac{1}{1 - \frac{\eta}{\xi}} V_a(\eta) \phi_a^+(\xi) \end{aligned}$$

Using commutator relations $[V_a^{-1}(\eta), \phi_a^-(\xi)] = -\frac{1}{1-\frac{\eta}{\xi}} V_a^{-1}(\eta)$ in the last item we obtain

$$\begin{aligned} U_1 &= \sum_{a \neq b} \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_b^-(\xi) \phi_a^+(\xi) - \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) \\ &\quad + \sum_{a > b} \oint_{\eta, \xi} \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) \phi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) \\ &\quad + \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^-(\xi) \phi_a^+(\xi) - \sum_a \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) \frac{1}{(1 - \frac{\eta}{\xi})^2} \phi_a^+(\xi) \\ U_1 &= \sum_{a,b} \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_b^-(\xi) \phi_a^+(\xi) - \sum_a \oint \frac{d\xi}{\xi\eta} \phi_a^-(\xi) (\phi_a^+(\xi))' - \sum_a \oint \frac{d\xi}{\xi} \phi_a^-(\xi) \phi_a^+(\xi) + \end{aligned}$$

$$\begin{aligned}
& \sum_{a>b} \oint_{\eta,\xi} \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) \phi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) V_a^{-1}(\eta) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) \\
U_2 = & \sum_{a,b,c} \oint \frac{d\xi d\eta d\gamma}{\xi\eta\gamma} \frac{\frac{\gamma}{\eta}}{1-\frac{\gamma}{\eta}} \frac{\frac{\eta}{\xi}}{1-\frac{\eta}{\xi}} \phi_a^-(\gamma) V_a^{-1}(\gamma) \phi_c^-(\eta) V_c(\gamma) V_c^{-1}(\eta) \phi_b^-(\xi) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) \\
& + \sum_{a,b,c} \oint \frac{d\xi d\eta d\gamma}{\xi\eta\gamma} \frac{\frac{\eta}{\xi}}{1-\frac{\eta}{\xi}} \phi_a^-(\gamma) V_a^{-1}(\gamma) V_c(\gamma) \phi_c^-(\eta) V_c^{-1}(\eta) \phi_b^-(\xi) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) \\
& - s \sum_{a,b} \oint \frac{d\xi d\eta}{\xi\eta} \frac{\frac{\eta}{\xi}}{1-\frac{\eta}{\xi}} \phi_a^-(\eta) V_a^{-1}(\eta) \phi_b^-(\xi) V_b(\eta) V_b^{-1}(\xi) V_a(\xi) \\
& + \frac{1}{3} \sum_{a,b,c} \oint \frac{d\xi d\eta d\gamma}{\xi\eta\gamma} \phi_a^-(\gamma) V_a^{-1}(\gamma) V_c(\gamma) \phi_c^-(\eta) V_c^{-1}(\eta) V_b(\eta) \phi_b^-(\xi) V_b^{-1}(\xi) V_a(\xi) \\
& - \frac{2}{3} \sum_{a,b} \oint \frac{d\xi d\eta}{\xi\eta} \phi_a^-(\eta) V_a^{-1}(\eta) V_b(\eta) \phi_b^-(\xi) V_b^{-1}(\xi) V_a(\xi) + \frac{s^2 - p_0 + 1}{3} p_0
\end{aligned}$$

A.3 Calculations for classical systems

Here we provide a calculation for the Poisson brackets in the classical limit.

$$\{\varphi_a(x), \varphi_b(y)\} = \delta'(x/y) \delta_{a,b}.$$

Calculate Poisson bracket with the Hamiltonian $\{\phi_a^-(x), \mathcal{H}\}$, $\{\phi_a^+(x), \mathcal{H}\}$:

$$\begin{aligned}
\{\varphi_a^-(x), \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) (\varphi_a^+(\xi))^2\} &= -2 \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) \varphi_a^+(\xi) \sum_{n>0} n \frac{\xi^n}{x^n} \\
\{\varphi_a^+(x), \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) (\varphi_a^+(\xi))^2\} &= \oint_{\xi} \frac{d\xi}{\xi} (\varphi_a^+(\xi))^2 \sum_{n>0} n \frac{x^n}{\xi^n} = x \frac{d}{dx} (\varphi_a^+(x))^2 \\
\{\varphi_a^-(x), \oint_{\xi} \frac{d\xi}{\xi} (\varphi_a^-(\xi))^2 \varphi_a^+(\xi)\} &= - \oint_{\xi} \frac{d\xi}{\xi} (\varphi_a^-(\xi))^2 \sum_{n>0} n \frac{\xi^n}{x^n} = x \frac{d}{dx} (\varphi_a^-(x))^2 \\
\{\varphi_a^+(x), \oint_{\xi} \frac{d\xi}{\xi} (\varphi_a^-(\xi))^2 \varphi_a^+(\xi)\} &= 2 \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) \varphi_a^+(\xi) \sum_{n>0} n \frac{x^n}{\xi^n}
\end{aligned}$$

From these four formulas we obtain

$$\{\varphi_a(x), \oint_{\xi} \frac{d\xi}{\xi} ((\varphi_a^-(\xi))^2 \varphi_a^+(\xi) + \varphi_a^-(\xi) (\varphi_a^+(\xi))^2)\} = x \frac{d}{dx} (\varphi_a(x))^2.$$

Summing the following two equations

$$\{\varphi_a^-(x), \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) (\varphi_a^+(\xi))'\} = - \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) \xi \frac{d}{d\xi} \left(\sum_{n>0} n \frac{\xi^n}{x^n} \right) = - \left(x \frac{d}{dx} \right)^2 \varphi_a^-(x)$$

$$\{\varphi_a^+(x), \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) (\varphi_a^+(\xi))'\} = \oint_{\xi} \frac{d\xi}{\xi} (\varphi_a^+(\xi))' \left(\sum_{n>0} n \frac{x^n}{\xi^n} \right) = \left(x \frac{d}{dx} \right)^2 \varphi_a^+(x)$$

we obtain

$$\{\varphi_a(x), \oint_{\xi} \frac{d\xi}{\xi} \varphi_a^-(\xi) (\varphi_a^+(\xi))'\} = \left(x \frac{d}{dx} \right)^2 (\varphi_a^+(x) - \varphi_a^-(x)).$$

The next items are the following

$$\{\varphi_a^-(x), \oint_{\eta} \frac{d\eta}{\eta} \varphi_a^-(\eta) \varphi_b^-(\eta) \varphi_a^+(\eta)\} = - \oint_{\eta} \frac{d\eta}{\eta} \varphi_a^-(\eta) \varphi_b^-(\eta) \sum_{n>0} n \frac{\eta^n}{x^n} = x \frac{d}{dx} (\varphi_a^-(x) \varphi_b^-(x))$$

$$\{\varphi_a^+(x), \oint_{\eta} \frac{d\eta}{\eta} \varphi_a^-(\eta) \varphi_b^-(\eta) \varphi_a^+(\eta)\} = \oint_{\eta} \frac{d\eta}{\eta} \varphi_a^+(\eta) \varphi_b^-(\eta) \sum_{n>0} n \frac{x^n}{\eta^n} = x \frac{d}{dx} (\varphi_a^+(x) \varphi_b^-(x))^+$$

$$\{\varphi_a^+(x), \oint_{\eta} \frac{d\eta}{\eta} \varphi_b^-(\eta) \varphi_a^-(\eta) \varphi_b^+(\eta)\} = x \frac{d}{dx} (\varphi_b^+(x) \varphi_b^-(x))^+$$

Poisson brackets with the the double integral are:

$$\begin{aligned} & \{\varphi_a^-(x), \oint_{\eta, \xi} \frac{d\xi d\eta}{\xi \eta} \varphi_a^-(\eta) \varphi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) \mathcal{V}_a^{-1}(\eta) \mathcal{V}_b(\eta) \mathcal{V}_b^{-1}(\xi) \mathcal{V}_a(\xi)\} \\ &= \oint_{\eta, \xi} \frac{d\xi d\eta}{\xi \eta} \left(\frac{1}{1 - \frac{\eta}{x}} - \frac{1}{1 - \frac{\xi}{x}} \right) \varphi_a^-(\eta) \varphi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) \mathcal{V}_a^{-1}(\eta) \mathcal{V}_b(\eta) \mathcal{V}_b^{-1}(\xi) \mathcal{V}_a(\xi) \\ &= \oint_{\eta} \frac{d\eta}{\eta} \frac{1}{1 - \frac{\eta}{x}} \mathcal{V}_a^{-1}(\eta) \mathcal{V}_b(\eta) \varphi_a^-(\eta) \left(\eta \frac{d}{d\eta} (\varphi_b^-(\eta) \mathcal{V}_b^{-1}(\eta) \mathcal{V}_a(\eta))^{\eta+} - \eta \frac{d}{d\eta} (\varphi_b^-(\eta) \mathcal{V}_b^{-1}(\eta) \mathcal{V}_a(\eta))^{\eta-} \right) \\ &\quad - \oint_{\xi} \frac{d\xi}{\xi} \frac{1}{1 - \frac{\xi}{x}} \varphi_b^-(\xi) \mathcal{V}_b^{-1}(\xi) \mathcal{V}_a(\xi) \left(\xi \frac{\partial}{\partial \xi} (\varphi_a^-(\xi) \mathcal{V}_a^{-1}(\xi) \mathcal{V}_b(\xi))^{\xi+} - \xi \frac{\partial}{\partial \xi} (\varphi_a^-(\xi) \mathcal{V}_a^{-1}(\xi) \mathcal{V}_b(\xi))^{\xi-} \right) \\ &= \left(\varphi_a^-(x) \mathcal{V}_a^{-1}(x) \mathcal{V}_b(x) x \frac{\partial}{\partial x} \left((\varphi_b^-(x) \mathcal{V}_b^{-1}(x) \mathcal{V}_a(x))^+ - (\varphi_b^-(x) \mathcal{V}_b^{-1}(x) \mathcal{V}_a(x))^- \right) \right)^- \\ &\quad - \left(\varphi_b^-(x) \mathcal{V}_b^{-1}(x) \mathcal{V}_a(x) x \frac{\partial}{\partial x} \left((\varphi_a^-(x) \mathcal{V}_a^{-1}(x) \mathcal{V}_b(x))^+ - (\varphi_a^-(x) \mathcal{V}_a^{-1}(x) \mathcal{V}_b(x))^- \right) \right)^-. \\ & \{\varphi_a^+(x), \oint_{\eta, \xi} \frac{d\xi d\eta}{\xi \eta} \varphi_a^-(\eta) \varphi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) \mathcal{V}_a^{-1}(\eta) \mathcal{V}_b(\eta) \mathcal{V}_b^{-1}(\xi) \mathcal{V}_a(\xi)\} \\ &= \oint_{\eta, \xi} \frac{d\xi d\eta}{\xi \eta} \sum_{n>0} n \frac{x^n}{\eta^n} \varphi_b^-(\xi) \sum_k k \left(\frac{\xi^k}{\eta^k} + \frac{\eta^k}{\xi^k} \right) \mathcal{V}_a^{-1}(\eta) \mathcal{V}_b(\eta) \mathcal{V}_b^{-1}(\xi) \mathcal{V}_a(\xi) \\ &= \oint_{\eta} \frac{d\eta}{\eta} \sum_{n>0} n \frac{x^n}{\eta^n} \mathcal{V}_a^{-1}(\eta) \mathcal{V}_b(\eta) \left(\eta \frac{d}{d\eta} (\varphi_b^-(\eta) \mathcal{V}_b^{-1}(\eta) \mathcal{V}_a(\eta))^{\eta+} - \eta \frac{d}{d\eta} (\varphi_b^-(\eta) \mathcal{V}_b^{-1}(\eta) \mathcal{V}_a(\eta))^{\eta-} \right) \end{aligned}$$

$$= x \frac{\partial}{\partial x} \left(\mathcal{V}_a^{-1}(x) \mathcal{V}_b(x) x \frac{\partial}{\partial x} \left((\varphi_b^-(x) \mathcal{V}_b^{-1}(x) \mathcal{V}_a(x))^+ - (\varphi_b^-(x) \mathcal{V}_b^{-1}(x) \mathcal{V}_a(x))^- \right) \right)^+.$$

The final answer is given in proposition 6. Both sides of the equality $\frac{dL}{dt} = [M, L]$ are equal to

$$\begin{aligned} & \sum_a \mathcal{V}_a (\varphi_a^+)^2 (\varphi_a^- \mathcal{V}_a^{-1} f)^+ + \sum_a \mathcal{V}_a z \frac{\partial \varphi_a^+}{\partial z} (\varphi_a^- \mathcal{V}_a^{-1} f)^+ + 2 \sum_a \mathcal{V}_a \left(z \frac{\partial}{\partial z} (\varphi_a^- \varphi_a^+)^- \mathcal{V}_a^{-1} f \right)^+ \\ & - \sum_a \mathcal{V}_a \left(\left(z \frac{\partial}{\partial z} \right)^2 (\varphi_a^-) \mathcal{V}_a^{-1} f \right)^+ - \sum_a \mathcal{V}_a \left((\varphi_a^+)^2 \varphi_a^- \mathcal{V}_a^{-1} f \right)^+ - \sum_a \mathcal{V}_a \left(z \frac{\partial \varphi_a^+}{\partial z} \varphi_a^- \mathcal{V}_a^{-1} f \right)^+ \\ & + \sum_a \mathcal{V}_a \left(z \frac{\partial (\varphi_a^-)^2}{\partial z} \mathcal{V}_a^{-1} f \right)^+ + 2 \sum_{a,b} \mathcal{V}_a (\varphi_b^- \varphi_b^+)_+ (\varphi_a^- \mathcal{V}_a^{-1} f)^+ - 2 \sum_{a,b} \mathcal{V}_a ((\varphi_b^- \varphi_b^+)_+ \varphi_a^- \mathcal{V}_a^{-1} f)^+ \\ & + 2 \sum_{a \neq b} \mathcal{V}_b \left(z \frac{\partial}{\partial z} (\varphi_b^- \mathcal{V}_{ab}) (\varphi_a^- \mathcal{V}_a^{-1} f)^+ \right)^+. \end{aligned}$$

A.4 Hamiltonians via q -determinant

Since $d_i = D_i + \beta \sum_{j < i} K_{ij}$, $H_1 = \sum_i d_i = \sum_i D_i + \beta \sum_{i,j < i} K_{ij} = \sum_i D_i + \frac{\beta}{2} \sum_{i \neq j} K_{ij}$. Here we use the relation $K_{ij} f = P_{ij} f = \sum_{a,b} E_i^{ab} E_j^{ba} f$ on functions $f \in M_N^S$.

$$H_1 = \sum_i D_i + \frac{\beta}{2} \sum_{i \neq j} P_{ij} = \sum_i D_i + \frac{\beta}{2} \sum_{i \neq j, a, b} E_i^{ab} E_j^{ba} = \sum_i D_i + \frac{\beta}{2} \sum_{a, b} \mathbb{E}^{ab} \mathbb{E}^{ba} - \frac{\beta}{2} s p_0.$$

In the last equality we use $\sum_i E_i^{ab} E_i^{ba} = s p_0$, where s and $p_0 = \sum_a p_{0,a}$ is the number of spins and particles, respectively.

$$H_2 = \sum_i D_i^2 + \beta \sum_{i,j < i} K_{ij} D_i + \beta \sum_{i,j < i} D_i K_{ij} + \beta^2 \sum_{i,j < i, k < i} K_{ij} K_{ik}.$$

Using relations (7) in the third item we obtain

$$H_2 = \sum_i D_i^2 + \beta \sum_{i \neq j} K_{ij} D_i + \beta^2 \sum_{i,j < i, k < i} K_{ij} K_{ik}.$$

The last item can be simplified using commutator relations $E_i^{ab} E_j^{cd} = \delta_{ij} (\delta_{bc} E_i^{ad} - \delta_{ad} E_i^{bc})$:

$$\begin{aligned} \sum_{i,j < i, k < i} K_{ij} K_{ik} &= \frac{1}{3} \sum_{i \neq j, i \neq k, j \neq i} P_{ij} P_{ik} + \frac{p_0(p_0 - 1)}{2} = \frac{1}{3} \sum_{i \neq j, i \neq k, j \neq i, a, b, c, d} E_i^{ab} E_j^{ba} E_i^{cd} E_k^{dc} + \frac{p_0(p_0 - 1)}{2} = \\ &= \frac{1}{3} \sum_{i \neq j, i \neq k, j \neq i, a, b, c} E_i^{ab} E_j^{bc} E_k^{ca} + \frac{p_0(p_0 - 1)}{2} = \end{aligned}$$

$$= \frac{1}{3} \sum_{i,j,k,a,b,c} E_i^{ab} E_j^{bc} E_k^{ca} - \frac{2s}{3} \sum_{i,j,a,b} E_i^{ab} E_j^{ba} - \frac{1}{3} \sum_{i,j,a,b} E_i^{aa} E_j^{bb} + \frac{s^2 p_0}{3} + \frac{p_0}{3} + \frac{p_0(p_0 - 1)}{2}.$$

Thus we obtain

$$H_2 = \sum_i D_i^2 + \beta \sum_{i,j,a,b} E_i^{ab} E_j^{ba} D_j - \beta s \sum_i D_i + \frac{\beta^2}{3} \sum_{a,b,c} \mathbb{E}^{ab} \mathbb{E}^{bc} \mathbb{E}^{ca} - \frac{2s\beta^2}{3} \sum_{a,b} \mathbb{E}^{ab} \mathbb{E}^{ba} + \beta^2 \frac{2s^2 + p_0 - 1}{6} p_0.$$

Now we show how to obtain the first elements of the q -determinant

$$q\text{-det } T(u) = \sum_{\sigma \in S_m} (-1)^{\text{sgn}(\sigma)} T^{\sigma(1),1}(u) T^{\sigma(2),2}(u-1) \dots T^{\sigma(m),m}(u-m+1) = 1 + \frac{\Delta_0}{u} + \frac{\Delta_1}{u^2} + \frac{\Delta_2}{u^3} + \dots$$

where $T^{ab}(u) = \delta^{ab} + \frac{T_0^{ab}}{u} + \frac{T_1^{ab}}{u^2} + \frac{T_2^{ab}}{u^3} + \dots$, $\Delta_0 = \sum_a T_0^{aa}$, and

$$\begin{aligned} \Delta_1 &= - \sum_{a>b} T_0^{ab} T_0^{ba} + \sum_{a<b} T_0^{aa} T_0^{bb} + \sum_a T_1^{aa} + \sum_{a>1} (a-1) T_0^{aa} = -\frac{1}{2} \sum_{a,b} T_0^{ab} T_0^{ba} + \frac{1}{2} \sum_{a,b} T_0^{aa} T_0^{bb} \\ &\quad - \frac{1}{2} \sum_{a>b} (T_0^{aa} - T_0^{bb}) + \sum_a T_1^{aa} + \sum_{a>1} (a-1) T_0^{aa} = \sum_a T_1^{aa} - \frac{1}{2} \sum_{a,b} T_0^{ab} T_0^{ba} + \frac{1}{2} (\Delta_0^2 + (s-1) \Delta_0), \\ \Delta_2 &= \sum_a T_2^{aa} + \sum_{a>1} 2(a-1) T_1^{aa} + \sum_{a<b} (T_0^{aa} T_1^{bb} + T_1^{aa} T_0^{bb}) - \sum_{a>b} (T_0^{ab} T_1^{ba} + T_1^{ab} T_0^{ba}) + \\ &\quad \sum_{a<b} (a+b-2) T_0^{aa} T_0^{bb} - \sum_{a>b} (a+b-2) T_0^{ab} T_0^{ba} + \sum_{a>1} (a-1)^2 T_0^{aa} + \sum_{a>b>c, \sigma \in S_3} (-1)^{\text{sgn}(\sigma)} T_0^{\sigma(c)c} T_0^{\sigma(b)b} T_0^{\sigma(a)a}. \end{aligned}$$

Using the relation $[T_1^{ab}, T_0^{cd}] = \delta_{cb} T_1^{ad} - \delta_{ad} T_1^{cb}$ we obtain

$$\begin{aligned} \Delta_2 &= \sum_a T_2^{aa} - \sum_{a,b} T_0^{ab} T_1^{ba} + (s-1) \sum_{a>1} T_1^{aa} + \sum_{a,b} T_0^{aa} T_1^{bb} + \\ &\quad + \sum_{a<b} (a+b-2) T_0^{aa} T_0^{bb} - \sum_{a>b} (a+b-2) T_0^{ab} T_0^{ba} + \sum_{a>1} (a-1)^2 T_0^{aa} + \frac{1}{6} \sum_{a,b,c} T_0^{aa} T_0^{bb} T_0^{cc} \\ &\quad + \frac{1}{3} \sum_{a,b,c} T_0^{ab} T_0^{bc} T_0^{ca} - \frac{1}{2} \sum_{a,b,c} T_0^{cc} T_0^{ab} T_0^{ba} + \frac{1}{6} \sum_{a,b} T_0^{aa} T_0^{bb} - \frac{1}{3} \sum_{a,b} T_0^{ab} T_0^{ba} + \frac{2-s}{6} \sum_a T_0^{aa} T_0^{aa} \\ &\quad + \frac{1}{3} \sum_{a>b} (3a+3b-1-4s) T_0^{ab} T_0^{ba} - \sum_{b>a} (a+b-1-s) T_0^{aa} T_0^{bb} + \frac{1}{3} \sum_{a>b} -2(s-a)(T_0^{bb} - T_0^{aa}) \\ &= \sum_a T_2^{aa} - \sum_{a,b} T_0^{ab} T_1^{ba} + \frac{1}{3} \sum_{a,b,c} T_0^{ab} T_0^{bc} T_0^{ca} + (s-1) \sum_{a>1} T_1^{aa} + \Delta_0 \sum_a T_1^{aa} - \Delta_0 \frac{1}{2} \sum_{a,b} T_0^{ab} T_0^{ba} \end{aligned}$$

$$\begin{aligned}
& +(s-1) \sum_{a < b} T_0^{aa} T_0^{bb} + \frac{5-4s}{3} \sum_{a > b} T_0^{ab} T_0^{ba} + \frac{1}{6} \sum_{a,b} T_0^{aa} T_0^{bb} - \frac{1}{3} \sum_{a,b} T_0^{ab} T_0^{ba} + \frac{2-s}{6} \sum_a T_0^{aa} T_0^{aa} \\
& + \sum_{a > 1} (a-1)^2 T_0^{aa} + \frac{1}{3} \sum_a (a-s)(s-a-1) T_0^{aa} + \frac{1}{3} \sum_a 2(s-a)(a-1) T_0^{aa} + \frac{1}{6} \Delta_0^3.
\end{aligned}$$

After these calculations we obtain the following formula for Δ_2 :

$$\begin{aligned}
\Delta_2 &= \sum_a T_2^{aa} - \sum_{a,b} T_0^{ab} T_1^{ba} + \frac{1}{3} \sum_{a,b,c} T_0^{ab} T_0^{bc} T_0^{ca} + \Delta_0 \sum_a T_1^{aa} - \Delta_0 \frac{1}{2} \sum_{a,b} T_0^{ab} T_0^{ba} \\
& (s-1) \sum_{a > 1} T_1^{aa} + \frac{3-4s}{6} \sum_{a,b} T_0^{ab} T_0^{ba} + \frac{1}{6} \Delta_0^3 + \frac{3s-2}{6} \Delta_0^2 + \frac{(s-1)(2s-1)}{6} \Delta_0, \\
\Delta_2 &= \sum_a T_2^{aa} - \sum_{a,b} T_0^{ab} T_1^{ba} + \frac{1}{3} \sum_{a,b,c} T_0^{ab} T_0^{bc} T_0^{ca} + s \sum_{a > 1} T_1^{aa} - \frac{2s}{3} \sum_{a,b} T_0^{ab} T_0^{ba} + \\
& + \Delta_0 \Delta_1 - \Delta_1 - \frac{1}{3} \Delta_0^3 + \frac{2}{3} \Delta_0^2 + \frac{(s^2-1)}{3} \Delta_0.
\end{aligned}$$

The Hamiltonians can be expressed in terms of Δ_i :

$$\begin{aligned}
\frac{H_1}{\beta} &= -\Delta_1 + \frac{1}{2} \Delta_0^2 - \frac{1}{2} \Delta_0, \\
\frac{H_2}{\beta^2} &= \Delta_2 - \Delta_0 \Delta_1 + \Delta_1 + \frac{1}{3} \Delta_0^3 - \frac{1}{2} \Delta_0^2 + \frac{1}{6} \Delta_0.
\end{aligned}$$

Acknowledgements

The authors thank M.L.Nazarov for valuable remarks and discussions, and I.Marshall and A.Schwarz for the careful proofreading. The research by M.M. was supported in part by the Simons Foundation and RFBR grant 16-01-00562. S.K appreciates the support of RSF grant, project 16-11-10316 dated 11.05.2016.

References

- [1] I. Andric, A. Jevicki and H. Levine, On the large-N limit in symplectic matrix models, Nucl. Phys. **B215** (1983) 307; I. Andric and V. Bardek, J. Phys. **A21** (1988) 2847
- [2] T. Arakawa, Drinfeld functor and finite-dimensional representations of the Yangian, Commun. Math. Phys. **205** (1999), 1–18.
- [3] H. Awata, Y. Matsuo, S. Odake, J. Shiraishi Collective field theory, Calogero-Sutherland model and generalized matrix models, Physics Letters B 347.1 (1995), 49-55.
- [4] Bernard, D., Gaudin, M., Haldane, F. D. M., Pasquier, V. Yang-Baxter equation in spin chains with long range interactions, J. Phys. A 26 (1993), 5219. arXiv preprint hep-th/9301084.
- [5] H. Awata, Y. Matsuo and T. Yamamoto, Collective field description of spin Calogero-Sutherland models, J. Phys. **A29** (1996) 3089-3098
- [6] Y. Kato and Y. Kuramoto Exact solution of the Sutherland model with arbitrary internal symmetry Phys. Rev. Lett. **74** 1222 (1995)
- [7] M.L. Nazarov and E. K. Sklyanin Sekiguchi-Debiard operators at infinity, Communications in Mathematical Physics 324.3 (2013): 831-849.
- [8] M.L. Nazarov and E.K. Sklyanin Integrable hierarchy of the quantum Benjamin-Ono equation Symmetry, Integrability and Geometry: Methods and Applications SIGMA **9** (2013), 078,
- [9] V. Pasquier A lecture on Calogero-Sutherland models, Integrable Models and Strings. Springer Berlin Heidelberg, 1994. 36-48. eprint hep-th/9405104,.
- [10] A.N. Sergeev, A.P. Veselov Calogero-Moser operators in infinite dimension, eprint arXiv:0910.1984. – 2009.
- [11] A. N. Sergeev, A. P. Veselov Dunkl operators at infinity and Calogero-Moser systems, IMRN. – 2015 ., eprint arXiv:1311.0853.
- [12] Uglov D. Yangian actions on higher level irreducible integrable modules of affine $\widehat{\mathfrak{gl}}_N$, eprint arXiv preprint math/9802048. – 1998.