

This is a repository copy of *Competition and habitat availability interact to structure arboreal ant communities across scales of ecological organization*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/203545/>

Version: Published Version

Article:

Adams, Benjamin J, Gora, Evan M, Donaldson-Matasci, Matina C et al. (2 more authors) (2023) Competition and habitat availability interact to structure arboreal ant communities across scales of ecological organization. *Proceedings of the Royal Society B: Biological Sciences*. 20231290. ISSN 1471-2954

<https://doi.org/10.1098/rspb.2023.1290>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Research



Cite this article: Adams BJ, Gora EM, Donaldson-Matasci MC, Robinson EJH, Powell S. 2023 Competition and habitat availability interact to structure arboreal ant communities across scales of ecological organization. *Proc. R. Soc. B* **290**: 20231290. <https://doi.org/10.1098/rspb.2023.1290>

Received: 12 June 2023
Accepted: 25 August 2023

Subject Category:
Ecology

Subject Areas:
ecology

Keywords:
community ecology, habitat filtering, competition, resource limitation, species coexistence

Author for correspondence:
Benjamin J. Adams
e-mail: b.maxadams@gmail.com

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6824120>.

Competition and habitat availability interact to structure arboreal ant communities across scales of ecological organization

Benjamin J. Adams¹, Evan M. Gora^{2,3}, Matina C. Donaldson-Matasci⁴, Elva J. H. Robinson⁵ and Scott Powell¹

¹Department of Biological Sciences, George Washington University, Washington, DC, USA

²Smithsonian Tropical Research Institute, Balboa, Panama

³Cary Institute of Ecosystem Studies, Millbrook, NY, USA

⁴Department of Biology, Harvey Mudd College, Claremont, CA 91711, USA

⁵Department of Biology, University of York, York, UK

BJA, 0000-0002-5857-175X; EMG, 0000-0002-0537-5835; MCD, 0000-0002-9877-384X; EJHR, 0000-0003-4914-9327; SP, 0000-0001-5970-8941

Understanding how resource limitation and biotic interactions interact across spatial scales is fundamental to explaining the structure of ecological communities. However, empirical studies addressing this issue are often hindered by logistical constraints, especially at local scales. Here, we use a highly tractable arboreal ant study system to explore the interactive effects of resource availability and competition on community structure across three local scales: an individual tree, the nest network created by each colony and the individual ant nest. On individual trees, the ant assemblages are primarily shaped by availability of dead wood, a critical nesting resource. The nest networks within a tree are constrained by the availability of nesting resources but also influenced by the co-occurring species. Within individual nests, the distribution of adult ants is only affected by distance to interspecific competitors. These findings demonstrate that resource limitation exerts the strongest effects on diversity at higher levels of local ecological organization, transitioning to a stronger effect of species interactions at finer scales. Collectively, these results highlight that the process exerting the strongest influence on community structure is highly dependent on the scale at which we examine the community, with shifts occurring even across fine-grained local scales.

1. Introduction

Resource-mediated habitat filtering and competition are expected to interact in the structuring of ecological communities [1–3]. The resources available in a habitat set the foundation for which species can live there, based on the fundamental niche and minimum resources required for long-term persistence of each species [4,5]. For each species that finds suitable resources, their persistence in the community can then be challenged by competition, potentially leading to exclusion [6–8]. Considerable work has focused on detecting the signatures of habitat filtering, competition and their interactions at large spatial scales [9,10]. Such work summarizes broad hierarchical patterns across a diversity of habitats, with habitat filtering generally being more important at large scales but with the balance shifting to competition at finer scales. However, the direct mechanisms underlying patterns of diversity are frequently obscured at broader spatial scales [11,12]. At the other extreme, local spatial scales provide opportunities to directly link known resource usage and competitive interactions among individuals with observable community patterns [13,14].

Nevertheless, studies that address the relative importance of resource availability and competition in structuring communities across multiple local scales are rare (but see work with communities in pitcher plants [15] and rock pools [16]). Filling this knowledge gap is critical for understanding the extent to which local processes and their interactions scale to influence the structure of ecological communities [14,17].

At local spatial scales, the influences of resource availability and biotic interactions on community structure are thought to be spatially dependent and complex [14]. Available resources often occur in discrete patches that limit access for species in the community, especially if dispersal to new resource patches is challenging [18,19]. Although frequently simplified to pairwise, linear interactions between species [20], communities generally exist as an assemblage of multiple species differentially interacting across space [21,22]. Indeed, how species use and partition resources in space is often complex and variable at local scales, but one common and understudied context is when individuals acquire resources and grow via establishing and expanding networked patches or nodes of resources. Examples of network patterns of resource acquisition and use span such disparate systems as fungal hyphae and root networks [23] and the foraging and multi-nest networks of social insect colonies [24–26]. Competition in these cases then plays out via interactions between abutting or intertwined networks, with any pattern of network growth facing an array of competitive pressures from multiple species and locations simultaneously. How resource distribution, network growth and competitive interactions shape local-scale patterns of coexistence should thus be highly dependent on fine-scale spatial relationships. Systems that incorporate these layers of local community interactions should then be ideal for addressing the broader knowledge gap of how local processes scale to community-level influences.

Arboreal ant communities are particularly tractable systems for studying the interactions between resource availability and competition in the context of growth via discrete expansion of resource networks [27]. Individual trees function as the primary habitat patches for arboreal ants to colonize and compete over. This is true even within dense forest environments, due to the phenomena of crown shyness and because arboreal ants rarely leave the crowns of trees [28,29]. As a result, arboreal ant communities within an individual tree typically act as isolated ‘island-like’ communities following classic species–area relationships [30–32]. Within a tree, resident colonies of ants also compete fiercely for a limited number of pre-existing nesting cavities [33–36]. Each colony must acquire and defend multiple nest sites, without the capacity to make more themselves, in order to grow and successfully reproduce [37,38], building a nest network as they do so. Finally, individual nests, which represent the finest spatial scale, vary in properties of quality [39] and defensibility [38,40], as well as in the competitive pressures they face [41]. Colonies must make collective decisions about which individual cavities they use within the resource-limited and highly competitive environment they occupy, and how they allocate colony members to a nest to maximize overall colony growth and reproduction [38,39,41].

Here, we use an arboreal ant study system, where growth is via network expansion, to explore the interactive effects of resource availability and competition on community structure at multiple spatial scales. More specifically, we focus

on the following three local scales: (i) a discrete resource patch, represented by the whole tree, (ii) resource networks built by organisms, represented by the within-tree nest networks of the resident ant colonies, and (iii) an individual resource within a network, represented by the individual nest. Our central hypothesis is that resource availability has an overarching influence on local community structure, and that species interactions emerge as more important at finer-grained local scales. We tested this hypothesis by surveying arboreal ant communities, quantifying nest site availability, mapping the spatial distribution of nests and quantifying the contents of individual nests. At the patch scale, we evaluated how the availability of resources and the competitive context on whole trees influences the ant community. At the scale of the resource network, we mapped nest networks and evaluated how these are shaped by tree characteristics, ant species and competitive context. Finally, at the scale of an individual resource or node within a resource network, we examined how competition and habitat limitations influenced the populations of adult ants and brood within individual nests. We expect that resource availability will be more important than competition for determining ant community metrics, such as species richness and nest abundance, measured at the scale of an individual tree. By contrast, we expect competition to determine nest selection and ant distribution, with less competitive ant species selecting nest sites and distributing adult ants and brood further away from competitors. Taken together, these detailed data across multiple spatial scales provide an integrative approach to identifying how local resource availability and competition shape community structure, including richness, composition and physical location within a habitat.

2. Methods

(a) Study site and focal species

We conducted all field work at the Dagny Johnson Key Largo Hammock Botanical State Park in Key Largo, Florida, USA (25.178°N, 80.366°W; hereafter Dagny State Park). The Florida Keys are marked by a mild subtropical climate with mean monthly temperatures ranging between 17.9°C and 31.9°C with approximately 101.2 cm of precipitation annually (<https://climatecenter.fsu.edu/products-services/data/1981-2010-normals/key-west>). Dagny State Park was established in 1982 and hosts the largest remnant in the United States of West Indian hardwood hammock forest [42–44].

The hardwood hammock forest of the Florida Keys present a novel opportunity to work in an arboreal ant system that overcomes many of the logistical constraints of other arboreal ant communities. For example, in the highly diverse tropical forest habitats where most arboreal ant research has focused, felling trees [45] or specialized equipment and training [46–48] are required simply to access the nesting ecology of the ants. Across all tropical habitats, including those that are more accessible, community diversity of more than a hundred species and tree-level diversity of 20 species or more [30,32,40,49] remains a challenge for understanding detailed species interaction or resource requirement. By contrast, the hardwood hammock forest of the Florida Keys is a species-rich ecosystem of conservation concern [43] that has a low and easily accessible canopy (does not exceed 10 m and is frequently less than 6 m; electronic supplementary material, figure S1) [42]. While the arboreal ant diversity is reduced compared with tropical habitats, tropical arboreal ant genera are still well-represented in the full community [50], and all species rely on the same nesting resource for

growth and reproduction: hollow cavities in dead stems that are often the abandoned feeding tunnels of wood-boring beetles [51]. Capturing all fine-grained local-scale interactions for even this reduced community of Florida Keys hammock forest could be overwhelming logistically, but most ant communities are dominated by a smaller subset of especially abundant species that capture an array of competitive interactions. Our surveys revealed four especially common species in the hammock forest system that will be our focus here. In addition to high colony incidence across surveyed trees, including frequent co-occurrence (below), these four species also represented contrasting ecology within the larger community. Thus, while this study is not an exhaustive study of the full arboreal ant community in the system, it uses the most abundant players in the community that span an array of potentially generalizable interactions and outcomes. Our four focal species are as follows: (i) *Pseudomyrmex ejectus* and (ii) *Pseudomyrmex simplex*, which are established native species that share similar niche space [52–54], (iii) *Pseudomyrmex gracilis*, which is a disruptive non-native with invasive potential [55,56] and (iv) *Cephalotes varians*, a native species with known defence specialization in its use of nesting resources [57–59].

(b) Identifying ant colonies and nest locations

We used a combination of multiple baiting and hand-collecting methods [38,60] to document ants foraging and nesting on 176 individual poisonwood trees (*Metopium toxiferum*) in relatively open areas of the hammock forest, in which individual trees are typically physically isolated from other trees. We specifically targeted trees that were not embedded in the larger forest canopy to eliminate any potential connectivity between neighbouring crowns and ensure that the only ants foraging at baits were nesting within the tree [32,61]. Baits (a combination of approx. 140 g of canned chicken and approx. 60 ml of honey with urine added as an additional attractant for *C. varians*) were placed throughout the entire crown of each tree at 12.00 and were examined and refreshed until 21.00 in order to document activity of both the diurnal (*P. ejectus*, *P. gracilis* and *P. simplex*) and nocturnal species (*C. varians*). We selected a subset of the trees ($n=31$) based on ease of access to the entire crown of the tree and a stratified sampling of tree sizes, and we then tracked foraging ants back to their nests. This method was used to locate all nests of all four of our focal species on each tree and to look for aggressive interactions among conspecifics at baits to ensure all conspecific ants within a tree were from the same colony [38]. Voucher specimens of all ants found at baits or on any other part of the tree were collected and stored in 95% ethanol to be identified in the lab using keys and voucher specimens [62]. It is worthwhile to note that, within the genera of interest, only our focal species of *Cephalotes* is found in the system, and that only our three focal species of the genus *Pseudomyrmex* were found on our study trees in the hammock forest, even though other members of the genus are found in the Florida Keys more generally [50].

(c) Measurements across local spatial scales

(i) Resource patch scale: whole tree

For each tree included in the initial survey, GPS coordinates were recorded, and the diameter of the trunk at 10 cm above the ground was measured. Although diameter at breast height (1.3 m above the ground) is a more common measure, nearly all of the trees in this forest branch below breast height (average distance to first branch = 83.6 cm; range = 10–173 cm in the present study). We used the diameter measured at 10 cm to calculate basal area ($BA = \pi(D/2)^2$) as a proxy for tree size and an estimate of the total resource patch size for the local ant community [30,63].

For each tree, we also quantified the total amount of dead wood (i.e. the nest resources available to the ant community)

in the crown using three methods that required increasing degrees of time and effort in the field but provide increasing resolution of the total resource availability. First, using visual surveys conducted by at least two individuals, we estimated the total per cent dieback for each tree crown to the nearest 5% and took the average between the two when different [64]. We multiplied this percentage by the basal area to produce a weighted proxy of available resources that accounted only for dead wood (i.e. the actual nesting resource). We also counted the total number of dead stems present in each tree crown. Finally, we quantified the total volume of dead wood in each tree by measuring every piece of dead wood by hand and calculating individual stem volume using Newton's Formula ($L(A_{\text{base}} + 4A_{\text{middle}} + A_{\text{tip}})/6$) which was then summed for the whole tree [65].

(ii) Resource network scale: nest network

In ant nest networks, reducing the number of nodes or junctions that an individual ant has to traverse may be more important for travel time than reducing physical distances between nests [24,66,67]. Therefore, we measured all possible paths between every nest and all other nests in a tree, recording all intervening junctions between nest pairs (e.g. branching forks in a tree stem, vines crossing a tree stem, two stems crossing each other or leaves from one stem touching another branch) to generate two distance measurements: the shortest physical distance between nest pairs and the smallest number of junctions between nest pairs. Physical distance was measured as the minimum distance an ant needs to walk between two nests (in cm) and 'junction distance' was measured as the fewest number of junctions encountered along any path between two nests. These measurements each produce two types of nest networks: (i) a community-wide nest network for a tree that includes all ant nests and (ii) an intraspecific nest network for each resident ant species that connects only nests within a colony together.

(iii) Individual resource scale: individual nests

For every stem containing an ant nest, we calculated stem volume using Newton's Formula as described above and then destructively harvested the stem at the end of the study. To harvest, we visited each nest at a time of day when each target species was not active, sealed all nest entrances and then removed the entire stem from the tree. This ensured all colony members were in their respective nests at the time of collection, and none escaped subsequently. Collected nests were shipped overnight back to the lab where they were immediately frozen at -20°C . All nests were dissected in the lab and the contents were quantified. Specifically, for each nest we confirmed the identity of the resident species and counted all eggs, larvae, worker pupae, soldier pupae, queen pupae, workers, soldiers, alate queens, dealate queens and males.

(d) Statistical analyses

(i) Whole tree analyses

We tested how tree characteristics shaped the arboreal ant community at the scale of an individual tree. We used linear regression to explore whether ant species richness or total ant nest count across all species in a tree were best predicted by each of tree basal area, crown dieback-weighted basal area, total dead stem count or total dead wood volume. The four predictor variables were highly correlated (correlation coefficient > 0.62) so we separately modelled each predictor for both species richness and total nest count (eight total models) and evaluated model fit by comparing Akaike Information Criterion (AIC) values (electronic supplementary material, table S1).

Individual ant species could also respond differently to tree characteristics and to the presence of other ant species on a tree. To explore this, we created four separate linear models each of eight different response variables: a binary presence/absence variable for each of the four focal ant species (four response variables) or the total nest count in a tree for each species (four response variables). All models included four predictors; three variables reporting total nest counts for the non-focal species of the analysis along with one of the four tree characteristic metrics listed above (32 models in total). For the presence/absence tests, we used generalized linear models with binomial errors and a log link function [68]. We reduced the models using backwards stepwise AIC selection [69] and AIC comparisons were used to determine best-fit model. Final models for all 32 tests are provided in electronic supplementary material, table S2.

(ii) Nest network analyses

We explored how the network of all ant nests within a tree was shaped by tree-level characteristics and the composition of the local ant community. We constructed six linear models with one of two response variables: the average physical distance or the average junction distance between any two nests in a tree. All models included five predictors; four predictors representing number of nests occupied by each of the four focal species, and a fifth predictor related to one of three metrics of dead wood availability within each tree (weighted basal area, dead twig count or total dead wood volume). To explore how the nest networks of each focal species responded to tree characteristics and the presence of other ant species, we evaluated another 24 models with the same set of predictors, with the response variables as either the average physical distance or the junction distance among nests of the same species in each tree (intraspecific nest distances). We reduced the models using backwards stepwise AIC selection and AIC comparisons were used to determine best-fit model. The final models for each of the tests are provided in electronic supplementary material, table S3.

Individual ant species could also display species-specific spatial nesting patterns. To examine these patterns, we explored differences in pairwise distance between all focal nests within each tree. We expanded the initial dataset by adding 19 more trees ($n = 50$ trees) with complete information on the ant community and nest networks, but lacking complete dead wood data. We documented 365 intraspecific pairwise nest distances split among the four focal ant species (e.g. distance between two *C. varians* nests, two *P. ejectus* nests, two *P. gracilis* nests or two *P. simplex* nests). We used two linear mixed models with physical or junction distance between two nests as the response variable, nest pair category as the predictor variable (four levels; one for each species) and Tree ID as a random grouping factor. We used a Tukey's *post hoc* test to explore differences in means per category.

(iii) Individual nest analyses

We first explored whether the mean volume of occupied stems differed among the four focal species and from the mean volume of unoccupied dead stems on a tree. We chose volume because nest quality is generally determined by cavity volume as it dictates how much space there is for colony growth (e.g. [39]). We fit stem volume as a function of species nest occupancy using a mixed-effects ANOVA, where species nest occupancy was a categorical variable with five levels (unoccupied, *C. varians*, *P. ejectus*, *P. gracilis* and *P. simplex*). Tree ID was included as a random grouping factor, and we used Tukey's *post hoc* tests to explore any differences among ant species nest selection.

We next explored whether the contents of a nest were predicted by stem volume, nesting ant species identity, and distance to the nearest nest of each of the four focal ant species

(eight predictors; two for each species to account for two different distance measurements) using zero-inflated generalized linear mixed models with negative binomial errors and a log link function. [41]. In each model, we included either the total count of the combination of all adult ants and brood, only adult ants or only brood as the response variable. We selected these metrics as nest defensibility is determined by defensive strategies of individual ant species [38,40,41,51] and competitor pressure depends on the neighbourhood of enemies trying to usurp the nest for themselves [41]. Ants will also differentially move their brood and redeploy adult ants based on perceived threat or nest defensibility. We started with 24 models (three response variables with all models including stem volume, nest ant species identity and one of eight distance measures). Model reduction and AIC comparison resulted in all models reducing to only ant species identity and the interaction between species identity and distance to the nearest *P. gracilis* nest as the best-fit models. Tree ID was treated as a random grouping variable for all models. We used a Tukey's *post hoc* test to explore any pairwise differences in nest contents between ant species.

All statistical tests were performed in the R environment version 4.2.2 [70] including packages *lme4* [71], *lmerTest* [72] and *glmmTMB* [73]. In all models, metrics of tree size, species richness, network distances, stem volumes and individual ant counts were log transformed to meet model assumptions where necessary and to match the expectation of a log-log linear relationship between species richness and area measurements [74]. Finally, we confirmed normality for all parametric models using Shapiro-Wilk tests on model residuals and performed residual diagnostics to confirm models conformed to model assumption using *DHARMa* [75].

3. Results

(a) Resource patch scale: whole tree

At the scale of discrete resource patches, whole trees with more nesting habitat had more ant nests and ant species, but species richness was not related to tree size. Specifically, trees with larger basal areas and more dead stems had more ant nests, but only dead stem count predicted variation in ant species richness. Basal area alone did not predict the number of ant species (figure 1; electronic supplementary material, table S1). Of the three metrics of dead wood availability, the number of dead stems in a tree was the best predictor of both species richness and total nests (electronic supplementary material, table S1).

The four focal ant species responded differently to nesting resource availability and potential species interactions. *Cephalotes varians* was influenced only by nesting resource availability, with its likelihood of being present on a tree higher in trees with more dead wood, with weighted basal area specifically functioning as the best predictor (electronic supplementary material, figure S2 and table S2). By contrast, the two native small-bodied *Pseudomyrmex*, *P. ejectus* and *P. simplex*, were influenced only by the presence of other ant species. These two species generally did not co-occur, but when both were present in a tree, the number of nests of the two species were negatively associated (electronic supplementary material, figure S3). Additionally, *P. ejectus* frequently co-occurred with *C. varians*, whereas *P. simplex* had a lower frequency of occurrence in trees that also hosted the non-native *P. gracilis* (electronic supplementary material, figure S4a, S4b and table S2). *P. gracilis* was influenced only by resource availability, establishing more nests




Figure 1. The relationships between arboreal ant species richness and total nests across tree basal area (*a,c*) or the total count of dead stems in a tree (*b,d*). Regression lines with 95% CI (shaded region) are included when there is a significant relationship (electronic supplementary material, table S1). Axes are on log-scales.

in trees with a higher volume of dead wood (electronic supplementary material, figure S5 and table S2).

(b) Resource network scale: nest network

For the resource network scale, both habitat availability and the presence of specific ant species shaped the community-wide nest network formed by all resident colonies on a tree. Specifically, both the physical distance and junction distance between any two nests in a tree increased with increasing dead wood availability, with dead stem volume acting as the best predictor for physical distance (figure 2*a*) and dead stem count as the best predictor for junction distance (figure 2*b*; electronic supplementary material, table S3). In addition, the average physical distances between ant nests in a tree were higher in trees with more *Cephalotes* nests (figure 2*c*).

The intraspecific nest networks of individual ant species responded differently to resource versus competitor pressures. The nest networks of *P. ejectus* and *P. simplex* were more spread out in trees with greater dead wood volume (both physical and junction distances for *P. ejectus* and physical distances for *P. simplex*; figure 3*a,b*; electronic supplementary material, table S3). *Cephalotes* and the two native *Pseudomyrmex* species also responded to competitor abundance. Specifically, the average intraspecific nest distances for both *C. varians* and *P. simplex* was smaller in trees with greater numbers of *P. gracilis* nests (figure 3*c,d*; electronic supplementary material, table S3). *Pseudomyrmex ejectus* also had a more clustered nest network in trees with

more *C. varians* nests (figure 3*e*; electronic supplementary material, table S3), whereas *P. simplex* had a less clustered nest network in trees with *C. varians*. Neither resource variables nor other ant species influenced the intraspecific distances among *P. gracilis* nests.

Measuring all intraspecific pairwise nest distances also revealed species-specific nesting patterns. *Cephalotes varians* and *P. ejectus* nests were, on average, more closely clustered together in space (both physical and junction distances) than the nests of *P. gracilis* or *P. simplex* (electronic supplementary material, figure S6; global tests— $F_{3,355} > 7.62$, $p < 0.0006$; Tukey Honestly Significant Difference (HSD)— $z > 2.67$, $p < 0.04$).

(c) Individual resource scale: individual nests

At the scale of an individual resource, defined by an individual ant nest within a nest network, *C. varians*, *P. ejectus* and *P. gracilis* all nested in stems that were of similar size and were larger than the average unoccupied dead stem on a tree (electronic supplementary material, figure S7). By contrast, *P. simplex* nested in smaller stems that were similar to average size of unoccupied dead stems on a tree (electronic supplementary material, figure S7; Tukey HSD— $z > 2.94$, $p < 0.02$). Exploring nest contents revealed that *C. varians* had more adult ants per nest than *P. ejectus* or *P. gracilis* (electronic supplementary material, figure S8; Tukey HSD— $z > 3.01$, $p < 0.02$). In addition, *C. varians* and the two native *Pseudomyrmex* ants showed consistent patterns for how they distributed brood and adult ants relative to their proximity




Figure 2. Statistically significant relationships between the community-wide distances between nests and the amount of dead wood or the number of ant nests in a tree. The shaded region around the regression lines indicates the 95% CI. Axes are on log-scales.

to a *P. gracilis* nest (electronic supplementary material, table S4). Specifically, *C. varians* had fewer total ants and brood in nests that were only a few junctions from the nearest *P. gracilis* nest (figure 4a), and *P. simplex* had fewer total ants and brood in nests that were physically closer to *P. gracilis* nests (figure 4b). By contrast, *P. ejectus* had more ants and brood in nests closer to nests of *P. gracilis* measured by both physical and junction distance (figure 4c,d).

4. Discussion

Our results broadly support our central hypothesis that resource availability has an overarching influence on local community structure, and that species interactions emerge as more important at finer-grained local scales. Specifically, we observed that availability of dead wood was the primary driver of ant diversity at the scale of a tree, that the nest network was shaped by interactions between resources and competition, and that the distribution of ants within a nest was entirely driven by competition. Resource limitations and competitive species interactions are frequently proposed as contrasting drivers of diversity in arboreal ant communities [40,61,76,77]. By incorporating multiple spatial scales into this current study, we are able to demonstrate that both processes are acting in tandem and that the strength of their effects is scale dependent. These findings provide rare empirical support for the theoretical and computational framework of habitat characteristics imposing limitations on local diversity prior to [78,79] or in concert with [80,81] species interactions. The results also further highlight the

need to match observations to the scale at which interactions occur, to avoid masking competition and other biotic interactions [13].

Species–area relationships are common among taxa and across spatial scales [82], but it is typically unclear what specific resources are underpinning these relationships [83]. At the scale of a whole tree, representing a discrete resource patch, we did not detect a direct relationship between species richness and tree size (basal area) despite this relationship being a common feature of arboreal ant communities [30–32,45]. We instead detected a species–resource relationship between ant species richness and dead wood availability, suggesting that nest site availability is the specific habitat limitation underlying area-based relationships within this arboreal ant community. We expect that this trend is widespread among arboreal ant communities, and we predict that where species–area relationships exist between tree size and ant species richness, incorporating measurements of nesting resources would better predict diversity patterns. Ultimately, habitable patch area is a proxy for a broad series of scale-dependent resources and ecological processes ranging from likelihood of encounter during dispersal, available food and nest sites and proximity to competitors [11,84]. These patterns suggest that habitat limitations on a community can be masked when fine-scale resource availability is not considered [13].

Resource availability provides a foundation for determining local diversity and community structure [14,18] but species interactions and behaviours can mediate the final outcome and dynamics [1,3,21]. We demonstrate that the arboreal ants in this forest follow these general trends in




Figure 3. Statistically significant relationships between intraspecific nest distances (physical or junction distance) for different focal ant species and dead wood availability or the number of nests of other ant species in a tree. The shaded region around the regression lines indicates the 95% CI. Distance and volume measures are on a log-scale.

terms of their resource networks, represented by networked nests that each colony occupies. For example, *C. varians* is a nest defence specialist that typically clusters its nests in a laboratory setting [25] and has a soldier caste that uses an armoured head dish to barricade the colony's nest entrances [59]. In a natural setting, we demonstrate that, compared with commonly co-occurring species, *C. varians* not only has the most clustered intraspecific nest network, as would be indicative of a species prioritizing defensibility, but also

further shrinks its network in the presence of the aggressive, non-native competitor *P. gracilis*. By contrast, *P. gracilis* disperses its nests broadly across a tree crown, as expected of a non-native under less competitive pressure [3,85,86]. In addition, *P. simplex* has a broadly dispersed nest network but has a significantly contracted nest network in the presence of *P. gracilis*. Collectively, these observations lend additional support to the idea that *P. gracilis* is using its widely dispersed nesting strategy to limit nest acquisition




Figure 4. Statistically significant relationships between the contents of ant nests versus the distance to the nearest nest of *P. gracilis*. The shaded region around the regression lines indicates the 95% CI. Axes are on log-scale.

by other members of the community and that less aggressive native species shrink their nest networks in response to this competitor.

Biotic interactions at finer spatial scales can also have meaningful impact on species growth within the community, even when it is not reflective in measurements of species richness or composition. However, the subtle impact of species interactions on growth patterns are frequently impossible to detect without extensive multi-year studies tracking individuals through time [87]. Here, we were able to collect data across local spatial scales, including at the fine-grained local scale of individual resources via distances between nests and the distribution of ants among nests. These data allowed us to demonstrate in a snapshot that the aggressive, non-native *P. gracilis* exerts competitive pressure on *C. varians* and *P. simplex* that limits the spatial extent and pattern of colony growth. More specifically, both ants have more clustered networks in trees with *P. gracilis* and tend to have fewer ants and brood in nests near *P. gracilis*. By contrast, *P. ejectus* tends to have more ants and brood in nests nearest to *P. gracilis*. Considering *P. ejectus* and *P. simplex* exhibit almost complete competitive exclusion, the distribution of *P. ejectus* ants in nests near to *P. gracilis* could arise from a form of competitive release [88,89] wherein *P. gracilis* limits *P. simplex* allowing for *P. ejectus* to better perform nearer to *P. gracilis*. Experimental manipulations of the ant community would be necessary to confirm

these observations, but being able to detect these patterns further highlights the value of multi-scale collection regimens for local community ecology data.

The outcome of multispecies interactions on the diversity and stability of ecological communities is notoriously difficult to understand and predict [6,22,90]. The majority of work on the subject is carried out in laboratory or mesocosm experiments [19,91], in plant systems where individuals can be more easily tracked [92,93] or via simulations [94]. The arboreal ant community of the Florida Keys hammock forest exhibits considerable utility in parsing the outcome of multispecies interactions in a complex but manageable animal community of conservation concern. The results of this study suggest that while nest site availability is the main determinant of ant species richness and abundance at the scale of a tree, competitive interactions between species shape the spatial distribution of nests within trees and ants within nests. Experiments modifying nest site availability via artificial nest additions [32,33,40,61] and modifying community structure via relocating ants species among trees [35] could provide further evidence for the outcomes recorded here. The arboreal ant community of the Florida Keys presents an opportunity to explore ecological processes across multiple scales of ecological organization in a system that is both accessible and amenable to experimental manipulations. Ultimately, the key to determining the drivers

of diversity is matching observations to the scale where interactions occur.

Ethics. The fieldwork was conducted under permit numbers 03271915, 02122025, and 02152135 from the Florida Department of Environmental Protection, Division of Recreation and Parks.

Data accessibility. All data included in this project are available via Dryad Digital Repository: <https://doi.org/10.5061/dryad.h70rxwdpr> [95].

Supplementary material is available online [96].

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. B.J.A.: conceptualization, data curation, formal analysis, investigation, methodology, project administration, supervision, validation, visualization, writing—original draft, writing—review and editing; E.M.G.: data curation, formal analysis, investigation, methodology, visualization, writing—original draft, writing—review and editing; M.C.D.-M. and E.J.H.R.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, resources, validation, writing—original draft, writing—review and editing; S.P.: conceptualization, data curation, formal analysis, funding acquisition, investigation,

methodology, project administration, resources, software, supervision, validation, visualization, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. All authors declare no conflict or competing interests.

Funding. This work was supported by National Science Foundation grants IOS-1755406 awarded to S.P., and IOS-1755425 awarded to M.C.D.-M., E.J.H.R., with additional support for undergraduate students Lucille Martin and Parisa Tabiatnejad provided by Harlan Research Fellowships from the George Washington University Department of Biological Sciences.

Acknowledgements. Jackie Fiore, Parisa Tabiatnejad and Melody Weber assisted with data collection in the field. Maggie Connolly, Rohit Kotha, Lucille Martin, Casey Robarts and Christopher Webb helped with nest dissections and counts of nest contents. Chloé Debyser and Valentin Lecheval assisted with code for computing nest spatial distances. We thank Trudy Ferraro and Paul Rice along with the other staff of the Florida Department of Environmental Protection and the Florida State Parks for access to sites and logistic support in Key Largo.

References

- Briones O, Montana C, Ezcurra E. 1998 Competition intensity as a function of resource availability in a semiarid ecosystem. *Oecologia* **116**, 365–372. (doi:10.1007/s004420050599)
- Rappart DJ, Turner JE. 1977 Economic models in ecology: the economics of resource allocation provide a framework for viewing ecological processes. *J. Kans. Entomol. Soc.* **195**, 367–373.
- Neumann K, Pinter-Wollman N. 2022 The effect of resource availability on interspecific competition between a native and an invasive ant. *Phil. Trans. R. Soc. B* **377**, 20210146. (doi:10.1098/rstb.2021.0146)
- Colwell RK, Rangel TF. 2009 Hutchinson's duality: the once and future niche. *Proc. Natl Acad. Sci. USA* **106**, 19 651–19 658. (doi:10.1073/pnas.0901650106)
- Hutchinson GE. 1957 Concluding remarks. population studies: animal ecology and demography. In *Cold Spring Harbor symposia on quantitative biology*, pp. 415–427. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Tilman D. 1982 *Resource competition and community structure*. Princeton, NJ: Princeton University Press.
- Chesson P. 2000 Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* **31**, 343–366. (doi:10.1146/annurev.ecolsys.31.1.343)
- Hardin G. 1960 The competitive exclusion principle. *J. Kans. Entomol. Soc.* **131**, 1292–1297.
- Meynard CN, Lavergne S, Boulangeat I, Garraud L, Van Es J, Mouquet N, Thuiller W. 2013 Disentangling the drivers of metacommunity structure across spatial scales. *J. Biogeogr.* **40**, 1560–1571. (doi:10.1111/jbi.12116)
- Ricklefs RE, Schluter D. 1993 Species diversity: regional and historical influences. In *Species diversity in ecological communities* (eds RE Ricklefs, D Schluter), pp. 350–363. Chicago, IL: University of Chicago Press.
- Levin SA. 1992 The problem of pattern and scale in ecology: the Robert H. MacArthur award lecture. *Ecology* **73**, 1943–1967. (doi:10.2307/1941447)
- Kneitel JM, Chase JM. 2004 Trade-offs in community ecology: linking spatial scales and species coexistence. *Ecol. Lett.* **7**, 69–80. (doi:10.1046/j.1461-0248.2003.00551.x)
- Huston MA. 1999 Local processes and regional patterns: appropriate scales for understanding variation in the diversity of plants and animals. *Oikos* **86**, 393–401. (doi:10.2307/3546645)
- Baiser B, Buckley HL, Gotelli NJ, Ellison AM. 2013 Predicting food-web structure with metacommunity models. *Oikos* **122**, 492–506. (doi:10.1111/j.1600-0706.2012.00005.x)
- Miller TE, Kneitel JM. 2005 Inquiline communities in pitcher plants as a prototypical metacommunity. In *Metacommunities: spatial dynamics and ecological communities* (eds Marcel Holyoak, Mathew Leibold, Robert Holt), pp. 122–145. Chicago, IL: University of Chicago Press.
- Vanschoenwinkel B, De Vries C, Seaman M, Brendonck L. 2007 The role of metacommunity processes in shaping invertebrate rock pool communities along a dispersal gradient. *Oikos* **116**, 1255–1266. (doi:10.1111/j.0030-1299.2007.15860.x)
- Kaspari M, Weiser MD. 1999 The size-grain hypothesis and interspecific scaling in ants. *Func. Ecol.* **13**, 530–538. (doi:10.1046/j.1365-2435.1999.00343.x)
- Leibold MA *et al.* 2004 The metacommunity concept: a framework for multi-scale community ecology. *Ecol. Lett.* **7**, 601–613. (doi:10.1111/j.1461-0248.2004.00608.x)
- Cadotte MW, Mai DV, Jantz S, Collins MD, Keele M, Drake JA. 2006 On testing the competition-colonization trade-off in a multispecies assemblage. *Am. Nat.* **168**, 704–709. (doi:10.1086/508296)
- Levine JM, Bascompte J, Adler PB, Allesina S. 2017 Beyond pairwise mechanisms of species coexistence in complex communities. *Nature* **546**, 56–64. (doi:10.1038/nature22898)
- Chesson P. 1994 Multispecies competition in variable environments. *Theor. Popul. Biol.* **45**, 227–276. (doi:10.1006/tpbi.1994.1013)
- Huisman J, Weissing FJ. 2001 Fundamental unpredictability in multispecies competition. *Am. Nat.* **157**, 488–494. (doi:10.1086/319929)
- Bebber DP, Hynes J, Darrah PR, Boddy L, Fricker MD. 2007 Biological solutions to transport network design. *Proc. R. Soc. B* **274**, 2307–2315. (doi:10.1098/rspb.2007.0459)
- Cook Z, Franks DW, Robinson E.J.H. 2014 Efficiency and robustness of ant colony transportation networks. *Behav. Ecol. Sociobiol.* **68**, 509–517. (doi:10.1007/s00265-013-1665-8)
- Chang J, Powell S, Robinson E.J.H, Donaldson-Matasci MC. 2021 Nest choice in arboreal ants is an emergent consequence of network creation under spatial constraints. *Swarm Intell.* **15**, 7–30. (doi:10.1007/s11721-021-00187-5)
- Lecheval V, Larson H, Burns DD, Ellis S, Powell S, Donaldson-Matasci MC, Robinson E.J.H. 2021 From foraging trails to transport networks: how the quality-distance trade-off shapes network structure. *Proc. R. Soc. B* **288**, 20210430. (doi:10.1098/rspb.2021.0430)
- Plowman NS, Mottl O, Novotny V, Idigil C, Philip FJ, Rimandai M, Klimes P. 2020 Nest microhabitats and tree size mediate shifts in ant community structure across elevation in tropical rainforest canopies. *Ecography* **43**, 431–442. (doi:10.1111/ecog.04730)
- Camargo RX, Oliveira PS. 2012 Natural history of the Neotropical arboreal ant, *Odontomachus hastatus*: nest sites, foraging schedule, and diet. *J. Insect Sci.* **12**, 48. (doi:10.1673/031.012.4801)

29. Ng FSP. 1977 Shyness in trees. *Nat. Malaysiana* **2**, 35–37.
30. Adams BJ, Schnitzer SA, Yanoviak SP. 2017 Trees as islands: canopy ant species richness increases with the size of liana-free trees in a Neotropical forest. *Ecography* **40**, 1067–1075. (doi:10.1111/ecog.02608)
31. Cuissi RG, Lasmar CJ, Moretti TS, Schmidt FA, Fernandes WD, Falleiros AB, Schoereder JH, Ribas CR. 2015 Ant community in natural fragments of the Brazilian wetland: species–area relation and isolation. *J. Insect Conserv.* **19**, 531–537. (doi:10.1007/s10841-015-9774-5)
32. Powell S, Costa AN, Lopes CT, Vasconcelos HL. 2011 Canopy connectivity and the availability of diverse nesting resources affect species coexistence in arboreal ants. *J. Anim. Ecol.* **80**, 352–360. (doi:10.1111/j.1365-2656.2010.01779.x)
33. Philpott SM, Foster PF. 2005 Nest-site limitation in coffee agroecosystems: artificial nests maintain diversity of arboreal ants. *Ecol. Appl.* **15**, 1478–1485. (doi:10.1890/04-1496)
34. Cerdá X, Retana J. 1998 Interference interactions and nest usurpation between two subordinate ant species. *Oecologia* **113**, 577–583. (doi:10.1007/s004420050411)
35. Mottl O, Yombai J, Fayle TM, Novotný V, Klimeš P. 2020 Experiments with artificial nests provide evidence for ant community stratification and nest site limitation in a tropical forest. *Biotropica* **52**, 277–287. (doi:10.1111/btp.12684)
36. Camarota F, Powell S, Melo AS, Priest G, Marquis RJ, Vasconcelos HL. 2016 Co-occurrence patterns in a diverse arboreal ant community are explained more by competition than habitat requirements. *Ecol. Evol.* **6**, 8907–8918. (doi:10.1002/ece3.2606)
37. Debout G, Schatz B, Elias M, Mckey D. 2007 Polydomy in ants: what we know, what we think we know, and what remains to be done. *Biol. J. Linn. Soc.* **90**, 319–348. (doi:10.1111/j.1095-8312.2007.00728.x)
38. Powell S. 2009 How ecology shapes caste evolution: linking resource use, morphology, performance and fitness in a superorganism. *J. Evol. Biol.* **22**, 1004–1013. (doi:10.1111/j.1420-9101.2009.01710.x)
39. Powell S, Dornhaus A. 2013 Soldier-based defences dynamically track resource availability and quality in ants. *Anim. Behav.* **85**, 157–164. (doi:10.1016/j.anbehav.2012.10.020)
40. Camarota F, Vasconcelos HL, Marquis RJ, Powell S. 2020 Revisiting ecological dominance in arboreal ants: how dominant usage of nesting resources shapes community assembly. *Oecologia* **194**, 151–163. (doi:10.1007/s00442-020-04748-z)
41. Powell S, Donaldson-Matasci M, Woodrow-Tomizuka A, Dornhaus A. 2017 Context-dependent defences in turtle ants: resource defensibility and threat level induce dynamic shifts in soldier deployment. *Func. Ecol.* **31**, 2287–2298. (doi:10.1111/1365-2435.12926)
42. Ross MS, O'Brien JJ, Flynn LJ. 1992 Ecological site classification of Florida Keys terrestrial habitats. *Biotropica* **24**, 488–502. (doi:10.2307/2389011)
43. Karim A, Main MB. 2009 Habitat fragmentation and conservation strategies for a rare forest habitat in the Florida Keys archipelago. *Urban Ecosystems* **12**, 359–370. (doi:10.1007/s11252-009-0089-8)
44. Wilder GJ, Sprunt SV, Duquesnel JA, Kolterman SF. 2014 A floristic inventory of Dagny Johnson Key Largo Hammock Botanical State Park and immediately adjacent lands (Monroe County), Florida, USA. *J. Bot. Res. Inst. Texas* **8**, 227–251.
45. Klimes P, Idigel C, Rimandai M, Fayle TM, Janda M, Weiblen GD, Novotny V. 2012 Why are there more arboreal ant species in primary than in secondary tropical forests? *J. Anim. Ecol.* **81**, 1103–1112. (doi:10.1111/j.1365-2656.2012.02002.x)
46. Basset Y, Novotny V, Miller SE, Kitching RL. 2003 Methodological advances and limitations in canopy entomology. In *Arthropods of tropical forests: spatio-temporal dynamics and resource use in the canopy* (ed. Yves Basset), pp. 7–16. Cambridge, UK: Cambridge University Press.
47. Antoniazzi R, Guevara R, García-Franco J, Janda M, Leponce M, Dáttilo W. 2020 Environmental drivers of ant dominance in a tropical rainforest canopy at different spatial scales. *Ecol. Entomol.* **46**, 440–450. (doi:10.1111/een.12988)
48. Leponce M, Dejean A, Mottl O, Klimes P. 2021 Rapid assessment of the three-dimensional distribution of dominant arboreal ants in tropical forests. *Insect Conserv. Diversity* **14**, 426–438. (doi:10.1111/icad.12486)
49. Longino JT, Colwell RK. 2020 The arboreal ants of a Neotropical rain forest show high species density and comprise one third of the ant fauna. *Biotropica* **52**, 675–685. (doi:10.1111/btp.12772)
50. Moreau CS, Deyrup MA, Davis LR. 2014 Ants of the Florida Keys: species accounts, biogeography, and conservation (Hymenoptera: Formicidae). *J. Insect Sci.* **14**, 295. (doi:10.1093/jisesa/ieu157)
51. Priest GV, Camarota F, Powell S, Vasconcelos HL, Marquis RJ. 2021 Ecosystem engineering in the arboreal realm: heterogeneity of wood-boring beetle cavities and their use by cavity-nesting ants. *Oecologia* **196**, 427–439. (doi:10.1007/s00442-021-04934-7)
52. Ward PS. 1985 The Nearctic species of the genus *Pseudomyrmex* (Hymenoptera: Formicidae). Las especies neárticas del género *Pseudomyrmex* (Hymenoptera: Formicidae). *Quaest. Entomol.* **21**, 209–246.
53. Larsen A, Philpott SM. 2010 Twig-nesting ants: the hidden predators of the coffee berry borer in Chiapas, Mexico. *Biotropica* **42**, 342–347. (doi:10.1111/j.1744-7429.2009.00603.x)
54. Gonthier DJ, Ennis KK, Philpott SM, Vandermeer J, Perfecto I. 2013 Ants defend coffee from berry borer colonization. *BioControl* **58**, 815–820. (doi:10.1007/s10526-013-9541-z)
55. Deyrup M, Davis L, Cover S. 2000 Exotic ants in Florida. *Trans. Am. Entomol. Soc.* **126**, 293–326.
56. Wetterer JK, Hill JG, MacGown JA, Booher DB. 2022 Populations of the graceful twig ant, *Pseudomyrmex gracilis*, now bridge the distribution gap between their native range in Texas and non-native range in Florida, USA. *Trans. Am. Entomol. Soc.* **148**, 51–58.
57. Wilson EO. 1976 A social ethogram of the neotropical arboreal ant *Zacryptocerus varians* (Fr. Smith). *Anim. Behav.* **24**, 354–363. (doi:10.1016/S0003-3472(76)80043-7)
58. De Andrade ML, Urbani CB. 1999 Diversity and adaptation in the ant genus *Cephalotes*, past and present. *Staatliches Museum für Naturkunde* **271**, 899.
59. Powell S, Price SL, Kronauer DJ. 2020 Trait evolution is reversible, repeatable, and decoupled in the soldier caste of turtle ants. *Proc. Natl Acad. Sci. USA* **117**, 6608–6615. (doi:10.1073/pnas.1913750117)
60. Yanoviak SP, Kaspari M. 2000 Community structure and the habitat templet: ants in the tropical forest canopy and litter. *Oikos* **89**, 259–266. (doi:10.1034/j.1600-0706.2000.890206.x)
61. Adams BJ, Schnitzer SA, Yanoviak SP. 2019 Connectivity explains local ant community structure in a Neotropical forest canopy: a large-scale experimental approach. *Ecology* **100**, e02673.
62. Ward PS. 1989 Systematic studies on pseudomyrmecine ants: revision of the *Pseudomyrmex oculatus* and *P. subtilissimus* species groups, with taxonomic comments on other species. *Quaest. Entomol.* **25**, 393–468.
63. O'Brien ST, Hubbell SP, Spiro P, Condit R, Foster RB. 1995 Diameter, height, crown, and age relationship in eight neotropical tree species. *Ecology* **76**, 1926–1939. (doi:10.2307/1940724)
64. Stolte K, Barbara C, Campbell S, Gillespie A. 2002 *Forest health indicators: forestry inventory and analysis program*. Research Triangle Park, NC: USDA Forest Service, Southern Research Station.
65. Harmon ME, Sexton J. 1996 Guidelines for Measurements of Woody Detritus in Forest Ecosystems. U.S. LTER Network Office: University of Washington, Seattle, WA, USA.
66. Gordon DM. 2012 The dynamics of foraging trails in the tropical arboreal ant *Cephalotes goniodontus*. *PLoS ONE* **7**, e50472. (doi:10.1371/journal.pone.0050472)
67. Gordon DM. 2017 Local regulation of trail networks of the arboreal turtle ant, *Cephalotes goniodontus*. *Am. Nat.* **190**, E156–E169. (doi:10.1086/693418)
68. Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White J-SS. 2009 Generalized linear mixed models: a practical guide for ecology and evolution. *Trends Ecol. Evol.* **24**, 127–135. (doi:10.1016/j.tree.2008.10.008)
69. Johnson JB, Omland KS. 2004 Model selection in ecology and evolution. *Trends Ecol. Evol.* **19**, 101–108. (doi:10.1016/j.tree.2003.10.013)
70. RCoreTeam. 2023 *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
71. Bates D, Maechler M, Bolker B, Walker S. 2014 lme4: Linear mixed-effects models using Eigen and S4. R package version **1**, 1–23.
72. Kuznetsova A, Brockhoff PB, Christensen RHB. 2017 lmerTest package: tests in linear mixed effects models. *J. Stat. Softw.* **82**, 1–26. (doi:10.18637/jss.v082.i13)
73. Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ,

- Machler M, Bolker BM. 2017 glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal* **9**, 378–400. (doi:10.32614/RJ-2017-066)
74. Preston FW. 1962 The canonical distribution of commonness and rarity: Part I. *Ecology* **43**, 185–215. (doi:10.2307/1931976)
75. Hartig F. 2022 DHARMA: residual diagnostics for hierarchical (multi-level/mixed) regression models. R package version 0.4.5. *Computer software*. See <https://CRAN.R-project.org/package=DHARMA>.
76. Dejean A, Compin A, Delabie JH, Azémar F, Corbara B, Leponce M. 2019 Biotic and abiotic determinants of the formation of ant mosaics in primary Neotropical rainforests. *Ecol. Entomol.* **44**, 560–570. (doi:10.1111/een.12735)
77. Ribas CR, Schoereder JH. 2002 Are all ant mosaics caused by competition? *Oecologia* **131**, 606–611. (doi:10.1007/s00442-002-0912-x)
78. Connell JH, Slatyer RO. 1977 Mechanisms of succession in natural communities and their role in community stability and organization. *Am. Nat.* **111**, 1119–1144. (doi:10.1086/283241)
79. Meiners SJ, Cadotte MW, Fridley JD, Pickett ST, Walker LR. 2015 Is successional research nearing its climax? New approaches for understanding dynamic communities. *Func. Ecol.* **29**, 154–164. (doi:10.1111/1365-2435.12391)
80. D'Amen M, Mod HK, Gotelli NJ, Guisan A. 2018 Disentangling biotic interactions, environmental filters, and dispersal limitation as drivers of species co-occurrence. *Ecography* **41**, 1233–1244. (doi:10.1111/ecog.03148)
81. Ulrich W, Zaplata MK, Winter S, Schaaf W, Fischer A, Soliveres S, Gotelli NJ. 2016 Species interactions and random dispersal rather than habitat filtering drive community assembly during early plant succession. *Oikos* **125**, 698–707. (doi:10.1111/oik.02658)
82. Rosenzweig ML. 1995 *Species diversity in space and time*. Cambridge, UK: University Press.
83. Turner WR, Tjørve E. 2005 Scale-dependence in species-area relationships. *Ecography* **28**, 721–730. (doi:10.1111/j.2005.0906-7590.04273.x)
84. Lomolino MV. 2000 Ecology's most general, yet protean pattern: the species-area relationship. *J. Biogeogr.* **27**, 17–26. (doi:10.1046/j.1365-2699.2000.00377.x)
85. Holway DA. 1999 Competitive mechanisms underlying the displacement of native ants by the invasive Argentine ant. *Ecology* **80**, 238–251. (doi:10.1890/0012-9658(1999)080[0238:CMUTDO]2.0.CO;2)
86. Warren RJ, Reed K, Mathew A, Krupp K, Goodman M, Archibald K, Spiering DJ. 2019 Release from intraspecific competition promotes dominance of a non-native invader. *Biol. Invas.* **21**, 895–909. (doi:10.1007/s10530-018-1868-z)
87. Clutton-Brock T, Sheldon BC. 2010 Individuals and populations: the role of long-term, individual-based studies of animals in ecology and evolutionary biology. *Trends Ecol. Evol.* **25**, 562–573. (doi:10.1016/j.tree.2010.08.002)
88. Chase JM, Abrams PA, Grover JP, Diehl S, Chesson P, Holt RD, Richards SA, Nisbet RM, Case TJ. 2002 The interaction between predation and competition: a review and synthesis. *Ecol. Lett.* **5**, 302–315. (doi:10.1046/j.1461-0248.2002.00315.x)
89. Paine RT. 1966 Food web complexity and species diversity. *Am. Nat.* **100**, 65–75. (doi:10.1086/282400)
90. Weigelt A, Schumacher J, Walther T, Bartelheimer M, Steinlein T, Beyschlag W. 2007 Identifying mechanisms of competition in multi-species communities. *J. Ecol.* **95**, 53–64. (doi:10.1111/j.1365-2745.2006.01198.x)
91. Rittmann BE, Manem JA. 1992 Development and experimental evaluation of a steady-state, multispecies biofilm model. *Biotechnol. Bioeng.* **39**, 914–922. (doi:10.1002/bit.260390906)
92. Austin M, Groves R, Fresco L, Kaye P. 1985 Relative growth of six thistle species along a nutrient gradient with multispecies competition. *J. Ecol.* **73**, 667–684. (doi:10.2307/2260503)
93. Kempel A, Chrobock T, Fischer M, Rohr RP, van Kleunen M. 2013 Determinants of plant establishment success in a multispecies introduction experiment with native and alien species. *Proc. Natl Acad. Sci. USA* **110**, 12 727–12 732. (doi:10.1073/pnas.1300481110)
94. Huisman J, Weissing FJ. 2001 Biological conditions for oscillations and chaos generated by multispecies competition. *Ecology* **82**, 2682–2695. (doi:10.1890/0012-9658(2001)082[2682:BCFOAC]2.0.CO;2)
95. Adams BJ, Gora EM, Donaldson-Matasci MC, Robinson EJH, Powell S. 2023 Data from: Competition and habitat availability interact to structure arboreal ant communities across scales of ecological organization. *Dryad Digital Repository*. (doi:10.5061/dryad.h70rxwdpr)
96. Adams BJ, Gora EM, Donaldson-Matasci MC, Robinson EJH, Powell S. 2023 Competition and habitat availability interact to structure arboreal ant communities across scales of ecological organization. Figshare. (doi:10.6084/m9.figshare.c.6824120)