



UNIVERSITY OF LEEDS

This is a repository copy of *Induced subgraphs and tree decompositions II. Toward walls and their line graphs in graphs of bounded degree.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/204535/>

Version: Accepted Version

Article:

Abrishami, T., Chudnovsky, M., Dibek, C. et al. (4 more authors) (2024) Induced subgraphs and tree decompositions II. Toward walls and their line graphs in graphs of bounded degree. *Journal of Combinatorial Theory: Series B*, 164. pp. 371-403. ISSN 0095-8956

<https://doi.org/10.1016/j.jctb.2023.10.005>

© 2023, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

INDUCED SUBGRAPHS AND TREE DECOMPOSITIONS

II. TOWARD WALLS AND THEIR LINE GRAPHS IN GRAPHS OF BOUNDED DEGREE

TARA ABRISHAMI^{*†}, MARIA CHUDNOVSKY^{*†}, CEMIL DIBEK^{*‡}, SEPEHR HAJEBI[§],
PAWEŁ RZAŻEWSKI[¶], SOPHIE SPIRKL^{§||}, AND KRISTINA VUŠKOVIĆ^{**}

ABSTRACT. This paper is motivated by the following question: what are the unavoidable induced subgraphs of graphs with large treewidth? Aboulker et al. made a conjecture which answers this question in graphs of bounded maximum degree, asserting that for all k and Δ , every graph with maximum degree at most Δ and sufficiently large treewidth contains either a subdivision of the $(k \times k)$ -wall or the line graph of a subdivision of the $(k \times k)$ -wall as an induced subgraph. We prove two theorems supporting this conjecture, as follows.

1. For $t \geq 2$, a t -theta is a graph consisting of two nonadjacent vertices and three internally vertex-disjoint paths between them, each of length at least t . A t -pyramid is a graph consisting of a vertex v , a triangle B disjoint from v and three paths starting at v and vertex-disjoint otherwise, each joining v to a vertex of B , and each of length at least t . We prove that for all k, t and Δ , every graph with maximum degree at most Δ and sufficiently large treewidth contains either a t -theta, or a t -pyramid, or the line graph of a subdivision of the $(k \times k)$ -wall as an induced subgraph. This affirmatively answers a question of Pilipczuk et al. asking whether every graph of bounded maximum degree and sufficiently large treewidth contains either a theta or a triangle as an induced subgraph (where a $theta$ means a t -theta for some $t \geq 2$).
2. A *subcubic subdivided caterpillar* is a tree of maximum degree at most three whose all vertices of degree three lie on a path. We prove that for every Δ and subcubic subdivided caterpillar T , every graph with maximum degree at most Δ and sufficiently large treewidth contains either a subdivision of T or the line graph of a subdivision of T as an induced subgraph.

1. INTRODUCTION

All graphs in this paper are finite and simple. Let $G = (V(G), E(G))$ be a graph. A *tree decomposition* (T, β) of G consists of a tree T and a map $\beta : V(T) \rightarrow 2^{V(G)}$, with the following properties:

- (i) For every $v \in V(G)$, there exists $t \in V(T)$ such that $v \in \beta(t)$.

^{*}PRINCETON UNIVERSITY, PRINCETON, NJ, USA

[§]DEPARTMENT OF COMBINATORICS AND OPTIMIZATION, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA

[¶]WARSAW UNIVERSITY OF TECHNOLOGY, POLAND/UNIVERSITY OF WARSAW, POLAND. THIS WORK IS A PART OF PROJECT CUTACOMBS THAT HAS RECEIVED FUNDING FROM THE EUROPEAN RESEARCH COUNCIL (ERC) UNDER THE EUROPEAN UNION'S HORIZON 2020 RESEARCH AND INNOVATION PROGRAMME (GRANT AGREEMENT No. 714704).

^{**}SCHOOL OF COMPUTING, UNIVERSITY OF LEEDS, UK. PARTIALLY SUPPORTED BY DMS-EPSRC GRANT EP/V002813/1.

[†] SUPPORTED BY NSF GRANT DMS-1763817 AND NSF-EPSRC GRANT DMS-2120644.

[‡] SUPPORTED BY NSF GRANT DMS-1763817.

^{||} WE ACKNOWLEDGE THE SUPPORT OF THE NATURAL SCIENCES AND ENGINEERING RESEARCH COUNCIL OF CANADA (NSERC), [FUNDING REFERENCE NUMBER RGPIN-2020-03912]. CETTE RECHERCHE A ÉTÉ FINANÇÉE PAR LE CONSEIL DE RECHERCHES EN SCIENCES NATURELLES ET EN GÉNIE DU CANADA (CRSNG), [NUMÉRO DE RÉFÉRENCE RGPIN-2020-03912].

Date: September 28, 2023.




FIGURE 1. The 6-by-6 square grid (left) and the 6-by-6 wall $W_{6 \times 6}$ (right).

- (ii) For every $v_1 v_2 \in E(G)$, there exists $t \in V(T)$ such that $v_1, v_2 \in \beta(t)$.
- (iii) For every $v \in V(G)$, the subgraph of T induced on the set $\beta^{-1}(v) = \{t \in V(T) \mid v \in \beta(t)\}$ is connected.

The *width* of the tree decomposition (T, β) is $\max_{v \in V(T)} |\beta(v)| - 1$. The *treewidth* of a graph G , denoted by $\text{tw}(G)$, is the minimum width of a tree decomposition of G . Treewidth, originally introduced by Robertson and Seymour in their study of graph minors, is widely considered to be an important graph parameter, both from a structural [22] and algorithmic [11] point of view. Roughly, the treewidth of a graph measures how “close to a tree” it is: trees have treewidth one, and in general, the larger the treewidth of a graph, the less “tree-like”, and hence the more complicated it is. So it is natural to ask how one would certify whether a graph is of large treewidth, and in particular, what can we say about the unavoidable substructures emerging in graphs of large treewidth. As an example, for each k , the k -by- k square grid is a planar graph of maximum degree three and with treewidth k , and the $(k \times k)$ -wall, denoted by $W_{k \times k}$, is the k -by- k hexagonal grid, which is planar graph with maximum degree three and with treewidth k (the formal definition is provided at the end of Subsection 1.1; see Figure 1). Every subdivision of $W_{k \times k}$ is also a graph of treewidth k . The Grid Theorem of Robertson and Seymour gives a complete characterization of the unavoidable minors, and the unavoidable subgraphs of graphs with large treewidth.

Theorem 1.1 (Robertson and Seymour [20]). *For every $k \geq 1$, every graph of sufficiently large treewidth has minor isomorphic to the k -by- k square grid, or equivalently, a subdivision of $W_{k \times k}$ as a subgraph.*

While tree decompositions and classes of graphs with bounded treewidth are central concepts in the study of graphs with forbidden minors [22], the problem of connecting tree decompositions with forbidden induced subgraphs had largely remained uninvestigated until very recently. In accordance, this work is a step toward understanding the unavoidable induced subgraphs of graphs with large treewidth. Formally, let us say a family \mathcal{F} of graphs is *useful* if there exists c such that every graph G with $\text{tw}(G) > c$ contains a member of \mathcal{F} as an induced subgraph. Then our work is motivated by the goal of characterizing useful families. For instance, Lozin and Razgon [18] have recently proved the following theorem, which gives a complete description of all finite useful families. Given a graph F , the *line graph* $L(F)$ of F is the graph with vertex set $E(F)$, such that two vertices of $L(F)$ are adjacent if the corresponding edges of G share an end.

Theorem 1.2 (Lozin and Razgon [18]). *Let \mathcal{F} be finite family of graphs. Then \mathcal{F} is useful if and only if it contains a complete graph, a complete bipartite graph, a forest in which each component has at most three leaves, and the line graph of such a forest.*

In fact, it is easy to see that the complete graph K_t has treewidth $t - 1$ and the complete bipartite graph $K_{t,t}$ has treewidth t . Also, as mentioned above, every subdivision of $W_{k \times k}$ is also of treewidth k , and crucially, no two non-isomorphic subdivisions of $W_{k \times k}$ are induced subgraphs of each other. The line graph of a subdivision of $W_{k \times k}$ is another example of a graph with large treewidth. Note that $L(W_{k \times k})$ does not contain $W_{k \times k}$ as an induced subgraph. In summary, if a family of graphs is useful, then it contains a complete graph, a complete bipartite graph, and for

some k , an induced subgraph of every subdivision of $W_{k \times k}$, and an induced subgraph of the line graph of every subdivision of $W_{k \times k}$. Therefore, it would be natural to ask whether the converse of the latter statement is also true:

Question 1.3. *Let \mathcal{F} be a family of graphs containing a complete graph, a complete bipartite graph, and for some k , an induced subgraph of every subdivision of $W_{k \times k}$, and an induced subgraph of the line graph of every subdivision of $W_{k \times k}$. Then is \mathcal{F} useful?*

It turns out that the answer to Question 1.3 is negative. To elaborate on this, we need a couple of definitions. By a *hole* in a graph we mean an induced cycle of length at least four, and an *even hole* is a hole on an even number of vertices. For graphs G and F , we say that G is F -free if G does not contain an induced subgraph isomorphic to F . If \mathcal{F} is a family of graphs, a graph G is \mathcal{F} -free if G is F -free for every $F \in \mathcal{F}$. It is not difficult to show that for large enough k , subdivisions of $W_{k \times k}$, line graphs of subdivisions of $W_{k \times k}$, and the complete bipartite graph $K_{k,k}$ all contain even holes. Therefore, the following theorem provides a negative answer to Question 1.3.

Theorem 1.4 (Sintiari and Trotignon [23]). *For every integer $\ell \geq 1$, there exists an (even hole, K_4)-free graph G_ℓ such that $\text{tw}(G_\ell) \geq \ell$.*

Observing that graphs G_ℓ in Theorem 1.4 have vertices of arbitrarily large degree, the following conjecture was made (and proved for the case $\Delta \leq 3$) in [1]:

Conjecture 1.5 (Aboulker, Adler, Kim, Sintiari and Trotignon [1]). *For every $\Delta > 0$ there exists c_Δ such that even-hole-free graphs with maximum degree at most Δ have treewidth at most c_Δ .*

Conjecture 1.5 was proved in [10] by three of the authors of the present paper. More generally, it is conjectured in [1] that there is an affirmative answer to Question 1.3 in the bounded maximum degree case (note that bounded maximum degree automatically implies that a large complete graph and a large complete bipartite graph are excluded).

Conjecture 1.6 (Aboulker, Adler, Kim, Sintiari and Trotignon [1]). *For every $\Delta > 0$ there is a function $f_\Delta : \mathbb{N} \rightarrow \mathbb{N}$ such that every graph with maximum degree at most Δ and treewidth at least $f_\Delta(k)$ contains a subdivision of $W_{k \times k}$ or the line graph of a subdivision of $W_{k \times k}$ as an induced subgraph.*

(We remark that, while the present paper was under review, Conjecture 1.6 was proved using a different method [17]. However, the techniques developed here provide foundation to a significant body of future work [2, 3, 4, 5, 6, 7, 8, 9].)

In [1] it is proved for proper minor-closed classes of graphs (in which case the bound on the maximum degree is not needed anymore).

Theorem 1.7 (Aboulker, Adler, Kim Sintiari and Trotignon [1]). *For every graph H there is a function $f_H : \mathbb{N} \rightarrow \mathbb{N}$ such that every graph of treewidth at least $f_H(k)$ and with no H -minor contains a subdivision of $W_{k \times k}$ or the line graph of a subdivision of $W_{k \times k}$ as an induced subgraph.*

In this paper we prove several theorems supporting Conjecture 1.6. In order to state our main results, we need a few more definitions.

A *path* is a graph P with vertex set $\{p_1, \dots, p_k\}$ and edge set $\{p_1p_2, p_2p_3, \dots, p_{k-1}p_k\}$. We write $P = p_1 - \dots - p_k$, and we say p_1 and p_k are the *ends* of P . The *length* of the path P is the number of edges in P . We say that P is a path *from* p_1 *to* p_k , where p_1 and p_k are the vertices of degree one in P . The *interior* of P is denoted P^* and is defined as $P \setminus \{p_1, p_k\}$.

Let G be a graph and let $X, Y \subseteq V(G)$ be disjoint. Then, X is *complete to* Y if for every $x \in X$ and $y \in Y$, we have $xy \in E(G)$, and X is *anticomplete to* Y if there are no edges from X to Y in G .

The *claw* is the graph with vertex set $\{a, b, c, d\}$ and edge set $\{ab, ac, ad\}$. For nonnegative integers t_1, t_2, t_3 , an S_{t_1, t_2, t_3} , also called a *long claw* or a *subdivided claw*, consists of a vertex v and three paths P_1, P_2, P_3 , where P_i is of length t_i , with one end v , such that $V(P_1) \setminus \{v\}$, $V(P_2) \setminus \{v\}$, and $V(P_3) \setminus \{v\}$ are pairwise disjoint and anticomplete to each other. Note that for every t , every subdivision of $W_{k \times k}$ for large enough k contains $S_{t, t, t}$ as an induced subgraph. Our first result is the following.

Theorem 1.8. *Let Δ, t, k be positive integers. There exists $c_{k, t, \Delta}$ such that for every $S_{t, t, t}$ -free graph G with maximum degree Δ and no induced subgraph isomorphic to the line graph of a subdivision of $W_{k \times k}$, we have $\text{tw}(G) \leq c_{k, t, \Delta}$.*

A *theta* is a graph consisting of three internally vertex-disjoint paths $P_1 = a \cdots b$, $P_2 = a \cdots b$, and $P_3 = a \cdots b$ of length at least 2, such that no edges exist between the paths except the three edges incident with a and the three edges incident with b . A *t-theta* is a theta such that each of P_1, P_2, P_3 has length at least t . A *pyramid* is a graph consisting of three paths $P_1 = a \cdots b_1$, $P_2 = a \cdots b_2$, and $P_3 = a \cdots b_3$ of length at least 1, two of which have length at least 2, pairwise vertex-disjoint except at a , and such that $b_1 b_2 b_3$ is a triangle and no edges exist between the paths except those of the triangle and the three edges incident with a . A *t-pyramid* is a pyramid such that each of P_1, P_2, P_3 has length at least t .

Note that the complete bipartite graph $K_{2,3}$ is in fact a theta. Also, for large enough k , every subdivision of $W_{k \times k}$ contains a theta as an induced subgraph, and the line graph of every subdivision of $W_{k \times k}$ contains a triangle. Therefore, the following theorem gives another reason why the answer to Question 1.3 is negative.

Theorem 1.9 (Sintiari and Trotignon [23]). *For every integer $\ell \geq 1$, there exists a (theta, triangle)-free graph G_ℓ such that $\text{tw}(G_\ell) \geq \ell$.*

In analogy to the situation with Theorem 1.4, the graphs G_ℓ in Theorem 1.9 contain vertices of arbitrary large degree. So it is asked in [19] whether (theta, triangle)-free graphs of bounded maximum degree have bounded treewidth (while it is proved in [19] that (theta, triangle, $S_{t, t, t}$)-free graphs, without a bound on the maximum degree, have bounded treewidth). We give an affirmative answer to this question. Indeed, our second result, the following, establishes a far-reaching generalization of this question. It also generalizes Theorem 1.8, and strongly addresses Conjecture 1.6.

Theorem 1.10. *Let Δ, t, k be positive integers with $t \geq 2$. Then, there exists $c_{t, k, \Delta}$ such that for every (t-theta, t-pyramid)-free graph G with maximum degree Δ and no induced subgraph isomorphic to the line graph of a subdivision of $W_{k \times k}$, we have $\text{tw}(G) \leq c_{t, k, \Delta}$.*

A tree T is a *subdivided caterpillar* if there is a path P in T such that P contains every vertex of T of degree at least three in T . The *spine* of T is the shortest path containing all vertices of degree at least three in T . A *leg* of a subdivided caterpillar T is a path in T from a vertex of degree one in T to a vertex of degree at least three in T . A graph G is *subcubic* if every vertex of G has degree at most three.

Note that for every subcubic subdivided caterpillar T and for large enough k , every subdivision of $W_{k \times k}$ contains a subdivision of T as an induced subgraph, and the line graph of every subdivision of $W_{k \times k}$ contains the line graph of a subdivision of T as an induced subgraph. Our third result is the following.

Theorem 1.11. *Let Δ be a positive integer and let T be a subcubic subdivided caterpillar. There exists $c_{T, \Delta}$ such that for every graph G with maximum degree Δ and no induced subgraph isomorphic to a subdivision of T or the line graph of a subdivision of T , we have $\text{tw}(G) \leq c_{T, \Delta}$.*

Let us now roughly discuss the proofs. Usually, to prove that a certain graph family has bounded treewidth, one attempts to construct a collection of “non-crossing decompositions,”

which roughly means that the decompositions “cooperate” with each other, and the pieces that are obtained when the graph is simultaneously decomposed by all the decompositions in the collection “line up” to form a tree structure. Such collections of decompositions are called “laminar.” In all the cases above, there is a natural family of decompositions to turn to, sharing a certain structural property: all the decompositions arise from removing from the graph the neighborhood of a small connected subgraph. Unfortunately, these natural decompositions are very far from being non-crossing, and therefore they cannot be used in traditional ways to get tree-decompositions. What turns out to be true, however, is that, due to the bound on the maximum degree of the graph, these collections of decompositions can be partitioned into a bounded number of laminar collections (where the bound on the number of collections depends on the maximum degree and on the precise nature of the decomposition). We will explain how to make use of this fact in Section 2.

Structure of the paper. We begin in Section 1.1 with a review of relevant definitions and notation. In Section 1.2, we define an important graph parameter tied to treewidth called separation number. In Section 2 we prove Theorem 2.10, which summarizes our main proof method. In Section 3, we bound the treewidth of claw-free graphs with no line graph of a subdivision of a wall, and in Section 4, we apply the results of Section 3 to prove Theorem 1.8. In Section 5, we prove Theorem 1.10, and in Section 6, we prove Theorem 1.11.

1.1. Definitions and Notation. Let G be a graph. In this paper, we use vertex sets and their induced subgraphs interchangeably. Let H be a graph. We say that $X \subseteq V(G)$ is an H in G if X is isomorphic to H . We say that G contains H if there exists $X \subseteq V(G)$ such that X is an H in G .

The *open neighborhood* of a vertex $v \in V(G)$, denoted $N(v)$, is the set of all vertices adjacent to v . The *degree* of $v \in V(G)$ is the size of its open neighborhood. A graph G has *maximum degree* Δ if the degree of every vertex $v \in V(G)$ is at most Δ . The *closed neighborhood* of a vertex $v \in V(G)$ is denoted $N[v]$ and is defined as $N[v] = N(v) \cup \{v\}$. Let $X \subseteq V(G)$. The *open neighborhood of X* , denoted $N(X)$, is the set of all vertices of $G \setminus X$ with a neighbor in X . The *closed neighborhood of X* is denoted $N[X]$ and is defined as $N[X] = N(X) \cup X$.

A set $X \subseteq V(G)$ is *connected* if for every $x, y \in X$, there is a path P in X from x to y . A set $C \subseteq V(G)$ is a *cutset* of a connected graph G if $G \setminus C$ is not connected. A set D is a *connected component of G* if D is inclusion-wise maximal such that $D \subseteq V(G)$ and D is connected.

Let $u, v \in V(G)$ and let $X \subseteq V(G)$. The *distance between u and v* is the length of a shortest path from u to v in G . The *distance between u and X* is the length of a shortest path from u to a vertex $x \in X$ in G . We denote by $N^d(v)$ the set of vertices at distance exactly d from v in G , and by $N^d[v]$ the set of vertices at distance at most d from v in G . Similarly, we denote by $N^d[X]$ the set of vertices of distance at most d from X in G . The *diameter* of a connected set $X \subseteq V(G)$ is the maximum distance in G between two vertices of X .

A *clique* is a set $K \subseteq V(G)$ such that every pair of vertices in K is adjacent. An *independent set* is a set $I \subseteq V(G)$ such that every pair of vertices in I is non-adjacent. The *clique number of G* , denoted $\omega(G)$, is the size of a largest clique in G . The *independence number of G* , denoted $\alpha(G)$, is the size of a largest independent set in G .

A *weight function on G* is a function $w : V(G) \rightarrow \mathbb{R}$ that assigns a non-negative real number to every vertex of G . A weight function is *normal* if $w(V(G)) = 1$. Unless otherwise specified, we assume all weight functions are normal. We denote by w^{\max} the maximum weight of a vertex; i.e. $w^{\max} = \max_{v \in V(G)} w(v)$.

Finally, let us include the precise definition of a wall. The $(n \times m)$ -*wall*, denoted $W_{n \times m}$, is the graph G with vertex set

$$V(G) = \{(1, 2j - 1) \mid 1 \leq j \leq m\}$$

$$\begin{aligned} & \cup \{(i, j) \mid 1 < i < n, 1 \leq j \leq 2m\} \\ & \cup \{(n, 2j - 1) \mid 1 \leq j \leq m, \text{ if } n \text{ is even}\} \\ & \cup \{(n, 2j) \mid 1 \leq j \leq m, \text{ if } n \text{ is odd}\} \end{aligned}$$

and edge set

$$\begin{aligned} E(G) = & \{(1, 2j - 1), (1, 2j + 1) \mid 1 \leq j \leq m - 1\} \\ & \cup \{(i, j), (i, j + 1) \mid 2 \leq i < n, 1 \leq j < 2m\} \\ & \cup \{(n, 2j), (n, 2j + 2) \mid 1 \leq j < m \text{ if } n \text{ is odd}\} \\ & \cup \{(n, 2j - 1), (n, 2j + 1) \mid 1 \leq j < m \text{ if } n \text{ is even}\} \\ & \cup \{(i, j), (i + 1, j) \mid 1 \leq i < n, 1 \leq j \leq 2m, i, j \text{ odd}\} \\ & \cup \{(i, j), (i + 1, j) \mid 1 \leq i < n, 1 \leq j \leq 2m, i, j \text{ even}\}. \end{aligned}$$

Again, see Figure 1 for an example.

1.2. Balanced separators and treewidth. Treewidth is tied to a parameter called the separation number. Let G be a graph, let $S \subseteq V(G)$, let k be a positive integer, and let $c \in [\frac{1}{2}, 1)$. A set $X \subseteq V(G)$ is a $(k, S, c)^*$ -separator if $|X| \leq k$ and for every component D of $G \setminus X$, it holds that $|D \cap S| \leq c|S|$. The *separation number* $\text{sep}_c^*(G)$ is the minimum k such that G has a $(k, S, c)^*$ -separator for every $S \subseteq V(G)$. The following lemma states that the separation number gives an upper bound for the treewidth of a graph.

Lemma 1.12 (Harvey and Wood [16]). *For every $c \in [\frac{1}{2}, 1)$ and every graph G , we have $\text{tw}(G) + 1 \leq \frac{1}{1-c} \text{sep}_c^*(G)$.*

Now, we redefine $(k, S, c)^*$ -separators using weight functions. Given a normal weight function w on a graph G and a constant $c \in [\frac{1}{2}, 1)$, a set $X \subseteq V(G)$ is a (w, c) -balanced separator of G if $w(D) \leq c$ for every component D of $G \setminus X$.

We call a weight function w on G a *uniform weight function* if there exists $Y \subseteq V(G)$ such that $w(v) = \frac{1}{|Y|}$ if $v \in Y$, and $w(v) = 0$ if $v \notin Y$. Lemma 1.12 implies the following:

Lemma 1.13. *Let $c \in [\frac{1}{2}, 1)$ and let G be a graph. If G has a (w, c) -balanced separator of size at most k for every uniform weight function w , then $\text{tw}(G) \leq \frac{1}{1-c}k$.*

Proof. We prove that $\text{sep}_c^*(G) \leq k$. Let $S \subseteq V(G)$ and let w_S be the weight function on G such that $w_S(v) = \frac{1}{|S|}$ if $v \in S$, and $w_S(v) = 0$ otherwise. Since w_S is a uniform weight function, it follows that G has a (w_S, c) -balanced separator X such that $|X| \leq k$. Let D be a component of $G \setminus X$, so $w(D) \leq c$. Consequently, $|D \cap S| \leq c|S|$, and so X is a $(k, S, c)^*$ -separator. Therefore, $\text{sep}_c^*(G) \leq k$, and the result follows from Lemma 1.12. ■

Lemma 1.13 implies that if for some fixed $c \in [\frac{1}{2}, 1)$, G has a balanced separator of size k for every weight function w , then the treewidth of G is bounded by a function of k . The next lemma states the converse.

Lemma 1.14 (Cygan, Fomin, Kowalik, Lokshantov, Marx, Pilipczuk and Pilipczuk [14]). *If $\text{tw}(G) \leq k$, then G has a (w, c) -balanced separator of size at most $k + 1$ for every normal weight function w and for every $c \in [\frac{1}{2}, 1)$.*

Together, Lemmas 1.13 and 1.14 show that treewidth is tied to the size of balanced separators. In this paper, we rely on balanced separators to prove that graphs have bounded treewidth. In what follows, we will often assume that G has no (w, c) -balanced separator of size d for some normal weight function w , $c \in [\frac{1}{2}, 1)$, and positive integer d , since otherwise, in light of Lemma 1.13, we are done.

2. CENTRAL BAGS AND FORCERS

A *separation* of a graph G is a triple (A, C, B) with $A, C, B \subseteq V(G)$ such that A, C , and B are pairwise disjoint, $A \cup C \cup B = V(G)$, and A is anticomplete to B . If A and B are both non-empty, then C is a cutset of G . If $S = (A, C, B)$ is a separation, we write $A(S) = A$, $C(S) = C$, and $B(S) = B$.

Separations provide a way to organize the structure of connected graphs around important cutsets. To that end, it is often useful to characterize the relationship between two separations of a graph. Here, we define two relations between graph separations. Two separations $S_1 = (A_1, C_1, B_1)$ and $S_2 = (A_2, C_2, B_2)$ of G are *non-crossing* if (possibly exchanging the roles of A_1 and B_1 , and of A_2 and B_2) $A_1 \cap C_2 = \emptyset$, $A_2 \cap C_1 = \emptyset$, and $A_1 \cap A_2 = \emptyset$; and *loosely non-crossing* if $A_1 \cap C_2 = \emptyset$ and $A_2 \cap C_1 = \emptyset$. The two non-crossing properties are illustrated in Figure 2. Note that if two separations are non-crossing, then they are also loosely non-crossing. A collection \mathcal{S} of separations is *(loosely) laminar* if the separations of \mathcal{S} are pairwise (loosely) non-crossing.

	A_1	C_1	B_1
A_2	\emptyset	\emptyset	
C_2	\emptyset		
B_2			

(A) Non-crossing

	A_1	C_1	B_1
A_2		\emptyset	
C_2	\emptyset		
B_2			

(B) Loosely non-crossing

FIGURE 2. Illustrations of two separations $S_1 = (A_1, C_1, B_1)$ and $S_2 = (A_2, C_2, B_2)$ being (A) non-crossing and (B) loosely non-crossing.

The notion of non-crossing separations was used in [21] as part of the study of tree decompositions. Let (T, β) be a tree decomposition of G . For $X \subseteq V(T)$, let $\beta(X) = \bigcup_{x \in X} \beta(x)$. Let $t_1 t_2 \in E(T)$ be an edge of T . Let T_1 and T_2 be the components of $T \setminus \{t_1 t_2\}$ containing t_1 and t_2 , respectively. Let $C = \beta(t_1) \cap \beta(t_2)$, let $A = \beta(V(T_1)) \setminus C$, and let $B = \beta(V(T_2)) \setminus C$. Then, it follows from the properties of tree decompositions that (A, C, B) is a separation of G . Up to symmetry between A and B , we say that (A, C, B) is the *separation of G corresponding to the edge $t_1 t_2$ of T* . Let $\mathcal{S}(T, \beta)$ be the collection of separations corresponding to the edges of T . It is shown in [21] that for every tree decomposition (T, β) of G , it holds that $\mathcal{S}(T, \beta)$ is laminar, and conversely, for every laminar collection \mathcal{S} of separations of G , there exists a tree decomposition (T, β) of G such that $\mathcal{S} = \mathcal{S}(T, \beta)$. Therefore, there is a one-to-one correspondence between the laminar collections of separations of a graph G and the tree decompositions of G . This correspondence can be used to show that a graph has bounded treewidth: instead of attempting to bound the treewidth directly, one can instead study its separations.

In this paper we modify the approach above and use loosely laminar collections of separations (in fact, a slight variant of that). We then introduce a tool that reduces the problem of bounding the treewidth of a graph G to the problem of bounding the treewidth of a certain induced subgraph β of G . This is summarized in Theorem 2.10, but we try to explain and motivate our lemmas and definitions as we go.

Let G be a graph and let $w : V(G) \rightarrow \mathbb{R}$ be a normal weight function on G . Let $\varepsilon \in (0, \frac{1}{2}]$. We say that a separation (A, C, B) is ε -*skewed* if $w(A) < \varepsilon$ or if $w(B) < \varepsilon$. For the remainder of the paper, we assume by convention that if S is ε -skewed for some $\varepsilon \in (0, \frac{1}{2}]$, then $w(A(S)) < \varepsilon$. We have now broken the symmetry between A and B . Let us say that separations (A_1, C_1, B_1) and (A_2, C_2, B_2) are *A-loosely non-crossing* if $A_1 \cap C_2 = A_2 \cap C_1 = \emptyset$ and *A-non-crossing* if $A_1 \cap C_2 = A_2 \cap C_1 = A_1 \cap A_2 = \emptyset$ (we also sometimes use “A-loosely crossing” and “A-crossing” to mean “not A-loosely non-crossing” and “not A-non-crossing.”) We define A-laminar and A-loosely laminar collections of separations similarly. Note that if (A_1, C_1, B_1) and (A_2, C_2, B_2) are

A -loosely non-crossing ε -skewed separations, then every component D of $A_1 \cup A_2$ is a component of A_1 or a component of A_2 , and therefore $w(D) < \varepsilon$.

Let G be a graph, w a normal weight function on G , $c \in [\frac{1}{2}, 1)$ and d an integer such that G has no (w, c) -balanced separator of size at most d (this will be our standard set up, in view of Lemma 1.13). Let $S = (A, C, B)$ be a separation of G with $|C| \leq d$. Since C is not a (w, c) -balanced separator of G , we deduce that $w(A) > c$ or $w(B) > c$. Therefore, S is $(1 - c)$ -skewed, so by our convention $w(A) < 1 - c$, and consequently $w(B) > c$.

Henceforth, we assume that all collections of separations are ordered, and we refer to these ordered collections as *sequences of separations*. Let G be a connected graph, let w be a normal weight function on G , and let \mathcal{S} be an A -loosely laminar sequence of ε -skewed separations of G . The *central bag* for \mathcal{S} , denoted $\beta_{\mathcal{S}}$, is defined as follows:

$$\beta_{\mathcal{S}} = \bigcap_{S \in \mathcal{S}} B(S) \cup C(S).$$

We also define a weight function $w_{\mathcal{S}} : \beta_{\mathcal{S}} \rightarrow [0, 1]$ on $\beta_{\mathcal{S}}$ as follows. Let $\mathcal{S} = (S_1, \dots, S_k)$, and assume that $C(S_i) \neq \emptyset$ for all $S_i \in \mathcal{S}$. We equip the sequence \mathcal{S} with an *anchor map* $\text{anchor}_{\mathcal{S}} : \mathcal{S} \rightarrow V(G)$, such that $\text{anchor}_{\mathcal{S}}(S_i) \in C(S_i)$ for every $S_i \in \mathcal{S}$. We call $\text{anchor}_{\mathcal{S}}(S)$ the *anchor* for S . Since \mathcal{S} is loosely laminar, it holds that $\text{anchor}_{\mathcal{S}}(S) \in \beta_{\mathcal{S}}$ for every $S \in \mathcal{S}$ (this is proven in (i) of Lemma 2.1). For $v \in \beta_{\mathcal{S}}$, let $a(v) = \{i \text{ s.t. } v \text{ is the anchor for } S_i\}$. Let $w^*(A_i) = w(A_i \setminus \bigcup_{1 \leq j < i} A_j)$. Then, we let $w_{\mathcal{S}}(v) = w(v) + \sum_{i \in a(v)} w^*(A_i)$ for all $v \in \beta_{\mathcal{S}}$. Thus, the anchor for a separation S is a way to record the weight of $A(S)$ in $\beta_{\mathcal{S}}$.

We state the next few lemmas in slightly greater generality than what we need here, in order to be able to use them in future work. The following lemma gives important properties of central bags and their weight functions.

Lemma 2.1. *Let $c \in [\frac{1}{2}, 1)$ and let d be a positive integer. Let G be a connected graph, let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let \mathcal{S} be an A -loosely laminar sequence of separations of G such that $C(S)$ is connected and $|C(S)| \leq d$ for all $S \in \mathcal{S}$, and let $\beta_{\mathcal{S}}$ be the central bag for \mathcal{S} . Then,*

- (i) $C(S) \subseteq \beta_{\mathcal{S}}$ for every $S \in \mathcal{S}$,
- (ii) $\beta_{\mathcal{S}}$ is connected,
- (iii) $w_{\mathcal{S}}(\beta_{\mathcal{S}}) = 1$.

Proof. Let $\mathcal{S} = (S_1, \dots, S_k)$, and let $S_i = (A_i, C_i, B_i)$ for all $1 \leq i \leq k$. Since \mathcal{S} is an A -loosely laminar sequence of separations, we have $C_i \cap A_j = \emptyset$ for all $1 \leq i, j \leq k$. Since $V(G) \setminus \beta_{\mathcal{S}} \subseteq \bigcup_{1 \leq i \leq k} A_i$, it follows that $C_i \cap (V(G) \setminus \beta_{\mathcal{S}}) = \emptyset$. Therefore, $C_i \subseteq \beta_{\mathcal{S}}$ for all $1 \leq i \leq k$. This proves (i).

Let D be a connected component of $\beta_{\mathcal{S}}$. Let $I = \{i : C_i \cap D \neq \emptyset\}$. Since $C_i \subseteq \beta_{\mathcal{S}}$ and C_i is connected, it follows that $C_i \subseteq D$ for all $i \in I$. Since $N(A_i) \subseteq C_i$, we deduce that $D \cup \bigcup_{i \in I} A_i$ contains a connected component of G . Since G is connected, we have $D \cup \bigcup_{i \in I} A_i = V(G)$, and so $D = \beta_{\mathcal{S}}$. This proves (ii).

For $1 \leq i \leq k$, let $v_i = \text{anchor}_{\mathcal{S}}(S_i)$. Note that

$$\begin{aligned} w_{\mathcal{S}}(\beta_{\mathcal{S}}) &= \sum_{v \in \beta_{\mathcal{S}} \setminus \{v_1, \dots, v_k\}} w_{\mathcal{S}}(v) + \sum_{v \in \{v_1, \dots, v_k\}} w_{\mathcal{S}}(v) \\ &= \sum_{v \in \beta_{\mathcal{S}}} w(v) + \sum_{1 \leq i \leq k} w \left(A_i \setminus \bigcup_{1 \leq j < i} A_j \right) \\ &= \sum_{v \in V(G)} w(v), \end{aligned}$$

where the last equality holds since for all $v \notin \beta_{\mathcal{S}}$, we have $v \in A_i$ for some $1 \leq i \leq k$. Since $w(G) = 1$, it follows that $w_{\mathcal{S}}(\beta_{\mathcal{S}}) = 1$. This proves (iii). \blacksquare

Lemma 2.2. *Let $c \in [\frac{1}{2}, 1)$, and let Δ, d, t be positive integers with $d \geq 1 + \Delta + \dots + \Delta^t$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let \mathcal{S} be an A -loosely laminar sequence of separations such that for every $S \in \mathcal{S}$, it holds that $C(S)$ is connected and has diameter at most t . Let $\beta_{\mathcal{S}}$ be the central bag for \mathcal{S} . Then, $\beta_{\mathcal{S}}$ has no $(w_{\mathcal{S}}, c)$ -balanced separator of size at most $d(1 + \Delta + \dots + \Delta^t)^{-1}$.*

Proof. Suppose for a contradiction that $\beta_{\mathcal{S}}$ has a $(w_{\mathcal{S}}, c)$ -balanced separator Y of size at most $d(1 + \Delta + \dots + \Delta^t)^{-1}$. Since G has no (w, c) -balanced separator of size at most d , it follows that $N^t[Y]$ is not a (w, c) -balanced separator of G of size at most d . Since $|N^t[Y]| \leq |Y|(1 + \Delta + \dots + \Delta^t) \leq d$, it follows that there exists a connected component X of $G \setminus N^t[Y]$ such that $w(X) > c$. Let Q_1, \dots, Q_ℓ be the connected components of $\beta_{\mathcal{S}} \setminus Y$. Let D_1, \dots, D_m be the connected components of $G \setminus \beta_{\mathcal{S}}$. Let $\mathcal{I} = \{i \text{ s.t. } Q_i \cap X \neq \emptyset\}$ and $\mathcal{J} = \{j \text{ s.t. } D_j \cap X \neq \emptyset\}$. Since \mathcal{S} is A -loosely laminar, it holds that for every $1 \leq j \leq m$, there exists $S \in \mathcal{S}$ such that $D_j \subseteq A(S)$. Let $\mathcal{S} = (S_1, \dots, S_k)$. For $j \in \mathcal{J}$, let $f(j)$ be minimum such that $D_j \subseteq A(S_{f(j)})$, let $S(j) = S_{f(j)}$, and let $v(j) = \text{anchor}_{\mathcal{S}}(S(j))$.

(1) *For all $j \in \mathcal{J}$, it holds that $C(S(j)) \cap Y = \emptyset$.*

Suppose $C(S(j)) \cap Y \neq \emptyset$ for some $j \in \mathcal{J}$. By assumption, $C(S(j))$ has diameter at most t , so $C(S(j)) \subseteq N^t[Y]$. Then, $N(D_j) \subseteq N^t[Y]$, and so $D_j = X$. By the choice of d and since $D_j \subseteq A(S(j))$, it follows that $w(D_j) \leq w(A(S(j))) < 1 - c \leq c$, a contradiction. This proves (1).

Suppose that $|\mathcal{I}| = 0$. It follows that $|\mathcal{J}| \neq 0$, so let $j \in \mathcal{J}$. Now, $D_j \subseteq A(S(j))$ and, since $\mathcal{I} = \emptyset$, it holds that $C(S(j)) \subseteq Y$, contradicting (1). Now, suppose that $|\mathcal{I}| \geq 2$. Assume Q_1, Q_2 are such that $Q_1 \cap X \neq \emptyset$ and $Q_2 \cap X \neq \emptyset$. Since Q_1 and Q_2 are distinct connected components of $\beta_{\mathcal{S}} \setminus Y$, it follows that there exists $j \in \mathcal{J}$ such that $N(D_j) \cap Q_1 \neq \emptyset$ and $N(D_j) \cap Q_2 \neq \emptyset$. Since $N(D_j) \subseteq C(S(j))$, it follows that $C(S(j)) \cap Q_1 \neq \emptyset$ and $C(S(j)) \cap Q_2 \neq \emptyset$. Since $C(S(j))$ is connected, it holds that $C(S(j)) \cap Y \neq \emptyset$, contradicting (1).

Therefore, $|\mathcal{I}| = 1$. Let $\mathcal{I} = \{i\}$. It follows that $C(S(j)) \subseteq Q_i$ for all $j \in \mathcal{J}$, and, in particular, $v(j) \in Q_i$ for all $j \in \mathcal{J}$. Let $a(v) = \{t \text{ s.t. } v = \text{anchor}_{\mathcal{S}}(S_t)\}$. Now,

$$\begin{aligned} w_{\mathcal{S}}(Q_i) &= w(Q_i) + \sum_{v \in Q_i} \sum_{t \in a(v)} w^*(A(S_t)) \\ &\geq w(Q_i) + \sum_{j \in \mathcal{J}} w^*(A(S(j))) \\ &\geq w(Q_i) + \sum_{j \in \mathcal{J}} w(D_j) \geq w(X). \end{aligned}$$

But $w_{\mathcal{S}}(Q_i) \leq c$, since Y is a $(w_{\mathcal{S}}, c)$ -balanced separator of $\beta_{\mathcal{S}}$, and so $w(X) \leq c$, a contradiction. This proves Lemma 2.2. \blacksquare

Lemma 2.2 shows that if G is a graph with large treewidth, then the central bag $\beta_{\mathcal{S}}$ for a well-behaved sequence \mathcal{S} of separations of G also has large treewidth. Our goal is to construct a sequence of separations \mathcal{S} such that bounding the treewidth of $\beta_{\mathcal{S}}$ is easier than bounding the treewidth of G . We discuss how to find such a sequence later in this section. However, our candidate sequences of separations are usually not A -loosely laminar. Therefore, we would like a generalization of Lemma 2.2 that holds for sequences of separations that are not necessarily A -loosely laminar. We do this by defining the dimension of sequences of separations. Let G be

a graph and let \mathcal{S} be a sequence of separations of G . The *dimension* of \mathcal{S} , denoted $\dim(\mathcal{S})$, is the minimum number of laminar sequences of separations with union \mathcal{S} . Clearly, $\dim(\mathcal{S}) = 1$ if and only if \mathcal{S} is laminar.

Next, we need the notion of a canonical separation. Let G be a graph, let $X \subseteq V(G)$, and let us fix an ordering (v_1, \dots, v_n) of $V(G)$. The ordering defines a lexicographic order on the subsets of $V(G)$. The *canonical separation for X* , denoted $S_X = (A_X, C_X, B_X)$, is defined as follows: B_X is the largest-weight connected component of $G \setminus N[X]$, $C_X = X \cup (N[X] \cap N(B))$, and $A_X = V(G) \setminus (B_X \cup C_X)$. If there is more than one largest weight connected component of $G \setminus N[X]$, we choose the lexicographically minimum largest-weight component. Note that the definition of canonical separation is compatible with the convention that if a separation S is ε -skewed, then $w(A) < \varepsilon$. The set X is called the *center* of the separation S_X and is denoted $\text{center}(S_X)$. A separation S is called a *canonical separation* if there exists $X \subseteq V(G)$ such that $S = S_X$. For the remainder of the paper, if S_X is a canonical separation, then we assume that the anchor for S_X is contained in $\text{center}(S_X)$.

Let \mathcal{S} be an A -loosely laminar sequence of canonical separations, and let $S_1, S_2 \in \mathcal{S}$. Suppose $B(S_1) \cup C(S_1) \subseteq B(S_2) \cup C(S_2)$. Then, $\beta_{\mathcal{S} \setminus S_2} = \beta_{\mathcal{S}}$, so S_2 is an ‘‘unnecessary’’ member of \mathcal{S} with respect to the central bag. We say that S_1 is a *shield for S_2* if $B(S_1) \cup C(S_1) \subseteq B(S_2) \cup C(S_2)$.

Lemma 2.3. *Let $c \in [\frac{1}{2}, 1)$ and let Δ, d, t be positive integers with $d \geq 1 + \Delta + \dots + \Delta^{t+1}$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let $S_X = (A_X, C_X, B_X)$ be a canonical separation of G such that C_X has diameter at most t . Then, $w(B_X) > c$.*

Proof. Since $X \subseteq C_X \subseteq N[X]$ and C_X has diameter at most t , it follows that $N[X]$ has diameter at most $t + 1$, and so $|N[X]| \leq 1 + \Delta + \dots + \Delta^{t+1} \leq d$. Suppose $w(B_X) \leq c$. By definition, B_X is a largest-weight connected component of $G \setminus N[X]$, so $w(D) \leq c$ for every connected component D of $G \setminus N[X]$. But now $N[X]$ is a (w, c) -balanced separator of G of size at most d , a contradiction. This proves the lemma. \blacksquare

Let \mathcal{S} be a sequence of separations. We say \mathcal{S} is *primordial* if for every distinct $S_1, S_2 \in \mathcal{S}$, it holds that S_1 is not a shield for S_2 . Note that S_1 is a shield for S_1 . Also, we say \mathcal{S} is (a, t) -good if every vertex $v \in V(G)$ is the anchor for at most a separations in \mathcal{S} and if $C(S)$ has diameter at most t for every $S \in \mathcal{S}$.

Lemma 2.4. *Let $c \in [\frac{1}{2}, 1)$ and let a, Δ, d, t be positive integers with $d \geq 1 + \Delta + \dots + \Delta^{t+1}$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let \mathcal{S} be an (a, t) -good primordial laminar sequence of canonical separations. Then, \mathcal{S} is A -laminar.*

Proof. Suppose $S, S' \in \mathcal{S}$ are such that S, S' are A -crossing. By Lemma 2.3, it holds that $w(B(S)) > c$ and $w(B(S')) > c$, so it follows that $B(S) \cap B(S') \neq \emptyset$. Therefore, since S, S' are non-crossing but A -crossing, we may assume up to symmetry between S and S' that $A(S') \cap C(S) = A(S') \cap B(S) = B(S) \cap C(S') = \emptyset$. But now $B(S) \cup C(S) \subseteq B(S') \cup C(S')$, so S is a shield for S' , a contradiction. \blacksquare

The following lemma extends the idea of a central bag to sequences of separations of bounded dimension.

Lemma 2.5. *Let $c \in [\frac{1}{2}, 1)$ and let d, Δ, k, a, t be positive integers with $d \geq (1 + \Delta + \dots + \Delta^{t+1})(1 + \Delta + \dots + \Delta^t)^k$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let \mathcal{S} be an (a, t) -good sequence of canonical separations of G with $\dim(\mathcal{S}) = k$. Assume that $\text{center}(S)$ is connected for every $S \in \mathcal{S}$. Then, there exists a sequence $\mathcal{S}_1, \dots, \mathcal{S}_k$ of A -laminar sequences of separations, with $\mathcal{S}^* = \mathcal{S}_1 \cup \dots \cup \mathcal{S}_k$ and $\beta = \bigcap_{S \in \mathcal{S}^*} B(S) \cup C(S)$, such that the following hold:*

- (i) $\mathcal{S}^* \subseteq \mathcal{S}$,
- (ii) For all $S \in \mathcal{S} \setminus \mathcal{S}^*$, there exists $S' \in \mathcal{S}^*$ such that either S' is a shield for S or $\text{center}(S) \cap A(S') \neq \emptyset$,
- (iii) β is connected,
- (iv) There is a normal weight function $w_{\mathcal{S}}$ on β such that β has no $(w_{\mathcal{S}}, c)$ -balanced separator of size at most $d(1 + \Delta + \dots + \Delta^t)^{-k}$.

Proof. Let $\mathcal{S}'_1, \dots, \mathcal{S}'_k$ be a partition of \mathcal{S} into laminar sequences. First, we will prove inductively that there exists an A -laminar sequence \mathcal{S}_i for every $i \in \{1, \dots, k\}$, with $\mathcal{S}_i \subseteq \mathcal{S}'_i$, and $\beta_i = \bigcap_{S \in \mathcal{S}_1 \cup \dots \cup \mathcal{S}_i} B(S) \cup C(S)$, such that β_i is connected and there exists a normal weight function w_i on β_i such that β_i has no (w_i, c) -balanced separator of size at most $d(1 + \Delta + \dots + \Delta^{t+1})^{-i}$. This will prove (iii) and (iv).

Let $T_1 = \{B(S) \cup C(S) : S \in \mathcal{S}'_1\}$, and let \mathcal{S}_1 be the sequence formed by adding, for every inclusion-wise minimal $Y \in T_1$, a separation $S \in \mathcal{S}'_1$ such that $B(S) \cup C(S) = Y$. Notice that by construction, no separation in \mathcal{S}_1 is a shield for another separation in \mathcal{S}_1 , so \mathcal{S}_1 is primordial. By Lemma 2.4, \mathcal{S}_1 is A -laminar. Let β_1 be the central bag for \mathcal{S}_1 (so $\beta_1 = \bigcap_{S \in \mathcal{S}_1} B(S) \cup C(S)$), and let w_1 be the weight function $w_{\mathcal{S}_1}$ on β_1 . Note that since \mathcal{S} is (a, t) -good and $d \geq 1 + \Delta + \dots + \Delta^{t+1}$, it follows that $C(S) \leq d$ for all $S \in \mathcal{S}$. By Lemmas 2.1 and 2.2, β_1 is connected and β_1 has no (w_1, c) -balanced separator of size $d(1 + \Delta + \dots + \Delta^t)^{-1}$. This proves the base case.

Suppose we have a sequence $\mathcal{S}_1, \dots, \mathcal{S}_i$ with $\mathcal{S}_i \subseteq \mathcal{S}'_i$ and $\beta_i = \bigcap_{S \in \mathcal{S}_1 \cup \dots \cup \mathcal{S}_i} B(S) \cup C(S)$, such that β_i is connected and there exists a normal weight function w_i on β_i such that β_i has no (w_i, c) -balanced separator of size at most $d(1 + \Delta + \dots + \Delta^t)^{-i}$. Let S be a separation of G and let H be an induced subgraph of G . We define $S \cap H$ as the separation of H given by $(A(S) \cap H, C(S) \cap H, B(S) \cap H)$. If \mathcal{S} is a sequence of separations, we define $\mathcal{S} \cap H = \{S \cap H \text{ s.t. } S \in \mathcal{S}\}$. Let $\mathcal{S}''_{i+1} = \{S \in \mathcal{S}'_{i+1} \text{ s.t. } \text{center}(S) \subseteq \beta_i\}$. Let $T_{i+1} = \{B(S) \cup C(S) : S \in \mathcal{S}''_{i+1}\}$, and let \mathcal{S}_{i+1} be the sequence formed by adding, for every inclusion-wise minimal $Y \in T_{i+1}$, a separation $S \in \mathcal{S}''_{i+1}$ such that $B(S) \cup C(S) = Y$. Note that since the anchors for canonical separations are contained in their centers, it follows that restricting the anchor map for \mathcal{S}'_{i+1} to β_i is a valid anchor map for \mathcal{S}''_{i+1} . Note also that by construction, \mathcal{S}_{i+1} is primordial. By Lemma 2.4, \mathcal{S}_{i+1} is A -laminar. Let β_{i+1} be the central bag for $\mathcal{S}_{i+1} \cap \beta_i$, and let w_{i+1} be the weight function on β_{i+1} . Then,

$$\begin{aligned} \beta_{i+1} &= \bigcap_{S \in \mathcal{S}_{i+1} \cap \beta_i} B(S) \cup C(S) \\ &= \bigcap_{S \in \mathcal{S}_{i+1}} (B(S) \cup C(S)) \cap \beta_i \\ &= \bigcap_{S \in \mathcal{S}_1 \cup \dots \cup \mathcal{S}_{i+1}} B(S) \cup C(S). \end{aligned}$$

From the choice of \mathcal{S}''_{i+1} , it holds that $C(S)$ is connected for all $S \in \mathcal{S}''_{i+1}$. Note that since \mathcal{S} is (a, t) -good and $d \geq (1 + \Delta + \dots + \Delta^t)^k$, it follows that $|C(S)| \leq 1 + \Delta + \dots + \Delta^t \leq \frac{d}{(1 + \Delta + \dots + \Delta^t)^i}$. By Lemmas 2.1 and 2.2, it follows that β_{i+1} is connected and that β_{i+1} has no (w_{i+1}, c) -balanced separator of size $d(1 + \Delta + \dots + \Delta^t)^{-(i+1)}$. This completes the induction. Let $\beta = \beta_k$, let $w_{\mathcal{S}} = w_k$, and let $\mathcal{S}^* = \mathcal{S}_1 \cup \dots \cup \mathcal{S}_k$. Then, $\beta = \bigcap_{S \in \mathcal{S}^*} B(S) \cup C(S)$, β is connected, and $w_{\mathcal{S}}$ is a normal weight function on β such that β has no $(w_{\mathcal{S}}, c)$ -balanced separator of size at most $d(1 + \Delta + \dots + \Delta^{t+1})^{-k}$. This proves (iii) and (iv).

By construction, $\mathcal{S}^* \subseteq \mathcal{S}$, which proves (i). It remains to prove (ii). Let $S \in \mathcal{S} \setminus \mathcal{S}^*$, let $\beta_0 = G$ and let $\mathcal{S}''_1 = \mathcal{S}'_1$, and assume $S \in \mathcal{S}'_i$ for some $1 \leq i \leq k$. Suppose $S \notin \mathcal{S}''_i$. Then, $\text{center}(S) \not\subseteq \beta_{i-1}$, so it follows that there exists $S' \in \mathcal{S}_1 \cup \dots \cup \mathcal{S}_{i-1}$ such that $\text{center}(S) \cap A(S') \neq \emptyset$.

Now, assume $S \in \mathcal{S}_i'' \setminus \mathcal{S}_i$. Because there exists $Y \in T_i$ with $Y \subseteq B(S) \cup C(S)$, it follows that there exists $S' \in \mathcal{S}_i$ such that S' is a shield for S . This proves (ii). \blacksquare

Previously, central bags were defined for A -laminar sequences of separations. Here, we define central bags for sequences of separations of bounded dimension: we call β as in Lemma 2.5 a *central bag* for \mathcal{S} , $w_{\mathcal{S}}$ the *weight function* on β , and $\mathcal{S}_1, \dots, \mathcal{S}_k$ the *central bag generator* for \mathcal{S} . Next, we show how to construct useful sequences of separations of bounded dimension. A sequence \mathcal{S} of separations is *strongly laminar* if $C(S_1) \cap C(S_2) = \emptyset$ for all distinct $S_1, S_2 \in \mathcal{S}$. The following lemma states that under certain conditions, a strongly laminar sequence is laminar.

Lemma 2.6. *Let $c \in [\frac{1}{2}, 1)$ and let Δ, d be positive integers with $d \geq 1 + \Delta + \dots + \Delta^{t+1}$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let \mathcal{S} be a strongly laminar sequence of canonical separations such that $C(S)$ is connected and $|C(S)| \leq d$ for every $S \in \mathcal{S}$. Then, \mathcal{S} is laminar.*

Proof. Let $S_1, S_2 \in \mathcal{S}$. Since $C(S_2)$ is connected, it follows that either $C(S_2) \subseteq B(S_1)$ or $C(S_2) \subseteq A(S_1)$. Similarly, either $C(S_1) \subseteq B(S_2)$ or $C(S_1) \subseteq A(S_2)$. Suppose that $C(S_2) \subseteq B(S_1)$ and $C(S_1) \subseteq A(S_2)$. Then, $C(S_2) \cap A(S_1) = \emptyset$ and $C(S_1) \cap B(S_2) = \emptyset$. Since G is connected, it follows that $B(S_2) \cap A(S_1) = \emptyset$, and thus S_1 and S_2 are non-crossing. The other cases follow by symmetry. \blacksquare

Let X be a connected graph, and let $\mathcal{X} = \{Y \subseteq V(G) : Y \text{ is an } X \text{ in } G\}$. Let $\mathcal{S}_X = \{S_Y : Y \in \mathcal{X}\}$ be the sequence of canonical separations with centers in \mathcal{X} . We call \mathcal{S}_X the *X -covering sequence* for G . The following lemma shows that if \mathcal{S}_X is (a, t) -good, then \mathcal{S}_X has bounded dimension.

Lemma 2.7. *Let $c \in [\frac{1}{2}, 1)$ and let a, d, t, Δ be positive integers with $d \geq 1 + \Delta + \dots + \Delta^{t+1}$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let X be a connected graph and let \mathcal{S}_X be the X -covering sequence. Suppose \mathcal{S}_X is (a, t) -good. Then, $\dim(\mathcal{S}_X) \leq a(1 + \Delta + \dots + \Delta^{2t}) + 1$.*

Proof. Let H be a graph with $V(H) = \{C(S) : S \in \mathcal{S}_X\}$. Two vertices $C_1, C_2 \in V(H)$ are adjacent if $C_1 \cap C_2 \neq \emptyset$ in G . If $C_1 \cap C_2 \neq \emptyset$, then $C_1 \cup C_2$ has diameter at most $2t$, so $\text{anchor}_{\mathcal{S}_X}(S_1)$ has distance at most $2t$ from $\text{anchor}_{\mathcal{S}_X}(S_2)$. Because G has maximum degree Δ , there are at most $1 + \Delta + \dots + \Delta^{2t}$ vertices in $N^{2t}[v]$ for every $v \in V(G)$. Since every vertex $v \in V(G)$ is the anchor for at most a separations in \mathcal{S}_X , it follows that for each $S_1 \in \mathcal{S}_X$ there are at most $a(1 + \Delta + \dots + \Delta^{2t})$ separations S_2 such that $C(S_1) \cap C(S_2) \neq \emptyset$. Therefore, H has maximum degree $a(1 + \Delta + \dots + \Delta^{2t})$, so $\chi(H) \leq a(1 + \Delta + \dots + \Delta^{2t}) + 1$.

Let $\gamma = a(1 + \Delta + \dots + \Delta^{2t}) + 1$. Let $\chi : V(H) \rightarrow \{1, \dots, \gamma\}$ be a coloring of H , and let $\mathcal{S}_i = \{S : \chi(C(S)) = i\}$ for all $1 \leq i \leq \gamma$. Now, \mathcal{S}_i is strongly laminar for all $1 \leq i \leq \gamma$, and so by Lemma 2.6, \mathcal{S}_i is laminar for all $1 \leq i \leq \gamma$. Therefore, $\mathcal{S}_1, \dots, \mathcal{S}_\gamma$ is a partition of \mathcal{S}_X into γ laminar sequences, so $\dim(\mathcal{S}_X) \leq a(1 + \Delta + \dots + \Delta^{2t}) + 1$. \blacksquare

Let G and X be graphs, let \mathcal{S}_X be the X -covering sequence in G , and let β_X be a central bag for \mathcal{S}_X . For certain graphs X , we can restrict the properties of β_X in helpful ways. To do this, we use structures called *forcers*. Let G be a graph, and let $X, Y \subseteq V(G)$ such that $X \cap Y = \emptyset$. We say that X *breaks* Y if for every component D of $G \setminus N[X]$ we have that $Y \not\subseteq N[D]$. A graph F is an *X -forcer* for G if for every $Y \subseteq V(G)$ such that Y is an F in G , there exists $X' \subset Y$ such that X' is an X in G and X' breaks $Y \setminus X'$. For a class \mathcal{C} of graphs, we say that F is an *X -forcer* for \mathcal{C} if F is an X -forcer for every $G \in \mathcal{C}$.

Lemma 2.8. *Let G, X , and F be graphs such that F is an X -forcer for G . Let H be an F in G . Then, there exists X' which is an X in G such that $X' \subset H$ and $A_{X'} \cap H \neq \emptyset$.*

Proof. Since H is an F in G and F is an X -forcer, there exists $X' \subset H$ such that X' is an X in H and X' breaks $H \setminus X'$. Note that $B_{X'}$ is a component of $G \setminus N[X']$ such that $B_{X'} \cup (C_{X'} \setminus X') \subseteq N[B_{X'}]$. Therefore, $H \setminus X' \not\subseteq B_{X'} \cup C_{X'}$, and so $H \cap A_{X'} \neq \emptyset$. ■

Lemma 2.9. *Let $c \in [\frac{1}{2}, 1)$ and let Δ, d, a, t, k be positive integers with $d \geq (1 + \Delta + \dots + \Delta^{t+1})(1 + \Delta + \dots + \Delta^t)^k$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let X be a connected graph, let \mathcal{S}_X be the X -covering sequence for G , and assume \mathcal{S}_X is (a, t) -good and $\dim(\mathcal{S}_X) = k$. Let β_X be a central bag for \mathcal{S}_X . Then, if F is an X -forcer for G , then β_X is F -free.*

Proof. Suppose for a contradiction that $H \subseteq \beta_X$ is an F in G and F is an X -forcer for G . By Lemma 2.8, there exists $X' \subset H$ such that X' is an X in G and $A_{X'} \cap H \neq \emptyset$. Let $\mathcal{S}_1, \dots, \mathcal{S}_k$ be the central bag generator for \mathcal{S}_X and let $\mathcal{S}^* = \mathcal{S}_1 \cup \dots \cup \mathcal{S}_k$. Since $\beta_X \subseteq B(S) \cup C(S)$ for all $S \in \mathcal{S}^*$, it follows that $S_{X'} \notin \mathcal{S}^*$. Then, by (ii) of Lemma 2.5, there exists $S' \in \mathcal{S}^* \mathcal{S}_1 \cup \dots \cup \mathcal{S}_k$ such that either S' is a shield for $S_{X'}$ or $X' \cap A(S') \neq \emptyset$. If $X' \cap A(S') \neq \emptyset$, then $H \not\subseteq \beta_X$, a contradiction. Therefore, S' is a shield for $S_{X'}$. But now $A(S_{X'}) \subseteq A(S')$, and $\beta_X \subseteq B(S') \cup C(S')$, so $H \not\subseteq \beta_X$, a contradiction. ■

We now summarize what we have proved so far, as follows.

Theorem 2.10. *Let \mathcal{C} be a class of graphs with maximum degree at most Δ closed under taking induced subgraphs. Let t, N be integers, and let X be a connected graph with $|V(X)| < t$. Let \mathcal{F} be a set of graphs such that F is an X -forcer for \mathcal{C} for every $F \in \mathcal{F}$. Let $\gamma(x) = 1 + \Delta + \Delta^2 + \dots + \Delta^x$. If $\text{tw}(H) < N$ for every \mathcal{F} -free graph H in \mathcal{C} , then $\text{tw}(G) \leq 2N\gamma(t+1)^{\Delta^{t^2}\gamma(2t)+1}$ for every $G \in \mathcal{C}$.*

Proof. Let $G \in \mathcal{C}$ and suppose that $\text{tw}(G) > 2N\gamma(t)^{\Delta^{t^2}\gamma(2t)+1}$. We may assume that G is connected. By Lemma 1.13 there exists a uniform weight function w on G such that G has no $(w, \frac{1}{2})$ -balanced separator of size at most $N\gamma(t+1)^{\Delta^{t^2}\gamma(2t)+1}$. Let \mathcal{S}_X be the X -covering sequence for G . Since $|V(X)| < t$, X is connected, and G has maximum degree Δ , it follows that every vertex of G belongs to at most $\binom{\Delta^t}{t} \leq \Delta^{t^2}$ copies of X . Also, since $|V(X)| < t$ and X is connected, for every $S \in \mathcal{S}_X$, $C(S)$ has diameter at most t . Therefore \mathcal{S}_X is (Δ^{t^2}, t) -good. By Lemma 2.7 we have that $\dim(\mathcal{S}_X) \leq \Delta^{t^2}\gamma(2t) + 1$. Let β_X be a central bag for \mathcal{S}_X . Now by Lemma 2.5 there is a normal weight function w_X on β_X such that β_X has no $(w_X, \frac{1}{2})$ -balanced separator of size at most N . But by Lemma 2.9, β_X is \mathcal{F} -free, and therefore $\text{tw}(\beta_X) < N$, contrary to Lemma 1.14. ■

Next, we give a useful application of the results of this section. While it is not used in this paper, it is an important tool for future applications of the central bag method.

A *clique cutset* of a connected graph G is a set $C \subseteq V(G)$ such that C is a clique in G and $G \setminus C$ is not connected. Let K be a clique cutset in G , so in particular, $K \neq \emptyset$. The *canonical separation for K* , denoted $S_K = (A_K, C_K, B_K)$, is defined as follows: B_K is the lexicographically minimum largest-weight connected component of $G \setminus K$, $C_K = K$, and $A_K = V(G) \setminus (B_K \cup C_K)$. A separation is called a *clique separation* if it is the canonical separation for some clique cutset K of G . Let \mathcal{C} be a primordial sequence of clique separations such that for every clique separation S_K of G , it holds that S_K has a shield in \mathcal{C} . We call \mathcal{C} a *clique covering* of G . The next lemma states that \mathcal{C} is A -loosely laminar.

Lemma 2.11. *Let Δ, d be positive integers with $d > \Delta$ and let $c \in [\frac{1}{2}, 1)$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and suppose G has no (w, c) -balanced separator of size at most d . Let \mathcal{C} be a clique covering of G . Then, \mathcal{C} is A -loosely laminar.*

Proof. Suppose there is $S_K, S_{K'} \in \mathcal{C}$ such that S_K and $S_{K'}$ are A -loosely crossing. We may assume that $A_K \cap C_{K'} \neq \emptyset$. Since $C_{K'}$ is a clique and A_K is anticomplete to B_K , it follows that $C_{K'} \cap B_K = \emptyset$. Since B_K is connected and $A_{K'}$ is anticomplete to $B_{K'}$, it follows that $A_{K'} \cap B_K = \emptyset$. Since G has no (w, c) -balanced separator of size $\Delta+1$, it holds that $w(B_K) > \frac{1}{2}$ and $w(B_{K'}) > \frac{1}{2}$, so $B_K \cap B_{K'} \neq \emptyset$. If $C_K \cap A_{K'} = \emptyset$, then S_K is a shield for $S_{K'}$, a contradiction, so $C_K \cap A_{K'} \neq \emptyset$. Since C_K is a clique and $A_{K'}$ is anticomplete to $B_{K'}$, it follows that $C_K \cap B_{K'} = \emptyset$. Since $B_{K'}$ is connected, it holds that $B_{K'} \cap A_K = \emptyset$, so $B_K = B_{K'}$. Now, $C_K \cap C_{K'}$ is a cutset of G separating $B_K = B_{K'}$ from $A_K \cup A_{K'}$, and so, since G is connected, $C_K \cap C_{K'} \neq \emptyset$. But $S_{K''} = (A_K \cup A_{K'}, C_K \cap C_{K'}, B_K = B_{K'})$ is a shield for both S_K and $S_{K'}$, a contradiction. ■

Now, we prove two important results about the central bag for \mathcal{C} .

Theorem 2.12. *Let Δ, d be positive integers with $d > \Delta$ and let $c \in [\frac{1}{2}, 1)$. Let G be a connected graph with maximum degree Δ , let w be a normal weight function on G , and assume that G has no (w, c) -balanced separator of size at most d . Let \mathcal{C} be a clique covering of G , let $\beta_{\mathcal{C}}$ be the central bag for \mathcal{C} , and let $w_{\mathcal{C}}$ be the weight function on $\beta_{\mathcal{C}}$. Then:*

- (i) $\beta_{\mathcal{C}}$ has no $(w_{\mathcal{C}}, c)$ -balanced separator of size $d(1 + \Delta)^{-1}$, and
- (ii) $\beta_{\mathcal{C}}$ has no clique cutset.

Proof. Since cliques have diameter one, it follows from Lemmas 2.11 and 2.2 that $\beta_{\mathcal{C}}$ has no $(w_{\mathcal{C}}, c)$ -balanced separator of size $d(1 + \Delta)^{-1}$. This proves (i).

Next, we prove (ii). Suppose $\beta_{\mathcal{C}}$ has a clique cutset K . Let D_1 and D_2 be two connected components of $\beta_{\mathcal{C}} \setminus K$. By Lemma 2.1 (i), $N(A)$ is a clique for every connected component A of $G \setminus \beta_{\mathcal{C}}$, so we deduce that D_1 and D_2 are in different connected components of $G \setminus K$. Therefore, $\beta_{\mathcal{C}}$ intersects two connected components of $G \setminus K$, so $\beta_{\mathcal{C}} \cap A_K \neq \emptyset$. Since $\beta_{\mathcal{C}} \subseteq B_{K'} \cup C_{K'}$ for all $S_{K'} \in \mathcal{C}$, it follows that $S_K \notin \mathcal{C}$ and S_K does not have a shield in \mathcal{C} , a contradiction. This proves (ii). ■

3. TREewidth OF CLAW-FREE GRAPHS

In this section, we prove that for every k , every claw-free graph with bounded maximum degree and with no induced subgraph isomorphic to the line graph of a $(k \times k)$ -wall has bounded treewidth. Our proof relies on a structural description of claw-free graphs due to the second author and Seymour. In particular, the theorem we apply here is a straightforward corollary of the main result of [12]. To state this theorem, we need a couple of definitions from [12].

Given a graph H , a set F of unordered pairs of vertices of H is called a *valid set* for H if every vertex of H belongs to at most one member of F . For a graph H and a valid set F of H , we say that a graph G is a *thickening* of (H, F) if for every $v \in V(H)$ there is a nonempty subset $X_v \subseteq V(G)$, all pairwise disjoint and with union $V(G)$, for which the following hold.

- For each $v \in V(H)$, the set X_v is a clique of G ,
- if $u, v \in V(H)$ are adjacent in H and $\{u, v\} \notin F$, then X_u is complete to X_v in G ,
- if $u, v \in V(H)$ are non-adjacent in H and $\{u, v\} \notin F$, then X_u is anticomplete to X_v in G ,
- if $\{u, v\} \in F$, then X_u is neither complete nor anticomplete to X_v in G .

Let Σ be a circle and let $I = \{I_1, \dots, I_k\}$ be a collection of subsets of Σ , such that each I_i is homeomorphic to the interval $[0, 1]$, no two of I_1, \dots, I_k share an endpoint, and no three of them have union Σ . Let H be a graph whose vertex set is a finite subset of Σ , and distinct vertices $u, v \in V(H)$ are adjacent precisely if $u, v \in I_i$ for some $i = 1, \dots, k$. The graph H is called a *long circular interval graph*. Let F' be the set of all pairs $\{u, v\}$ such that $u, v \in V(H)$ are distinct endpoints of I_i for some i and there exists no $j \neq i$ for which $u, v \in I_j$. Also, let $F \subseteq F'$. Then, for every such H and F , every thickening G of (H, F) is called a *fuzzy long circular interval graph*.

Given a graph G , a *strip-structure* of G is a pair (H, η) , where H is a graph with no isolated vertices and possibly with loops or parallel edges, and η is a function mapping each $e \in E(H)$ to a subset $\eta(e)$ of $V(G)$, and each pair (e, u) consisting of an edge $e \in E(H)$ and an end u of e to a subset $\eta(e, u)$ of $\eta(e)$, with the following specifications.

- (S1) The sets $(\eta(e) : e \in E(H))$ are non-empty and partition $V(G)$.
- (S2) For each $v \in V(H)$, the union of sets $\eta(e, v)$ for all $e \in E(H)$ incident with v is a clique of G . In particular, $\eta(e, v)$ is a clique of G for all $e \in E(H)$ and $v \in V(H)$ an end of e .
- (S3) For all distinct $e_1, e_2 \in E(H)$, if $x_1 \in \eta(e_1)$ and $x_2 \in \eta(e_2)$ are adjacent, then there exists $v \in V(H)$ with v an end of both e_1 and e_2 , such that $x_i \in \eta(e_i, v)$ for $i = 1, 2$.

We say a strip-structure (H, η) is *non-trivial* if $|E(H)| \geq 2$. The following can be derived from Theorem 7.2 in [12].

Theorem 3.1 (Corollary of Theorem 7.2 from [12]). *Let G be a connected claw-free graph. Then one of the following holds.*

- We have $\alpha(G) \leq 3$.
- G is a fuzzy long circular interval graph.
- G admits a non-trivial strip structure (H, η) , such that for every $e \in E(G)$ with ends u and v ,
 - either $\alpha(\eta(e)) \leq 4$ or $\eta(e)$ is a fuzzy long circular interval graph; and
 - there exists a path P_e in $\eta(e)$ (possibly of length zero) with an end in $\eta(e, u)$ and an end in $\eta(e, v)$ whose interior is disjoint from $\eta(e, u) \cup \eta(e, v)$.

To begin with, we show that every fuzzy long circular interval graph with bounded maximum degree has bounded treewidth. Indeed, the proof is almost immediate from the following well-known fact about *chordal* graphs, i.e. graphs with no induced cycle of length at least four.

Theorem 3.2 (folklore). *A graph G is chordal if and only if it admits a tree decomposition (T, β) where for every $t \in V(T)$, the set $\beta(t)$ is a clique of G . Consequently, if G is chordal, then $\text{tw}(G) = \omega(G) - 1$.*

Theorem 3.3. *Let G be a fuzzy long circular interval graph of maximum degree at most Δ . Then we have $\text{tw}(G) \leq 4\Delta + 3$.*

Proof. Suppose that G is a thickening of (H, F) , where H is a long circular interval graph with Σ, I_1, \dots, I_k as in the definition, and F is a valid set for H as in the definition. Let G^* be the graph with $V(G^*) = V(G)$ and

$$E(G^*) = E(G) \cup \left(\bigcup_{\{u,v\} \in F} \{ab : a \in X_u, b \in X_v\} \right).$$

Then G^* is a long circular interval graph (the same interval representation Σ, I_1, \dots, I_k works for G^* , as well). In addition, we may easily observe that

- $\omega(G^*) \leq 2\omega(G) \leq 2(\Delta + 1)$;
- for all $i = 1, \dots, k$, the set $C_i = \bigcup_{u \in V(H) \cap I_i} X_u$ is a clique of G^* ; and
- for all $i = 1, \dots, k$, the graph $G - C_i$ is a chordal.

By Theorem 3.2 and the third bullet above, $G - C_1$ admits a tree decomposition (T, β) of width $\omega(G^*) - 1$. Now, for every $t \in V(T)$, let $\beta^*(t) = \beta(t) \cup C_1$. Then it is readily seen that (T, β^*) is a tree decomposition of G^* of width $\omega(G^*) + |C_1| - 1 \leq 2\omega(G^*) - 1$, where the last inequality follows from the the second bullet above. Hence, since G is a subgraph of G^* , we have $\text{tw}(G) \leq \text{tw}(G^*) \leq 2\omega(G^*) - 1 \leq 4\Delta + 3$, where the last inequality follows from the first bullet above. This proves Theorem 3.3. ■

The following is an easy observation.

Observation 3.4. *Let H be a graph and H' be a subdivision of H . Then $\text{tw}(H) = \text{tw}(H')$.*

We also use Theorem 1.1 with an explicit value of $f(k)$. In fact, a considerable amount of work has been devoted to understanding the order of magnitude of $f(k)$, and as of now, the following result of Chuzhoy and Tan provides the best known bound.

Theorem 3.5 ([13]). *There exist universal constants c_1 and c_2 such that for every integer k , every graph with no subgraph isomorphic to a subdivision of the $(k \times k)$ -wall has treewidth at most $c_1 k^9 \log^{c_2} k$.*

Now we are in a position to prove the main result of this section.

Theorem 3.6. *Let Δ, k be integers and c_1 and c_2 be as in Theorem 3.5. Let*

$$w(\Delta, k) = \max\{c_1 k^9 \log^{c_2} k (\Delta + 1)^2, 6(\Delta + 1)\} - 1.$$

Then for every claw-free graph G of maximum degree at most Δ and with no induced subgraph isomorphic to the line graph of a subdivision of the $(k \times k)$ -wall, we have $\text{tw}(G) \leq w(\Delta, k)$.

Proof. We may assume that G is connected, and so we may apply Theorem 3.1. Note that if we allow for trivial strip-structures, then the first two bullets of Theorem 3.1 will be absorbed into the first dash of the third bullet. In other words, we have

- (2) G admits a (possibly trivial) strip structure (H, η) , such that for every $e \in E(G)$ with ends u and v ,
- either $\alpha(\eta(e)) \leq 4$ or $\eta(e)$ is a fuzzy long circular interval graph; and
 - if (H, η) is non-trivial, then there exists a path P_e in $\eta(e)$ (possibly of length zero) with an end in $\eta(e, u)$ and an end in $\eta(e, v)$ whose interior is disjoint from $\eta(e, u) \cup \eta(e, v)$.

We also deduce:

- (3) For every $e \in E(H)$ and every $v \in V(H)$ incident with e , we have $|\eta(e, v)| \leq \Delta + 1$.

By (S2), $\eta(e, v)$ is a clique of G . So from G being of maximum degree at most Δ , we have $|\eta(e, v)| \leq \Delta + 1$. This proves (3).

- (4) For every $v \in V(H)$, the number of edges $e \in E(H)$ incident with v for which $\eta(e, v) \neq \emptyset$ is at most $\Delta + 1$.

For otherwise by (S2), the union of sets $\eta(e, v)$ for all $e \in E(H)$ with $v \in e$ contains a clique of G of size at least $\Delta + 2$, which is impossible. This proves (4).

- (5) H admits a tree decomposition (T_0, β_0) of width at most $c_1 k^9 \log^{c_2} k$.

If $|E(H)| \leq 1$, then we are done. So we may assume that (H, η) is non-trivial. Let the paths $\{P_e : e \in E(H)\}$ be as promised in the second bullet of (2), and let H^- be the graph obtained from H by removing its loops. Then, using (S1), (S2) and (S3) from the definition of a strip structure, one may observe that $G' = G[\bigcup_{e \in E(H^-)} V(P_e)]$ is isomorphic to the line graph of a subdivision H' of H^- . Now, since G has no induced subgraph isomorphic to the line graph of a subdivision of the $(k \times k)$ -wall, neither does G' , and so H' has no subgraph isomorphic to a subdivision of the $(k \times k)$ -wall. Thus, by Theorem 3.5, we have $\text{tw}(H') \leq c_1 k^9 \log^{c_2} k$, and so

by Observation 3.4, we have $\text{tw}(H^-) \leq c_1 k^9 \log^{c_2} k$. This, along with the fact that every tree decomposition of H^- is also a tree decomposition of H , proves (5).

(6) For every $e \in E(H)$, $\eta(e)$ admits a tree decomposition (T_e, β_e) of width at most $4(\Delta + 1)$.

Note that $\eta(e)$ is of maximum degree at most Δ . So if $\alpha(\eta(e)) \leq 4$, then we have $\text{tw}(\eta(e)) \leq |\eta(e)| \leq \alpha(\eta(e))(\Delta + 1) \leq 4(\Delta + 1)$, as desired. Otherwise, by the first bullet of (2), $\eta(e)$ is a fuzzy long circular interval graph, and so by Theorem 3.3, we have $\text{tw}(G) \leq 4\Delta + 3$. This proves (6).

Let (T_0, β_0) be as in (5), and for every $e \in E(H)$, let (T_e, β_e) be as promised by (6). We assume T_0 and T_e 's have mutually disjoint vertex sets and edge sets. Now, we construct a tree T as follows. For every $e \in E(H)$ with ends u and v , choose a vertex $s_e \in \beta_0^{-1}(u) \cap \beta_0^{-1}(v)$, which exists by definition of tree decomposition, and pick $t_e \in V(T_e)$ arbitrarily. Let $V(T) = V(T_0) \cup (\bigcup_{e \in E(H)} V(T_e))$, and $E(T) = \{s_e t_e : e \in E(H)\} \cup E(T_0) \cup (\bigcup_{e \in E(H)} E(T_e))$. We also define $\beta : V(T) \rightarrow 2^{V(G)}$ as follows. Let $t \in V(T)$. If $t \in V(T_0)$, then

$$\beta(t) = \bigcup_{u \in \beta_0(t)} \bigcup_{e \in E(H): u \text{ is an end of } e} \eta(e, u).$$

Otherwise, if $t \in V(T_e)$ for some $e \in E(H)$ with ends u and v , then $\beta(t) = \beta_e(t) \cup \eta(e, u) \cup \eta(e, v)$.

(7) (T, β) is a tree decomposition of G .

By (S1), for every vertex $x \in V(G)$, there exists $e \in E(H)$ such that $x \in \eta(e)$, and so (T_e, β_e) being a tree decomposition of $\eta(e)$, there exists $t \in V(T_e) \subseteq V(T)$ with $x \in \beta_e(t) \subseteq \beta(t)$.

Also, by (S3), for every edge $x_1 x_2 \in E(G)$, either $x_1 x_2 \in E(\eta(e))$ for some $e \in E(H)$, or there exists $v \in V(H)$ and $e_1, e_2 \in E(H)$ with v an end of e_1 and e_2 such that $x_i \in \eta(e_i, v)$ for $i = 1, 2$. In the former case, since (T_e, β_e) is a tree decomposition of $\eta(e)$, there exists $t \in V(T_e) \subseteq V(T)$ with $x_1, x_2 \in \beta_e(t) \subseteq \beta(t)$. In the latter case, since (T_0, β_0) is a tree decomposition of H , there exists $t \in V(T_0) \subseteq V(T)$ with $v \in \beta_0(t)$. Therefore, for $i = 1, 2$, we have

$$x_i \in \eta(e_i, v) \subseteq \bigcup_{e \in E(H): v \text{ is an end of } e} \eta(e, v) \subseteq \bigcup_{u \in \beta_0(t)} \bigcup_{e \in E(H): u \text{ is an end of } e} \eta(e, u) = \beta(t),$$

and so $x_1, x_2 \in \beta(t)$.

It remains to show that for every $x \in V(G)$, the graph $T|\beta^{-1}(x)$ is connected. By (S1), there exists a unique edge $e \in E(H)$ with ends u and v , with $x \in \eta(e)$. First, suppose that either $x \in \eta(e, u)$ or $x \in \eta(e, v)$, say the former. Then we have $\beta^{-1}(x) = \beta_0^{-1}(u) \cup V(T_e)$. Also, since $s_e \in \beta_0^{-1}(u)$ and $t_e \in V(T_e)$, we have $E(T|\beta^{-1}(x)) = \{s_e t_e\} \cup E(T_0|\beta_0^{-1}(u)) \cup E(T_e)$. Now, from (T_0, β_0) being a tree decomposition of H , we deduce that $T_0|\beta_0^{-1}(u)$ is connected, and so $T|\beta^{-1}(x)$ is connected, as well.

Next, suppose that $x \in \eta(e) \setminus (\eta(e, u) \cup \eta(e, v))$. Then we have $\beta^{-1}(x) = \beta_e^{-1}(x)$. So from (T_e, β_e) being a tree decomposition of $\eta(e)$, we deduce that $T|\beta^{-1}(x) = T_e|\beta_e^{-1}(x)$ is connected. This proves (7).

Now, let $t \in V(T)$. If $t \in V(T_0)$, then by (3), (4) and (5), we have

$$|\beta(t)| = \sum_{u \in \beta_0(t)} \sum_{e \in E(H): u \text{ is an end of } e} |\eta(e, u)| \leq |\beta_0(t)|(\Delta + 1)^2 \leq c_1 k^9 \log^{c_2} k (\Delta + 1)^2 \leq w(\Delta, k) + 1.$$

Also, if $t \in V(T_e)$ for some $e \in E(H)$, then by (3) and (6), we have

$$|\beta(t)| \leq |\beta_e(t)| + |\eta(e, u)| + |\eta(e, v)| \leq 6(\Delta + 1) \leq w(\Delta, k) + 1.$$

Hence, by (7), (T, β) is a tree decomposition of G of width at most $w(\Delta, k)$. This proves Theorem 3.6. \blacksquare

4. LONG CLAWS AND LINE GRAPHS OF WALLS

Here, we apply the results of Sections 3 to prove Theorem 1.8, that excluding a long claw and the line graphs of all subdivisions of $W_{k \times k}$ gives bounded treewidth.

Let t_1, t_2, t_3 be integers, with $t_1 \geq 0$ and $t_2, t_3 \geq 1$. Recall from the introduction that a *long claw*, also called a *subdivided claw*, denoted S_{t_1, t_2, t_3} , is a vertex v and three paths P_1, P_2, P_3 , of length t_1, t_2 , and t_3 , respectively, with one end v , such that $P_1 \setminus \{v\}$, $P_2 \setminus \{v\}$, and $P_3 \setminus \{v\}$ are pairwise disjoint and anticomplete to each other. We call P_1, P_2, P_3 the *paths* of S_{t_1, t_2, t_3} . The vertex v is called the *root* of S_{t_1, t_2, t_3} . For two graphs H_1, H_2 , we denote by $H_1 + H_2$ the graph with vertex set $V(H_1) \cup V(H_2)$ and edge set $E(H_1) \cup E(H_2)$. We start with a lemma.

Lemma 4.1. *Let t_1, t_2, t_3 be positive integers with $t_1 \geq 2$. Let G be an S_{t_1, t_2, t_3} -free graph. Then, $S_{t_1-1, t_2, t_3} + K_1$ is an S_{t_1-2, t_2, t_3} -forcer for G .*

Proof. Let H be an S_{t_1-1, t_2, t_3} in G , and let $u \in V(G)$ be anticomplete to H , so that $H \cup \{u\}$ is an $S_{t_1-1, t_2, t_3} + K_1$. Let $H = P_1 \cup P_2 \cup P_3$, where $P_1 = v-x_1-\dots-x_{t_1-1}$, $P_2 = v-y_1-\dots-y_{t_2}$, and $P_3 = v-z_1-\dots-z_{t_3}$. Let $X = H \setminus x_{t_1-1}$. Let D be a connected component of $G \setminus N[X]$. Suppose $u, x_{t_1-1} \in N[D]$. It follows that there exists a path $P = x_{t_1-1}-p_1-\dots-p_k-u$ from x_{t_1-1} to u with $P^* \subseteq D$, so X is anticomplete to P^* . Then, $H \cup \{p_1\}$ is isomorphic to S_{t_1, t_2, t_3} , a contradiction. Therefore, X breaks $\{u, x_{t_1-1}\}$, and it follows that $S_{t_1-1, t_2, t_3} + K_1$ is an S_{t_1-2, t_2, t_3} -forcer for G . \blacksquare

Now we can prove Theorem 1.8, which we restate.

Theorem 4.2. *Let Δ, t_1, t_2, t_3, k be positive integers with $t = t_1 + t_2 + t_3$. Let \mathcal{C} be the class of all S_{t_1, t_2, t_3} -free graph with maximum degree Δ and no induced subgraph isomorphic to the line graph of a subdivision of $W_{k \times k}$. There exists an integer $N_{k, t, \Delta}$ such that $\text{tw}(G) \leq N_{k, t, \Delta}$ for every $G \in \mathcal{C}$.*

Proof. The proof is by induction on $t_1 + t_2 + t_3$. If $t_1 = t_2 = t_3 = 1$, the result follows from Theorem 3.6. Thus we may assume that $t_1 \geq 2$. By Theorem 2.10 and Lemma 4.1, it is enough to find a bound on the treewidth of $(S_{t_1-1, t_2, t_3} + K_1)$ -free graphs in \mathcal{C} .

Let $H \in \mathcal{C}$ be $(S_{t_1-1, t_2, t_3} + K_1)$ -free. By the inductive hypothesis we may assume that there exists $X \subseteq V(H)$ such that X is an S_{t_1-1, t_2, t_3} in H . Since H does not contain $S_{t_1-1, t_2, t_3} + K_1$, it follows that $V(H) \subseteq N[X]$, and therefore $\text{tw}(H) \leq |V(H)| \leq t\Delta$. \blacksquare

5. t -THETAS, t -PYRAMIDS, AND LINE GRAPHS OF WALLS

In this section, we prove Theorem 1.10, that for all k, t , excluding t -thetas, t -pyramids, and the line graphs of all subdivisions of $W_{k \times k}$ in graphs with bounded degree gives bounded treewidth. The proof involves an application of Theorem 4.2. We also need the following lemma.

Lemma 5.1. *Let x_1, x_2, x_3 be three distinct vertices of a graph G . Assume that H is a connected induced subgraph of $G \setminus \{x_1, x_2, x_3\}$ such that H contains at least one neighbor of each of x_1, x_2, x_3 , and that subject to these conditions $V(H)$ is minimal subject to inclusion. Then, one of the following holds:*

- (i) *For some distinct $i, j, k \in \{1, 2, 3\}$, there exists P that is either a path from x_i to x_j or a hole containing the edge $x_i x_j$ such that*
 - $H = P \setminus \{x_i, x_j\}$, and
 - either x_k has at least two non-adjacent neighbors in H or x_k has exactly two neighbors in H and its neighbors in H are adjacent.
- (ii) *There exists a vertex $a \in H$ and three paths P_1, P_2, P_3 , where P_i is from a to x_i , such that*

- $H = (P_1 \cup P_2 \cup P_3) \setminus \{x_1, x_2, x_3\}$, and
 - the sets $P_1 \setminus \{a\}$, $P_2 \setminus \{a\}$ and $P_3 \setminus \{a\}$ are pairwise disjoint, and
 - for distinct $i, j \in \{1, 2, 3\}$, there are no edges between $P_i \setminus \{a\}$ and $P_j \setminus \{a\}$, except possibly $x_i x_j$.
- (iii) There exists a triangle $a_1 a_2 a_3$ in H and three paths P_1, P_2, P_3 , where P_i is from a_i to x_i , such that
- $H = (P_1 \cup P_2 \cup P_3) \setminus \{x_1, x_2, x_3\}$, and
 - the sets P_1, P_2 and P_3 are pairwise disjoint, and
 - for distinct $i, j \in \{1, 2, 3\}$, there are no edges between P_i and P_j , except $a_i a_j$ and possibly $x_i x_j$.

Proof. For some distinct $i, j, k \in \{1, 2, 3\}$, let P be a path from x_i to x_j with $V(P^*) \subseteq V(H)$ (in the graph where the edge $x_i x_j$ is deleted if it exists). Such a path exists since x_i and x_j have neighbors in H and H is connected. Assume that x_k has neighbors in P^* . Then, by the minimality of $V(H)$, we have $H = P^*$. If x_k has two non-adjacent neighbors in P^* , or x_k has two neighbors in P^* and its neighbors in P^* are adjacent, then outcome (i) holds. If x_k has a unique neighbor in P^* , then outcome (ii) holds. Thus, we may assume that x_k is anticomplete to P^* .

Let Q be a path with $Q \setminus \{x_k\} \subseteq H$ from x_k to a vertex $w \in H \setminus P$ (so $x_k \neq w$) with a neighbor in P^* . Such a path exists since x_k has a neighbor in H , x_k is anticomplete to P^* , and H is connected. By the minimality of $V(H)$, we have $V(H) = (V(P) \cup V(Q)) \setminus \{x_1, x_2, x_3\}$ and no vertex of $Q \setminus w$ has a neighbor in P^* . Moreover, by the argument of the previous paragraph, we may assume that x_i and x_j are anticomplete to $Q \setminus \{x_k\}$.

Now, if w has a unique neighbor in P^* , then outcome (ii) holds. If w has two neighbors in P^* and its neighbors in P^* are adjacent, then outcome (iii) holds. Therefore, we may assume that w has two non-adjacent neighbors in P^* . Let y_i and y_j be the neighbors of w in P^* that are closest in P^* to x_i and x_j , respectively. Let R be the subpath of P^* from y_i to y_j . Now, the graph H' induced by $((P \cup Q) \setminus R^*) \setminus \{x_1, x_2, x_3\}$ is a connected induced subgraph of $G \setminus \{x_1, x_2, x_3\}$ and it contains at least one neighbor of x_1, x_2 , and x_3 . Moreover, $H' \subset H$ since $R^* \neq \emptyset$. This contradicts the minimality of $V(H)$. \blacksquare

Now we are ready to prove Theorem 1.10, which we restate.

Theorem 5.2. *Let Δ, t, k be positive integers with $t \geq 2$. Let \mathcal{C} be the class of graphs of maximum degree Δ with no t -theta, no t -pyramid, and no induced subgraph isomorphic to the line graph of a subdivision of $W_{k \times k}$. There exists an integer $M_{k,t,\Delta}$ such that $\text{tw}(G) \leq M_{k,t,\Delta}$ for every $G \in \mathcal{C}$.*

Proof. We start by proving a result about the existence of forcers for \mathcal{C} .

- (8) $S_{t,t,t}$ is an $S_{t-1,t-1,t-1}$ -forcer for \mathcal{C} .

Let $G \in \mathcal{C}$, and let Y be an $S_{t,t,t}$ in G , let r be the root of Y , let x, y, z be the leaves of Y , and let $X = Y \setminus \{x, y, z\}$. Let D be a connected component of $G \setminus N[X]$, and suppose $\{x, y, z\} \subseteq N[D]$. Let $Z \subseteq D$ be an inclusion-wise minimal connected subset of D such that x, y, z each have a neighbor in Z . By Lemma 5.1, one of three cases holds. If case (ii) or case (iii) holds, then it is clear that $Y \cup Z$ is either a t -theta or a t -pyramid, so we may assume case (i) holds. Then, up to symmetry between x, y , and z , the subgraph of G induced on $Z \cup \{x, z\}$ is a path from x to z . Suppose y has two non-adjacent neighbors in Z . Let p, q in Z be the first and last neighbors of y in Z , such that x, p, q, z appear in x - Z - z in that order. Then G contains a theta between r and y through r - Y - y , r - Y - x - Z - p - y , and r - Y - z - Z - q - y . Since each of the paths of the theta contains a path of Y , it follows that every path of the theta has length at least t , a contradiction. Therefore, y has exactly two adjacent neighbors p, q in Z such that

x, p, q, z appear in x - Z - z in that order. But now G contains a pyramid from r to $\{y, p, q\}$ through r - Y - y , r - Y - x - Z - p , and r - Y - z - Z - q . Since each of the paths of the pyramid contains a path of Y , it follows that every path of the pyramid has length at least t , a contradiction. Therefore, X breaks $\{x, y, z\}$, so $S_{t,t,t}$ is an $S_{t-1,t-1,t-1}$ -forcer for G . This proves (8).

Now by Theorem 2.10, the result follows immediately from Theorem 4.2. \blacksquare

6. SUBCUBIC SUBDIVIDED CATERPILLARS AND THEIR LINE GRAPHS

In this section, we prove Theorem 1.11, that excluding a subdivided subcubic caterpillar and its line graph in graphs with bounded degree gives bounded treewidth. The proof uses Theorem 4.2 to get a structural result involving a family of induced subgraphs called (k, t) -creatures. We begin with the following lemma.

Lemma 6.1. *Let $\Delta > 0$ and $t > 0$ be integers, G be a graph and P be an induced path in G of length at least $t(1 + \Delta) - 1$. Also, let $z \in G \setminus P$ have at least one and at most Δ neighbors in P . Then there exists a subpath $P' = p'_0 \cdots p'_t$ of P of length t where $N(z) \cap P' = \{p'_0\}$.*

Proof. Suppose not. Let $P = p_0 \cdots p_\ell$, where $\ell \geq t(1 + \Delta) - 1$. Also, let $|N(z) \cap P| = j \leq \Delta$, and $0 \leq i_1 < \cdots < i_j \leq \ell$ satisfy $N(z) \cap P = \{p_{i_k} : k = 1, \dots, j\}$. If the subpath $p_{i_1} \cdots p_0$ of P is of length at least t , then $P' = p_{i_1} \cdots p_{i_1-t}$ satisfies Lemma 6.1, a contradiction. So $p_{i_1} \cdots p_0$ is of length at most $t - 1$. Similarly, $p_{i_j} \cdots p_\ell$ is of length at most $t - 1$. As a result, $j \geq 2$.

Now, if for some $k \in \{1, \dots, j - 1\}$, the subpath $p_{i_k} \cdots p_{i_{k+1}}$ of P is of length at least $t + 1$, then $P' = p_{i_k} \cdots p_{i_k+t}$ satisfies the lemma. Thus, for all $k \in \{1, \dots, j - 1\}$, $p_{i_k} \cdots p_{i_{k+1}}$ is of length at most t . But then P is of length at most $2(t - 1) + t(j - 1) = t(j + 1) - 2 \leq t(1 + \Delta) - 2$, which is impossible. This proves Lemma 6.1. \blacksquare

Next, we define creatures properly. For integers $k > 0$ and $t \geq 0$, a (k, t) -creature in a graph G is a pair $\Xi = (J, \mathcal{P})$, where

- J is a connected subset of G .
- \mathcal{P} is a collection of k mutually vertex-disjoint and anticomplete induced paths in $G \setminus J$, each of length t .
- For every $P \in \mathcal{P}$, an end v of P , called the P -joint of Ξ , satisfies the following:
 - v has a neighbor in J , and
 - $P \setminus v$ is anticomplete to J .

We also use Ξ to denote the set $J \cup (\bigcup_{P \in \mathcal{P}} P)$.

Lemma 6.2. *Let $\Delta, k, t \geq 0$ be integers and let G be a graph of maximum degree at most Δ with no (k, t) -creature. Let $X \subseteq G$ be an $S_{t+1,t+1,t+1}$ in G and let $x \in X$ be a leaf of X . Then the connected component of $G \setminus (N[X \setminus \{x\}] \setminus \{x\})$ containing x has no $(k - 1, t(1 + \Delta))$ -creature.*

Proof. Suppose not. Since $S_{t+1,t+1,t+1}$ is a $(3, t)$ -creature, it follows that $k \geq 4$. So we may choose a $(k - 1, t(1 + \Delta))$ -creature $\Xi = (J, \{P_1, \dots, P_{k-1}\})$ in the component C of $G \setminus (N[X \setminus \{x\}] \setminus \{x\})$ containing x with P_i -joint v_i for $i = 1, \dots, k - 1$, and an induced path L in C from x to some vertex $z \in N[\Xi]$, such that no vertex in $L \setminus \{z\}$ belongs to $N[\Xi]$. Let u be the root of X and let P, Q, R be the paths of X , with x an end of P . We deduce the following.

(9) z has a neighbor in $\bigcup_{i=1}^{k-1} P_i$.

Suppose for a contradiction that z is anticomplete to $\bigcup_{i=1}^{k-1} P_i$, and so $z \in N[J]$. Note that the path z - L - x - P - u has length at least t (since P does), and so we may choose a subpath P' of z - L - x - P - u containing z and of length equal to t . Also, for each $i \in \{1, \dots, k - 1\}$, since P_i is of length $t(1 + \Delta) \geq t$, we may choose a subpath P'_i of P_i containing v_i and of length equal to

t . But then $(J, \{P'_i : i \in \{1, \dots, k-1\}\} \cup \{P'\})$ is a (k, t) -creature in G , a contradiction. This proves (9).

Let I be the set of all indices $i \in \{1, \dots, k-1\}$ for which z has a neighbor in P_i . By (9), we have $I \neq \emptyset$, and so we may select an element $i_0 \in I$. Let $J' = J \cup P_{i_0} \cup L \cup P$. Note that J' is connected, and $u, z \in J'$.

(10) *For each $i \in I \setminus \{i_0\}$, there exists a subpath $P'_i = p_0^i \cdots p_t^i$ of P_i of length t , such that p_0^i is the only neighbor of z in P'_i , and P'_i is anticomplete to $J' \setminus \{z\}$.*

If $t = 0$, then $P_i = \{v_i\}$, z is adjacent to v_i , and so $P'_i = P_i$ satisfies (10). Thus, we may assume that $t \geq 1$, and so P_i is of length at least one. By Lemma 6.1 applied to $P_i \setminus v_i$ and z , we obtain a subpath $P'_i = p_0^i \cdots p_t^i$ of $P_i \setminus v_i$ (and so of P_i) of length t , such that p_0^i is the only neighbor of z in P'_i . Also, by the choice of X, Ξ and L , we obtain that $P_i \setminus \{v_i\}$ is anticomplete to $J' \setminus \{z\}$. Therefore, $P'_i \subseteq P_i \setminus \{v_i\}$ is anticomplete to $J' \setminus \{z\}$, as well. This proves (10).

Now, for each $i \in I \setminus \{i_0\}$, let P'_i be as promised in (10). Moreover, for each $i \in \{1, \dots, k-1\} \setminus (I \setminus \{i_0\})$, there exists a subpath P'_i of P_i containing v_i and of length equal to t , as P_i is of length $t(1 + \Delta) \geq t$. But then by (10) and the choice of X, Ξ and L , $(J', \{P'_i : i \in \{1, \dots, k-1\} \setminus \{i_0\}\} \cup \{Q \setminus u, R \setminus u\})$ is a (k, t) -creature in G , a contradiction. This proves Lemma 6.2. \blacksquare

Lemma 6.3. *Let $\Delta > 0$ and $\ell > 1$ be integers, and K be a connected graph of maximum degree at most Δ with $|K| \geq 1 + \sum_{i=0}^{\ell-2} \Delta^i$. Then K contains an induced path on at least ℓ vertices.*

Proof. Since G has maximum degree at most Δ , then for every $i \geq 0$ and every vertex $v \in G$, the set of vertices in G at distance i from v is of size at most Δ^i . Therefore, G has a vertex at distance at least $\ell - 1$ from v . This proves Lemma 6.3. \blacksquare

Recall from the introduction that a tree T is a *subcubic subdivided caterpillar* if it is of maximum degree at most three, and there exists a path $P \subseteq T$ such that P contains every vertex of T of degree three. The *spine* of T is the shortest path containing all vertices of degree at least three in T . A *leg* of a subdivided caterpillar T is a path in T from a leaf to a vertex of degree three in T whose all internal vertices are of degree two.

Theorem 6.4. *Let T be a subcubic subdivided caterpillar and let $\Delta > 0$ be an integer. Then there exist k, t such for every graph G of maximum degree at most Δ , if G contains a (k, t) -creature, then G contains a subdivision of T or the line graph of a subdivision of T .*

Proof. Note that if T has no vertex of degree three, then it is a path, and so setting $k = 1$ and $t = |T|$, we are done. So we may assume that S has a spine S with $|S| = s \geq 1$. We define $\ell = 6s^3 + s^2 - 1 > 1$ and $k = 1 + \Delta + \Delta^2 + \dots + \Delta^{\ell-2}$. Also, for every leaf u of T , let U_u be the leg of T having u as one of its end. Let t be the maximum length of U_u taken over all leaves $u \in T \setminus S$ of T . We claim that the values of k, t defined as above satisfy the theorem. Suppose not. Then there exists a graph G of maximum degree at most Δ , containing a (k, t) -creature but not containing a subdivision of T or the line graph of a subdivision of T .

(11) *We may choose H and \mathcal{P} with the following specifications.*

- H is a connected induced subgraph of G .
- \mathcal{P} is a collection of k mutually vertex-disjoint and anticomplete induced paths in G , each of length at least t .
- For every $P \in \mathcal{P}$, there is an end of P , denoted by v_P , with $P \cap H = \{v_P\}$ and $N(P \setminus \{v_P\}) \cap H = \{v_P\}$.

Note that there exists a (k, t) -creature $\Xi = (J, \mathcal{P})$ in G . For every $P \in \mathcal{P}$, let v_P be the P -joint

of Ξ . Let $H = (J \cup \{v_P : P \in \mathcal{P}\})$. From the definition of a (k, t) -creature, it follows directly that H and \mathcal{P} satisfy the above three bullets. This proves (11).

We choose H and \mathcal{P} satisfying (11) and with $|H|$ as small as possible. For every $P \in \mathcal{P}$, let v_P be as in the third bullet of (11). Let $A = \{v_P : P \in \mathcal{P}\}$ and $J = H \setminus A$.

(12) *Every vertex in $v \in J$ is a cut-vertex of H .*

For otherwise $H \setminus v$ and \mathcal{P} satisfy (11), violating the minimality of H . This proves (12).

For every vertex $v \in H$, let us say v is *redundant* if $v \in J$ and $N(v) \cap H$ is a stable set in H of size exactly two. Otherwise, we say v is *irredundant*.

(13) *There exists an induced path Q_1 in H containing at least $6s^3 + s^2 - 1$ irredundant vertices.*

For every redundant vertex $v \in H$ with $N(v) \cap H = \{x, z\}$, by *suppressing* v , we mean removing v from H and adding the edge xz to the resulting graph, while we refer to the reverse operation as *unsuppressing* v . Let K be the graph obtained from H by repeatedly suppressing redundant vertices until none is left. Note that the maximum degree of K does not exceed that of H , which in turn does not exceed Δ , as H is an induced subgraph of G . Also, we have $A \subseteq K$, and so $|K| \geq |A| = k = 1 + \sum_{i=0}^{\ell-2} \Delta^i$. Thus, by Lemma 6.3, K contains an induced path Q_0 on at least $\ell = 6s^3 + s^2 - 1$ vertices. Therefore, after unsuppressing all redundant vertices of H , we obtain an induced path Q_1 in H where every vertex in Q_0 is an irredundant vertex of Q_1 . This proves (13).

Henceforth, let Q_1 be as guaranteed in (13).

(14) *There exists an induced path Q_2 in H such that $Q_2 \subseteq J$ and Q_2 contains at least $6s^2 + s - 1$ irredundant vertices.*

Let $A_1 = Q_1 \cap A$ and $B_1 = Q_1 \cup (\bigcup_{P \in \mathcal{P}: v_P \in A_1} P)$. If $|A_1| \geq s$, then B_1 contains a subdivision of T , and hence so does G , a contradiction. It follows that $|A_1| \leq s - 1$. As a result, $Q_1 \setminus A_1$ has at most s connected components, and by (13), $Q_1 \setminus A_1$ contains at least $6s^3 + s^2 - s$ irredundant vertices. Therefore, there exists a connected component Q_2 of $Q_1 \setminus A_1$ (hence $Q_2 \subseteq J$) containing at least $6s^2 + s - 1$ irredundant vertices. This proves (14).

Henceforth, let Q_2 be as promised in (14). Note that by (9), every vertex in Q_2 is a cut-vertex of H . We say a vertex $x \in Q_2$ is *docile* if there exists a connected component of $H \setminus x$, denoted by D_x , such that $D_x \cap Q_2 = \emptyset$. The following is immediate from the definition.

(15) *Let $x \in Q_2$ be a docile vertex. Then*

- D_x is anticomplete to $Q_2 \setminus \{x\}$;
- $N(x) \cap D_x \neq \emptyset$; and
- for every docile vertex $y \in Q_2 \setminus \{x\}$, D_x is anticomplete to D_y in G .

Also, we deduce:

(16) *For every docile vertex $x \in G$, there exists a (possibly not unique) path $P_x \in \mathcal{P}$ with $v_{P_x} \in D_x$.*

Otherwise $H \setminus D_x$ and \mathcal{P} satisfy (11), violating the minimality of H . This proves (16).

(17) *There is a subpath Q_3 of Q_2 which has at least $6s$ irredundant vertices and no docile vertices.*

Let D be the set of all docile vertices in Q_2 . For every $x \in D$, let P_x be as in (16). Then by the second bullet in (15), we may choose a shortest path W_x in D_x from v_{P_x} to some vertex

$w_x \in N(x) \cap D_x$ (so W_x is induced and $W_x \setminus w_x$ is disjoint from $N(x) \cap D_x$). Let $B_2 = Q_2 \cup (\bigcup_{x \in D} (W_x \cup P_x))$. If $|D| \geq s$, then by the first and the third bullets of (15), B_2 contains a subdivision of T , and hence so does G , a contradiction. So $|D| \leq s-1$. It follows that $Q_2 \setminus D$ has at most s connected components, and by (14), $Q_2 \setminus D$ contains at least $6s^2$ irredundant vertices. Therefore, there exists a connected component Q_3 of $Q_2 \setminus D$ containing at least $6s$ irredundant vertices. This proves (17).

From now on, let Q_3 be as obtained in (17), $r = |Q_3|$, and $Q_3 = q_1 \cdots q_r$. By (17), we have $r \geq 6s \geq 6$. For every $i \in \{2, \dots, r-1\}$, we denote by L_i and R_i the components of $Q_2 \setminus q_i$ containing q_{i-1} and q_{i+1} , respectively. Since q_i is not docile, the vertex-set of every connected component of $H \setminus q_i$ contains either L_i or R_i . Also, by (12), q_i is a cut-vertex of H . So $H \setminus q_i$ has exactly two distinct connected components λ_i and ρ_i , such that $L_i \subseteq \lambda_i$ and $R_i \subseteq \rho_i$. For every $i \in \{1, \dots, r-1\}$, let us say i is a *bump* if there exists a connected component of $H \setminus \{q_i, q_{i+1}\}$, denoted by μ_i , such that $\mu_i \cap Q_2 = \emptyset$. From this definition, we immediately deduce the following.

(18) *Let $i \in \{1, \dots, r-1\}$ be a bump. Then*

- μ_i is anticomplete to $Q_2 \setminus \{q_i, q_{i+1}\}$;
- $(N(q_i) \cup N(q_{i+1})) \cap \mu_i \neq \emptyset$; and
- for every bump $j \in \{1, \dots, r-1\} \setminus \{i\}$, μ_i is anticomplete to μ_j in G .

Also, we have:

(19) *For every bump $i \in \{1, \dots, r-1\}$, there is a (possibly not unique) path $P_i \in \mathcal{P}$ with $v_{P_i} \in \mu_i$.*

Otherwise $H - \mu_i$ and \mathcal{P} satisfy (11), violating the minimality of H . This proves (19).

(20) *For every $i \in \{2, \dots, r-1\}$, if q_i is irredundant, then either $i-1$ or i is a bump.*

Since q_i is irredundant, it has a neighbor $w \in H \setminus \{q_{i-1}, q_i, q_{i+1}\}$. It follows that either $w \in \lambda_i \setminus \{q_{i-1}\}$ or $w \in \rho_i \setminus \{q_{i+1}\}$. Assume the former holds. Note that q_i separates w from R_i in H . Also, if there exists a path M in $H \setminus q_{i-1}$ from w to some vertex in $x \in L_{i-1}$, then q_i - w - M - x is a path in $H \setminus q_{i-1}$ from $q_i \in R_{i-1}$ to $x \in L_{i-1}$, a contradiction. Thus, q_{i-1} separates w from L_{i-1} in H . As a result, $\{q_{i-1}, q_i\}$ separates w from $Q_2 \setminus \{q_{i-1}, q_i\}$, and so the connected component of $H \setminus \{q_{i-1}, q_i\}$ containing w is anticomplete to $Q_2 \setminus \{q_{i-1}, q_i\}$. But then $i-1$ is a bump. Similarly, if $w \in \rho_i \setminus \{q_{i+1}\}$, then i is a bump. This proves (20).

Let $i \in \{1, \dots, r-1\}$ be a bump, and $P_i \in \mathcal{P}$ be as in (19). By the second bullet of (18), we may choose a shortest path Z_i in μ_i from v_{P_i} to some vertex $z_i \in (N(q_i) \cup N(q_{i+1})) \cap \mu_i$ (so Z_i is induced and $Z_i \setminus z_i$ is disjoint from $N(q_i) \cup N(q_{i+1})$). We say i is a bump of *type 1* if $z_i \in N(q_i) \setminus N(q_{i+1})$, of *type 2* if $z_i \in N(q_{i+1}) \setminus N(q_i)$, and of *type 3* if $z_i \in N(q_i) \cap N(q_{i+1})$. Note that every bump is of type 1, 2 or 3.

By (17), there exists $I \subseteq \{2, \dots, r-1\}$ with $|I| \geq 6s-2$ such that q_i is irredundant for all $i \in I$. Therefore, by (20), there exists a $I' \subseteq \{1, \dots, r-1\}$ with $|I'| \geq 3s-1$ such that every $i \in I'$ is a bump. Consequently, there exists $I'' \subseteq I$ with $|I''| \geq s$ such that all elements of I'' are bumps of the same type. Now, let $B_3 = Q_2 \cup (\bigcup_{i \in I''} (Z_i \cup P_i))$. If either all elements of I'' are of type 1 or all elements of I'' are of type 2, then B_3 contains a subdivision of T , which is impossible. Otherwise, all elements of I'' are of type 3. But then B_3 contains the line graph of a subdivision of T , a contradiction. This proves Theorem 6.4. \blacksquare

Next we prove a lemma.

Lemma 6.5. *Let Δ, b, k, t be positive integers where $k \geq 3$. Let \mathcal{C} be the class of graphs with maximum degree Δ that do not contain a (k, t) -creature or the line graph of a subdivision of $W_{b \times b}$. There exists $R_{b, t, k, \Delta}$ such that $\text{tw}(G) \leq R_{b, t, k, \Delta}$ for every $G \in \mathcal{C}$.*

Proof. Let $t_i = t(1 + \Delta)^{k-i}$. Let \mathcal{C}_i be the class of graphs with maximum degree Δ that do not contain an (i, t_i) -creature and have no induced subgraph isomorphic to the line graph of a subdivision of $W_{b \times b}$. We will prove by induction that there exists $R_{b,t,k,i,\Delta}$ such that $\text{tw}(G) \leq R_{b,t,k,i,\Delta}$ for every $G \in \mathcal{C}_i$. Since S_{t_3,t_3,t_3} is a $(3, t_3)$ -creature, for $i = 3$ the result follows from Theorem 4.2. Next we prove a result about the existence of forcers in graphs in \mathcal{C}_i .

(21) $S_{t_{i+1},t_{i+1},t_{i+1}} + H$ is a $S_{t_i,t_{i+1},t_{i+1}}$ -forcer for \mathcal{C}_i for every $(i-1, t_{i-1})$ -creature H .

Let $G \in \mathcal{C}_i$ and let H be an $(i-1, t_{i-1})$ -creature. Let Y be an $S_{t_{i+1},t_{i+1},t_{i+1}} + H$ in G , let $Y' = Y \setminus H$, let $x \in Y'$ be a leaf of Y' , and let $X = Y' \setminus \{x\}$. Let D be a connected component of $G \setminus N[X]$. Suppose $x \in N[D]$. Then, by Lemma 6.2, it follows that D has no $(i-1, t_{i-1})$ -creature. Since H is anticomplete to Y' , we have that $H \not\subseteq N[D]$. Therefore, X breaks $\{x\} + H$, so $S_{t_{i+1},t_{i+1},t_{i+1}} + H$ is a $S_{t_i,t_{i+1},t_{i+1}}$ -forcer for G . This proves (21).

By Theorem 2.10, it is now enough to bound the treewidth of $\{(S_{t_{i+1},t_{i+1},t_{i+1}} + H) : H \text{ is an } (i-1, t_{i-1})\text{-creature}\}$ -free graphs in \mathcal{C}_i . Let F be a graph with no (i, t_i) -creature. If F is $S_{t_{i+1},t_{i+1},t_{i+1}}$ -free, the result follows from Theorem 4.2. Thus, let $Q \subseteq V(F)$ be an $S_{t_{i+1},t_{i+1},t_{i+1}}$ in F . Then, $F \setminus N[Q]$ has no $(i-1, t_{i-1})$ -creature, so by the inductive hypothesis, we deduce that $\text{tw}(F \setminus N[Q]) \leq R_{b,t,k,i-1,\Delta}$. But $|Q| = 3t_i + 4$, and therefore $|N[Q]| \leq (3t_i + 4)\Delta$. Consequently, $\text{tw}(F) \leq R_{b,t,k,i-1,\Delta} + (3t_i + 4)\Delta$, and we can set $R_{b,t,k,i,\Delta} = R_{b,t,k,i-1,\Delta} + (3t_i + 4)\Delta$. ■

We can now prove Theorem 1.11, which we restate.

Theorem 6.6. *Let Δ be a positive integer and let T be a subcubic subdivided caterpillar. Let \mathcal{C} be the class of graphs with maximum degree at most Δ which do not contain a subdivision of T or the line graph of a subdivision of T . Then there exists $R_{\Delta,T}$ such that $\text{tw}(G) \leq R_{\Delta,T}$ for every $G \in \mathcal{C}$.*

Proof. Let $G \in \mathcal{C}$. By Theorem 6.4, there exist integers k, t such that if $G \in \mathcal{C}$ then G does not contain a (k, t) -creature.

Next we observe:

(22) *Let $G \in \mathcal{C}$. Then G does not contain the line graph of a subdivision of $W_{|T| \times |T|}$.*

Let H be the line graph of a subdivision of $W_{|T| \times |T|}$. Then, H contains the line graph of a subdivision of T . It follows that if G contains H , then G contains the line graph of a subdivision of T , a contradiction. This proves (22).

Now the result follows from Lemma 6.5. ■

REFERENCES

- [1] Aboulker, P., Adler, I., Kim, E. J., Sintuari, N. L. D., and Trotignon, N. “On the tree-width of even-hole-free graphs.” *European J. Combinatorics* 98 (2021), 103394.
- [2] Abrishami, T., Alecu, B., Chudnovsky, M., Hajebi S., and Spirkl S. “Induced subgraphs and tree decompositions X. Toward logarithmic treewidth in even-hole-free graphs.” *arXiv:2307.13684*, 2023.
- [3] Abrishami, T., Alecu, B., Chudnovsky, M., Hajebi S., and Spirkl S. “Tree independence number for (even hole, diamond, pyramid)-free graphs.” *arXiv:2305.16258*, 2023.
- [4] Abrishami, T., Alecu, B., Chudnovsky, M., Hajebi S., Spirkl, S., and Vušković, K. “Induced subgraphs and tree decompositions V. One neighbor in a hole.” *arXiv:2205.04420*, 2022.
- [5] Abrishami, T., Chudnovsky, M., Dibek, C., and Rzażewski, P. “Polynomial-time algorithm for Maximum Independent Set in bounded-degree graphs with no long induced claws.” *arXiv:2107.05434*, 2022.
- [6] Abrishami, T., Chudnovsky, M., Dibek, C., and Vušković, K. “Separators and submodular functions in perfect graphs.” *arXiv:2110.00108*, 2021.
- [7] Abrishami, T., Chudnovsky, M., Hajebi S., and Spirkl S. “Induced subgraphs and tree decompositions III. Three paths configurations and logarithmic treewidth.” *Advances in Combinatorics*, 2022:6, 29pp.
- [8] Abrishami, T., Chudnovsky, M., Hajebi S., and Spirkl, S. “Induced subgraphs and tree decompositions VI. Graphs with 2-cutsets.” *arXiv:2207.05538*, 2022.

- [9] T. Abrishami, M. Chudnovsky, S. Hajebi, and S. Spirkl. “Induced subgraphs and tree decompositions IV. (Even-hole, diamond, pyramid)-free graphs.” *Electronic Journal of Combinatorics* **30**, 2 (2023).
- [10] Abrishami, T., Chudnovsky, M., and Vušković, K. “Induced Subgraphs and Tree decompositions I. Even-hole-free graphs of bounded degree” *J. Combin. Theory Ser. B*, 157 (2022), 144 – 175.
- [11] Bodlaender, H. L. “Dynamic programming on graphs with bounded treewidth.” Springer, Berlin, Heidelberg, (1988), pp. 105 – 118.
- [12] Chudnovsky, M., Seymour, P. “Claw-free graphs. V. Global structure.” *J. Combin. Theory Ser. B* 98 (6) (2008), 1373 – 1410.
- [13] Chuzhoy, J., and Tan, Z. “Towards tight(er) bounds for the Excluded Grid Theorem.” *J. Combin. Theory Ser. B*, 146 (1) (2021), 219 – 265.
- [14] Cygan, M., Fomin, F. V., Kowalik, L., Lokshtanov, D., Marx, D., Pilipczuk, M., Pilipczuk, M., and Saurabh, S. *Parameterized algorithms*. Springer, (2015).
- [15] Fomin, F., Golovach, P., and Thilikos, D. “Contraction obstructions for treewidth.” *J. Combin. Theory Ser. B*, 101 (5) (2011) 302 – 314.
- [16] Harvey, D.J. and Wood, D.R. “Parameters Tied to Treewidth.” *J. Graph Theory* 84 (4) (2017), 364 – 385.
- [17] T. Korhonen. “Grid Induced Minor Theorem for Graphs of Small degree.” *J. Combin. Theory Ser. B* **160** (2023), 206 – 214.
- [18] V. Lozin and I. Razgon. “Tree-width dichotomy.” *European J. Combinatorics* 103 (2022): 103517.
- [19] Pilipczuk, M., Sintiari, N.L.D., Thomassé, S., and Trotignon, N. “(Theta, triangle)-free and (even hole, K_4)-free graphs. Part 2: bounds on treewidth.” *J. Graph Theory* 97 (4) (2021), 624 – 641.
- [20] Robertson, N. and Seymour, P. “Graph minors. V. Excluding a planar graph.” *J. Combin. Theory Ser. B*, 41 (1) (1996), 92 – 114.
- [21] Robertson, N. and Seymour, P. “Graph minors. X. Obstructions to tree-decomposition.” *J. Combin. Theory Ser. B* 52, 2 (1991), 153 – 190.
- [22] Robertson, N., and Seymour, P. “Graph minors. XVI. Excluding a non-planar graph.” *J. Combin. Theory, Ser. B*, **89** (2003), 43 – 76.
- [23] Sintiari, N. L. D., and Trotignon, N. “(Theta, triangle)-free and (even hole, K_4)-free graphs. Part 1 : Layered wheels.” *J. Graph Theory* 97 (4) (2021), 475 – 509.