

Assessing the size and uncertainty of remaining carbon budgets

Received: 5 August 2022

Accepted: 20 September 2023

Published online: 30 October 2023

 Check for updates

Robin D. Lamboll ¹✉, Zebedee R. J. Nicholls ^{2,3}, Christopher J. Smith ^{3,4,5}, Jarmo S. Kikstra ^{1,3,6}, Edward Byers ³ & Joeri Rogelj ^{1,3,6}

The remaining carbon budget (RCB), the net amount of CO₂ humans can still emit without exceeding a chosen global warming limit, is often used to evaluate political action against the goals of the Paris Agreement. RCB estimates for 1.5 °C are small, and minor changes in their calculation can therefore result in large relative adjustments. Here we evaluate recent RCB assessments by the IPCC and present more recent data, calculation refinements and robustness checks that increase confidence in them. We conclude that the RCB for a 50% chance of keeping warming to 1.5 °C is around 250 GtCO₂ as of January 2023, equal to around six years of current CO₂ emissions. For a 50% chance of 2 °C the RCB is around 1,200 GtCO₂. Key uncertainties affecting RCB estimates are the contribution of non-CO₂ emissions, which depends on socioeconomic projections as much as on geophysical uncertainty, and potential warming after net zero CO₂.

The remaining carbon budget (RCB) is the net amount of carbon dioxide (CO₂) humans can still emit while keeping global warming below a given limit with a given probability, taking into account the effect of other anthropogenic climate forcers^{1,2}. The concept is key when considering the speed of decarbonization required to meet the goal of the Paris Agreement to keep global warming to well below 2 °C relative to pre-industrial levels and pursuing efforts to limit it to below 1.5 °C (ref. 3). Many approaches to equitable international climate action involve estimating the global RCB and dividing it among nations according to various principles of equity^{4,5}. However the RCB for the Paris-relevant temperature targets (generally interpreted as a 50% chance of keeping global warming below 1.5 °C and anywhere from a 66% to 90% chance of 2.0 °C (ref. 6)) is small compared with the uncertainty in their values. This means subtle updates to the assessments can substantially affect the values, which makes their use challenging.

Previous work shows that the temperature rise is, to first order, not strongly dependent on when carbon emissions occur, only on their cumulative sum^{7–13}; however, the RCB is strongly dependent on both how much and when different types of non-CO₂ emissions occur^{14–19}. As a result, the RCB requires some set of scenarios describing co-evolutions of CO₂ and other emissions to estimate.

In the Working Group I (WG1) report for the IPCC Sixth Assessment Report (AR6)¹, a set of values were established using the approach presented in refs. 19,20, decomposing the RCB into CO₂ and non-CO₂ parts. The CO₂ part was assessed analytically by integrating information from multiple lines of evidence, while the non-CO₂ part was assessed using a reduced-complexity climate model (or emulator), MAGICC 7.5.1^{21–23}, calibrated to the IPCC AR6 assessment²⁴. The impact of non-CO₂ emissions on the RCB was estimated by fitting a linear trend to the relationship between future non-CO₂ and future total warming at net zero CO₂ emissions for available scenarios in the database accompanying the IPCC Special Report on Global Warming of 1.5 °C (SR1.5)²⁵. Following an update to historical data, an updated version of MAGICC (7.5.3) was available and used in the WG3 report^{26,27}.

The WG3 report discusses how updates to the non-CO₂ contribution at the time of net zero reduce the 1.5 °C RCB by about 100 GtCO₂ (about one-fifth) relative to estimates reported in WG1, although it did not tabulate values. It also makes comparisons between the RCB and the cumulative emissions until net zero of scenarios meeting a given temperature goal, which it finds approximately consistent with each other, although with less consistency for 1.5 °C of global warming than for higher levels. While this 20% change in the RCB estimate

¹Centre for Environmental Policy, Imperial College London, London, UK. ²School of Geography, Earth and Atmospheric Sciences, The University of Melbourne, Melbourne, Victoria, Australia. ³International Institute for Applied Systems Analysis, Laxenburg, Austria. ⁴Met Office Hadley Centre, Exeter, UK. ⁵School of Earth and Environment, University of Leeds, Leeds, UK. ⁶Grantham Institute – Climate Change and Environment, Imperial College London, London, UK. ✉e-mail: r.lamboll@imperial.ac.uk




Fig. 1 | How the carbon budgets are calculated. **a**, Schematic of how different factors contribute to the remaining carbon budget, based on ref. 20. **b**, Non-CO₂ contribution to warming after 2010–2019 for updated MAGICC, FaIR and the average of these values for each scenario in the AR6 database when it reaches net

zero CO₂. The bar at the top left indicates the median warming expected from 100 GtCO₂. We plot both the linear fit to the given quantiles and the quantiles of QRW fits to the averaged data points.

is small compared with the overall uncertainty and with past updates between the IPCC Fifth Assessment Report⁸ and the SR1.5¹⁹, it is politically important and warrants investigation.

In this Article, we update the RCB calculations fully and include results from an additional simple climate model calibrated for use in the latest IPCC report, FaIR^{24,28}. We then present improvements to the calculation methodology and examine the impact of these changes. We assess the RCBs through six contributing factors following ref. 20 and present the results of various changes in calculation that lead to updated values. Where not otherwise mentioned, RCBs are listed for keeping warming to the specified warming limits with 50% probability.

Sources of uncertainty

The main contributing factors in the budget calculation assessed by WG1¹ are transient climate response to cumulative CO₂ emissions (TCRE, the temperature rise per unit carbon emitted^{2,29}), historical warming (assessed human-induced global average temperature rise at present relative to pre-industrial levels), unrepresented Earth system feedbacks (ESFs), zero-emissions commitment (ZEC, the CO₂-based warming that continues after CO₂ emissions reach and are kept at net zero), warming from non-CO₂ emissions relative to the historical period and recent emissions. The equation combining these can be found in Methods, and a schematic of the equation is found in Fig. 1a. The values used for these variables can be found in Table 1.

Each of these factors comes with uncertainties, and the nature of and relationships between these uncertainties are complex. For example, while by default, uncertainty ranges are assumed to be normally distributed, other distributions for the range of TCRE are possible³⁰, as well as subtle nonlinearities³¹. Earth system feedbacks that are not included in the majority of Earth system models are notoriously difficult to quantify^{1,20,29}. Relationships between these distributions probably exist but are under-researched; we discuss these in the Supplementary Information. Pre-industrial temperatures are also somewhat uncertain; the IPCC considers them known to only 0.2 °C accuracy, consisting of uncertainty both in the relevant period and in what historical temperatures were. However, knowing exactly what pre-industrial temperatures were is irrelevant when considering future impacts of climate change, and this uncertainty can shrink if a

Table 1 | Table of values defining CO₂ contribution to warming

Name	Value	Discussion
TCRE	0.27–0.63 °C per 1,000 GtCO ₂ (1.0–2.3 °C per 1,000 PgC)	We investigate normal (default), positive-only normal and log-normal distributions.
Historical warming	1.07 °C 2010–2019	
ESF	26 ± 97 GtCO ₂	
ZEC	0 ± 0.19 °C	Based on ref. 33. We also consider an asymmetric distribution, where negative values are set to 0, and 0 ± 0.3, based on ref. 35.
Recent emissions	325 GtCO ₂	Emissions from 2015 to 2021, estimated from ref. 45.

The default assumptions are all following ref. 1 except where specified.

more recent historical benchmark is used with a predefined offset. The important point is that the same definition of historical temperature is used as in IPCC impact assessments and was intended by the Paris Agreement. In this way, we can call it a definitional uncertainty rather than uncertainty in future risk profile. This is explored further in the Supplementary Information. A similar definitional unclarity applies to the separation between CO₂ emissions from human-managed land-use changes and natural feedbacks included in the definition of TCRE. The problem can be removed by using a consistent database’s definition for all terms.

In principle, ZEC can influence our calculation both when it is positive and when it is negative. In practice, a negative ZEC may be fully realized only after peak warming occurs and therefore be less relevant for limiting maximum warming. Thus, while the assessed distribution of ZEC is a Gaussian based around zero, for very delayed ZEC impacts, the effective impact of ZEC for our calculation may be defined only by the positive part of this distribution. A recent model intercomparison project on ZEC (ZEC-MIP³²) indicates that for gradually declining emissions, some of the value identified as ZEC under the idealized conditions of an abrupt stop in emissions will be realized

Table 2 | Absolute and relative changes in remaining carbon budgets at 50, 66 and 90% exceedance probabilities on changing single aspects of the calculation from the default update

Temperature (°C)	Change	Relative 50% (%)	Absolute 50% (GtCO ₂)	Absolute 66% (GtCO ₂)	Absolute 90% (GtCO ₂)
1.5	Include permafrost in MAGICC results	-0.9	-3	-3	-3
1.5	Log-normal TCRE distribution	7.1	22	10	10
1.5	Positive-only normal TCRE distribution	2.0	6	6	8
1.5	Maximum non-CO ₂ warming	-17.8	-56	-53	-54
1.5	Non-CO ₂ warming at peak average total temperature, only NZ scenarios	-15.9	-50	-48	-48
1.5	Non-CO ₂ warming at peak total temperature	-7.5	-24	-23	-23
1.5	Non-CO ₂ warming at peak total temperature, only NZ scenarios	-17.4	-55	-52	-53
1.5	Non-CO ₂ warming at preharmonized NZ	1.9	6	6	6
1.5	Non-CO ₂ normalized 2010–2019	-1	-3	-3	-3
1.5	Recent emissions	36.8	117	117	117
1.5	Use QRW for non-CO ₂ fit	-6.9	-22	-21	-21
1.5	Use SR1.5 database	-0.9	-3	-3	-3
1.5	ZEC only impacts if positive	-38.8	-123	-43	0
1.5	ZEC standard deviation 0	1.3	4	102	327
1.5	ZEC standard deviation 0.3	-0.5	-1	-93	-319
2	Include permafrost in MAGICC results	-0.6	-8	-7	-6
2	Log-normal TCRE distribution	8.5	105	62	-8
2	Positive-only normal TCRE distribution	0.9	11	9	13
2	Maximum Non-CO ₂ warming	-7	-86	-77	-66
2	Non-CO ₂ warming at peak average total temperature, only NZ scenarios	-2.6	-32	-29	-24
2	Non-CO ₂ warming at peak total temperature	-4.8	-59	-53	-45
2	Non-CO ₂ warming at peak total, only NZ scenarios	-2.7	-33	-29	-25
2	Non-CO ₂ warming at preharmonized NZ	-0.6	-7	-7	-5
2	Non-CO ₂ normalized 2010–2019	-0.3	-4	-4	-3
2	Recent emissions	9.5	117	117	117
2	Use QRW for non-CO ₂ fit	1.5	19	16	14
2	Use SR1.5 database	-6.4	-78	-70	-60
2	ZEC only impacts if positive	-12.6	-155	-99	-19
2	ZEC standard deviation 0	0.3	4	57	172
2	ZEC standard deviation 0.3	-0.3	-4	-70	-221

NZ scenarios are scenarios reaching net zero after harmonization.

before net zero is reached³³, and this is replicated by simple climate models such as FaIR³⁴. This means that a negative ZEC could result in a budget increase, but by how much is uncertain. Depending on the other characteristics of the pathway, the time taken for ZEC to materialize may also reduce its impacts; if the scenario has decreasing non-CO₂ warming, this can mask a positive ZEC, and vice versa. Typically ZEC measured until 50 years after emissions stop is used in RCB estimates¹, but the peak non-CO₂ warming in MAGICC and FaIR is typically much earlier. A pre-net zero negative ZEC may also mean that some low level of CO₂ is emitted after peak temperature has been reached and does not affect the value of the peak. To further complicate matters, ZEC-MIP suggests that the uncertainty in ZEC depends on future warming whereas the IPCC provides a ZEC assessment at only one level of cumulative carbon emissions. It reports a central value for ZEC after 1,000 PgC of cumulative carbon emissions of zero with an assessed likely range of ± 0.3 °C in ref. 35; we include this uncertainty

as a robustness case. This estimate is for 2 °C of initial warming, so it is probably a little high for the 1.5 °C budget³³.

Despite this uncertainty, we can set bounds on the impact these considerations might have. We explore the impact of ignoring negative ZEC in our calculation as an upper bound on ZEC occurring too late to prevent peak warming from exceeding the predefined global warming limit. Table 2 indicates that this would have a very substantial effect, reducing the 50% 1.5 °C budget by over a third. This is the largest single impact explored here. While this is a high estimate of the impact and indicates an impact that might materialize only in the decades after net zero CO₂ is reached, it emphasizes that an increased understanding of ZEC would be very valuable to improve the accuracy of our budgets. Symmetrically increasing the uncertainty of ZEC has only very minor impact on the median budgets but substantially reduces the budget for a 66% chance of limiting warming to 1.5 °C or 2 °C. Reducing it would increase the 66% budget.




Fig. 2 | The impact of model and scenario family on carbon budgets. We use scenarios from the AR6 database to estimate non-CO₂ warming at peak warming by interpolating between scenarios from the same model with the same SSP (except for All AR6, where we interpolate between all scenarios). **a**, The impact of SSP family on non-CO₂ warming for IMAGE 3.0.1 scenarios (the only model with

a complete set of SSPs). **b**, Budgets for 1.5 °C for different models and scenarios for models where there are at least three scenarios. All AR6 scenarios similarly interpolates the non-CO₂ warming between all AR6 scenarios. Box plots show median and 25th–75th percentile range; whiskers show the 10th–90th percentile values, with seven points per SSP/model group.

Non-CO₂ warming contribution

Estimating RCBs requires an estimate of how much non-CO₂ emissions will contribute to warming. This requires estimates of both how much we will emit of many different species over time and what impact they will have on the climate^{36,37}. It therefore combines sociopolitical with geophysical uncertainty, which requires more complicated models than discussed so far. In an attempt to capture future socioeconomic developments, we use the AR6 scenario database³⁸, the most comprehensive current database of global emissions projections from different socioeconomic models. For assessing the geophysical uncertainty, we use two climate emulators. Full details of our emulator and database choices can be found in Methods. In the AR6 WG1 report, budgets were calculated with the emulator MAGICC and the SRL5 database²⁵; we explore adding FaIR and look at the impact of different versions of these models.

A version update to MAGICC (from version 7.5.1 to 7.5.3) reduced the 1.5 °C RCB by over 100 GtCO₂ (equivalent to roughly 0.05 °C in terms of temperature) due to a change in the historic aerosol emissions used to calibrate the model. A similar, although smaller, effect occurred when the FaIR model was updated. After combining the budgets, we find that the net effect of the updates is a 22% reduction of the 50% 1.5 °C RCB and a 13% reduction of the 66% 2 °C RCB. The budgets before and after updating are compared in Supplementary Fig. 1 and indicate uncertainties of around 100 GtCO₂ in the 1.5 °C budget and 200 GtCO₂ in the 2 °C budget from the geophysical impact of non-CO₂ emissions. This is broadly in line with the within-emulator uncertainty over non-CO₂ warming once the emulators are averaged, which gives 120 GtCO₂ for 1.5 °C and 210 GtCO₂ for 2 °C. Details of how we use these emulators to calculate non-CO₂ contributions are presented in Methods.

Previous estimates have assumed a linear relationship between additional temperature increase until peak warming and the non-CO₂ warming contribution until then. We investigate the impact of non-linear relationships, fitting a local quantile regression function called quantile rolling windows (QRW, described in Methods) as seen in Fig. 1. While the median QRW line deviates substantially from the linear relationship for higher degrees of total warming, for the 1.5 °C and 2 °C

budgets, the impact of allowing for a nonlinear relationship is less than 7% of the total budgets (Table 2).

The details of this are worth investigating. Projected non-CO₂ warming in pathways meeting 2.5 °C (at the lower end of a ‘current policy’ trajectory^{39,40}) are 0.30 °C, compared with 0.13 °C in 1.5 °C-compliant pathways—a difference expected to be equal to the warming of over 350 GtCO₂. For 2 °C-compliant pathways, this non-CO₂ warming decreases to 0.22 °C, or around 190 GtCO₂ cooler than the 2.5 °C pathway. Subtle changes in calculation methodology or non-CO₂ mitigation effort could result in similarly large changes to the budget. The total non-CO₂ contribution is tabulated in the Supplementary Information.

Normally, RCBs are calculated using all scenarios available in a particular database because there is no particular reason to favour one model or family of scenario above another. However, it is also instructive to consider how each individual model and scenario family represents the relationship between total and non-CO₂ warming. In the AR6 database, only the IMAGE model has at least three results for all of the widely used family of scenarios known as shared socioeconomic pathways (SSPs). The SSPs, numbered one to five, represent different population, urbanization and education storylines with differing levels of challenges to mitigation of and adaptation to climate change, influencing GHG emissions and global warming projections⁴¹. We can estimate how the relationship between non-CO₂ warming for a given total temperature rise depends on a specific set of global socioeconomic assumptions by interpolating between individual scenarios in the same SSP group, as shown in Fig. 2a. Interestingly, Fig. 2a shows that for each SSP ‘world’ of scenarios, there is a highly nonlinear relationship between non-CO₂ warming and peak total warming. However, if the SSP world is unknown, the overall trend is approximately linear (dashed line). As expected from earlier literature looking at deep mitigation scenarios⁴², non-CO₂ warming changes little with total warming for low total warming, but changes rapidly after some threshold. This threshold differs markedly between different SSP implementations. The different thresholds make the average fit to all SSP scenarios within the IMAGE model very linear; similar coincidences cause the linear approximation to be relatively good for the whole scenario collection.




Fig. 3 | Relationship between total and non-CO₂ warming over time in each emulator. a, MAGICC. b, FaIR. Five pathways from different models in AR6, representative of types of scenario that reach net zero with different pathways, are highlighted, with markers indicating how different definitions

of when to take non-CO₂ warming will affect the results. The same scenarios are highlighted in both plots in the same colour. Actual net zero refers to net zero in the final emissions data, whereas original net zero is for the scenario before harmonization.

In general, scenarios are designed to limit global warming to below a certain limit. Such scenarios aim to limit all GHG emissions, often modelled by applying a CO₂-equivalent price to all GHGs. Intuitively, one therefore expects a monotonic relationship between total warming and warming from non-CO₂ GHGs. However, clear limits to reducing non-CO₂ GHGs to zero have been identified, as insufficient mitigation measures have been identified to fully eliminate them for some activities such as agriculture⁴³. Aerosol forcing, which cools Earth and generally reduces as CO₂ emissions do, also reduces the strength of the expected correlation. Typically, this floor of non-CO₂ warming is already achieved in pathways that limit warming below 2.0 °C and is not markedly reduced further when aiming to limit warming further to lower levels⁴². This minimum floor of non-CO₂ emissions determines to a large degree the non-CO₂ warming expected around the time CO₂ emissions reach net zero. Importantly, this minimum floor level can differ substantially both between models and between model configurations, for example, depending on assumptions about future socioeconomic development, what mitigation options are possible in a model or how land systems are treated. While the 17–83% uncertainty range in the fit to scenario data corresponds to a change in budgets of only around 100 GtCO₂, many individual scenarios lie several times this outside this range, as seen in Fig. 1b.

We also investigate the impact of model and SSP scenario family on RCBs (Fig. 2b). Similar plots for the SRI.5 database can be found in Supplementary Fig. 2. While results are clearly different for each combination, no clear trends emerge, assuaging concerns that over-representation of a few models or scenario families in the AR6 database might systematically bias the RCB calculations. This concern is also assuaged by the small impact of changing between the AR6 and SRI.5 databases (<1% change for the 50% 1.5 °C budget and 6% change at 2 °C; Table 2), which have very different distributions of scenarios. We find that the standard deviation between the 50% 1.5 °C budgets calculated with different single model–SSP combinations are around 120 GtCO₂ with scenarios from the AR6 database. The range of values across all model–SSP combinations is from 490 GtCO₂ to a minimum value of 80 GtCO₂. Carrying out the same analysis with the scenarios available in the SRI.5 database results in similar values. This emphasizes that, depending on how successfully non-CO₂ emissions are reduced, the 1.5 °C RCB can change by a factor of around two and that a more precise RCB estimate needs to be conditional on the non-CO₂ pathway to net zero. Equally, the use of RCBs to assess the global warming performance of pathways can be made more accurate

if these sorts of conditional RCBs are used for comparison instead of generic central estimates.

Timing of non-CO₂ warming

The RCB is properly defined as the cumulative CO₂ emissions until annual net CO₂ emissions become zero. However, in virtually all pathways, CO₂ is the only major GHG to reach net zero. Residual emissions of other long-lived GHGs mean that Earth may continue to warm after reaching net zero. In practice, most scenarios that reach net zero CO₂ in our scenario databases then achieve net negative CO₂ emissions, and these negative emissions soon cancel out the warming from other forcers. Furthermore, both emulators used in this study have slightly negative ZECs (despite being calibrated to the IPCC AR6 assessment, which reports that the assessed value of ZEC is close to zero but with low confidence in the sign^{32,33,35}). This negative ZEC in the emulators usually prevents rises in median temperature in net zero scenarios to the end of the century. These facts defang but do not resolve the question of when we should measure the non-CO₂ warming.

Our default definition of non-CO₂ warming is the non-CO₂ contribution to warming at the time CO₂ emissions become net zero, consistent with recent IPCC RCB estimates¹. It has the benefit of decoupling the time used for determining non-CO₂ warming from the temporal evolution of the emulator's temperature response. This, for example, reduces the impact of the emulator's negative ZEC. It is, however, not the right choice of timing to ensure a given temperature is not exceeded because it does not estimate the non-CO₂ contribution at the time of peak temperature. We therefore consider variations on this assumption, described in detail in Methods and plotted for a few scenarios in Fig. 3. We find that while in some scenarios different approaches will get very similar results, in other scenarios results may differ by more than 0.1 °C. Some alternative approaches that can be considered are the non-CO₂ warming at the time of the model-reported net zero date (the date of net zero before emissions were harmonized to be consistent with recent emissions⁴⁴), the maximum possible non-CO₂ warming at any point over the twenty-first century and the non-CO₂ warming at maximum total temperature. The impact of changing between these measures is investigated in Table 2. Most pathways do not reach net zero and therefore do not contribute to the calculation in the first two approaches. It will generally improve results to also exclude them from other approaches since these scenarios do not reach their peak temperature during the twenty-first century and so do not have well-defined RCBs.




Fig. 4 | Plots of the changes to the carbon budget from each modification of the calculation. a, RCB for 50% chance of 1.5 °C. **b**, RCB for 50% chance of 2 °C. **c**, RCB for a range of temperatures (displaying only WGI and updated budgets). Uncertainty intervals indicate 33rd and 66th percentile budgets considering uncertainty distribution in CO₂ warming factors. Our default update corresponds to the changes until the use of QRW. In **a** and **b**, lighter blue represents an increase and red represents a decrease in the budget with each step.

The maximum non-CO₂ warming is an upper bound on the non-CO₂ term (which is negative in the equation for the RCB) rather than a fair estimate. The original (before harmonization to recent observations) net

zero test functions as a robustness check against any distorting impact of harmonization on pathways. Table 2 shows that the influence of this standard operation is minor. The most appropriate estimate of non-CO₂ warming comes from the estimates of non-CO₂ emissions at the time of peak warming since this is the deciding point for whether the scenario exceeds a particular limit. To combine the evidence that comes from the non-CO₂ warming estimates of MAGICC and FaIR, the temperature trends of the two emulators should be averaged before a maximum is found because otherwise the estimates may come from different years. Furthermore, viewing the two estimates as the true value plus an error term, averaging first and then finding the maximum allows more opportunities for error cancellation. We therefore consider average-first non-CO₂ warming at peak total temperature the best estimate of the marginal effect of non-CO₂ warming on the peak temperature. It is generally higher than the average non-CO₂ warming at net zero and hence decreases the 50% 1.5 °C RCB by 16% (Table 2). We use this technique in our ‘recommended update’. The temperature limit indicated by this non-CO₂ contribution is generally temporary and before peak CO₂ warming is reached; hence, the older practice (continued in our ‘default update’) of taking the contribution at net zero might be justified. The default update simply incorporates new data into the pre-existing methodology; the recommended update includes calculation methodology changes.

Comparison of recommended result with AR6 WGI results

The RCB factors updated from the AR6 WGI report to the approach we recommend can be summarized as follows: more recent emissions were included; the version of the climate emulator MAGICC was updated and calculations from FaIR were included; the database of scenarios was changed from SRI.5 to AR6; the non-CO₂ trend was found using QRW instead of a linear trend; and the non-CO₂ warming is taken at the time of peak total warming from scenarios that reach net zero instead of at the time of net zero. As seen in Fig. 4, recent emissions, recalibrating MAGICC and the addition of FaIR had the largest impact. The difference between recommended and previous budgets is small by 2 °C, and the updated RCB for higher degrees of warming is larger for temperature rises above 2.2 °C. A diagram of budgets with different MAGICC and FaIR versions can be found in Supplementary Fig. 1. Including a variety of emulators increases the robustness of the estimate as does making non-CO₂ assumptions explicit; applying a nonlinear relationship for estimating non-CO₂ warming as a function of total warming, choice of time for non-CO₂ warming and the database of scenarios is less impactful.

After making all these changes, our best (50%) RCB estimate starting after 2022 is 250 GtCO₂ (17–83% range from uncertainty in the impact of CO₂; –170 to 840 GtCO₂) for the RCB for limiting warming to 1.5 °C. The same uncertainty in the modelled impact of non-CO₂ forcing gives a range of 160–280 GtCO₂. We can combine these uncertainties using a generalized extreme values functional fit to the quantiles (described in Methods), resulting in a range of –200 to 830 GtCO₂. Note that the skew on non-CO₂ uncertainty lowers both limits. For 2 °C, we have 1,220 GtCO₂ (650–2,270 GtCO₂ from CO₂; 600–2,240 GtCO₂ including non-CO₂ uncertainty). For limiting warming to 2 °C with 66% or 90% probability, the RCBs are estimated at 940 and 500 GtCO₂, respectively. With 40 GtCO₂ emitted in 2022⁴⁵, this is roughly equivalent to 23 and 12 years of current CO₂ emissions for a 66% or 90% chance, respectively, of limiting warming to 2 °C and 6 years of emissions for a 50% chance of 1.5 °C. Translated into linear paths to net zero, this implies reaching global net zero CO₂ emissions around 2070, 2050 and 2035.

Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41558-023-01848-5>.

References

1. Canadell, J. G. et al. in *Climate Change 2021: The Physical Science Basis* (eds Masson-Delmotte, V. et al.) Ch. 5 (Cambridge Univ. Press, 2021); <https://doi.org/10.1017/9781009157896.007>
2. IPCC in *Climate Change 2021: The Physical Science Basis* (eds Masson-Delmotte, V. et al.) Annex VII (Cambridge Univ. Press, 2021); <https://doi.org/10.1017/9781009157896.022>
3. *Adoption of the Paris Agreement* FCCC/CP/2015/L.9/Rev.1 (UNFCCC, 2015); https://unfccc.int/sites/default/files/english_paris_agreement.pdf
4. van den Berg, N. J. et al. Implications of various effort-sharing approaches for national carbon budgets and emission pathways. *Climatic Change* **162**, 1805–1822 (2020).
5. Lahn, B. A history of the global carbon budget. *Wiley Interdiscip. Rev. Clim. Change* **11**, 636 (2020).
6. Schleussner, C.-F., Ganti, G., Rogelj, J. & Gidden, M. J. An emission pathway classification reflecting the Paris Agreement climate objectives. *Commun. Earth Environ.* **3**, 135 (2022).
7. Meinshausen, M. et al. Greenhouse-gas emission targets for limiting global warming to 2°C. *Nature* **458**, 1158–1162 (2009).
8. Collins, M. et al. *Long-term Climate Change: Projections, Commitments and Irreversibility* (Cambridge Univ. Press, Cambridge, 2013).
9. Tokarska, K. B. et al. Recommended temperature metrics for carbon budget estimates, model evaluation and climate policy. *Nat. Geosci.* **12**, 964–971 (2019).
10. Allen, M. R. et al. Warming caused by cumulative carbon emissions towards the trillionth tonne. *Nature* **458**, 1163–1166 (2009).
11. MacDougall, A. H. The oceanic origin of path-independent carbon budgets. *Sci. Rep.* **7**, 10373 (2017).
12. Zickfeld, K., Eby, M., Matthews, H. D. & Weaver, A. J. Setting cumulative emissions targets to reduce the risk of dangerous climate change. *Proc. Natl Acad. Sci. USA* **106**, 16129–16134 (2009).
13. Matthews, H. D., Gillett, N. P., Stott, P. A. & Zickfeld, K. The proportionality of global warming to cumulative carbon emissions. *Nature* **459**, 829–832 (2009).
14. Rogelj, J., Meinshausen, M., Schaeffer, M., Knutti, R. & Riahi, K. Impact of short-lived non-CO₂ mitigation on carbon budgets for stabilizing global warming. *Environ. Res. Lett.* **10**, 075001 (2015).
15. Rogelj, J. et al. Differences between carbon budget estimates unravelled. *Nat. Clim. Change* **6**, 245–252 (2016).
16. Tokarska, K. B., Gillett, N. P., Arora, V. K., Lee, W. G. & Zickfeld, K. The influence of non-CO₂ forcings on cumulative carbon emissions budgets. *Environ. Res. Lett.* **13**, 034039 (2018).
17. Millar, R. J. et al. Emission budgets and pathways consistent with limiting warming to 1.5°C. *Nat. Geosci.* **10**, 741–747 (2017).
18. Matthews, D. H. et al. An integrated approach to quantifying uncertainties in the remaining carbon budget. *Commun. Earth Environ.* **2**, 7 (2021).
19. Rogelj, J. et al. in *Special Report on Global Warming of 1.5°C* (eds Masson-Delmotte, V. et al.) 93–174 (WMO, 2018); <http://www.ipcc.ch/report/sr15/>
20. Rogelj, J., Forster, P. M., Kriegler, E., Smith, C. J. & Séférian, R. Estimating and tracking the remaining carbon budget for stringent climate targets. *Nature* **571**, 335–342 (2019).
21. Meinshausen, M., Raper, S. C. B. & Wigley, T. M. L. Emulating coupled atmosphere–ocean and carbon cycle models with a simpler model, MAGICC6—Part 1: model description and calibration. *Atmos. Chem. Phys.* **11**, 1417–1456 (2011).
22. Meinshausen, M. et al. The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500. *Geosci. Model Dev.* **13**, 3571–3605 (2020).
23. Nicholls, Z. et al. Reduced Complexity Model Intercomparison Project Phase 2: Synthesizing Earth System Knowledge for Probabilistic Climate Projections. *Earths Future* **9**, 2020–001900 (2021).
24. Forster, P. et al. in *Climate Change 2021: The Physical Science Basis* (eds Masson-Delmotte, V. et al.) 923–1054 (Cambridge Univ. Press, Cambridge, 2021); <https://doi.org/10.1017/9781009157896.009>
25. Huppmann, D., Rogelj, J., Kriegler, E., Krey, V. & Riahi, K. A new scenario resource for integrated 1.5°C research. *Nat. Clim. Change* <https://doi.org/10.1038/s41558-018-0317-4> (2018).
26. Riahi, K. et al. in *Climate Change 2022: Mitigation of Climate Change* (eds Shukla, P. R. et al.) Ch. 3 (Cambridge Univ. Press, 2022).
27. Guivarch, C. et al. in *Climate Change 2022: Mitigation of Climate Change* (eds Shukla, P. R. et al.) Annex iii (Cambridge Univ. Press, 2022).
28. Smith, C. J. et al. FAIR v1.3: a simple emissions-based impulse response and carbon cycle model. *Geosci. Model Dev.* **11**, 2273–2297 (2018).
29. MacDougall, A. H. Estimated effect of the permafrost carbon feedback on the zero emissions commitment to climate change. *Biogeosciences* **18**, 4937–4952 (2021).
30. Spafford, L. & MacDougall, A. H. Quantifying the probability distribution function of the transient climate response to cumulative CO₂ emissions. *Environ. Res. Lett.* **15**, 034044 (2020).
31. Nicholls, Z. R. J., Gieseke, R., Lewis, J., Nauels, A. & Meinshausen, M. Implications of non-linearities between cumulative CO₂ emissions and CO₂-induced warming for assessing the remaining carbon budget. *Environ. Res. Lett.* **15**, 074017 (2020).
32. Jones, C. D. et al. The Zero Emissions Commitment Model Intercomparison Project (ZECMIP) contribution to C4MIP: quantifying committed climate changes following zero carbon emissions. *Geosci. Model Dev.* **12**, 4375–4385 (2019).
33. MacDougall, A. H. et al. Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂. *Biogeosciences* **17**, 2987–3016 (2020).
34. Koven, C. D., Sanderson, B. M. & Swann, A. L. S. Much of zero emissions commitment occurs before reaching net zero emissions. *Environ. Res. Lett.* **18**, 14017 (2023).
35. Lee, J.-Y. et al. in *Climate Change 2021: The Physical Science Basis* (eds Masson-Delmotte, V. et al.) 553–672. (Cambridge Univ. Press, 2021); <https://doi.org/10.1017/9781009157896.006>
36. Mengis, N. & Matthews, H. D. Non-CO₂ forcing changes will likely decrease the remaining carbon budget for 1.5°C. *NPJ Clim. Atmos. Sci.* **3**, 19 (2020).
37. Jenkins, S. et al. Quantifying non-CO₂ contributions to remaining carbon budgets. *NPJ Clim. Atmos. Sci.* **4**, 47 (2021).
38. Byers, E. et al. AR6 Scenarios Database. Zenodo <https://doi.org/10.5281/zenodo.7197970> (2022).
39. Rogelj, J. et al. Credibility gap in net-zero climate targets leaves world at high risk. *Science* **380**, 1014–1016 (2023).
40. *Emissions Gap Report 2022* (UNEP, 2022); <https://www.unep.org/resources/emissions-gap-report-2022>
41. Riahi, K. et al. The shared socioeconomic pathways and their energy, land use, and greenhouse gas emissions implications: an overview. *Glob. Environ. Change* **42**, 153–168 (2017).
42. Rogelj, J. et al. Scenarios towards limiting global mean temperature increase below 1.5°C. *Nat. Clim. Change* **8**, 325–332 (2018).
43. Nabuurs, G.-J. et al. in *Climate Change 2022: Mitigation of Climate Change* (eds Shukla, P. R. et al.) Ch. 7 (Cambridge Univ. Press, 2022); <https://doi.org/10.1017/9781009157926.009>
44. Kikstra, J. S. et al. The IPCC Sixth Assessment Report WGIII climate assessment of mitigation pathways: from emissions to global temperatures. *EGUsphere* <https://doi.org/10.5194/gmd-15-9075-2022> (2022).

45. Friedlingstein, P. et al. Global carbon budget 2022. *Earth Syst. Sci. Data* **14**, 4811–4900 (2022).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the

source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023, corrected publication 2023

Methods

The equation for the RCB B for a temperature T is expressed as

$$B = (T - ZEC - \delta T_{\text{nonCO}_2}(T) - \delta T_{\text{historic}}) / \text{TCRE} - \text{ESF}(T) - E_{\text{recent}}, \quad (1)$$

for $\delta T_{\text{historic}}$ the historical warming, $\delta T_{\text{nonCO}_2}$ the non-CO₂ warming, ESF the CO₂ emitted from any Earth system feedbacks otherwise not covered by the TCRE uncertainty and E_{recent} emissions that occurred too recently to be accounted for in the period of historical warming. Values for these can be found in Table 1 and a schematic is in Fig. 1a. This equation is used for all parts of this study, and we propose to modify only the distributions of individual terms. Diagrams showing the relationships between distributions of TCRE and distributions of RCB can be found in Supplementary Fig. 3. Robustness checks using emulators that would be sensitive to correlations between variables are presented in Supplementary Fig. 4.

For each temperature target, 10 million values for ESF, ZEC and TCRE are drawn from the relevant distributions (Table 1), assuming independence between each of the estimates, combined with the best estimate of non-CO₂ warming contribution for this level of peak warming and plugged into equation (1). Quantiles of the resulting budgets are then calculated. Where the normal distribution is used to capture the uncertainty in TCRE, it is possible to obtain a negative TCRE value. This would be an odd assumption that often results in a negative budget. However, this negative budget is the lower bound rather than the upper bound for emissions reaching that temperature target. The probability of a negative TCRE is less than 1% with our distribution based on the IPCC AR6 assessment¹; for this reason and for visual clarity, graphs such as Supplementary Fig. 3 do not depict the top and bottom 1% of results for any distribution. The impact of replacing these negative TCRE draws with a very small positive value (a positive-only normal distribution) is investigated in Table 2 and found to be minor for 50%–90% budgets; it is only relevant to the extreme-probability budgets. ESF is expressed as CO₂ emitted per degree of warming and is also given by a normal distribution of values multiplied by future warming; its impact is small for budgets below 2.5 °C, so we do not consider robustness checks of this. The emissions from 2020 to 2022 were not included in the WG1 budgets and were recently evaluated as amounting to 121 GtCO₂ (ref. 45), although changes in the estimates of emissions in earlier years mean that our recent emissions change by 131 GtCO₂.

For the non-CO₂ components of projections, we default to (and recommend using) the AR6 scenario database³⁸ but also investigate the use of the SRI.5 database²⁵ for comparison with previous IPCC RCBs. The emissions scenarios in both databases are vetted to ensure that key emissions species and socioeconomic variables are within reasonable ranges in the recent past and near future, then harmonized to match historical emissions precisely and infilled with any missing emissions⁴⁴.

The emissions scenarios from the AR6 and SRI.5 databases are then run through reduced-complexity climate model emulators. For climate emulators, we use runs from both MAGICC 7.5.3²¹ and FaIR 1.6.2²⁸. By default, we use versions of the emulators calibrated to assessments in the AR6 WG1 report²⁴, but we also compare these results with runs using older model versions and pre-AR6 calibrations (MAGICC 7.5.1 and FaIR 1.3.4) for robustness checks. Both versions of MAGICC also have the option of including a module designed to mimic the effects of permafrost thawing; the impact of turning this option on is also investigated. Note that this affects only the relationship between total warming and non-CO₂ emissions as the feedback of permafrost melting on the warming per unit of CO₂ is included in the ESF.

The non-CO₂ warming contribution is calculated slightly differently in FaIR and MAGICC. In FaIR, we calculate the warming from only anthropogenic emissions and the warming from the same scenarios with only anthropogenic CO₂ emissions. We subtract the average temperatures in the period 2010–2019 from each case, and the difference between these values is then the non-CO₂ contribution to warming.

In MAGICC, we do three experiments for each scenario: one with all emissions and natural climate forcers, one with anthropogenic forcers only and one with anthropogenic CO₂ emissions only. The difference between the all anthropogenic forcers and anthropogenic CO₂-only experiments is the non-CO₂ contribution to warming. We use a different approach to MAGICC when processing FaIR data because by default, FaIR includes the effects of a substantial solar cycle in future temperatures, which we avoid including. Precalculated MAGICC and FaIR results for all these cases are included in the codebase for running these calculations.

In all cases, the peak temperature in the emulator until 2100 is compared with the non-CO₂ warming at various times, depending on the non-CO₂ peaking definition (see discussion in main text). The default peaking choice, in keeping with previous work, is the non-CO₂ warming at the time the scenario actually reaches net zero in the harmonized emissions, but we also consider the time it originally reached net zero CO₂ before CO₂ emissions were harmonized to recent historical data (non-CO₂ warming at original net zero CO₂); the non-CO₂ warming at peak total warming, either in all cases or restricted to scenarios that make net zero (after harmonization); the non-CO₂ warming at the time of peak total warming in MAGICC, conditional on meeting net zero (after harmonization); and, very conservatively, the maximum non-CO₂ warming experienced in the twenty-first century. While non-CO₂ warming at net zero CO₂ is defined only for emissions trajectories that reach net zero CO₂ (either before or after harmonization), it can also be calculated for scenarios that never reach net zero. These scenarios typically are either high warming, and hence less relevant for low-warming calculations, or nearly reach zero, meaning the difference between approaches is smaller.

Whichever value of non-CO₂ warming is used, the rest of the calculation is the same. If both MAGICC and FaIR are used, the peak warming and non-CO₂ warming are averaged before the fit to the relationship is made, as seen in Fig. 1b.

There are several ways of fitting a relationship of non-CO₂ warming contribution to total warming at peak. The default method is a linear trend, which fits a straight line to the points using quantile regression to find the 50th percentile. This is preferred to a least-squares fit, which would be more influenced by extreme points. Alternatively, this fit can be performed using a QRW method, which weights points according to $1/(1 + \Delta x^2)$ for Δx the distance along the x axis, normalized by a value proportional to the total range of x values. With this weighting, weighted quantiles are evaluated at ten points equally spaced across the x axis, and results for points in between are linearly interpolated. See ref. 46 for more details on this method. This technique is reasonably similar to calculating the rolling quantiles of points but with smoother behaviour and defined over a wider x -axis interval. A third technique is linear interpolation, which is appropriate only when few data points are available. We linearly interpolate between these known points to find the non-CO₂ warming corresponding to this total temperature rise, with total temperatures outside the known range assumed to have non-CO₂ warming equal to the closest point.

For runs where only a single model/scenario family is used, we filter the database for each specific model and then look for cases with at least three scenarios with names starting ‘SSP*n*’ for n between 1 and 5. We calculate the non-CO₂ component using the non-CO₂ warming at peak total warming of these cases, not filtering out scenarios that do not reach net zero to avoid a lack of data. Linear interpolation is used to make the fit.

To combine errors from CO₂ and non-CO₂ physical uncertainty, we fit a generalized extreme value distribution (GEV) to the quantiles of the data for each at each relevant temperature, following the approach of ref. 47. For the CO₂ error (the largest contribution), we fit the GEV to the 0.1, 0.17, 0.33, 0.5, 0.66, 0.83 and 0.9 quantiles of standard runs. For the non-CO₂ uncertainty, we make runs using the 0.17, 0.5 or 0.83 non-CO₂ warming temperature quantiles for each scenario. Fits to the

scenario warming are made as before and budgets calculated. A GEV fit to the 50% probability budget as a function of the non-CO₂ quantile is made, and to convert this into an uncertainty, we subtract that value at 0.5 non-CO₂ warming. The combined error is then determined by adding a million draws from the two GEVs together and calculating the relevant quantiles of the result.

Data availability

The required statistics from runs of MAGICC and FaIR are available from Zenodo: <https://doi.org/10.5281/ZENODO.8332951> ref. 48.

Code availability

The original code used to generate the data is available from Zenodo with the data⁴⁸. Updates to the codebase are available from <https://github.com/Rlamboll/AR6CarbonBudgetCalc>.

References

46. Lamboll, R. D., Nicholls, Z. R. J., Kikstra, J. S., Meinshausen, M. & Rogelj, J. Silicone v1.0.0: an open-source Python package for inferring missing emissions data for climate change research. *Geosci. Model Dev.* **13**, 5259–5275 (2020).
47. Possolo, A., Merktas, C. & Bodnar, O. Asymmetrical uncertainties. *Metrologia* **56**, 045009 (2019).
48. Lamboll, R., Nicholls, Z. & Rogelj, J. Carbon budget calculator. Zenodo <https://doi.org/10.5281/ZENODO.8332951> (2023).

Acknowledgements

R.D.L. and J.R. received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 820829 (CONSTRAIN), and J.R. and Z.R.J.N. also under grant

agreement no. 101003536 (ESM2025). J.S.K. and E.B. acknowledge funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 821471 (ENGAGE) and funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 951542) (GENIE). C.J.S. was supported by a NERC/IIASA Collaborative Research Fellowship (NE/TO09381/1).

Author contributions

R.D.L., J.S.K. and J.R. conceived the experiments. R.D.L., Z.R.J.N. and C.J.S. wrote and executed code to collect data. R.D.L., E.B., J.S.K. and J.R. analysed the data. All authors contributed to writing the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41558-023-01848-5>.

Correspondence and requests for materials should be addressed to Robin D. Lamboll.

Peer review information *Nature Climate Change* thanks Nadine Mengis, Antti-Ilari Partanen and Benjamin Sanderson for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.