



This is a repository copy of *Electromagnetic self-force on a charged particle on Kerr spacetime: equatorial circular orbits*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/207772/>

Version: Published Version

Article:

Torres, T. and Dolan, S.R. orcid.org/0000-0002-4672-6523 (2022) Electromagnetic self-force on a charged particle on Kerr spacetime: equatorial circular orbits. *Physical Review D*, 106 (2). 024024. ISSN 2470-0010

<https://doi.org/10.1103/physrevd.106.024024>

© 2022 American Physical Society. Reproduced in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Electromagnetic self-force on a charged particle on Kerr spacetime: Equatorial circular orbits

Theo Torres^{*} and Sam R. Dolan[†]

Consortium for Fundamental Physics, School of Mathematics and Statistics, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, United Kingdom

 (Received 23 September 2020; revised 28 May 2022; accepted 8 July 2022; published 18 July 2022)

We calculate the self-force acting on a charged particle on a circular geodesic orbit in the equatorial plane of a rotating black hole. We show by direct calculation that the dissipative self-force balances with the sum of the flux radiated to infinity and through the black hole horizon. Prograde orbits are found to stimulate black hole superradiance, though we confirm that the condition for floating orbits cannot be met. We calculate the conservative component of the self-force by application of the mode sum regularization method, and we present a selection of numerical results. We obtain the leading-order coefficients in post-Newtonian expansions of the dissipative and conservative components of the self-force, using an analytical method and numerical fitting, respectively. The self-force on the innermost stable circular orbits of the Kerr spacetime is calculated, and comparisons are drawn between the electromagnetic and gravitational self forces.

DOI: [10.1103/PhysRevD.106.024024](https://doi.org/10.1103/PhysRevD.106.024024)

I. INTRODUCTION

The first detection of gravitational waves in 2015 heralded the arrival of gravitational wave astronomy as an observational science [1]. Two years on, the near-coincident detection of gravitational waves (GWs) and gamma rays from a binary neutron-star merger confirmed that GWs travel at the speed of light [2]. This dual detection highlighted the potential of *multimessenger astronomy*, as the host galaxy was quickly located by telescopes working in the electromagnetic spectrum, enabling a new measurement of the expansion rate of the universe to be made, that is, an independent determination of the Hubble parameter [3]. By 2021, ground-based detectors had reported 90 gravitational wave events, detailed in the transient catalogues (GWTCs) 1, 2 and 3 [4]. With each doubling of sensitivity, we can expect that number to increase by a factor of approximately eight.

The mathematical modeling of compact binary inspirals, and their gravitational wave signatures, has progressed steadily over recent decades. Three leading approaches to modeling the gravitational two-body problem give complementary information, namely, post-Newtonian expansions [5], self-force calculations [6], and the simulations of numerical relativity [7]. The essential inspiral process can be understood from the leading-order term in the post-Newtonian expansion. A pair of compact bodies of masses M_1 and M_2 , on quasicircular orbits about the center of mass, will radiate gravitational waves predominantly in the

quadrupole mode ($\ell = 2$) at twice the orbital frequency [8,9]. Consequently, the binary system loses energy, and the GW frequency increases with a characteristic chirp profile,

$$f(t) \approx \frac{5}{8\pi} \left(\frac{5GM}{c^3} \right)^{-5/8} (t_0 - t)^{-3/8}, \quad (1)$$

where f is the wave frequency, c is the speed of light, t_0 is the time of collision, and $\mathcal{M} = (M_1 M_2)^{3/5} / (M_1 + M_2)^{1/5}$ is the *chirp mass* [10]. The spectrogram of the gravitational wave signal from the binary neutron star inspiral was found to track this chirp profile remarkably closely over the last ~ 100 seconds before merger [11], despite the fact that, formally, Eq. (1) arises only from the leading-order term of a post-Newtonian expansion for the radiated flux [9].

A natural question arises: does the catalogue of GW events from inspirals put strong constraints on beyond-GR theories, such as Einstein-Maxwell-dilaton theory [12,13], or dark matter scenarios featuring exotic charges [14]? A more prosaic, but related, question is whether the compact bodies could carry significant *electromagnetic* charges. In both scenarios, the existence of a radiative dipole mode ($\ell = 1$) will, in principle, change the character of the inspiral. However, numerical simulations [15] suggest that the gravitational-wave chirps observed from comparable mass binaries are compatible with non-negligible charge-to-mass ratios (e.g., up to ~ 0.3 for GW150914, for example [15]).

In future decades, space-based observatories will detect low-frequency gravitational waves from systems involving supermassive black holes. A key target for LISA are so-called extreme mass-ratio inspirals (EMRIs) with $M_1 \gg M_2$.

^{*}t.torres-vicente@sheffield.ac.uk

[†]s.dolan@sheffield.ac.uk

The scenario in which the central black hole (M_1) is charged was recently investigated using black hole perturbation theory [16,17]. In this work, we consider the complementary case, in which the central black hole (M_1) is uncharged, but the orbiting body (M_2) carries an electromagnetic charge q . We shall approach this problem from the perspective that the orbiting body experiences an electromagnetic self-force, at order q^2 , that causes a radiation reaction upon its motion. First, let us review the roots of this idea in the birth of quantum theory.

It is well-known that classical field theory is unable to satisfactorily account for the observed stability of the hydrogen atom. In the “planetary” version of the Rutherford atomic model [18], a pointlike electron orbits the atomic nucleus. The centripetal acceleration of the charged electron generates electromagnetic (EM) radiation at the orbital frequency of $\sim 10^{15}$ Hz and, consequently, a radiation-reaction force acts upon the electron, causing the rapid collapse of the atom within 10^{-8} s. Invoking the Abraham-Lorentz [19,20] force law,

$$\mathbf{F} = \frac{2}{3} \frac{e^2}{4\pi\epsilon_0 c^3} \dot{\mathbf{a}}, \quad (2)$$

nonrelativistic classical theory¹ implies that a pointlike electron on a quasicircular inspiral trajectory will generate EM radiation with the following “chirp” profile [cf. Eq. (1)]

$$f(t) \approx \frac{1}{4\pi\alpha} \sqrt{\frac{c}{a_0}} (t_0 - t)^{-1/2}. \quad (3)$$

Here α is the fine-structure constant and a_0 is the Bohr radius (see Appendix A for a derivation).

There is no experimental support for collapsing atoms and/or EM chirps, of course. To the contrary, experiments with electric discharges from the 1850s onwards show that atoms emit EM radiation at certain discrete frequencies [22]. Tension between theory and experiment led to the introduction of the Bohr-Rutherford atomic model [23], and on to quantum theory itself. However, the idea of an orbit driven by a radiation-reaction force has reemerged as a useful concept on a very different scale in the universe.

In this article we consider the radiation-reaction process for a charged particle orbiting a black hole of mass M , rather than a charged nucleus. We shall assume that the length-scales of the particle, such as its Compton wavelength, are substantially smaller than the curvature scale, so that classical field theory provides an adequate framework. One might expect that, since the gravitational force and the Coulomb force both follow inverse square-laws in the weak-field, the radiation reaction process will proceed in a broadly

similar fashion, producing a chirp frequency which scales with $(t_0 - t)^{-1/2}$ while $v \ll c$ and $r \gg GM/c^2$. However, an important difference that cannot be overlooked is that the spacetime of a black hole is curved, not flat.

The first expression for an EM self-force on a weakly curved spacetime was obtained by DeWitt-Morette and DeWitt [24] in 1964. The self-force on a particle of charge q on a vacuum spacetime characterized by a Newtonian potential $\Phi_N = GM/c^2 r \ll 1$ is given by

$$\mathbf{F}_{\text{self}} \approx \frac{q^2}{4\pi\epsilon_0 c^3} \left(\frac{2}{3} \frac{d\mathbf{g}}{dt} + \frac{GMc}{r^3} \hat{\mathbf{r}} \right), \quad (4)$$

where $\mathbf{g} = -c^2 \nabla \Phi_N$ is the Newtonian gravitational field. The first term in parentheses in Eq. (4) is the standard Abraham-Lorentz force, which leads to the dissipation of orbital energy, and thus to an analogue of Eq. (3). The second term is a *conservative* correction to the Newtonian force $m\mathbf{g}$, which is *not* present in flat spacetime. Analogous equations were obtained for scalar and gravitational self-forces in weakly-curved spacetimes in Ref. [25].

To move beyond the Newtonian/weak-field context, we must acknowledge several key differences between a point mass in Newtonian theory and a black hole in general relativity. First, there exists an innermost stable circular orbit (ISCO), inside of which circular orbits cannot be sustained. Second, orbital velocities are sizable ($v/c \sim 0.4$ at the Schwarzschild ISCO), necessitating a fully relativistic description. Third, the issue of regularization is more subtle in a curved space-time, and Dirac’s time-reversal approach (“half-advanced-minus-retarded”) breaks down and requires modification [26–28].

The conservative component of the EM self-force leads to a shift in the orbital energy and angular momentum, and to a shift in the ISCO radius and frequency. The dissipative component of the EM self-force leads to orbital decay, and to the possibility of two interesting phenomena: floating orbits, and synchrotron radiation. The possibility of *floating orbits*—orbits which do not decay—arises due to *super-radiance*, which allows a particle on a corotating orbit to stimulate the release of energy and angular momentum from a rotating black hole [29–31]. The possibility of synchrotron radiation arises from the high velocities on ISCO orbits, leading to the beaming of radiation in the direction of motion [32,33].

In 1960, DeWitt and Brehme [26] derived an expression for the self-force on a point electric charge (see Eq. (1.33) in Ref. [34]) that consists of two parts: a local term which depends on the external force and the local Ricci tensor [35], and a *tail integral*, which encapsulates the effect of radiation emitted at earlier times that reaches the particle after interacting with the spacetime curvature. Thus, self-force in curved spacetime is nonlocal in time, since it depends on the past history of the motion of the particle, as well as its current state.

¹For a fully relativistic treatment, one would instead start with the Abraham-Lorentz-Dirac equation [21]; but note that $v/c < 0.01$ for a pointlike electron at the Bohr radius.

Calculating the tail integral in practice is a technical challenge (though see [36]); fortunately, there are equivalent formulations available, as described in the review articles [34,6] (see also Ref. [37]). Prominent among these is the *mode sum regularization* (MSR) method introduced by Barack and Ori [38], which has been applied by numerous authors [39–48] for efficient and accurate calculations of the self-force. Schematically, a regularized self-force $\mathcal{F}_\mu^{\text{reg}}$ is obtained by subtracting *regularization parameters* $\mathcal{F}_\mu^{[-1]}$, $\mathcal{F}_\mu^{[0]}$, etc., from the ℓ modes of a “bare” force:

$$\mathcal{F}_\mu^{\text{reg}} = \sum_{\ell=0}^{\infty} (\mathcal{F}_\mu^{\text{bare},\ell} - \mathcal{F}_\mu^{[-1]}(2\ell+1) - \mathcal{F}_\mu^{[0]} - \dots). \quad (5)$$

The regularization parameters are obtained from a local analysis of the symmetric-singular Detweiler-Whiting field [27]. Happily, regularization parameters for the EM field have already been calculated for the Schwarzschild black hole by Barack and Ori [49] and for the Kerr black hole by Heffernan, Wardell and Ottewill [50–52], and we make use of these here.

The MSR method is suited to cases where the field equations allow for a complete decomposition into modes in such a way as to reduce the problem to the solution of ordinary differential equations. Fortunately, the field equations for an EM field on Kerr spacetime fall into this class, as shown by Teukolsky [53–55], and the Faraday tensor $F_{\mu\nu}$ can be fully reconstructed from Maxwell scalars of spin-weight ± 1 that satisfy second-order ODEs [56,57].

The article is organized as follows. Section II describes the formulation of the calculation, covering the spacetime and its geodesic orbits (II A); Maxwell’s equations in the Teukolsky formalism (II B); the distributional source terms due to the particle (II C); the mode solutions (II D) and the special cases of static modes and the monopole (II E); the dissipative self-force and fluxes (II F); and the conservative self-force (II G) calculated by projecting from spin-weighted spheroidal harmonics to spherical harmonics (II G 2) and by mode sum regularization (II G 3). Section III describes the implementation, addressing numerical issues (III A) and the validation of the results (III B). Results are given in Sec. IV for the dissipative (IV A) and conservative (IV B) aspects of the self-force. We conclude with a discussion in Sec. V.

We employ units in which the physical constants G , c and $4\pi\epsilon_0$ are equal to unity. The spacetime signature is $\{-++++\}$.

II. FORMULATION

A. Spacetime and geodesic orbits

1. Spacetime

The Kerr solution with mass M and angular momentum $J = aM$ expressed in Boyer-Lindquist coordinates $\{t, r, \theta, \phi\}$ has the line element

$$\begin{aligned} ds^2 &\equiv g_{\mu\nu} dx^\mu dx^\nu \\ &= -\frac{\Delta}{\Sigma} (dt - a \sin^2 \theta d\phi)^2 + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2 \\ &\quad + \frac{\sin^2 \theta}{\Sigma} ((r^2 + a^2) d\phi - a dt)^2, \end{aligned} \quad (6)$$

where $\Sigma \equiv r^2 + a^2 \cos^2 \theta$ and $\Delta \equiv r^2 - 2Mr + a^2$. When the condition $a^2 \leq M^2$ is satisfied, the Kerr solution corresponds to a black hole spacetime with two distinct horizons: an internal (Cauchy) horizon at $r_- = M - \sqrt{M^2 - a^2}$ and an external (event) horizon at $r_+ = M + \sqrt{M^2 - a^2}$. The angular velocity of the event horizon is [58]

$$\Omega_h = \frac{a}{2Mr_+}. \quad (7)$$

The inverse metric $g^{\mu\nu}$ can be written in terms of a null basis $\{l^\mu, n^\mu, m^\mu, \bar{m}^\mu\}$, where the overline denotes the complex conjugate, as

$$g^{\mu\nu} = -2l^{(\mu} n^{\nu)} + 2m^{(\mu} \bar{m}^{\nu)} \quad (8)$$

$$= \frac{\Delta_r}{\Sigma} l_+^{(\mu} l_-^{\nu)} + \frac{1}{\Sigma} m_+^{(\mu} m_-^{\nu)}. \quad (9)$$

Here we employ the Kinnersley tetrad [59],

$$l^\mu = l_+^\mu, \quad n^\mu = -\frac{\Delta_r}{2\Sigma} l_-^\mu, \quad m^\mu = \frac{1}{\sqrt{2}(r+ia\cos\theta)} m_+^\mu, \quad (10)$$

written in terms of a non-normalized null basis

$$\begin{aligned} l_\pm^\mu &\equiv [\pm(r^2 + a^2)/\Delta, 1, 0, \pm a/\Delta], \\ m_\pm^\mu &\equiv [\pm ia \sin \theta, 0, 1, \pm i \csc \theta] = \bar{m}_\mp^\mu. \end{aligned} \quad (11)$$

The legs l_\pm^μ are aligned with the two principal null directions of the spacetime. The inner products of the tetrad l_\pm^μ and m_\pm^μ are

$$g_{\mu\nu} l_+^\mu l_-^\nu = \frac{2\Sigma}{\Delta}, \quad g_{\mu\nu} m_+^\mu m_-^\nu = 2\Sigma, \quad (12)$$

with all others zero.

2. Circular equatorial geodesic orbits

Let $x_p^\mu(\tau)$ denote the particle’s worldline, with tangent vector $u^\mu \equiv \frac{dx_p^\mu}{d\tau}$ satisfying $g_{\mu\nu} u^\mu u^\nu = -1$. In the absence of forces $x_p^\mu(\tau)$ is a geodesic, satisfying $u^\nu \nabla_\nu u^\mu = 0$. Geodesic orbits on the Kerr spacetime are characterized by three constants of motion: energy $E = -u_\mu \xi_{(t)}^\mu$, azimuthal angular momentum $L = u_\mu \xi_{(\phi)}^\mu$ and Carter constant $Q = Q^{\mu\nu} u_\mu u_\nu$, where $\xi_{(t)}^\mu = (\partial_t)^\mu$ and $\xi_{(\phi)}^\mu = (\partial_\phi)^\mu$ are

Killing vectors and $Q^{\mu\nu}$ is the Killing tensor. For a circular orbit in the equatorial plane at Boyer-Lindquist radius r_0 ,

$$E = \frac{1 - 2\nu^2 + \tilde{a}\nu^3}{\sqrt{1 - 3\nu^2 + 2\tilde{a}\nu^3}}, \quad L = r_0\nu \frac{1 - 2\tilde{a}\nu^3 + \tilde{a}^2\nu^4}{\sqrt{1 - 3\nu^2 + 2\tilde{a}\nu^3}}, \quad Q = 0, \quad (13)$$

where $\tilde{a} = a/M$ and $\nu = \sqrt{M/r_0}$. Explicitly, the equatorial circular geodesic orbit has $x_p^\mu(\tau) = [t(\tau), r_0, 0, \Omega t(\tau)]$ and $u^\mu = u^t[1, 0, 0, \Omega]$, where

$$\Omega = \frac{\nu^3}{M(1 + \tilde{a}\nu^3)}, \quad u^t = \frac{1 + \tilde{a}\nu^3}{\sqrt{1 - 3\nu^2 + 2\tilde{a}\nu^3}}, \quad (14)$$

We adopt the convention [60] that L and Ω are always positive and $a > 0$ ($a < 0$) for prograde (retrograde) orbits.

The innermost stable circular orbit (ISCO) is at the radius

$$r_{\text{isco}}/M = 3 + Z_2 \mp \sqrt{(3 - Z_1)(3 + Z_1 + 2Z_2)} \quad (15)$$

where $Z_1 = 1 + (1 - \tilde{a}^2)^{1/3}[(1 + \tilde{a})^{1/3} + (1 - \tilde{a})^{1/3}]$ and $Z_2 = \sqrt{3\tilde{a}^2 + Z_1^2}$ and the upper (lower) sign in Eq. (15) corresponds to prograde (retrograde) motion [61,62].

B. Maxwell's equations and the Teukolsky formalism

The electromagnetic field equations in their standard covariant form are

$$\nabla_\nu F^{\mu\nu} = 4\pi J^\mu, \quad \nabla_{[\mu} F_{\nu\sigma]} = 0, \quad (16)$$

where $F^{\mu\nu}$ is the Faraday tensor and J^μ is a vector field representing a four-current that is divergence-free ($\nabla_\mu J^\mu = 0$). It is convenient to introduce a complexified version of the Faraday tensor, $\mathbb{F}^{\mu\nu} = F^{\mu\nu} - i\tilde{F}^{\mu\nu}$, where $\tilde{}$ denotes the Hodge dual, i.e., $\tilde{F}^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\sigma\gamma}F_{\sigma\gamma}$. The complexified tensor is self-dual by virtue of the property $\tilde{\mathbb{F}}^{\mu\nu} = i\mathbb{F}^{\mu\nu}$. The field equations (16) then reduce to a single tensorial equation

$$\nabla_\nu \mathbb{F}^{\mu\nu} = 4\pi J^\mu. \quad (17)$$

The six degrees of freedom of $F^{\mu\nu}$ are encapsulated in 3 complex Maxwell scalars,

$$\phi_0 \equiv F_{\mu\nu}l^\mu m^\nu, \quad \phi_2 \equiv F_{\mu\nu}\bar{m}^\mu n^\nu, \quad \phi_1 \equiv \frac{1}{2}F_{\mu\nu}(l^\mu n^\nu - m^\mu \bar{m}^\nu), \quad (18)$$

and the self-dual Faraday tensor is specified in terms of Maxwell scalars by

$$\mathbb{F}^{\mu\nu} = 4(\phi_0\bar{m}^{[\mu}n^{\nu]} + \phi_2l^{[\mu}m^{\nu]} + \phi_1(n^{[\mu}l^{\nu]} - \bar{m}^{[\mu}m^{\nu]})). \quad (19)$$

For future reference, we introduce rescaled quantities:

$$\Phi_{+1} \equiv \phi_0 = \frac{1}{\sqrt{2}Q}l_+^\mu m_+^\nu F_{\mu\nu}, \quad \Phi_{-1} \equiv 2\bar{Q}^2\phi_2 = \frac{\Delta}{\sqrt{2}Q}l_-^\mu m_-^\nu F_{\mu\nu}, \quad (20)$$

where $q = r + ia \cos \theta$.

Projecting (17) onto a null tetrad aligned with the principal null directions leads to four equations in Newman-Penrose form [54]

$$(D - 2\rho)\phi_1 - (\bar{\delta} + \pi - 2\alpha)\phi_0 = -2\pi J_l, \quad (21a)$$

$$(\delta - 2\tau)\phi_1 - (\Delta + \mu - 2\gamma)\phi_0 = -2\pi J_m, \quad (21b)$$

$$(D - \rho + 2\epsilon)\phi_2 - (\bar{\delta} + 2\pi)\phi_1 = -2\pi J_{\bar{m}}, \quad (21c)$$

$$(\delta - \tau + 2\beta)\phi_2 - (\Delta + 2\mu)\phi_1 = -2\pi J_n, \quad (21d)$$

where $D = l^\mu\partial_\mu$, $\Delta = n^\mu\partial_\mu$, $\delta = m^\mu\partial_\mu$ are directional derivatives, and $J_l = l^\mu J_\mu$, $J_n = n^\mu J_\mu$, etc., are projections of the four-current, and α , ρ , τ , π etc. are the Newman-Penrose coefficients associated with the null tetrad.

In 1973, Teukolsky [54] showed that one can obtain a decoupled equation for ϕ_0 , and also for ϕ_2 , by exploiting a commutation relation between first-order operators. After inserting the Newman-Penrose quantities for the Kinnersley tetrad, viz. $\kappa = \sigma = \nu = \lambda = 0$,

$$\rho = -1/(r - ia \cos \theta), \quad \beta = -\rho^* \cot \theta / 2\sqrt{2}, \quad (22a)$$

$$\pi = ia\rho^2 \sin \theta / \sqrt{2}, \quad \alpha = \pi - \beta^*,$$

$$\tau = -ia\rho\rho^* \sin \theta / \sqrt{2}, \quad \mu = \rho^2\rho^* \Delta_r / 2, \quad (22b)$$

$$\gamma = \mu + \frac{1}{4}\rho\rho^* \Delta', \quad \epsilon = 0,$$

one arrives at a master equation, Eq. (4.7) in Ref. [54]. This may be cast into the form [63]

$$[(\nabla_\mu \pm \Gamma_\mu)(\nabla^\mu \pm \Gamma^\mu) - 4\psi_2]\Phi_{\pm 1} = 4\pi T_{\pm 1}, \quad (23)$$

where ∇_μ denotes the covariant derivative on the Kerr spacetime, and here the so-called ‘‘connection vector’’ [63] is

$$\Gamma^\mu \equiv \frac{1}{\Sigma} \left[\frac{M(r^2 - a^2)}{\Delta} - (r + ia \cos \theta), r - M, 0, \frac{a(r - M)}{\Delta} + i \frac{\cos \theta}{\sin^2 \theta} \right] \quad (24)$$

and $\psi_2 = M/(r - ia \cos \theta)^3$ is the only nonvanishing Weyl scalar for the Kerr spacetime in the Kinnersley tetrad. The source terms in Eq. (23) are

$$T_{+1} = J_0 \equiv (\delta - \beta - \bar{\alpha} - 2\tau + \bar{\pi})J_l - (D - \epsilon + \bar{\epsilon} - 2\rho - \bar{\rho})J_m, \quad (25)$$

$$\frac{1}{2\bar{Q}^2}T_{-1} = J_2 \equiv (\Delta + \gamma - \bar{\gamma} + 2\mu + \bar{\mu})J_{\bar{m}} - (\bar{\delta} + \alpha + \bar{\beta} + 2\pi - \bar{\nu})J_n. \quad (26)$$

Remarkably, Eq. (23) admits separable solutions. The solution can be constructed from a sum over modes, with each mode in the form

$$\Phi_{\pm 1} = R_{\pm 1}(r)S_{\pm 1}(\theta)e^{-i\omega t + im\phi}. \quad (27)$$

In the vacuum case ($J^\mu = 0$), inserting Eq. (27) into Eq. (23) leads to homogeneous Teukolsky equations in Chandrasekhar's form,

$$\begin{aligned} (\Delta\mathcal{D}^\dagger\mathcal{D} - 2i\omega r - \lambda)P_{-1} &= 0, \\ (\Delta\mathcal{D}\mathcal{D}^\dagger + 2i\omega r - \lambda)P_{+1} &= 0, \end{aligned} \quad (28a)$$

$$\begin{aligned} (\mathcal{L}\mathcal{L}_1^\dagger + 2a\omega \cos\theta + \lambda)S_{-1} &= 0, \\ (\mathcal{L}^\dagger\mathcal{L}_1 - 2a\omega \cos\theta + \lambda)S_{+1} &= 0, \end{aligned} \quad (28b)$$

where $P_{+1} = \Delta R_{+1}$, $P_{-1} = R_{-1}$ and λ is the separation constant for $s = -1$ [57]. Here we have made use of directional derivatives along $\{l^\mu, l^\mu, m^\mu, m^\mu\}$, denoted by $\{\mathcal{D}, \mathcal{D}^\dagger, \mathcal{L}^\dagger, \mathcal{L}\}$, where

$$\mathcal{D} \equiv l^\mu_+ \partial_\mu = \partial_r - \frac{iK}{\Delta}, \quad \mathcal{L}^\dagger \equiv m^\mu_+ \partial_\mu = \partial_\theta - Q, \quad (29a)$$

$$\mathcal{D}^\dagger \equiv l^\mu_- \partial_\mu = \partial_r + \frac{iK}{\Delta}, \quad \mathcal{L} \equiv m^\mu_- \partial_\mu = \partial_\theta + Q, \quad (29b)$$

with $K \equiv \omega(r^2 + a^2) - am$ and $Q \equiv m \csc\theta - a\omega \sin\theta$. We assume that these operators act only on quantities with harmonic time dependence $\chi \equiv e^{-i\omega t + im\phi}$. Furthermore, let $\mathcal{L}_n = \mathcal{L} + n \cot\theta$ and $\mathcal{L}_n^\dagger = \mathcal{L}^\dagger + n \cot\theta$.

For consistency these functions must also satisfy the Teukolsky-Starobinsky identities,

$$\Delta\mathcal{D}\mathcal{D}P_{-1} = \mathcal{B}P_{+1}, \quad \mathcal{L}^\dagger\mathcal{L}_1^\dagger S_{-1} = \mathcal{B}S_{+1}, \quad (30a)$$

$$\Delta\mathcal{D}^\dagger\mathcal{D}^\dagger P_{+1} = \mathcal{B}P_{-1}, \quad \mathcal{L}\mathcal{L}_1 S_{+1} = \mathcal{B}S_{-1}, \quad (30b)$$

where $\mathcal{B} \equiv \sqrt{\lambda^2 + 4am\omega - 4a^2\omega^2}$.

The modes of the Maxwell scalar of zero spin-weight, ϕ_1 , can be constructed by applying differential operators to the modes of ϕ_0 and ϕ_2 [56]. From Chap. 8 in Chandrasekhar [59],

$$\begin{aligned} \phi_1^{\ell m} &= \frac{\chi}{\sqrt{2}(r - ia \cos\theta)^2} [g_{+1}(r)\mathcal{L}_1 S_{+1}(\theta) \\ &\quad - ia f_{-1}(\theta)\mathcal{D}P_{-1}(r)] \end{aligned} \quad (31a)$$

$$\begin{aligned} &= -\frac{\chi}{\sqrt{2}(r - ia \cos\theta)^2} [g_{-1}(r)\mathcal{L}_1^\dagger S_{-1}(\theta) \\ &\quad - ia f_{+1}(\theta)\mathcal{D}^\dagger P_{+1}(r)] \end{aligned} \quad (31b)$$

where

$$\mathcal{B}g_{+1}(r) = (r\mathcal{D} - 1)P_{-1}, \quad (32a)$$

$$\mathcal{B}g_{-1}(r) = (r\mathcal{D}^\dagger - 1)P_{+1}, \quad (32b)$$

$$\mathcal{B}f_{+1}(\theta) = (\cos\theta\mathcal{L}_1^\dagger + \sin\theta)S_{-1}, \quad (32c)$$

$$\mathcal{B}f_{-1}(\theta) = (\cos\theta\mathcal{L}_1 + \sin\theta)S_{+1}. \quad (32d)$$

C. Source terms

For a pointlike charge q on a geodesic orbit, the four-current is

$$J^\mu = q \int u^\mu(\tau) \delta^4(x^\mu - x^\mu_p(\tau)) (-g(x))^{-1/2} d\tau, \quad (33)$$

$$= \frac{q\hat{U}^\mu}{r_0^2} \delta(r - r_0) \delta(\theta - \pi/2) \delta(\phi - \Omega t). \quad (34)$$

On the second line we have inserted the expressions in Sec. II A 2 to specialize to a circular geodesic orbit in the equatorial plane ($\theta = \pi/2$). Here $\hat{U}^\mu \equiv u^\mu/u^t = [1, 0, 0, \Omega]$, with Ω defined in Eq. (14); projecting onto the Kinnersley tetrad yields

$$\hat{U}^\mu l_\mu = -(1 - a\Omega) = \hat{U}^\mu n_\mu \frac{2r_0^2}{\Delta},$$

$$\hat{U}^\mu m_\mu = \frac{i}{\sqrt{2}r_0} ((r_0^2 + a^2)\Omega - a). \quad (35)$$

The first task is to compute the source terms J_0 and J_2 in Eqs. (25) and (26). Here we must handle the distributional terms with some care (see Appendix E), noting that whereas $f(x)\delta(x - x_0) = f(x_0)\delta(x - x_0)$, on the other hand

$$f(x)\delta'(x - x_0) = f(x_0)\delta'(x - x_0) - f'(x_0)\delta(x - x_0), \quad (36)$$

where $f(x)$ is any differentiable function and x_0 is a constant. Using

$$\delta(\phi - \Omega t) = \frac{1}{2\pi} \sum_{m=-\infty}^{\infty} \chi_m, \quad \chi_m \equiv e^{im(\phi - \Omega t)}, \quad (37)$$

and evaluating on the equatorial plane at $r = r_0$ after employing (36) leads to

$$\begin{aligned} \Sigma J_0 = & \frac{-q}{2\pi\sqrt{2}r_0} \sum_m \left[(1 - a\Omega) \left(\partial_\theta - m(1 - a\Omega) + \frac{ia}{r_0} \right) \right. \\ & \left. + i((r_0^2 + a^2)\Omega - a) \left(\partial_r - \frac{im((r_0^2 + a^2)\Omega - a)}{\Delta_0} + \frac{1}{r_0} \right) \right] \chi_m \delta(r - r_0) \delta\left(\theta - \frac{\pi}{2}\right). \end{aligned} \quad (38)$$

At this point we employ the orthonormality of the spin-weighted spheroidal harmonics,

$$\int S_{\pm 1}^{\ell m}(\theta) S_{\pm 1}^{\ell' m}(\theta) d(\cos \theta) = \frac{1}{2\pi} \delta_{\ell \ell'}, \quad (39)$$

to establish that

$$\delta\left(\theta - \frac{\pi}{2}\right) = 2\pi \sum_{\ell=1}^{\infty} S_{\pm 1}^{\ell m}\left(\frac{\pi}{2}\right) S_{\pm 1}^{\ell m}(\theta), \quad (40)$$

$$\delta'\left(\theta - \frac{\pi}{2}\right) = 2\pi \sum_{\ell=1}^{\infty} -S_{\pm 1}^{\ell m'}\left(\frac{\pi}{2}\right) S_{\pm 1}^{\ell m}(\theta). \quad (41)$$

Hence

$$\begin{aligned} \Sigma J_0 = & \frac{-q}{\sqrt{2}r_0} \sum_{\ell m} S_{\pm 1}^{\ell m}(\theta) \chi_m \left\{ i((r_0^2 + a^2)\Omega - a) S_{\pm 1}^{\ell m}\left(\frac{\pi}{2}\right) \delta'(r - r_0) + \right. \\ & \left. - (1 - a\Omega) S_{\pm 1}^{\ell m'}\left(\frac{\pi}{2}\right) \delta(r - r_0) + \left[ir_0\Omega + m \left(\frac{((r_0^2 + a^2)\Omega - a)^2}{\Delta_0} - (1 - a\Omega)^2 \right) \right] S_{\pm 1}^{\ell m}\left(\frac{\pi}{2}\right) \delta(r - r_0) \right\}. \end{aligned} \quad (42)$$

From the form of (42), we see that the master equation Eq. (23) admits a separable solution

$$\Phi_{\pm 1} = \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{\ell} R_{\pm 1}^{\ell m} S_{\pm 1}^{\ell m} \chi_m \quad (43)$$

where

$$(\Delta \mathcal{D} \mathcal{D}^\dagger + 2im\Omega r - \lambda) P_{\pm 1}^{\ell m} = \mathcal{S} (+iBS_+ \delta'(r - r_0) + \{(mA^{(r)} + iA^{(i)})S_+ + CS'_+\} \delta(r - r_0)), \quad (44a)$$

$$(\Delta \mathcal{D}^\dagger \mathcal{D} - 2im\Omega r - \lambda) P_{-1}^{\ell m} = \mathcal{S} (-iBS_- \delta'(r - r_0) + \{(mA^{(r)} - iA^{(i)})S_- - CS'_-\} \delta(r - r_0)), \quad (44b)$$

where $P_{\pm 1}^{\ell m} = \Delta R_{\pm 1}^{\ell m}$ and $P_{-1}^{\ell m} = R_{-1}^{\ell m}$, and $S_{\pm 1} = S_{\pm 1}^{\ell m}(\frac{\pi}{2})$ and $S'_{\pm 1} = S_{\pm 1}^{\ell m'}(\frac{\pi}{2})$, and

$$\mathcal{S} = -\frac{4\pi q}{\sqrt{2}r_0}, \quad (45a)$$

$$B = \Delta_0((r_0^2 + a^2)\Omega - a), \quad (45b)$$

$$A^{(r)} = r_0(r_0((r_0^2 + a^2)\Omega^2 - 1) + 2M(1 - a\Omega)^2), \quad (45c)$$

$$A^{(i)} = a^2(2M - r_0)\Omega + 2a(r_0 - M) - r_0^3\Omega, \quad (45d)$$

$$C = -\Delta_0(1 - a\Omega). \quad (45e)$$

D. Mode solutions

The source terms in Eqs. (44) are distributions with support at $r = r_0$ only. Hence solutions to the inhomogeneous equations may be constructed from solutions to the homogeneous equations in the standard manner. Let $P_{\pm 1}^{\ell m, h}$ and $P_{\pm 1}^{\ell m, \infty}$ be a pair of solutions to Eq. (28) that satisfy the physical boundary conditions, that is, let $P_{\pm 1}^{\ell m, h}$ be ingoing at the future horizon, and let $P_{\pm 1}^{\ell m, \infty}$ be outgoing at future infinity. The inhomogeneous solution takes the form

$$P_{\pm 1}^{\ell m}(r) = \alpha_{\pm 1}^{\infty} P_{\pm 1}^{\ell m, \infty}(r) \Theta(r - r_0) + \alpha_{\pm 1}^h P_{\pm 1}^{\ell m, h}(r) \Theta(r_0 - r), \quad (46)$$

where $\Theta(\cdot)$ is the Heaviside step function, and $\alpha_{\pm 1}^{\infty}$ and $\alpha_{\pm 1}^h$ are complex coefficients to be determined. Inserting (46) into (44) yields the matrix equation

$$\begin{pmatrix} \alpha_{\pm 1}^{\ell m, \infty} \\ \alpha_{\pm 1}^{\ell m, h} \end{pmatrix} = \frac{1}{\Delta_0 W_{\pm}} \begin{pmatrix} -(P_{\pm 1}^{\ell m, h})' & P_{\pm 1}^{\ell m, h} \\ -(P_{\pm 1}^{\ell m, \infty})' & P_{\pm 1}^{\ell m, \infty} \end{pmatrix} \Big|_{r=r_0} \begin{pmatrix} \mathfrak{B}_{\pm} \\ \mathfrak{A}_{\pm} \end{pmatrix}. \quad (47)$$

where

$$W_{\pm} = P_{\pm 1}^{\ell m, h} \frac{dP_{\pm 1}^{\ell m, \infty}}{dr} - P_{\pm 1}^{\ell m, \infty} \frac{dP_{\pm 1}^{\ell m, h}}{dr}, \quad (48a)$$

$$\mathfrak{B}_{\pm} = \pm i S B S_{\pm}, \quad (48b)$$

$$\mathfrak{A}_{\pm} = \mathcal{S}\{(mA^{(r)} \pm i\tilde{A}^{(i)})S_{\pm} \pm CS'_{\pm}\}. \quad (48c)$$

Here \mathcal{S} , B , $A^{(r)}$ and C are defined in Eq. (45), and $\tilde{A}^{(i)} = r_0 \Delta_0 \Omega$.

E. Static modes and the monopole

1. $m = 0$ homogeneous modes

The $m = 0$ modes are static ($\omega = 0$). In this case we employ the homogeneous modes

$$P_{\pm 1}^{\ell 0, h} = \Delta \partial_r P_{\ell}(z), \quad P_{\pm 1}^{\ell 0, \infty} = \Delta \partial_r Q_{\ell}(z), \quad (49)$$

where $P_{\ell}(\cdot)$ and $Q_{\ell}(\cdot)$ are Legendre functions with the branch cut on the real axis from $-\infty$ to $+1$, and $z \equiv \Delta_r / (r_+ - r_-)$. The Wronskian is

$$W_{\pm} \equiv P_{\pm 1}^{\ell 0, h} \frac{dP_{\pm 1}^{\ell 0, \infty}}{dr} - P_{\pm 1}^{\ell 0, \infty} \frac{dP_{\pm 1}^{\ell 0, h}}{dr} = \frac{1}{2} (r_+ - r_-) \ell(\ell + 1). \quad (50)$$

The angular functions are

$$S_{\pm 1}^{\ell 0}(\theta) = \mp \sqrt{\frac{2\ell + 1}{4\pi\ell(\ell + 1)}} \frac{d}{d\theta} P_{\ell}(\cos\theta), \quad (51)$$

such that the normalization condition (39) holds.

2. Monopole mode

To complete the solution, we must now add ‘‘by hand’’ a nonradiative monopole mode which is responsible for the q/r part of the electric field far from the black hole.

The homogeneous vector potential

$$A_{(0)}^{\mu} \equiv \frac{qr}{2\Sigma} (l_+^{\mu} - l_-^{\mu}), \quad (52)$$

in Lorenz gauge ($\nabla_{\mu} A_{(0)}^{\mu} = 0$) generates a homogeneous Faraday tensor $F_{(0)}^{\mu\nu} = \nabla^{\mu} A^{\nu} - \nabla^{\nu} A^{\mu}$ that satisfies the vacuum equation $\nabla_{\nu} F_{(0)}^{\mu\nu} = 0$. It has the key properties that

$$F_{(0)}^{ir} = \frac{q}{r^2} + O(r^{-3}), \quad (53)$$

in the far-field and

$$\frac{1}{2} \int F_{(0)}^{\mu\nu} dS_{\mu\nu} = 4\pi q, \quad (54)$$

where the two-surface integral is taken over any ‘‘sphere’’ of constant Boyer-Lindquist coordinate r , or any closed surface enclosing the horizon. It is quick to verify that the Maxwell scalars ϕ_0 and ϕ_2 (but not ϕ_1) associated with the homogeneous solution are zero.

The inhomogeneous monopole mode,

$$F_{\text{mono}}^{\mu\nu} = \Theta(r - r_0) F_{(0)}^{\mu\nu}, \quad (55)$$

does *not* satisfy the vacuum equation; instead, $\nabla_{\nu} \mathbb{F}_{\text{mono}}^{\mu\nu} = 4\pi \mathbb{J}_{\text{mono}}^{\mu}$ where $\mathbb{F}_{\text{mono}}^{\mu\nu} \equiv F_{\text{mono}}^{\mu\nu} - i\tilde{F}_{\text{mono}}^{\mu\nu}$ and it is straightforward to show that

$$\mathbb{J}_{\text{mono}}^{\mu} = \frac{q}{4\pi} \frac{\Delta}{2\Sigma} \frac{1}{(r - ia \cos\theta)^2} (l_+^{\mu} - l_-^{\mu}) \delta(r - r_0). \quad (56)$$

Note that $\mathbb{J}_{\text{mono}}^{\mu}$ associated with the step in the monopole mode is not restricted to the particle worldline, but instead has support on the sphere at $r = r_0$. Although $\mathbb{J}_{\text{mono}}^{\mu}$ itself is not zero, a short calculation shows that there are no additional source terms for the Teukolsky equation (23), that is, $J_0^{\text{mono}} = J_2^{\text{mono}} = 0$. In other words, the inhomogeneous monopole is associated with a step in ϕ_1 , the Maxwell scalar of spin-weight zero, only.

The inhomogeneous monopole mode makes a contribution to the radial component of the self-force of

$$\begin{aligned}\mathcal{F}_r^{\text{mono}} &= q^2 u^t \frac{(r_0^2 - a^2 \cos^2 \theta)(1 - a\Omega \sin^2 \theta)}{\Sigma^2} \\ &\equiv \sum_{\ell=0}^{\infty} \mathcal{F}_r^{\ell, \text{mono}} Y_0^{\ell 0}(\theta).\end{aligned}\quad (57)$$

Evaluating at $\theta = \pi/2$ yields $\mathcal{F}_r^{\text{mono}} = q^2 u^t (1 - a\Omega)/r_0^2$.

F. Dissipative force and fluxes

1. Dissipative component of the self-force

The dissipative components of the self-force are the t and ϕ components of $\mathcal{F}_\mu \equiv qF_{\mu\nu}u^\nu$. From the symmetry of the Faraday tensor, it is straightforward to see that $\mathcal{F}_t = qF_{t\phi}\Omega u^t = -\mathcal{F}_\phi\Omega$ and in the following we will focus on the t component of the self-force. The $(t\phi)$ component of the Faraday tensor can be expressed in terms of the Maxwell scalars as:

$$\begin{aligned}F_{t\phi} &= \sqrt{2}\text{Re} \left[i \sin \theta (r - ia \cos \theta) \phi_2 \right. \\ &\quad \left. + \frac{i\Delta \sin \theta}{2\Sigma} (r + ia \cos \theta) \phi_0 \right].\end{aligned}\quad (58)$$

Evaluating the force on the particle's worldline, i.e., at $r = r_0$ and $\theta = \pi/2$, yields

$$\begin{aligned}\mathcal{F}_t &= \sqrt{2}qu^t\Omega\text{Re} \left[ir_0\phi_2 + \frac{i\Delta_0}{2r_0}\phi_0 \right] \\ &= \frac{q\Omega u^t}{\sqrt{2}r_0} \sum_{\ell m} \text{Re} \left[iP_{-1}^{\ell m}(r_0)S_{-1}^{\ell m}\left(\frac{\pi}{2}\right) + iP_{+1}^{\ell m}(r_0)S_{+1}^{\ell m}\left(\frac{\pi}{2}\right) \right] \\ &= \frac{q\Omega u^t}{\sqrt{2}r_0} \sum_{\ell m} \text{Re} \left[i((-1)^{\ell+m}P_{-1}^{\ell m}(r_0) + P_{+1}^{\ell m}(r_0))S_{+1}^{\ell m}\left(\frac{\pi}{2}\right) \right],\end{aligned}\quad (59)$$

where we have used the fact that $S_{-1}^{\ell m}(\frac{\pi}{2}) = (-1)^{l+m}S_{+1}^{\ell m}(\frac{\pi}{2})$.

2. Energy flux

For an electromagnetic field given by a Faraday tensor $F_{\mu\nu}$ with energy-momentum $T^{\mu\nu} = F^{\mu\alpha}F^{\nu\beta}g_{\alpha\beta} - \frac{1}{4}F^{\alpha\beta}F_{\alpha\beta}g^{\mu\nu}$, and a Killing vector K^μ , one can construct a current:

$$Y^\mu = T^{\mu\nu}K_\nu.\quad (62)$$

In vacuum, this current is divergence-free but in the presence of a source, which is the case of interest here, the current satisfies the following continuity equation:

$$\nabla_\mu Y^\mu = F^{\mu\nu}K_\mu J_\nu = \frac{\mathcal{F}_\mu K^\mu}{r_0^2 u^t} \delta(r - r_0) \delta\left(\theta - \frac{\pi}{2}\right) \delta(\phi - \Omega t).\quad (63)$$

Using Gauss' theorem

$$\int_V \nabla_\mu Y^\mu \sqrt{-g} d^4x = \int_{\partial V} Y^\mu d\Sigma_\mu\quad (64)$$

where V is a space-time volume with boundary ∂V that spans from the horizon to infinity, we can relate the force at the particle to the fluxes through the boundary. Since the system is stationary, only the fluxes at infinity and through the horizon contribute to the total flux (see Appendix B):

$$\frac{\mathcal{F}_a K^a}{u^t} = \Phi_\infty^K + \Phi_h^K,\quad (65)$$

where the superscript K correspond to the choice of Killing vector. As mentioned the link between the t and ϕ component of the force is trivial and we focus on the time component of the force which correspond to the choice $K^a = [1, 0, 0, 0]$. In the following we will drop the superscript K and keep in mind that we are considering the energy flux. In Appendix B we derive the expression for the energy flux at infinity and through the horizon in terms of the α coefficients defined in Eq. (46). Explicitly, the energy flux at infinity is

$$\Phi_\infty = \frac{1}{8\pi} \sum_{\ell m} |\alpha_{-1}^{\ell m, \infty}|^2,\quad (66)$$

and through the horizon,

$$\Phi_h = \frac{1}{8\pi} \sum_{\ell m} \frac{\omega}{2Mr_+ \tilde{\omega}} |\alpha_{+1}^{\ell m, h}|^2,\quad (67)$$

with $\tilde{\omega} = \omega - m\Omega_h$ and Ω_h as defined in Eq. (7).

G. Conservative force and regularization

1. Conservative component of the self-force

We compute here the conservative component of the self-force, i.e., \mathcal{F}_r , in terms of the Maxwell scalars. From the definition of the force, we have:

$$\mathcal{F}_r = qF_{r\mu}u^\mu = qu^t(F_{rt} + F_{r\phi}\Omega).\quad (68)$$

Using the expression of the Faraday tensor in terms of the Maxwell scalars,

$$F_{\mu\nu} = 2[\phi_2 l_{[\mu} m_{\nu]} + \phi_0 \bar{m}_{[\mu} n_{\nu]} + \phi_1 (n_{[\mu} l_{\nu]} + m_{[\mu} \bar{m}_{\nu]})] + \text{c.c.},\quad (69)$$

we get that

$$\frac{\mathcal{F}_r}{qu^t} = \sqrt{2}((a^2 + r^2)\Omega - a) \sin\theta \left[-\frac{i\phi_0}{4(r - ia \cos\theta)} + \frac{i\phi_2(r - ia \cos\theta)}{2\Delta} \right] + (1 - a\Omega \sin^2\theta)\phi_1 + \text{c.c.} \quad (70)$$

Inserting the mode decompositions (27) and (31a) and evaluating at $\phi = \Omega t$ yields

$$\begin{aligned} \mathcal{F}_r = qu^t \sum_{lm} \frac{\sqrt{2}((r^2 + a^2)\Omega - a)}{4\Delta(r - ia \cos\theta)} \sin(\theta) \left[-iP_{+1}^{lm} S_{+1}^{lm} + iP_{-1}^{lm} S_{-1}^{lm} \right] \\ + (1 - a\Omega \sin^2\theta) \frac{g_{+1}^{lm} \mathcal{L}_1 S_{+1}^{lm} - ia f_{-1}^{lm} \mathcal{D}P_{-1}^{lm}}{\sqrt{2}(r - ia \cos\theta)^2} + \text{c.c.} \end{aligned} \quad (71)$$

2. Projection onto scalar harmonics

Before the mode sum regularization procedure can be applied, it is necessary to project the spin-weighted spheroidal harmonics onto the scalar spherical harmonics. Using the results of Appendix C,

$$\mathcal{F}_r = qu^t \sum_{l,m,\hat{l},\hat{l}} \left\{ \frac{\sqrt{2}((r^2 + a^2)\Omega - a)}{4\Delta(r - ia \cos\theta)} \left[-iP_{+1}^{lm} C_{lm\hat{l}}^{+1} + iP_{-1}^{lm} C_{lm\hat{l}}^{-1} \right] \right. \quad (72)$$

$$\left. + (1 - a\Omega \sin^2\theta) \frac{\mathcal{B} g_{+1}^{lm} C_{lm\hat{l}}^{\mathcal{L}} - ia [C_{lm\hat{l}}^{\mathcal{L}} \cos\theta + C_{lm\hat{l}}^{+1}] \mathcal{D}P_{-1}^{lm}}{\sqrt{2}\mathcal{B}(r - ia \cos\theta)^2} \right\} Y_0^{\hat{l}m} + \text{c.c.} \quad (73)$$

with

$$C_{lm\hat{l}}^{+1} = (b_{+1}^m)_i^l (A_{+1}^m)_i^{\hat{l}} \quad (74a)$$

$$C_{lm\hat{l}}^{-1} = (b_{-1}^m)_i^l (A_{-1}^m)_i^{\hat{l}} \quad (74b)$$

$$C_{lm\hat{l}}^{\mathcal{L}} = (b_{+1}^m)_i^l \left[\sqrt{\hat{l}(\hat{l} + 1)} \delta_i^{\hat{l}} - am\Omega (A_{+1}^m)_i^{\hat{l}} \right]. \quad (74c)$$

Expanding (73) in $z = \cos\theta$, we have

$$\mathcal{F}_r = qu^t \sum_{\ell m} [{}_0\mathcal{F}_r^{\ell m} + {}_1f_r^{\ell m} z + {}_2f_r^{\ell m} z^2 + o(z^3)] Y_0^{\ell m}, \quad (75)$$

with

$$\begin{aligned} {}_0\mathcal{F}_r^{\ell m} = \sum_{\hat{l}} \frac{((r_0^2 + a^2)\Omega - a)}{\sqrt{2}\Delta_0 r_0} \text{Im} [P_{+1}^{lm} C_{lm\hat{l}}^{+1} - P_{-1}^{lm} C_{lm\hat{l}}^{-1}] \\ + \frac{\sqrt{2}(1 - a\Omega)}{r_0^2} \text{Re} \left[g_{+1}^{lm} C_{lm\hat{l}}^{\mathcal{L}} - \frac{ia C_{lm\hat{l}}^{+1} \mathcal{D}P_{-1}^{lm}}{\mathcal{B}} \right], \end{aligned} \quad (76)$$

and

$$\begin{aligned} {}_1f_r^{\ell m} = a \sum_{\hat{l}} \frac{((r_0^2 + a^2)\Omega - a)}{\sqrt{2}\Delta_0 r_0^2} \text{Re} [P_{+1}^{lm} C_{lm\hat{l}}^{+1} - P_{-1}^{lm} C_{lm\hat{l}}^{-1}] \\ - \frac{2\sqrt{2}(1 - a\Omega)}{r_0^3} \text{Im} [g_{+1}^{lm} C_{lm\hat{l}}^{\mathcal{L}} - \frac{ia}{\mathcal{B}} C_{lm\hat{l}}^{+1} \mathcal{D}P_{-1}^{lm}] \\ + \frac{2(1 - a\Omega)}{\sqrt{2}r_0^2} \text{Im} \left[\frac{C_{lm\hat{l}}^{\mathcal{L}} \mathcal{D}P_{-1}^{lm}}{\mathcal{B}} \right], \end{aligned} \quad (77)$$

$$\begin{aligned} {}_2f_r^{\ell m} = a^2 \sum_{\hat{l}} \frac{-((r_0^2 + a^2)\Omega - a)}{\sqrt{2}\Delta_0 r_0^3} \text{Im} [P_{+1}^{lm} C_{lm\hat{l}}^{+1} - P_{-1}^{lm} C_{lm\hat{l}}^{-1}] \\ - \frac{3\sqrt{2}(1 - a\Omega)}{r_0^4} \text{Re} \left[g_{+1}^{lm} C_{lm\hat{l}}^{\mathcal{L}} - \frac{ia C_{lm\hat{l}}^{+1} \mathcal{D}P_{-1}^{lm}}{\mathcal{B}} \right] \\ + \frac{2\sqrt{2}(1 - a\Omega)}{r_0^3} \text{Re} \left[\frac{C_{lm\hat{l}}^{\mathcal{L}} \mathcal{D}P_{-1}^{lm}}{\mathcal{B}} \right] \\ + a\Omega \sum_{\hat{l}} \frac{\sqrt{2}}{r_0^2} \text{Re} \left[g_{+1}^{lm} C_{lm\hat{l}}^{\mathcal{L}} - \frac{ia C_{lm\hat{l}}^{+1} \mathcal{D}P_{-1}^{lm}}{\mathcal{B}} \right], \end{aligned} \quad (78)$$

Finally, expanding $zY_0^{\ell m}$ and $z^2Y_0^{\ell m}$ using

$$\cos\theta Y_0^{\ell m} = \sum_{\ell_1} {}_1B_{\ell_1}^{\ell m} Y_0^{\ell_1 m}, \quad (79a)$$

$$\cos^2\theta Y_0^{\ell m} = \sum_{\ell_2} {}_2B_{\ell_2}^{\ell m} Y_0^{\ell_2 m}, \quad (79b)$$

where

$${}_1B_{\ell_1}^{\ell m} = (-1)^m \sqrt{(2\ell+1)(2\ell_1+1)} \times \begin{pmatrix} 1 & \ell & \ell_1 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & \ell & \ell_1 \\ 0 & m & -m \end{pmatrix} \quad (80)$$

$${}_2B_{\ell_2}^{\ell m} = (-1)^m \frac{2\sqrt{(2\ell+1)(2\ell_2+1)}}{3} \times \begin{pmatrix} 2 & \ell & \ell_2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 & \ell & \ell_2 \\ 0 & m & -m \end{pmatrix} + \frac{1}{3} \delta_{\ell, \ell_2} \quad (81)$$

leads to

$$\mathcal{F}_r = qu^t \sum_{\ell m} [{}_0\mathcal{F}_r^{\ell m} + {}_1\mathcal{F}_r^{\ell m} + {}_2\mathcal{F}_r^{\ell m} + o(z^3)] Y_0^{\ell m}, \quad (82a)$$

$$= \sum_{\ell=0}^{\infty} \mathcal{F}_r^{\ell} \quad (82b)$$

with

$${}_1\mathcal{F}_r^{\ell m} = \sum_{\ell_1} {}_1f_r^{\ell_1 m} {}_1B_{\ell_1}^{\ell m} \quad (83a)$$

$${}_2\mathcal{F}_r^{\ell m} = \sum_{\ell_2} {}_1f_r^{\ell_2 m} {}_2B_{\ell_2}^{\ell m}. \quad (83b)$$

3. Mode sum regularization

The regularization procedure is based on the subtraction of an appropriate singular component from the retarded field, in order to leave a finite regular field that is solely responsible for the self-force. The subtracted component must have the same singular structure as the retarded field in the vicinity of the particle, and must be sufficiently symmetric as to not contribute to the self-force (or at least, not in such a way that cannot be easily corrected for). Detweiler and Whiting identified an appropriate choice of the singular (S) field, based on a Green's function decomposition [27]. Subtracting this singular field is equivalent to regularizing at the level of the ℓ -mode sum [6,34,38–48].

In the electromagnetic case, Heffernan *et al.* [50–52] (see also Haas [64,65]) showed that subtracting the S field leads to a regularized force $\mathcal{F}_\mu^{\text{reg}}$ with a radial component in the form

$$\mathcal{F}_r^{\text{reg}} = \sum_{\ell=0}^{\infty} \mathcal{F}_r^{\text{reg}[n]\ell}, \quad \mathcal{F}_r^{\text{reg}[n]\ell} \equiv \mathcal{F}_r^{\ell} - \mathcal{F}_r^{[n]\ell}, \quad (84)$$

where $[n]$ denotes the order of the local expansion of the S field, and

$$\mathcal{F}_r^{[n]\ell} = (2\ell+1)\mathcal{F}_r^{[-1]} + \mathcal{F}_r^{[0]} + \frac{\mathcal{F}_r^{[2]}}{(2\ell-1)(2\ell+3)} + \dots + \mathcal{F}_r^{[n]}\mathcal{G}_{[n]}(\ell). \quad (85)$$

Here $n \geq 0$ is an even integer denoting the order, and $\mathcal{G}_{[n]}(\ell) \equiv 1/(2\ell+1-n)(2\ell+3-n)\dots(2\ell+1+n)$ is

defined for $n > 0$ such that $\sum_{\ell=0}^{\infty} \mathcal{G}_{[n]}(\ell) = 0$. Explicit expressions for the mode sum regularization parameters $\mathcal{F}_r^{[-1]}$, $\mathcal{F}_r^{[0]}$ and $\mathcal{F}_r^{[2]}$ are given in Eq. (2.54), (2.56), and (2.59) of Ref. [51] for the Kerr case, and $\mathcal{F}_r^{[4]}$ is given in Eq. (5.52) of Ref. [50] for the Schwarzschild case.

The regularized force in Eq. (85) should include the monopole piece given in Eq. (57).

III. IMPLEMENTATION

A. Numerics

1. Homogeneous solution to the Teukolsky equations

In order to compute the components of the self-force, we need to evaluate radial Teukolsky functions $P_{\pm 1}^{lm}(r)$ and spin-weighted spheroidal harmonics $S_{\pm 1}^{lm}(\theta)$ at the particle's location, that is $r = r_0$ and $\theta = \pi/2$. To do so, we use the BlackHolePerturbation toolkit [66]. The angular functions are computed using the *SpinWeightedSpheroidalHarmonics* package and the radial functions are computed with the *Teukolsky* package of the toolkit. The Teukolsky package implements the Mano-Suzuki-Takasugi (MST) method [67,68] to compute the homogeneous solution of the Teukolsky equations.

2. High- l tail contribution

Our approach to compute the self-force requires us to sum over spin-weighted spheroidal modes or scalar spherical modes. Ideally one would sum an infinite number of modes but in practice we can only compute a finite number of components, up to ℓ_{max} . In the case of the dissipative components of the self-force, the magnitude of the terms to be summed over decays exponentially, as can be seen in Fig. 4, and therefore the error from truncating the sum is negligible. However, for the regularized conservative part of the self-force, the terms in the sum decay as an inverse power of $L = \ell + 1/2$ instead of an exponential, and the associated error from neglecting the higher modes is sizable. To reduce this error, we estimate the contribution coming from the $\ell > \ell_{\text{max}}$ modes following the standard approach of [39] which we outline below.

In the large- ℓ regime, the modes of the regularized force in Eq. (84) are approximately

$$\mathcal{F}_r^{\text{reg}[n]\ell} \approx \frac{D_n}{L^n}, \quad (86)$$

where n denotes the regularization order (with $n = 6$ in the Schwarzschild case and $n = 4$ in the Kerr case) and D_n is a numerical coefficient to be determined by fitting to the high- ℓ modes. Figure 5 shows that Eq. (86) is a reasonable approximation for high values of ℓ . The contribution of the high- ℓ modes is then approximately

$$\sum_{\ell=\ell_{\max}+1}^{\infty} \mathcal{F}_r^{\text{reg}[n]\ell} \approx \sum_{\ell=\ell_{\max}+1}^{\infty} \frac{D_n}{L^n} = D_n \zeta(n, \ell_{\max} + 1), \quad (87)$$

where $\zeta(s, a)$ is the Hurwitz Zeta function.

3. Projection

In order to apply the mode-sum regularization procedure, we need to project the force onto the scalar spherical harmonics basis. The original quantities in the spin-weighted spheroidal harmonics (associated to the index l) are first projected onto the spin-weighted spherical harmonics (associated with the index \hat{l}) which are then expanded onto scalar spherical harmonics (associated with the index ℓ). For the subdominant terms, which are proportional to $\cos\theta$ and $\cos^2\theta$, one extra projection is needed (associated with the index ℓ_1 and ℓ_2). Due to the presence of the 3j-symbols, and their association to spin-weighted or scalar quantities, the summation indices satisfy

$$\ell - 2 \leq \ell_2 \leq \ell + 2 \quad (88)$$

$$\ell - 1 \leq \ell_1 \leq \ell + 1 \quad (89)$$

$$\ell - (1 + n) \leq \hat{l} \leq \ell + (1 + n) \quad (90)$$

$$|m| \leq l, \hat{l}, \ell, \ell_1, \ell_2 \quad (91)$$

where $n = 0, 1, 2$ when computing the dominant, subdominant or subsubdominant term.

Since one spin-weighted spheroidal mode couples to several scalar spherical modes, we first compute all spin-weighted spheroidal modes separately and then perform the sums. We start by summing over ℓ_1 or ℓ_2 if we are computing the subdominant contributions at fixed m , \hat{l} and l . We then sum over m modes with fixed \hat{l} and l and then we sum over \hat{l} modes at fixed l . All these sums performed at this point are finite and can be performed for any value of l . Finally we sum over l which in principle can take any nonzero integer values. In practice however, we sum over a finite number of l modes and estimate the contribution of the higher l as described above.

B. Validation

In order to validate our numerical code when computing the energy fluxes at infinity and through the horizon, we compare the total flux with the dissipative component of the self-force computed using (61). We check that the two quantities agree up to numerical accuracy according to (65). Furthermore, each flux is computed at $r = r_0^+$ and $r = r_0^-$ using different solutions to the homogeneous Teukolsky equations. We verify that the two fluxes obtained agree to numerical accuracy, meaning that our dissipative component of the self-force is continuous across the particle.

In the case of the conservative piece of the self-force, we do not have a conservation law to support our numerical code. To validate our numerical approach in this case, we first verify that the radial component of the self-force is continuous across the particle as in the conservative component case. We note that while \mathcal{F}_r is continuous across the particle, up to the expected precision, each spherical harmonic component \mathcal{F}_r^ℓ is discontinuous (for $a \neq 0$). We also observe that the sum of the even (odd) ℓ modes are independently continuous across the particle. Both features are likely due to the fact that we are only using a finite number of terms when expanding around $\cos\theta \approx 0$.

We observe that the bare modes, \mathcal{F}_r^ℓ , are well regularized using the regularization parameters found in the literature [50,51]. Finally, our result for the conservative self-force in the Schwarzschild case agrees with the results of Haas [64] (see Fig. 7).

IV. RESULTS

Below we present a selection of numerical results for the self-force. Where a dimensionless value is stated, e.g., $\tilde{\mathcal{F}}_r$, the physical value should be inferred by reinstating the dimensionful constants, e.g., $\mathcal{F}_r = (q^2/4\pi\epsilon_0)(c^2/GM)^2 \tilde{\mathcal{F}}_r$.

A. Dissipative effects

1. Total fluxes

Figure 1 shows the total energy flux Φ for a charged particle on a circular orbit about a black hole, as a function of orbital radius. The total flux is related to the self-force component \mathcal{F}_t by Eq. (65). In the large- r_0 limit, the flux approaches an asymptotic value of Φ_{Newt} , where

$$\Phi_{\text{Newt}} = \frac{2}{3} \beta \tilde{r}_0^{-4}, \quad \beta \equiv \frac{q^2 c^5}{4\pi\epsilon_0 G^2 M^2} \quad (92)$$

and

$$\tilde{r}_0 \equiv r_0 / (GM/c^2). \quad (93)$$

In Appendix A, it is shown that Φ_{Newt} results from combining Keplerian orbits with the Abraham-Lorentz force (2). In the discussion below, we omit the dimensionful factor β .

By fitting the numerical results in the weak-field region ($r_0 \gg M$), we infer that, for the flux at infinity, $\Phi_\infty \approx \Phi_{\text{Newt}}$ at leading order, with a linear-in- a contribution of $-\frac{8}{3} \tilde{a} \tilde{r}_0^{-11/2}$ at leading order (where $\tilde{a} = a/M$). For the horizon flux, we infer that $\Phi_h \approx \frac{8}{3} \tilde{r}_0^{-7}$ at leading order for the Schwarzschild case, with a linear-in- a contribution of $-\frac{2}{3} \tilde{a} \tilde{r}_0^{-11/2}$ at leading order in the Kerr case. Note that, for the horizon flux, the Kerr term begins at a *lower* order in the expansion in $1/\tilde{r}_0$ than the Schwarzschild term.

Alternatively, an expansion of the fluxes in inverse powers of $1/\tilde{r}_0$ can be obtained using the Mano-Suzuki-Takasugi (MST) formalism [67] by applying the approach of Kavanagh *et al.* [69,70] to our Eqs. (66) and (67). At leading orders, we find that the dipole flux is $\Phi_\infty^{(\ell=1)} = \frac{2}{3}\tilde{r}_0^{-4} - \frac{44}{15}\tilde{r}_0^{-5}$ and the quadrupole flux is $\Phi_\infty^{(\ell=2)} = \frac{8}{5}\tilde{r}_0^{-5}$, which sum to $\Phi_\infty = \frac{2}{3}\tilde{r}_0^{-4} - \frac{4}{3}\tilde{r}_0^{-5}$ through next-to-leading order. This expansion is consistent with the numerical data presented in Fig. 1.

Figure 2 shows the ratio of the flux through the horizon to the flux radiated away to infinity, for the three types of field (scalar, electromagnetic and gravitational). The scalar and electromagnetic cases are qualitatively similar, with radiation emitted principally in the dipole ($\ell = 1$) modes. For particles that are orbiting in the same sense as the black hole spin, superradiance can lead to a significant extraction of energy from the horizon. For $a = 0.99M$, the energy extracted from the hole is up to $\sim 26.5\%$ of that radiated away in the EM case, and up to $\sim 22.3\%$ in the scalar-field case. Since this ratio falls below the threshold for balance (100%), there are no floating orbits. In the gravitational case, radiation is emitted principally in the quadrupole ($\ell = 2$) modes, and the maximum ratio is smaller ($\sim 8.7\%$ for $a = 0.99M$). Again, there are no floating orbits.




FIG. 1. The radiated flux for an electromagnetically charged particle on a circular orbit at radius $r = r_0$ around a Kerr black hole of spin a . Here the total flux $\Phi_{\text{tot}} = \Phi_\infty + \Phi_h$ has been scaled by the Newtonian value Φ_{Newt} (see Eq. (92)). The solid lines correspond to prograde orbits ($a > 0$), while the dotted lines correspond to retrograde orbits ($a < 0$), and the color of the lines gives the magnitude of a . In each case, the minimum radius is the innermost stable circular orbit.

In the gravitational case, these results are consistent with those previously presented by Kapadia, Kennefick and Glampedakis [31].

Figure 3 shows the ratio of fluxes Φ_h/Φ_∞ for a particle on the innermost stable circular orbit (ISCO), as a function of the spin of the black hole. The ratio changes sign at $a = a_c \approx 0.359403M$. This is the value of a at which the angular frequency of the ISCO orbit [see Eq. (15)] matches the angular frequency of the event horizon Ω_h . For $a > a_c$, the (prograde) horizon frequency exceeds the orbital frequency. In this case, the electromagnetic field slows the rotation of the black hole, generating superradiance, leading to an extraction of flux from the event horizon and $\Phi_h/\Phi_\infty < 0$.

Figure 4 shows the multipolar structure of the flux generated by a particle at the ISCO for the scalar, electromagnetic and gravitational-wave cases. The lowest radiative multipole $\ell = \max(|s|, 1)$ generates the greatest flux at the horizon, and the low multipoles also dominate the flux at infinity. The plots show evidence for the expected exponential fall-off of the modal fluxes with $\ell + 1/2$.

B. Conservative effects

1. Schwarzschild case

Regularization. Figure 5 illustrates the application of the regularization procedure to the radial component of the self-force, in the $a = 0$ case. The unregularized (“bare”) modes scale with $L = \ell + 1/2$ in the large- ℓ limit. After subtracting $\mathcal{F}_r^{[-1]\ell}$ and $\mathcal{F}_r^{[0]\ell}$ as in Eq. (85), that is, removing the leading and subleading order regularization terms, one obtains modes that scale with L^{-2} . This is the minimum necessary to obtain a convergent sum. To reduce the error associated with the high- ℓ tail, and to demonstrate that our results match expectations, we removed a further two regularization terms, that is, we subtracted $\mathcal{F}_r^{[4]\ell}$, leaving a mode sum whose terms converge as L^{-6} in the large- ℓ regime, as shown in Fig. 5.

Weak field expansion. Using numerical data for the radial component of the self-force at large values of r_0 we infer a weak-field expansion in the form

$$\mathcal{F}_r(r_0) \approx \frac{q^2}{4\pi\epsilon_0 c^2} \frac{GM}{r_0^3} \left(1 + \frac{3}{2\tilde{r}_0} + \frac{\alpha_2 \log(\tilde{r}_0)}{\tilde{r}_0^2} + \frac{\alpha_3}{\tilde{r}_0^2} + o\left(\frac{1}{\tilde{r}_0^3}\right) \right). \quad (94)$$

where \tilde{r}_0 is the dimensionless radial coordinate introduced in Eq. (93). The coefficients α_2 and α_3 were estimated from summing over the first 15ℓ -modes, with data in two ranges (i) $1000 < r_0 < 1500$ and (ii) $900 < r_0 < 1000$, yielding

$$\alpha_2 = 1.249(2) \quad \alpha_2 = 1.231(1) \quad (95a)$$

$$\alpha_3 = 1.38(1) \quad \alpha_3 = 1.48(1). \quad (95b)$$




FIG. 2. The ratio of the energy flux falling onto the horizon, Φ_h , to the energy flux radiated to infinity, Φ_∞ , as a function of orbital radius r_0 , for various spin parameters \tilde{a} , and for the scalar, electromagnetic and gravitational cases. The solid lines correspond to prograde orbits ($a \geq 0$). The dotted lines on the second plot, corresponds to retrograde orbits ($a < 0$). The color of the lines gives the magnitude of a . Negative ratios arise due a negative flux from the horizon associated with superradiance. A value less than -1 would indicate the existence of floating orbits.

The numeral in parantheses is the confidence interval in the final digit quoted, which is specific to the particular dataset used for the fitting. The data supports the presence of a log term at sub-sub-leading order, but accurate estimates for α_2 and α_3 have not been obtained.

Figure 6 compares the weak-field expansion, Eq. (94), with numerical data for \mathcal{F}_r for $a = 0$. It shows that \mathcal{F}_r increases monotonically as r_0 decreases. Moreover, \mathcal{F}_r differs from the leading order term in Eq. (94) by no more

than a factor of ~ 1.44 across the range $[r_{\text{ISCO}}, \infty)$. Including successive terms in the expansion improves the agreement with the data; and Eq. (94) gives a relative error of $\sim 6\%$ at the ISCO.

Shifts in orbital parameters. The conservative self-force has the effect of shifting the orbital parameters from their geodesic values at order q^2 . For a circular orbit on the Schwarzschild spacetime, the fractional change in the orbital energy E , angular momentum J and frequency Ω is given by




FIG. 3. The ratio of fluxes Φ_h/Φ_∞ for a particle on the innermost stable circular orbit, as a function of a the spin of the black hole. Negative (positive) values of a correspond to retrograde (prograde) circular orbits. For $\Omega_h > \Omega$, there is a negative flux ($\Phi_h < 0$) from the horizon, a manifestation of superradiance.




FIG. 4. The multipolar structure of the flux radiated through the horizon (upper) and to infinity (lower) by a charged particle on a circular orbit at the ISCO of a Kerr black hole with $a = 0.99M$. The trendline indicates an exponential fall-off with multipole number ℓ .




FIG. 5. Scalar spherical modes of the radial component of the self-force \mathcal{F}_r , and regularization at various orders. Here we have chosen $a = 0$ (Schwarzschild) and $r_0/M = 20$. The blue dots are the values of the bare force, which grow linearly with $L = \ell + 1/2$ at large L . The solid black lines are guidelines to represent the decay of the regularized force.




FIG. 6. Comparison between the numerically-determined value of \mathcal{F}_r (solid) and the weak-field expansion in Eq. (94) (dashed), for the Schwarzschild ($a = 0$) case. Here the dashed lines $\mathcal{F}_r^{(n)}$ show truncated versions of (94), with n indicating the number of terms included.




FIG. 7. Fractional change of the energy (green), angular momentum (blue) and frequency (red) for a particle on a circular orbit as a function of the orbital radius r_0 in the Schwarzschild case. The inset shows the radial range presented in Fig. 12 of Haas [64]. Our results agree with [64] and provides the behavior of the fractional for a larger radial range.




FIG. 8. Scalar-spherical modes of the radial component of the self-force and their regularization at various orders in the Kerr case. Here we have chosen $r_0/M = 20$ and $a = \frac{1}{2}M$. The blue dots are the values of the bare force, which (at leading order) grow linearly with $L = \ell + 1/2$. The solid black lines are guidelines to indicate the power-law decay of the regularized modes.

$$\frac{\Delta \Omega}{\Omega_0} = -\frac{(r_0 - 3M)r_0}{2\mu M} \mathcal{F}_r, \quad (96a)$$

$$\frac{\Delta E}{E_0} = -\frac{r_0}{2\mu} \mathcal{F}_r, \quad (96b)$$

$$\frac{\Delta J}{J_0} = -\frac{(r_0 - 2M)r_0}{2\mu M} \mathcal{F}_r. \quad (96c)$$

Figure 7 shows the shift in E , J and Ω as a function of r_0 . In each case, the self-force leads to a reduction in E , J and Ω . The shifts for the Kerr case are given in Appendix D.

2. Kerr case

Figure 8 shows that the bare modes of the force, \mathcal{F}_r^ℓ defined in Eq. (82), are correctly regularized with the regularization parameters calculated by Heffernan *et al.* [51]. This is a nontrivial test of the formulation, and of the projection onto spherical harmonics. In the projection step, we find that it is necessary to expand to subsubleading order in $z = \cos \theta$ in Eq. (82) to achieve regularization at order $n = 2$, and to obtain a regularized force $\mathcal{F}_r^{\text{reg}[2]}$ which is well defined on the particle such that




FIG. 9. Radial component of the self-force (scaled by r_0^3) for various black hole spins, $a \in \{-0.99, 0.5, 0, 0.5, 0.99\}$. The solid lines correspond to prograde orbits ($a > 0$), while the dotted lines correspond to retrograde orbits ($a < 0$), and the color of the lines gives the magnitude of a . In each case, the minimum radius is the innermost stable circular orbit.




FIG. 10. The radial and time components of the self-force for a particle on the innermost stable circular orbit, as a function of a the spin of the black hole. Negative (positive) values of a correspond to retrograde (prograde) circular orbits.

its left-sided limit ($r \rightarrow r_0^-$) and right-sided limit ($r \rightarrow r_0^+$) are in agreement.

Figure 9 shows \mathcal{F}_r as a function of r_0 , for several values of the black hole spin parameter a/M . We observe that \mathcal{F}_r is everywhere positive (i.e., repulsive) and greater than q^2/r_0^3 . At fixed radius, \mathcal{F}_r is larger on the retrograde orbit than on the prograde orbit. The effect of black hole rotation increases as r_0 decreases, as expected.

By fitting the numerical data, we find a linear-in- a contribution to \mathcal{F}_r of $-3\tilde{a}\tilde{r}_0^{-9/2}$ at leading order.

Figure 10 shows the self-force on the ISCO, as a function of a/M . The conservative component, \mathcal{F}_r , is always positive (i.e., repulsive). The total flux is always positive, indicating that superradiance is insufficient for a floating orbit to arise. The magnitudes of \mathcal{F}_r and \mathcal{F}_t are largest on the corotating ISCO of a rapidly-rotating black hole. In the limit $a \rightarrow M$, the ISCO approaches $r_0 = M$.

Table I provides a selection of values of \mathcal{F}_r for circular orbits of radii $r_0 \in [r_{\text{isco}}, 50M]$, for the black hole spin parameters $a = 0, \pm 0.5M$ and $\pm 0.99M$.

TABLE I. *Radial component of self-force for circular equatorial geodesic orbits.* The dimensionless values in the table correspond to $\mathcal{F}_r/[(q^2/4\pi\epsilon_0)(c^2/GM)^2]$. The digit in parantheses is an estimate of the uncertainty in the final quoted digit. The ISCO radius, defined in Eq. (15), is $r_{\text{isco}}/M \in \{8.971861, 7.554585, 6.0, 4.233003, 1.454498\}$ (to 7 s.f.) for the cases $a/M \in \{-0.99, -0.5, 0, 0.5, 0.99\}$.

r_0/M	$\mathcal{F}_r(r_0)$				
	$a = -0.99$	$a = -0.5$	$a = 0$	$a = 0.5$	$a = 0.99$
r_{isco}/M	0.001967652(2)	0.003315094(1)	0.0066497(5)	0.019003(2)	0.479(1)
10	0.0013513595(1)	0.0012770754(1)	0.00120985(2)	0.00114927(1)	0.001093823(1)
20	0.000141150327(2)	0.00013867449(5)	0.00013624(1)	0.000133916(2)	0.0001316275(1)
50	0.000008332378(2)	0.000008296911(1)	0.000008261044(2)	0.000008225470(2)	0.000008190833(6)

V. DISCUSSION AND CONCLUSION

In this article, we have computed the electromagnetic self-force acting on a point charge—or, with caveats, on a charged compact body—on a circular geodesic lying in the equatorial plane of a rotating black hole. This represents the first EM self-force calculation on Kerr spacetime in a dynamical scenario (see below for static cases). Our results complement those already available for the *gravitational* self-force on Kerr [42,43,48,62,71–73], a topic which has received much attention due to its relevance in modeling extreme mass-ratio inspirals for gravitational wave detectors.

To compare the dissipative effects of the electromagnetic and gravitational self-forces, consider once more the inspiral of a particle or compact body of mass μ and charge q into a black hole of mass M , driven by the dissipative component of the self-force. From the chirp formulas (3) and (1), valid in the large- r_0 regime, an order-of-magnitude estimate of the merger timescale, starting with an orbit of radius r_0 , is

$$\tau_{\text{EM}} \sim \left(\frac{\pi\epsilon_0 GM_\odot^2}{Q^2} \right) \cdot \left(\frac{M}{\mu} \right) \cdot \left(\frac{r_0}{GM/c^2} \right)^2 \cdot \frac{r_0}{c}, \quad (97a)$$

$$\tau_{\text{grav}} \sim \left(\frac{5}{2^{1/3}32} \right) \cdot \left(\frac{M}{\mu} \right) \cdot \left(\frac{r_0}{GM/c^2} \right)^3 \cdot \frac{r_0}{c}. \quad (97b)$$

Here Q is the net charge density of the particle/compact body in Coulombs per solar mass, and we have made the assumption that $\mu \ll M$ to obtain (97b). Numerical evaluation of the first parantheses in Eq. (97a) yields $7.4 \times 10^{39} \text{ C}^2/Q^2$, and thus, for a compact body, an electromagnetically driven inspiral is much slower than a gravitationally-driven inspiral, unless the compact body can support implausibly high net charge densities of $Q \gtrsim 10^{18} \text{ C}$ per solar mass. On the other hand, for an elementary charged particle the converse is true, as $Q \approx 1.9 \times 10^{38} \text{ C}$ per solar mass for a proton, for instance. That is, for a charged elementary particle, the EM inspiral is more rapid and the gravitational wave flux is negligible; but nevertheless, the inspiral into a black hole is exceedingly slow due to the suppressing factor M/μ . Of course,

an elementary-particle-black-hole-inspiral scenario is rather artificial, not least because we have neglected all contents of the universe but two.

One key result of this work is a demonstration that the local dissipative component of the self-force \mathcal{F}_t exactly balances with the sum of the electromagnetic flux radiated to infinity and down the horizon of the black hole, in accord with Eq. (65), up to the expected numerical precision. Closer examination of the fluxes, in Figs. 1, 2 and 3, shows that superradiance is stimulated when the angular velocity of the black hole horizon exceeds the orbital angular velocity. However, we find that superradiance is not sufficient to support floating orbits, even at the ISCO (see also [31]).

A key difference between the electromagnetic self-force and the gravitational self-force is that the latter is *gauge-dependent* under small changes in the coordinate system at $O(\mu)$. More precisely, for circular orbits the dissipative component of the gravitational self-force—relating to the radiated fluxes—*can* be identified uniquely, but the conservative component cannot; it is coordinate-dependent. This means that it is *not* possible to directly compare \mathcal{F}_r between the electromagnetic and gravitational cases. Instead, one must look to the gauge-invariant consequences of the conservative component of self-force to make meaningful comparisons. For example, Fig. 7 shows the fractional change in the orbital energy, angular momentum and frequency at fixed r_0 due to the conservative component of the self-force.

One such gauge-invariant consequence, slightly beyond the scope of this work, is the shift in the ISCO at $O(q^2)$ that arises due to the conservative component of the self-force. This can be calculated by examining mildly eccentric orbits [74], or possibly by using a Hamiltonian approach with circular-orbit data as input [62]; a comparison with known results for the ISCO shift induced by the gravitational self-force would certainly be of interest. Another observable that could be compared directly is the self-force-induced shift in the advance of the periapsis of an eccentric bound orbit [71].

The new results presented in Sec. IV are mostly numerical in nature. We have inferred leading order terms in weak-field expansions by fitting the numerical data. A complementary approach is to apply the Mano-Suzuki-Takasugi (MST) formalism [67] to obtain analytical results in the form of post-Newtonian expansions (see e.g., [69,70]). This has been done successfully in the gravitational self-force case, for quantities such as fluxes [75–77], Detweiler’s redshift invariant [78], and the spin-precession invariant [79]. The MST method can be straightforwardly adapted from the $s = 2$ to the $s = 1$ case. Indeed, in Sec. IVA 1 we have applied this method to obtain the flux through subleading order in \tilde{r}_0^{-1} . A promising avenue for future work, therefore, is to apply the MST method to the formulae

herein to obtain very high-order expansions of (e.g.,) \mathcal{F}_t and \mathcal{F}_r in closed form.

It is worth noting that the calculation presented here is not fully self-consistent, in the sense that we have evaluated the self-force by assuming the past worldline of the particle is a *geodesic*, rather than a trajectory that has itself been accelerated by its own self-force. Introducing the “true” trajectory would introduce subdominant contributions to the force starting at $O(q^4)$. One challenge, for future investigation, is to evolve the orbit in a fully self-consistent manner under the action of the electromagnetic self-force. This has already been done successfully for the gravitational self-force [73,80].

The *electrostatic* self-force on a charged particle on Kerr was examined many years ago by Léauté and Linet [81], and later by Piazzese and Rizzi [82]. For the special case of a particle at rest on the symmetry axis $\theta = 0$ at $r = r_0$, the (repulsive, conservative) self-force is available in closed form [82],

$$\mathcal{F}_{\text{self}}^\mu = \frac{q^2(Mr_0 - a^2)}{(r_0^2 + a^2)^2} e_3^\mu, \quad (98)$$

where e_3^μ is a unit spacelike vector along the symmetry axis. It is notable that Eq. (98) does not depend on the sign of a , and thus frame-dragging effects are absent in this highly symmetric case. Here, we have established that \mathcal{F}_r has a linear-in- a contribution for geodesic orbits in the equatorial plane.

Two further avenues of enquiry suggest themselves. First, the self-force on the ISCO in the $a \rightarrow M$ extremal limit has been investigated in the gravitational self-force context [83], but not yet in the electromagnetic self-force context. Second, an additional physical effect which has not been examined here is the self-torque that would arise at $O(q^2)$ if the particle (or compact body) is endowed with a magnetic dipole moment. In other words, the force arising from the (regularized) magnetic field in the rest frame of the particle.

ACKNOWLEDGMENTS

With thanks to Barry Wardell and Niels Warburton for email correspondence and discussions, and to the anonymous referees for constructive feedback. With additional thanks to Chris Kavanagh for assistance in obtaining the next-to-leading order expansion of the flux in Sec. IVA 1. This work makes use of the Black Hole Perturbation Toolkit [66]. T. T. and S. D. acknowledge financial support from the Science and Technology Facilities Council (STFC) under Grant No. ST/T001038/1. S. D. acknowledges financial support from the European Union’s Horizon 2020 research and innovation programme under the H2020-MSCA-RISE-2017 Grant No. FunFiCO-777740.

APPENDIX A: DISSIPATIVE SELF-FORCE IN THE NEWTONIAN LIMIT

For circular orbits far from a black hole ($r_0 \gg GM/c^2$), the speed of the particle, $|v| = r_0\Omega = \sqrt{GM/r_0}$, is small in comparison with the speed of light c , and a leading-order Newtonian approximation for the flux (92) and the chirp formula (3) is obtained by combining the Abraham-Lorentz force (2) with circular orbits in Newtonian gravity.

The work done in unit time P upon a particle of charge q by the Abraham-Lorentz force (2) is

$$P = \mathbf{F} \cdot \mathbf{v} = \frac{2}{3} \frac{q^2}{4\pi\epsilon_0 c^3} \dot{\mathbf{a}} \cdot \mathbf{v}. \quad (\text{A1})$$

Inserting a fixed circular orbit with $\mathbf{r} = r_0 \hat{\mathbf{r}}$ and $\mathbf{v} \equiv \dot{\mathbf{r}} = r_0 \Omega \hat{\boldsymbol{\phi}}$ and $\mathbf{a} \equiv \dot{\mathbf{v}} = -r_0 \Omega^2 \hat{\mathbf{r}}$ and $\dot{\mathbf{a}} \equiv -r_0 \Omega^2 \dot{\hat{\boldsymbol{\phi}}}$, where $\hat{\mathbf{r}}$ and $\hat{\boldsymbol{\phi}}$ are unit vectors and $\Omega = \sqrt{GM/r_0^3}$ is the angular frequency of the orbit, yields

$$P = -\frac{2}{3} \frac{q^2}{4\pi\epsilon_0 c^3} r_0^2 \Omega^4 \quad (\text{A2})$$

$$= -\frac{2}{3} \frac{q^2}{4\pi\epsilon_0 c^3} \frac{G^2 M^2}{r_0^4}. \quad (\text{A3})$$

By conservation of energy, the flux radiated to infinity is equal and opposite to the work done on the particle by the Abraham-Lorentz force, that is, $\Phi_{\text{Newt}} = -P$, yielding Eq. (92).

For a particle of mass μ on a circular orbit under gravity, the sum of kinetic and (Newtonian) potential energies is $E = -GM\mu/2r_0$. We now allow the particle to gradually spiral inwards on a sequence of quasicircular orbits, by equating P with $\dot{E} = -GM\mu\dot{r}/2r^2$. This leads to

$$\frac{\dot{f}}{f^3} = \frac{8\pi^2 q^2}{4\pi\epsilon_0 c^3}, \quad (\text{A4})$$

where $f = \Omega/2\pi$ is the orbital frequency. Integrating with respect to time leads to

$$f(t) = \frac{1}{4\pi} \sqrt{\frac{4\pi\epsilon_0 c^3 \mu}{q^2} (t_0 - t)^{-1/2}}, \quad (\text{A5})$$

where the time of collision t_0 arises as the constant of integration. For the case of an electron of mass $\mu = m_e$ and charge $q = -e$, Eq. (A5) reduces to Eq. (3) once we insert the definition of the fine-structure constant $\alpha = e^2/(4\pi\epsilon_0 \hbar c)$ and the Bohr radius $a_0 = 4\pi\epsilon_0 \hbar^2/(m_e e^2)$.

APPENDIX B: ENERGY FLUX

1. Flux at infinity

At infinity the energy flux is given by

$$\Phi_R^K = \Delta t^{-1} \int T^{ab} K_b d\Sigma_a \quad (\text{B1})$$

where we have chosen $K^\mu = [1, 0, 0, 0]$ to be the Killing vector and $d\Sigma_\mu$ is defined by the condition $r \rightarrow \infty$ and is given by

$$d\Sigma_\mu = n_\mu d\Sigma, \quad (\text{B2})$$

with

$$n_\mu = \frac{[0, 1, 0, 0]}{\sqrt{g^{rr}}} \quad \text{and} \quad d\Sigma = |h|^{1/2} dt d\theta d\phi. \quad (\text{B3})$$

where $h_{\mu\nu}$ is the induced metric on the hypersurface define by $r \rightarrow \infty$. Since $|h| = g/g_{rr}$, we have

$$d\Sigma_\mu = [0, 1, 0, 0] \sqrt{-g} dt d\theta d\phi. \quad (\text{B4})$$

The flux is then given by

$$\Phi_R = \int T^r{}_t \sqrt{-g} d\theta d\phi. \quad (\text{B5})$$

The energy-momentum tensor can be expressed in terms of the Maxwell scalars as [54]

$$4\pi T_{\mu\nu} = -\{\phi_0 \phi_0^* n_\mu n_\nu + 2\phi_1 \phi_1^* [l_{(\mu} n_{\nu)} + m_{(\mu} m_{\nu)}^*] + \phi_2 \phi_2^* l_\mu l_\nu - 4\phi_0^* \phi_1 n_{(\mu} m_{\nu)} - 4\phi_1^* \phi_2 l_{(\mu} m_{\nu)} + 2\phi_2 \phi_0^* m_\mu m_\nu\} + \text{c.c.} \quad (\text{B6})$$

where parentheses denote symmetrization. With our choice of tetrad, we find that the relevant terms as $r \rightarrow \infty$ are

$$\lim_{r \rightarrow \infty} T^r{}_t = \frac{1}{2\pi} \left(\phi_2 \phi_2^* - \frac{\phi_0 \phi_0^*}{4} \right) \quad (\text{B7})$$

We recall that

$$\phi_0 = \sum_{l,m} {}_{+1}R_{lm}(r) S_{+1}^{\ell m}(\theta) e^{im(\phi - \Omega t)}, \quad (\text{B8})$$

$$2(r - ia \cos \theta)^2 \phi_2 = \sum_{l,m} {}_{-1}R_{lm}(r) S_{-1}^{\ell m}(\theta) e^{im(\phi - \Omega t)}, \quad (\text{B9})$$

as well as the fact that

$$\lim_{r \rightarrow \infty} r^2 |{}_{+1}R_{lm}(r)| = 0 \quad \text{and} \quad \lim_{r \rightarrow \infty} \frac{|{}_{-1}R_{lm}(r)|}{r} = |{}_{-1}\alpha_{lm}^\infty|. \quad (\text{B10})$$

Using the orthonormality properties of the spin-weighted spheroidal harmonics, we therefore get the energy flux radiated at infinity

$$\Phi_\infty = \lim_{r \rightarrow \infty} \int \frac{r^2 \sin \theta}{2\pi} \phi_2 \phi_2^* d\theta d\phi = \sum_{l,m} \frac{|\alpha_{lm}^\infty|^2}{8\pi} \quad (\text{B11})$$

2. Flux through the horizon

In order to evaluate the flux of energy through the horizon, we first need to modify the tetrad basis we have defined in Eq. (10) since it is singular at the horizon. Following Chap. 8 Sec. 76(b) of [59], we first perform a rotation of class III (according to Chandrashekar's convention):

$$l^\mu \rightarrow \frac{\Delta}{2(r^2 + a^2)} l^\mu = \left[\frac{1}{2}, \frac{\Delta}{2(r^2 + a^2)}, 0, \frac{a}{2(r^2 + a^2)} \right] \quad (\text{B12})$$

and

$$n^\mu \rightarrow \frac{2(r^2 + a^2)}{\Delta} n^\mu = \left[\frac{(r^2 + a^2)^2}{\Sigma \Delta}, -\frac{(r^2 + a^2)}{\Sigma}, 0, a \frac{(r^2 + a^2)}{\Sigma \Delta} \right]. \quad (\text{B13})$$

We then go to a Kerr-Schild frame via the coordinate transformation:

$$dv = dt + \frac{r^2 + a^2}{\Delta} dr \quad \text{and} \quad d\tilde{\phi} = d\phi + \frac{a}{\Delta} dr. \quad (\text{B14})$$

In this frame, the null vectors $l_{(HH)}^\mu$ and $n_{(HH)}^\mu$, where HH stands for Hartle-Hawking, are given by

$$l_{(HH)}^\mu = \left[1, \frac{\Delta}{2(r^2 + a^2)}, 0, \frac{a}{r^2 + a^2} \right], \quad (\text{B15})$$

$$n_{(HH)}^\mu = \left[0, -\frac{r^2 + a^2}{\Sigma}, 0, 0 \right]. \quad (\text{B16})$$

On the horizon, the vector $l_{(HH)}$ can be expressed in terms of the time and angular Killing vector $K_T^\mu = [1, 0, 0, 0]$ and $K_L^\mu = [0, 0, 0, 1]$ as

$$l_{(HH)}^\mu = K_T^\mu + \Omega_h K_L^\mu. \quad (\text{B17})$$

where $\Omega_h = a/2Mr_+$ is the angular frequency of the horizon. In this basis, which is well behaved at the horizon, the Maxwell scalars $\phi_0^{(HH)}$ is related to the Maxwell scalar ϕ_0 computed in the basis (10) via

$$\phi_0^{(HH)} = \frac{\Delta}{2(r^2 + a^2)} \phi_0. \quad (\text{B18})$$

The surface element $d\Sigma_a$ of the horizon, which is a null hypersurface, is given by

$$d\Sigma_\mu = l_\mu^{(HH)} d\sigma dt \quad (\text{B19})$$

where $d\sigma = 2Mr_+ \sin(\theta) d\theta d\phi$ is the elementary surface area of the event horizon. Therefore, the elementary flow of energy and angular momentum through the horizon are

$$\left(\frac{d^2 \Phi_h^T}{dt d\Omega} \right) = 2Mr_+ l_\mu^{(HH)} K_T^\nu T_\nu^\mu \quad (\text{B20})$$

$$\left(\frac{d^2 \Phi_h^L}{dt d\Omega} \right) = 2Mr_+ l_\mu^{(HH)} K_L^\nu T_\nu^\mu \quad (\text{B21})$$

Combining these with (B17) and using the fact that $\Phi^T = -\Omega \Phi^L$ yields

$$\left(\frac{d^2 \Phi_h^T}{dt d\Omega} \right) = \frac{2Mr_+ \omega}{\omega - m\Omega_h} T^{\mu\nu} l_\mu^{(HH)} l_\nu^{(HH)}. \quad (\text{B22})$$

By definition $T^{\mu\nu} l_\mu^{(HH)} l_\nu^{(HH)} = \phi_0^{(HH)} \phi_0^{*(HH)}/2\pi$, and therefore

$$\left(\frac{d^2 \Phi_h^T}{dt d\Omega} \right) = \frac{\omega}{8Mr_+ \tilde{\omega}} \frac{\Delta}{2\pi} \phi_0 \phi_0^*. \quad (\text{B23})$$

Finally, integrating over the surface element using the decomposition (B8), the orthonormality of the spin-weighted spheroidal harmonics, and the asymptotic behavior of the radial function near the horizon,

$$\Phi_h^T = \sum_{l,m} \frac{\omega}{16\pi Mr_+ \tilde{\omega}} |\alpha_{lm}^h|^2. \quad (\text{B24})$$

APPENDIX C: PROJECTION ONTO SCALAR SPHERICAL HARMONICS

To compute the physical conservative part of the self-force we apply the mode-sum regularization procedure. As a preliminary step before applying the regularization, one should decompose the radial force onto a basis of scalar spherical harmonics. Since the structure of the Kerr metric invited us to use spin-weighted spheroidal harmonics as a basis for the angular functions of our problem, we now need to project the spin-weighted spheroidal harmonics onto scalar spherical harmonics.

1. Projection of the spin-weighted spheroidal harmonics

a. From spin-weighted spheroidal harmonics to spin-weighted spherical harmonics

We first decompose the spin-weighted spheroidal harmonics S_s^{lm} onto the spin-weighted spherical harmonics Y_s^{lm} :

$$S_s^{lm}(\theta) = \sum_{\tilde{l}} (b_s^m)_{\tilde{l}} Y_{\tilde{l}}^{lm}(\cos \theta). \quad (\text{C1})$$

The coefficients $(b_s^m)_i^l$ are computed using the Black Hole Perturbation Toolkit [66].

b. From spin-weighted spherical harmonics to scalar spherical harmonics

We decompose the spin-weighted spherical harmonics in terms of spherical harmonics Y_0^{lm} ,

$$Y_{+1}^{lm}(z) = \sum_{\tilde{l}} \frac{(A_{+1}^m)_i^l}{\sqrt{1-z^2}} Y_{\tilde{l}}^{lm}(z), \quad (\text{C2})$$

$$Y_0^{lm}(z) = \sum_{\tilde{l}} \delta_{\tilde{l}}^l Y_{\tilde{l}}^{lm}(z), \quad (\text{C3})$$

$$Y_{-1}^{lm}(z) = \sum_{\tilde{l}} \frac{(A_{-1}^m)_i^l}{\sqrt{1-z^2}} Y_{\tilde{l}}^{lm}(z), \quad (\text{C4})$$

where $z = \cos \theta$ and the coefficients are given by

$$(A_{+1}^m)_i^l = (-1)^{m+1} \sqrt{2(2l+1)(2\tilde{l}+1)} \\ \times \begin{pmatrix} 1 & l & \tilde{l} \\ 0 & m & -m \end{pmatrix} \begin{pmatrix} 1 & l & \tilde{l} \\ 1 & -1 & 0 \end{pmatrix}, \quad (\text{C5a})$$

$$(A_{-1}^m)_i^l = (-1)^m \sqrt{2(2l+1)(2\tilde{l}+1)} \\ \times \begin{pmatrix} 1 & l & \tilde{l} \\ 0 & m & -m \end{pmatrix} \begin{pmatrix} 1 & l & \tilde{l} \\ -1 & 1 & 0 \end{pmatrix}. \quad (\text{C5b})$$

It follows from the properties of the Wigner 3j symbols that

$$(A_{+1}^m)_i^l = (-1)^{l+\tilde{l}} (A_{-1}^m)_i^{\tilde{l}}. \quad (\text{C6})$$

Combining the two decompositions, we can write the spin-weighted spheroidal harmonics as

$$S_s^{lm}(\theta) = \sum_{\tilde{l}, \tilde{m}} (b_s^m)_i^l (A_s^m)_i^{\tilde{m}} \frac{Y_{\tilde{l}}^{\tilde{m}}(z)}{\sqrt{1-z^2}}. \quad (\text{C7})$$

Note that due to the presence of the 3j-symbols in Eqs. (C5), the indices \tilde{l} and \hat{l} satisfy $\tilde{l}-1 \leq \hat{l} \leq \tilde{l}+1$.

2. Expansion of $\mathcal{L}_1 S_{+1}^{lm}(\theta)$

The definition of $\mathcal{L}_1 S_{+1}^{lm}$ is

$$\mathcal{L}_1 S_{+1}^{lm}(\theta) = \partial_\theta S_{+1}^{lm}(\theta) + \left(\frac{m}{\sin \theta} - a\omega \sin \theta \right) S_{+1}^{lm} + \frac{\cos \theta}{\sin \theta} S_{+1}^{lm}. \quad (\text{C8})$$

In order to project $\mathcal{L}_1 S_{+1}^{\ell m}$ onto scalar spherical harmonics, we first need to project $\partial_\theta S_{+1}^{\ell m}$.

The spherical harmonics of different spins are related by

$$\delta Y_s^{lm} = \sqrt{(l-s)(l+s+1)} Y_{s+1}^{lm}, \quad (\text{C9})$$

$$\bar{\delta} Y_s^{lm} = -\sqrt{(l+s)(l-s+1)} Y_{s-1}^{lm}, \quad (\text{C10})$$

where the spin-raising and spin-lowering operators δ and $\bar{\delta}$ are defined by

$$\delta f_s(\theta, \phi) = -(\sin \theta)^s \left[\frac{\partial}{\partial \theta} + \frac{i}{\sin \theta} \frac{\partial}{\partial \phi} \right] ((\sin \theta)^{-s} f_s) \quad (\text{C11})$$

$$\bar{\delta} f_s(\theta, \phi) = -(\sin \theta)^{-s} \left[\frac{\partial}{\partial \theta} - \frac{i}{\sin \theta} \frac{\partial}{\partial \phi} \right] ((\sin \theta)^s f_s). \quad (\text{C12})$$

We have that

$$\partial_\theta S_{+1}^{lm}(\theta) = \sum_{\tilde{l}} (b_{+1}^m)_i^l \partial_\theta Y_{+1}^{\tilde{l}m}(\cos \theta). \quad (\text{C13})$$

We can eliminate the derivative using the relation (C11) and the expression for $\bar{\delta}$, namely,

$$\bar{\delta} Y_{+1}^{lm}(z) = -(\sin \theta)^{-1} \left[\frac{\partial}{\partial \theta} - \frac{i}{\sin \theta} \frac{\partial}{\partial \phi} \right] (\sin \theta Y_{+1}^{lm}(z)) \quad (\text{C14})$$

$$= -\frac{1}{\sin \theta} [(\cos \theta + m) Y_{+1}^{lm}(z) + \sin \theta \partial_\theta Y_{+1}^{lm}(z)] \quad (\text{C15})$$

$$= -\sqrt{l(l+1)} Y_0^{lm}. \quad (\text{C16})$$

Therefore, we have that

$$\partial_\theta S_{+1}^{lm}(\theta) = \sum_{\tilde{l}} (b_{+1}^m)_i^l \left(\sqrt{\hat{l}(\hat{l}+1)} Y_0^{\hat{l}m}(\cos \theta) - \frac{(\cos \theta + m)}{\sin \theta} Y_{+1}^{\hat{l}m}(\cos \theta) \right). \quad (\text{C17})$$

Substituting (C17) into (C9), we get

$$\mathcal{L}_1 S_{+1}^{lm} = \sum_{\tilde{l}} (b_{+1}^m)_i^l \left[\sqrt{\hat{l}(\hat{l}+1)} Y_0^{\hat{l}m} - a m \Omega \sin \theta Y_{+1}^{\hat{l}m} \right] \quad (\text{C18})$$

$$= \sum_{\tilde{l}, \tilde{m}} (b_{+1}^m)_i^l \left[\sqrt{\hat{l}(\hat{l}+1)} \delta_{\tilde{l}}^{\hat{l}} - a m \Omega (A_{+1}^m)_i^{\tilde{l}} \right] Y_0^{\tilde{m}}. \quad (\text{C19})$$

APPENDIX D: SHIFTS IN ORBITAL PARAMETERS FROM CONSERVATIVE SELF-FORCE

On Kerr spacetime, the shifts in the energy and angular momentum at fixed r_0 are

$$\frac{\Delta E}{E_0} = -\frac{r_0 \Delta_0 (L_0 (r_0 - 2M) + 2aME_0)}{2E_0 \mathcal{X}} \mathcal{F}_r / \mu, \quad (\text{D1a})$$

$$\frac{\Delta L}{L_0} = -\frac{r_0 \Delta_0 (E_0 (r_0^3 + a^2 (r_0 + 2M)) - 2aML_0)}{2L_0 \mathcal{X}} \mathcal{F}_r / \mu, \quad (\text{D1b})$$

$$\mathcal{X} = (r_0^3 - 3Mr_0^2 - 2Ma^2)E_0 L_0 + aM((3r_0^2 + a^2)E_0^2 + L_0^2) \quad (\text{D1c})$$

and the shift in the angular velocity at fixed r_0 is

$$\frac{\Delta \Omega}{\Omega_0} = -\frac{\Delta_0^2 r_0^3 ((r_0^3 + a^2 (r_0 + 2M))E_0^2 - 4E_0 L_0 aM - L_0^2 (r_0 - 2M))}{2\mathcal{X}(E_0 r_0 (r_0^2 + a^2) - 2aM(L_0 - aE_0))(r_0 L_0 - 2M(L_0 - aE_0))} \mathcal{F}_r / \mu. \quad (\text{D2})$$

APPENDIX E: A DISTRIBUTIONAL IDENTITY

The Dirac delta distribution $\delta(x - x_0)$ is of compact support, with $\delta(x - x_0) = 0$ for $x \neq x_0$ and the fundamental property

$$\int_{x_0-\epsilon}^{x_0+\epsilon} h(x) \delta(x - x_0) dx = h(x_0). \quad (\text{E1})$$

Here $\epsilon > 0$, x_0 is a real constant and $h(x)$ is a test function that is continuous at x_0 (and, below, differentiable). Now consider $f(x)\delta'(x - x_0)$ appearing in Eq. (36), where $f(x)$ is a differentiable function and the prime denotes an ordinary derivative with respect to x . By application of integration by parts,

$$\begin{aligned} & \int_{x_0-\epsilon}^{x_0+\epsilon} h(x) f(x) \delta'(x - x_0) dx \\ &= - \int_{x_0-\epsilon}^{x_0+\epsilon} \{h'(x) f(x) + h(x) f'(x)\} \delta(x - x_0), \\ &= -h'(x_0) f(x_0) - h(x_0) f'(x_0). \end{aligned} \quad (\text{E2})$$

By contrast, the integral of $f(x_0)\delta'(x - x_0)$, where $f(x_0)$ is a constant rather than a function of x , has but a single term,

$$\int_{x_0-\epsilon}^{x_0+\epsilon} h(x) f(x_0) \delta'(x - x_0) dx = -h'(x_0) f(x_0). \quad (\text{E3})$$

Hence $f(x)\delta'(x - x_0)$ and $f(x_0)\delta'(x - x_0)$ are not equivalent. On the other hand,

$$\begin{aligned} & \int_{x_0-\epsilon}^{x_0+\epsilon} h(x) \{f(x_0) \delta'(x - x_0) - f'(x_0) \delta(x - x_0)\} dx \\ &= -h'(x_0) f(x_0) - h(x_0) f'(x_0), \end{aligned} \quad (\text{E4})$$

which matches Eq. (E2). This equivalence justifies the replacement made in Eq. (36).

-
- [1] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), *Phys. Rev. Lett.* **116**, 241102 (2016).
 - [2] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), *Phys. Rev. Lett.* **119**, 161101 (2017).
 - [3] LIGO Scientific, Virgo, 1M2H, Dark Energy Camera GW-E, DES, DLT40, Las Cumbres Observatory, VINROUGE, MASTER Collaborations *et al.*, *Nature (London)* **551**, 85 (2017).
 - [4] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA Collaborations), *arXiv:2111.03606*.
 - [5] L. Blanchet, *Living Rev. Relativity* **17**, 2 (2014).
 - [6] L. Barack and A. Pound, *Rep. Prog. Phys.* **82**, 016904 (2019).
 - [7] M. D. Duez and Y. Zlochower, *Rep. Prog. Phys.* **82**, 016902 (2019).
 - [8] A. Einstein, *Sitzber. Preuss. Akad. Wiss.* **47**, 831 (1915).
 - [9] P. Peters and J. Mathews, *Phys. Rev.* **131**, 435 (1963).
 - [10] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), *Ann. Phys. (Berlin)* **529**, 1600209 (2017).
 - [11] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), *Phys. Rev. Lett.* **119**, 161101 (2017).
 - [12] E. W. Hirschmann, L. Lehner, S. L. Liebling, and C. Palenzuela, *Phys. Rev. D* **97**, 064032 (2018).
 - [13] M. Khalil, N. Sennett, J. Steinhoff, J. Vines, and A. Buonanno, *Phys. Rev. D* **98**, 104010 (2018).

- [14] V. Cardoso, C. F. B. Macedo, P. Pani, and V. Ferrari, *J. Cosmol. Astropart. Phys.* **05** (2016) 054; 04 (2020) E01.
- [15] G. Bozzola and V. Paschalidis, *Phys. Rev. Lett.* **126**, 041103 (2021).
- [16] R. Zhu and T. Osburn, *Phys. Rev. D* **97**, 104058 (2018).
- [17] J. Y. J. Burton and T. Osburn, *Phys. Rev. D* **102**, 104030 (2020).
- [18] E. Rutherford, London, Edinburgh, Dublin Philos. Mag. J. Sci. **27**, 488 (1914).
- [19] M. Abraham, *Theorie der Elektrizität* (BG Teubner, Leipzig, 1905), Vol. II.
- [20] H. A. Lorentz, *Arch. Néerl.* **25**, 363 (1892).
- [21] P. A. M. Dirac, *Proc. R. Soc. A* **167**, 148 (1938).
- [22] A. L. Schawlow, *Rev. Mod. Phys.* **54**, 697 (1982).
- [23] C. Baily, *Eur. Phys. J. H* **38**, 1 (2013).
- [24] C. M. DeWitt and B. S. DeWitt, *Phys. Phys. Fiz.* **1**, 3 (1964).
- [25] M. J. Pfenning and E. Poisson, *Phys. Rev. D* **65**, 084001 (2002).
- [26] B. S. DeWitt and R. W. Brehme, *Ann. Phys. (N.Y.)* **9**, 220 (1960).
- [27] S. L. Detweiler and B. F. Whiting, *Phys. Rev. D* **67**, 024025 (2003).
- [28] S. E. Gralla, A. I. Harte, and R. M. Wald, *Phys. Rev. D* **80**, 024031 (2009).
- [29] W. H. Press and S. A. Teukolsky, *Nature (London)* **238**, 211 (1972).
- [30] V. Cardoso, S. Chakrabarti, P. Pani, E. Berti, and L. Gualtieri, *Phys. Rev. Lett.* **107**, 241101 (2011).
- [31] S. J. Kapadia, D. Kennefick, and K. Glampedakis, *Phys. Rev. D* **87**, 044050 (2013).
- [32] C. W. Misner, R. A. Breuer, D. R. Brill, P. L. Chrzanowski, H. G. Hughes, and C. M. Pereira, *Phys. Rev. Lett.* **28**, 998 (1972).
- [33] M. Davis, R. Ruffini, J. Tiomno, and F. Zerilli, *Phys. Rev. Lett.* **28**, 1352 (1972).
- [34] E. Poisson, A. Pound, and I. Vega, *Living Rev. Relativity* **14**, 7 (2011).
- [35] J. Hobbs, *Ann. Phys. (N.Y.)* **47**, 141 (1968).
- [36] B. Wardell, C. R. Galley, A. Zenginoğlu, M. Casals, S. R. Dolan, and A. C. Ottewill, *Phys. Rev. D* **89**, 084021 (2014).
- [37] N. Khusnutdinov, *Eur. Phys. J. Plus* **136**, 669 (2021).
- [38] L. Barack and A. Ori, *Phys. Rev. D* **61**, 061502 (2000).
- [39] L. Barack and N. Sago, *Phys. Rev. D* **75**, 064021 (2007).
- [40] L. Barack and N. Sago, *Phys. Rev. D* **81**, 084021 (2010).
- [41] S. Akcay, *Phys. Rev. D* **83**, 124026 (2011).
- [42] A. G. Shah, T. S. Keidl, J. L. Friedman, D.-H. Kim, and L. R. Price, *Phys. Rev. D* **83**, 064018 (2011).
- [43] A. G. Shah, J. L. Friedman, and T. S. Keidl, *Phys. Rev. D* **86**, 084059 (2012).
- [44] S. Akcay, N. Warburton, and L. Barack, *Phys. Rev. D* **88**, 104009 (2013).
- [45] S. R. Dolan and L. Barack, *Phys. Rev. D* **87**, 084066 (2013).
- [46] T. Osburn, E. Forseth, C. R. Evans, and S. Hopper, *Phys. Rev. D* **90**, 104031 (2014).
- [47] M. van de Meent and A. G. Shah, *Phys. Rev. D* **92**, 064025 (2015).
- [48] M. van de Meent, *Phys. Rev. D* **97**, 104033 (2018).
- [49] L. Barack and A. Ori, *Phys. Rev. D* **67**, 024029 (2003).
- [50] A. Heffernan, A. Ottewill, and B. Wardell, *Phys. Rev. D* **86**, 104023 (2012).
- [51] A. Heffernan, A. Ottewill, and B. Wardell, *Phys. Rev. D* **89**, 024030 (2014).
- [52] A. Heffernan, The self-force problem: Local behaviour of the Detweiler-Whiting singular field, Ph.D. thesis, University College, Dublin, 2012.
- [53] W. H. Press and S. A. Teukolsky, *Astrophys. J.* **185**, 649 (1973).
- [54] S. A. Teukolsky, *Astrophys. J.* **185**, 635 (1973).
- [55] S. A. Teukolsky and W. H. Press, *Astrophys. J.* **193**, 443 (1974).
- [56] S. Chandrasekhar, *Proc. R. Soc. A* **349**, 1 (1976).
- [57] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Oxford University Press, Oxford, 1983).
- [58] S. M. Carroll, *Spacetime and Geometry* (Cambridge University Press, Cambridge, England, 2019).
- [59] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Oxford University Press, New York, 1998), Vol. 69.
- [60] N. Warburton and L. Barack, *Phys. Rev. D* **81**, 084039 (2010).
- [61] J. M. Bardeen, W. H. Press, and S. A. Teukolsky, *Astrophys. J.* **178**, 347 (1972).
- [62] S. Isoyama, L. Barack, S. R. Dolan, A. Le Tiec, H. Nakano, A. G. Shah, T. Tanaka, and N. Warburton, *Phys. Rev. Lett.* **113**, 161101 (2014).
- [63] D. Bini, C. Cherubini, R. T. Jantzen, and R. J. Ruffini, *Prog. Theor. Phys.* **107**, 967 (2002).
- [64] R. Haas, arXiv:1112.3707.
- [65] P. Nolan, High accuracy calculations of self force gauge invariants for compact binaries, Ph.D. thesis, University College Dublin, 2015.
- [66] Black Hole Perturbation Toolkit (bhptoolkit.org).
- [67] S. Mano, H. Suzuki, and E. Takasugi, *Prog. Theor. Phys.* **95**, 1079 (1996).
- [68] S. Mano, H. Suzuki, and E. Takasugi, *Prog. Theor. Phys.* **96**, 549 (1996).
- [69] C. Kavanagh, A. C. Ottewill, and B. Wardell, *Phys. Rev. D* **92**, 084025 (2015).
- [70] C. Kavanagh, A. C. Ottewill, and B. Wardell, *Phys. Rev. D* **93**, 124038 (2016).
- [71] M. van de Meent, *Phys. Rev. Lett.* **118**, 011101 (2017).
- [72] M. van de Meent, *Phys. Rev. D* **94**, 044034 (2016).
- [73] M. Van De Meent and N. Warburton, *Classical Quantum Gravity* **35**, 144003 (2018).
- [74] L. Barack and N. Sago, *Phys. Rev. Lett.* **102**, 191101 (2009).
- [75] R. Fujita, *Prog. Theor. Phys.* **128**, 971 (2012).
- [76] R. Fujita, *Prog. Theor. Exp. Phys.* **2015**, 033E01 (2015).
- [77] C. Munna, *Phys. Rev. D* **102**, 124001 (2020).
- [78] D. Bini and T. Damour, *Phys. Rev. D* **91**, 064050 (2015).
- [79] D. Bini and T. Damour, *Phys. Rev. D* **91**, 064064 (2015).
- [80] N. Warburton, S. Akcay, L. Barack, J. R. Gair, and N. Sago, *Phys. Rev. D* **85**, 061501 (2012).
- [81] B. Léauté and B. Linet, *J. Phys. A* **15**, 1821 (1982).
- [82] F. Piazzese and G. Rizzi, *Gen. Relativ. Gravit.* **23**, 403 (1991).
- [83] S. E. Gralla, A. P. Porfyriadis, and N. Warburton, *Phys. Rev. D* **92**, 064029 (2015).