

This is a repository copy of European Society for Radio-therapy and Oncology and European Organisation for Research and Treatment of Cancer consensus on reirradiation: definition, reporting, and clinical decision making.

White Rose Research Online URL for this paper: https://eprints.whiterose.ac.uk/208224/

Version: Supplemental Material

Article:

Andratschke, N., Willmann, J., Appelt, A.L. orcid.org/0000-0003-2792-9218 et al. (16 more authors) (2022) European Society for Radio-therapy and Oncology and European Organisation for Research and Treatment of Cancer consensus on re-irradiation: definition, reporting, and clinical decision making. The Lancet Oncology, 23 (10). e469-e478. ISSN 1470-2045

https://doi.org/10.1016/S1470-2045(22)00447-8

© 2022, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: https://creativecommons.org/licenses/


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Re-irradiation Type I n-th RT


- Overlap of irradiated volumes
- With or without concern for toxicity from cumulative doses



- No overlap of irradiated volumes
- Concern for toxicity from cumulative doses

Repeat organ irradiation n-th RT

- No overlap of irradiated volumes
- No concern for toxicity from cumulative doses
- Target volumes in the same organ



Target volumes in different organs

doses