

**REPLY: Navigating Nutritional Nuances
in the Research of Heart Failure With
Preserved Ejection Fraction**



We are delighted that Drs Dev and Ahmed found our study¹ of interest. Their comments highlight the importance of malnutrition in patients with heart failure with preserved ejection fraction (HFpEF) and its potential to influence therapeutic interventions. Our study's main focus was to explore the underlying biological mechanisms responsible for skeletal muscle pathology in HFpEF, which we harnessed to identify therapeutic targets via combined caloric restriction (CR) with exercise.¹ We found that CR with exercise rescued muscle pathology in a rat model of HFpEF, whereas pharmacologic cardiac therapies did not.¹ That similar mechanisms underlying muscle growth were affected in both animals and patients with HFpEF suggests a conserved response that, as we highlighted, warrants further investigation.¹

In complex pathologies such as HFpEF, any therapeutic approach requires the optimization of multiple variables, and we welcome the suggestion that nutritional status should form part of this. Indeed, future studies incorporating comprehensive nutritional status assessments before recommending CR in patients with HFpEF seem essential. Unfortunately, physicians often overlook nutritional status, focusing primarily on pharmacologic therapy. Therefore, introducing controlled CR in clinical practice may have the additional benefit of allowing malnutrition to be routinely screened in these patients. However, the lack of standardized definitions for malnutrition and varied assessment tools currently pose a limitation.² In our study,¹ we noted that CR with exercise requires further investigation to ensure efficacy in patients with HFpEF.

Drs Dev and Ahmed further highlight difficulties in translating CR and exercise to the clinic in HFpEF and how malnutrition may limit applicability. Randomized controlled trials have shown that CR and exercise in obese, older patients with HFpEF is safe and feasible and improves muscle performance and quality of life.^{3,4} Unfortunately, malnutrition was not evaluated in these studies,^{3,4} although one may consider that a significant number of patients included would have been malnourished, on the basis of the prevalence cited by Drs Dev and Ahmed. Despite this, no unfavorable responses were reported.^{3,4} Although exercise can counteract frailty in malnourished adults,⁵ more studies are needed on

the effects of CR. Nevertheless, as more than one-half of patients with HFpEF are not diagnosed as malnourished, CR with exercise offers a feasible treatment for a significant proportion of this patient population.^{3,4}

Ever Espino-Gonzalez, PhD^a

Peter G. Tickle, PhD^a

Raffaele Altara, PhD^{b,c}

George W. Booz, PhD^c

Alessandro Cataliotti, MD, PhD^d

Lee D. Roberts, PhD^a

Stuart Egginton, PhD^a

*T. Scott Bowen, MSc, PhD^a

^aSchool of Biomedical Sciences

Faculty of Biological Sciences

University of Leeds

Leeds LS2 9JT, United Kingdom

E-mail: t.s.bowen@leeds.ac.uk

From the ^aUniversity of Leeds, Leeds, United Kingdom;

^bMaastricht University, Maastricht, the Netherlands;

^cUniversity of Mississippi Medical Center, Jackson, Mississippi, USA; and the

^dOslo University Hospital and the University of Oslo, Oslo, Norway.

<https://doi.org/10.1016/j.jacbts.2024.01.014>

© 2024 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

REFERENCES

1. Espino-Gonzalez E, Tickle Peter G, Altara R, et al. Caloric restriction rejuvenates skeletal muscle growth in heart failure with preserved ejection fraction. *J Am Coll Cardiol Basic Trans Science*. 2024;9(2):223-240.
2. Driggin E, Cohen LP, Gallagher D, et al. Nutrition assessment and dietary interventions in heart failure: JACC review topic of the week. *J Am Coll Cardiol*. 2022;79:1623-1635.
3. Kitzman DW, Brubaker P, Morgan T, et al. Effect of caloric restriction or aerobic exercise training on peak oxygen consumption and quality of life in obese older patients with heart failure with preserved ejection fraction: a randomized clinical trial. *JAMA*. 2016;315:36-46.
4. Brubaker PH, Nicklas BJ, Houston DK, et al. A randomized, controlled trial of resistance training added to caloric restriction plus aerobic exercise training in obese heart failure with preserved ejection fraction. *Circ Heart Fail*. 2023;16:e010161.
5. Fiatarone MA, O'Neill EF, Ryan ND, et al. Exercise training and nutritional supplementation for physical frailty in very elderly people. *N Engl J Med*. 1994;330:1769-1775.