



This is a repository copy of *Patient-reported outcomes and prognostic factors in ankle fracture-dislocation: a systematic review*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/212063/>

Version: Published Version

Article:

Ahmed, R. orcid.org/0000-0003-1758-2708, Ward, A.E. orcid.org/0000-0002-9609-830X and Thornhill, E. (2024) Patient-reported outcomes and prognostic factors in ankle fracture-dislocation: a systematic review. *Trauma*, 26 (2). pp. 113-123. ISSN 1460-4086

<https://doi.org/10.1177/14604086231183582>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Patient-reported outcomes and prognostic factors in ankle fracture-dislocation: A systematic review

Trauma
2024, Vol. 26(2) 113–123
© The Author(s) 2023



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/14604086231183582

journals.sagepub.com/home/tra



Ramsha Ahmed , Alex E Ward  and Elizabeth Thornhill

Abstract

Objectives: Ankle fractures have an incidence of around 90,000 per year in the United Kingdom. They affect younger patients following high energy trauma and, in the elderly, following low energy falls. Younger patients with pre-existing comorbidities including raised BMI or poor bone quality are also at risk of these injuries which impact the bony architecture of the joint and the soft tissues leading to a highly unstable fracture pattern, resulting in dislocation. At present, there is no literature exploring what effect ankle fracture-dislocations have on patients' quality of life and activities of daily living, with only ankle fractures being explored.

Methods: Relevant question formatting was utilised to generate a focused search. This was limited to studies specifically mentioning ankle injuries with a focus on ankle fracture-dislocations. The number of patients, fracture-dislocation type, length of follow up, prognostic factors, complications and outcome measures were recorded.

Results: Nine hundred and thirty-nine fractures were included within the studies. Eight studies looked at previously validated foot and ankle scores, two primarily focused on the American Orthopaedic Foot and Ankle Society score (AOFAS), three on the foot and ankle outcome score (FAOS), and one study on the Olerud–Molander score (OMAS). Patient, injury, and management factors were identified as being associated with poorer clinical outcomes.

Conclusions: Not only are age and BMI a risk factor for posttraumatic osteoarthritis but they were also identified as prognostic indicators for functional outcome in this review. Patients sustaining a concurrent fracture-dislocation were found to have poorer clinical outcomes, and the timing and success of reduction further influenced outcomes. This review found that the quality of reduction was directly related to the patients' functional outcomes post-follow up, and the risk of developing posttraumatic osteoarthritis, which was more frequent in patients sustaining Bosworth fractures, posterior malleolar fractures, and in patients with increasing age.

Level of evidence: IV.

Keywords

Ankle fracture, ankle fracture dislocation, open fracture, open reduction internal fixation, American Orthopaedic Foot and Ankle Society score, Olerud–Molander foot and ankle score, posttraumatic osteoarthritis, prognostic factor, outcome

Introduction

Ankle fractures have an incidence of approximately 90,000 per year in the United Kingdom. They have a bimodal distribution, affecting younger patients following high energy trauma and, in the elderly, following low energy falls. Younger patients with pre-existing comorbidities such as raised body mass index (BMI) or poor bone quality are also at risk of these injuries^{1,2} which not only affect the bony architecture of the joint but also the soft tissues, resulting in a highly unstable fracture pattern associated with dislocation. When the soft tissues fail, this may result in an open fracture and add to the risk profile.³

Current UK practice advises for all ankle fracture-dislocations to be reduced urgently within the emergency department (ED), minimising the risk of further soft tissue

Department of Trauma and Orthopaedics, Sheffield Teaching Hospitals, Sheffield, South Yorkshire, UK

Corresponding author:

Ramsha Ahmed, The University of Sheffield, Academic Department of Medical Education, Beech Hill Road, Sheffield, South Yorkshire S10 2RX, UK.

Email: rahmed9@sheffield.ac.uk

injury and oedema. The British Orthopaedic Association Standards for Trauma and Orthopaedics (BOAST) 12 guidance recommends a pre-reduction radiograph unless this would cause an unacceptable delay.⁴ Ankle fracture-dislocations are commonly managed with either primary open reduction and internal fixation (ORIF), or staged management with a period of external fixation prior to definitive surgery, allowing recovery of soft tissues. The definitive use of external fixation is indicated for patients with a poor soft tissue envelope, or those not medically fit for a second procedure.

Previous studies have illustrated an association between ankle fracture-dislocations and post-traumatic osteoarthritis (PTOA) due to imperfections in the articular surface and changing joint biomechanics. This can affect the individual's ability to return to work and increases the risk of further operative intervention in the form of ankle arthroplasty or fusion.⁵ Subsequently, the effect of changing the joint biomechanics increases the potential for further fusions, including those of the talocalcaneal and talonavicular joints.⁶

Existing literature reviews have identified prognostic factors associated with outcome progression in patients with non-displaced ankle fractures. These include osteochondral lesions resulting in a lower threshold for recurrent injury⁷ and early weight-bearing which improves functionality.⁸ This review aims to explore the current evidence for outcomes of ankle fracture-dislocations and the available prognostic evidence.

Methods

Data sources

This systematic review was written in accordance to the PRISMA guidelines (Figure 1). Relevant question formatting was utilised to generate a focused search using PICO: population, intervention, comparison, and outcome. This guided the selection of literature sources and provided a focus for the search. Our population, as well as search limitation, consisted of patients over the age of 16 who sustained ankle fracture-dislocations. Patients receiving operative intervention were compared to those who were managed conservatively to identify factors associated with poorer prognosis. The assessment of functional outcomes was derived using appropriate foot and ankle scoring systems, weight-bearing status, and radiographic evaluation. An extensive search strategy (Figure 1) was employed using Boolean operators. Computerised databases consisted of Wolters Kluwer OvidSP Database (1733 papers), Cochrane Library (4 papers), and EBSCOhost (5 papers).

The literature search was limited to studies specifically mentioning ankle injuries with a focus on ankle fracture-dislocations. The exclusion criteria consisted of studies

which did not utilise foot and ankle scoring systems to assess functional outcomes, no data on completion, and a focus solely on ankle fractures without dislocation. Articles reporting a mixed series of non-displaced ankle fractures and ankle fracture-dislocations were excluded if the qualitative and quantitative analysis between the two subgroups could not be distinguished. Papers focussing on surgical techniques were also excluded.

Study selection

During the screening stage, the title and abstract of prospective articles were assessed. Studies which either demonstrated the inclusion criteria, or insufficient detail, progressed to the successive stage. Full paper screening of these studies then took place with two authors (AEW and RA) assessing which studies met the inclusion criteria (Table 1). In a situation of disagreement, the final decision followed a discussion between both authors to reach a consensus. Of the initial 1742 papers, 9 met our inclusion criteria. Figure 1 demonstrates the screening process.

Data extraction

The data of the included studies were extracted by one author (RA) and verified by the second author (AEW). Only data from the published articles were included in the analysis with the authors not being contacted for further details. Studies were assigned their level of evidence as defined by the Centre for Evidence-Based Medicine.⁹ Relevant information including the population sample, pattern of fracture-dislocation, length of follow up, prognostic factors, complications and outcome measures were recorded.

Risk of bias in individual studies

During the full paper screening, obvious sources of bias were recorded (Figure 2) including funding and affiliations. These are discussed within the limitations section of this review. A specific bias screening tool was not utilised.

Data synthesis and statistics

Using methods identified by Stufkens et al., prognostic indicators were assigned a grade A to D which associates their clinical value.¹¹ These gradings are outlined in Table 2. Papers were compared using the functional outcomes scores, and the most frequently used functional outcome score was used to prevent duplication within a single paper utilising multiple clinical outcome scores.




Figure 1. PRISMA flow diagram.

Results

The included nine studies totalled 939 ankle fractures. Amongst these papers, three were Level III^{12–14} studies and the remainder were Level IV.^{15–20} (Table 3). Two papers focussed on prognostic factors associated with poorer outcomes in ankle fracture dislocations^{15,19} and three papers identified risk factors associated with PTOA.^{16,18,20}

Four of the included nine studies did not report the number of patients initially identified.^{14,15,18,20} Of those patients in the remaining five studies (n = 670), 86.71% (n = 581) completed follow-up.^{12,13,16,17,19} Six papers reported a follow-up range between 3 and 128 months.^{12,13,15–18}

Functional outcomes

Three studies used the American Orthopaedic Foot and Ankle Society (AOFAS) score to assess the clinical outcomes of different types of fracture-dislocation. The score is out of 100 and consists of nine questions, focussing on three areas: pain (40 points), function (50 points), and alignment (10 points). Scores of more than 90 show an ‘excellent’ result, more than 80 show ‘good’ result, but an acceptable result is debated.²¹

Cho et al.¹⁵ further subdivided the injuries as standard Danis–Weber B or C fracture-dislocations and Bosworth fracture-dislocations. These injuries are defined as ‘fracture-dislocation of the ankle with fixed displacement

of the proximal fragment of the distal fibular fracture behind the posterior tibial tubercle¹⁵ and are commonly caused by severe external rotation with supination and can be irreducible unless opened.²²

Table 1. Inclusion and exclusion criteria.

Inclusion criteria	Exclusion Criteria
Population <ul style="list-style-type: none"> • Patients over 16-year-old • Patients with ankle fracture dislocation/s 	<ul style="list-style-type: none"> • Studies with absent foot and ankle scores • Absent data on study completion • Ankle fracture/s with no dislocation/s • Mixed series of ankle fracture/s and ankle fracture dislocation/s • Studies with a sole focus on surgical techniques
Intervention <ul style="list-style-type: none"> • Operative management 	
Comparison <ul style="list-style-type: none"> • Conservative management 	
Outcome <ul style="list-style-type: none"> • Foot and ankle scores • Weightbearing status • Radiographic evaluation 	

Shou et al.¹⁸ and Wang et al.²⁰ examined the outcomes for the Log-splitter fracture-dislocation. Both assessed the typical pattern of injury caused by a vertical axial load and illustrated that when anatomically reduced intraoperatively, these injuries had a similar ‘acceptable outcome’ (75.33 ± 6.53 vs 75.05 ± 13.86).

Testa et al.¹⁹ compared the outcomes of ankle-fracture dislocations according to the Danis–Weber classification.²³ Weber B injuries (those at the level of the syndesmosis), and Weber C (those above the syndesmosis), both had a significant improvement at the end of the 12-month follow up ($78.67, 45\text{--}100$ vs $69.72, 35\text{--}100, p=0.03$). Patients did suffer with residual ankle pain, stiffness and swelling.

Table 2. Clinical relevance gradings.¹⁰

Stufkens grade	Description
A	Treatment options or prognosis are supported by strong evidence (consistent with Level I or II studies)
B	Treatment options or prognosis are supported by fair evidence (consistent with Level III or IV studies)
C	Treatment options or prognosis are supported by either conflicting or poor quality evidence (Level IV studies)
D	When insufficient evidence exists to make a recommendation




Figure 2. Risk of bias.¹⁰

Table 3. Overall outcomes.

Study, year, level of evidence	Included (n)	Type of injury	F/U range in months (mean)	Outcome parameters	Scores at final follow up	Complications	Conclusion
Cho BK et al. ¹⁵ 2018 IV	15 Bosworth 25 control	Bosworth fracture dislocations vs control (Bimalleolar and trimalleolar ankle fracture-dislocations. Weber classification (types B and C))	24–109 (mean 66.1 ± 28.8)	AOFAS 0–100 OMAS	AOFAS at final F/U Bosworth 87.8 ± 11.7 vs Control 89.1 ± 10.2. NS OMAS at final F/U Bosworth 88.4 ± 10.8 vs Control 89.8 ± 9.6. NS	Ankle stiffness, Non-union, Wound infection, skin necrosis, Compartment syndrome. All NS	Delayed surgical reduction and repeated attempts at closed reduction appear to be prognostic factors for poorer outcomes.
Lindsjö U ¹⁶ 1985 IV	Total 298 Weber B 229 Weber C 66	Weber classification (types B and C)	24–72 (Not known)	Association for the Study of Internal Fixation principles; excellent, good, acceptable, poor. Computer analysis (AID)	82% excellent and good, 8% acceptable, 10% poor	Posttraumatic osteoarthritis NS	Open reduction internal fixation seems to be the best method of treatment for obtaining accurate anatomical restoration and good function.
Sculco PK et al. ¹⁷ 2015 IV	Total 108 SER IV dislocated 35 SER IV non-dislocated 73	SER IV ankle fracture dislocation	12–52 (mean 21)	FAOS, 0–100; symptoms, pain, ADL, sport, QOL. Subtalar range of motion (ROM); normal inversion, normal eversion, normal plantarflexion, normal dorsiflexion.	Accuracy of articular reduction – dislocation 28 vs non-dislocation 71* FAOS pain – dislocation 84.0 vs non-dislocation 73.0* FAOS symptoms – dislocation 69.4 vs non-dislocation 75.9. NS FAOS ADL – dislocation 78.1 vs non-dislocation 87.7* FAOS sport – dislocation 55.1 vs non-dislocation 65.5. NS FAOS QOL – dislocation 53.4 vs non – dislocation 60.5. NS ROM normal inversion – dislocation 10 vs non-dislocation 42* ROM normal eversion – dislocation 15 vs non-dislocation 49* ROM normal plantarflexion – dislocation 10 vs non-dislocation 52* ROM normal dorsiflexion – dislocation 14 vs non-dislocation 43 NS	Post-operative infections, wound complications. All NS	Concurrent fracture dislocations are associated with worse radiographic and functional outcomes, without an increase in superficial or deep infections.
Shou K et al. ¹⁸ 2020 IV	Total 31 Intraoperative anatomical reduction 24	Log Splitter injury – High energy Transsyndesmotoc	13–26 (mean 22.9 ± 3.3)	ROM ankle joint, AOFAS, 0–100, BCS (good/fair/poor), K-L (0–4).	ROM plantarflexion – reduction 27.52 ± 3.72 vs non-reduction 21.24 ± 2.89* ROM dorsal expansion – reduction 12.47 ± 4.28 vs 9.04 ± 4.18*	Posttraumatic osteoarthritis. Reduction 8 vs non-reduction 6*	The quality of intraoperative reduction for the treatment of Log

Table 3. Continued.

Study, year, level of evidence	Included (n)	Type of injury	F/U range in months (mean)	Outcome parameters	Scores at final follow up	Complications	Conclusion
	Intraoperative non-anatomical reduction 7	ankle fracture dislocation			ROM eversion – reduction 11.51 ± 2.84 vs $8.93 \pm 3.91^*$ ROM inversion – reduction 22.63 ± 2.85 vs non-reduction $17.68 \pm 3.37^*$ AOFAS – reduction 75.33 ± 6.53 vs non-reduction $66.89 \pm 4.28^*$ K-L score – reduction 0.62 vs non-reduction 1.83* BCS – 16 good, 7 fair, 8 poor. For all patients.		Splitter injury is scientifically significant and can be utilised as a major factor to predict clinical outcome. Compared to cause and functional outcome, this may play a major role in functional outcome.
Tantigate T et al. ¹² 2019 III	Total 118 Fracture dislocation 33 Non-dislocation 85	Bimalleolar and trimalleolar ankle fracture-dislocations. Weber classification (types B and C)	12–76 (41 ± 12 vs 39 ± 14)	FAOS 0–100; symptom, pain, ADL, sport, QOL. Quality of reduction.	FAOS symptom – dislocation 73.21 vs non-dislocation 85.71. NS FAOS pain – dislocation 76.39 vs 91.67* FAOS ADL – dislocation 86.77 vs non-dislocation 95.59. NS FAOS sport – dislocation 55.00 vs non-dislocation 85.00. NS FAOS QOL – dislocation 50.00 vs non-dislocation 62.50. NS	Post-operative wound complications: surgical site infection, wound erythema, wound dehiscence. All NS	Ankle fracture dislocations are more prevalent in patients who are elderly, female, and diabetic.
Testa G et al. ¹⁹ 2019 IV	Total 48 Weber B 30 Weber C 18	Trimalleolar ankle fracture dislocations (Weber types B and C)	12 (Not known)	Visual Analogue Scores (VAS), Olerud–Molander Scores(OMAS).	VAS – Weber B 1.73 vs Weber C 3.06* OMAS – Weber B 78.67 vs Weber C 69.72*	Residual ankle pain, residual ankle stiffness, residual ankle swelling	Patients with one or more negative factors (aged over 61 years, BMI > 40, class ASA > 1, type C fracture according to Danis–Weber or a fracture dislocation) could have worse outcomes
Wang Z et al. ²⁰ 2017 IV	Total 41 Typical 19 Untypical 22	Log Splitter fracture-dislocation (typical – vertical axial load, untypical –	Not Known (7.48 ± 5.56)	AOFAS, 0–100, BCS, range of motion (ROM, postoperative	AOFAS – typical 75.05 ± 13.86 vs untypical 81.55 ± 5.60 . NS ROM dorsal expansion – typical 23.84 ± 2.11 vs untypical $26.14 \pm 2.10^*$	Posttraumatic osteoarthritis* Post-operative infection,	The ORIF procedure may be an optimal approach to treat Log-splitter injuries.

(continued)

Table 3. Continued.

Study, year, level of evidence	Included (n)	Type of injury	F/U range in months (mean)	Outcome parameters	Scores at final follow up	Complications	Conclusion
		rotational mechanism)		tibiofibular width, (mm).	ROM plantarflexion – typical 25.58 ± 3.10 vs untypical $30.27 \pm 2.33^*$ ROM eversion – typical 22.11 ± 3.62 vs untypical $27.59 \pm 2.40^*$ ROM inversion – typical 22.84 ± 3.10 vs $28.32 \pm 3.68^*$ BCS – typical 6 good/10 fair/3 poor vs untypical 16 good/6 fair/0 poor. Postoperative tibiofibular width – typical 17.24 ± 3.30 vs untypical $9.69 \pm 2.55^*$	post-operative non-union. All NS	Patients with untypical injury had better fracture reduction and range of motion and low incidence rate of posttraumatic arthritis compared to patients with untypical injuries.
Warner S] et al. ¹³ 2015 III	Total 47 Fracture dislocation 20 Non-dislocation 27	PER IV ankle fracture dislocation	12–128 (30)	FAOS 0–100; (symptom, pain, ADL, QOL, ROM), radiographic imaging; articular malreduction (>2 mm joint surface gap), syndesmotric malreduction (>2 mm distance between anterior and posterior incisura).	FAOS symptoms – dislocation 46 vs non-dislocation 70* FAOS pain – dislocation 56 vs non-dislocation 82* FAOS ADL – dislocation 61 vs non-dislocation 84* FAOS sports – dislocation 37 vs non-dislocation 59 NS FAOS QOL – dislocation 25 vs non-dislocation 59* ROM dorsiflexion – dislocation 15 vs non-dislocation 18. NS ROM plantarflexion – dislocation 41 vs non-dislocation 49. NS ROM eversion – dislocation 21 vs non-dislocation 24. NS ROM inversion – dislocation 24 vs non-dislocation 30. NS Articular malreduction – dislocation 6 vs non-dislocation 3. NS Syndesmotric malreduction – dislocation 8 vs non-dislocation 10. NS	Post-operative infection, wound complication. All NS	PER VI fracture dislocations have a higher rate of articular malreduction and poorer functional outcomes than non-dislocated PER IV fractures
Wicks L et al. ¹⁴	Total 233 Group A 62	Lauge-Hansen classification	Not reported	OMAS, LEFS.	OMAS – group A 64.4 vs group B 72* LEFS – group A 48.1 vs group B 56.5. NS	Postoperative wound	The time taken to achieve a reduced

(continued)

Table 3. Continued.

Study, year, level of evidence	Included (n)	Type of injury	F/U range in months (mean)	Outcome parameters	Scores at final follow up	Complications	Conclusion
2018 III	Group B 51 Group C 110 analysis)					infection. NS	ankle mortise was significantly longer for patients who had a radiograph before manipulation with the potential for exacerbating soft tissue trauma.

Summary of studies (AOFAS, American Orthopaedic Foot and Ankle Society score; OMAS, Olerud–Molander score; FAOS, foot and ankle outcome score; VAS, visual analogue scale; BCS, Burwell–Charnley score; K–L, Kellergren–Lawrence criteria; LEFS, lower extremity functional scale; ROM, range of movement; QOL, quality of life; ADL, activities of daily living; NS, not significant; *, significant).

Wicks et al.¹⁴ assessed ankle fracture dislocations using the Lauge–Hansen classification²⁴ but the functional outcomes were not grouped according to this, but rather whether they underwent manipulation according to clinical groups alone or if they had a radiograph performed prior to reduction. This method was used to assess outcomes according to the BOAST 12 which suggests ‘*Reduction and splinting should be performed urgently for clinically deformed ankles. Radiographs should be obtained before reduction unless this will cause an unacceptable delay*’.⁴ Patients in group B (patients undergoing manipulation in the ED based on clinical grounds alone), had better outcomes than those in group A (patients receiving an X-ray before any attempt at ankle reduction) (72, 25–100 vs 64.4, 20–90, $p=0.039$).

Two papers assessed the injuries according to the Lauge–Hansen Classification.²⁴ The results were significant for the pronation-external rotation (PER) injuries studied by Warner et al.¹³ Of the 47 injuries assessed, the 20 fracture-dislocations illustrated significantly worse outcomes than the 27 fractures alone across the symptom, pain, ADL and quality of life (QOL) domains (Symptoms 70 ± 23.3 vs 46 ± 21.7 , $p=0.002$; Pain 82 ± 20.1 vs 56 ± 20.7 , $p \leq 0.001$; ADL 84 ± 21.6 vs 61 ± 23.3 , $p=0.002$; QOL 59 ± 29.4 vs 25 ± 21.3 , $p \leq 0.001$).

Fracture reduction as a prognostic indicator

Across the nine studies, a range of scores and measurements were used to assess whether the ankle fracture-dislocations were reduced successfully following operative intervention. Lindsjö¹⁶ judged the acceptability of the reduction following operative intervention but did not state what criteria were used; 90% of the fractures were satisfactorily reduced, and this was linked to better clinical outcomes on combined subjective clinical and examination findings (Stufkens Grade C).^{11,16}

Other studies did outline clear measurements the authors had used on postoperative films to identify whether the fracture had been effectively reduced. Warner et al.¹³ used the tibiofibular distance to judge how well the syndesmosis had been reduced, as well as judging the articular reduction on postoperative CT scans. Syndesmotic malreduction was common, in not only fracture-dislocations but also in fractures alone (44.4% vs 47.6%, $p=0.95$). Articular malreduction was more common in fracture-dislocations (33.3% vs 14.3%, $p=0.15$) (Stufkens Grade C).^{11,13}

Two studies used the Burwell–Charnley score to identify fractures that were unsatisfactorily reduced.^{19,20,25} This scoring system classifies fractures into those with anatomical, fair, and poor reduction based on the degree of postoperative displacement.⁶ Both studies found that the quality of intraoperative reduction was not only linked to significantly better short-term outcomes, as measured by the AOFAS, but also to significantly lower rates of early

post-traumatic arthritis after completion of the 2-year follow-up period (Stufkens Grade B).^{11,18,20}

Other prognostic factors associated with poorer outcomes

Patient, injury and management factors were identified as being associated with poorer clinical outcomes. Testa et al. found that patients with a BMI of over 40 as well as those above the age of 61 years experienced poorer outcomes (AOFAS) at the end of the 12-month follow-up period.¹⁹ The authors also found that patients with comorbidities which caused them to be assigned an American Society of Anaesthesiologists (ASA) grade of greater than one were more likely to have poorer clinical outcomes (Stufkens Grade C).^{11,19}

As for the fracture personality, unsurprisingly patients who had a concurrent fracture-dislocation were found to have poorer clinical outcomes than those with an ankle fracture alone (Stufkens Grade C).^{11,17,19} Patients with these injuries had poor reduction on radiographs during follow up (Stufkens Grade B).^{11,19,26} Patients with Weber C fracture-dislocations were also identified as having poorer clinical function owing to the greater disruption of the syndesmosis (Stufkens Grade C).^{11,27}

Within the ED, the timing and success of reduction were found to be important factors influencing outcomes. Wicks et al. found that patients with ‘frankly displaced ankle fracture-dislocations’ who waited longer for reduction, due to a prolonged waiting time for pre-reduction radiographs, had a significantly worse clinical function at the end of the study period. OMAS and FAOS were used to assess the functional outcome in this study (Stufkens Grade C).^{11,15} This was also true for patients who required multiple attempts to achieve adequate closed reduction (Stufkens Grade C).^{11,16} One study did demonstrate how prolonged reduction time for ankle fracture-dislocations leads to poorer functional outcomes.¹⁴

Complications associated with ankle-fracture dislocations

Soft tissue complications were reported across six of the nine studies.^{12–15,17,20} Primarily, the rates of superficial wound infections were described in five studies, but when compared to patients who suffered fractures without dislocation, there were no significant differences (Stufkens Grade D).^{11–15,17,20} Oral antibiotics and better surgical site hygiene were described as adequate solutions for the management of these infections.¹⁹

Other soft tissue complications included the impact of managing the patient in casts. Lindsjö et al. reported one patient suffering with a pressure sore and a second having

a hypersensitivity reaction to plaster of Paris (Stufkens Grade D).^{11,16}

Thromboembolic events were reported in two studies, although these were rare. Lindsjö¹⁶ described two patients (0.006%) who were treated for deep venous thrombosis (DVTs), both of which were mobile throughout their treatment in a cast. The use of chemical thromboprophylaxis was not discussed. According to Tantigate et al.,¹² two patients suffered from thromboembolic events in the non-dislocated cohort (2.4%), in comparison to the dislocated subgroup ($p=0.42$) which reported no patients suffering this complication (Stufkens Grade D).^{11,12}

Three studies focussed on factors associated with PTOA.^{16,18,20} Lindsjö¹⁶ found that the incidence of PTOA was significantly higher in patients over the age of 35 years and in those whose injury included a posterior malleolus fracture. The quality of reduction was also associated with the likelihood of developing PTOA, those with excellent or good reduction being less likely to suffer in the future.³

Discussion

Patients sustaining severe ankle fractures including PER and Weber C, have worse clinical outcomes when compared to patients sustaining less severe injuries.^{11,13,27} It is therefore no surprise that fracture dislocations have worse outcomes than fractures alone.^{15,19}

The quality of fracture reduction in the included studies was directly related to the patient's functional outcomes at the end of follow up.^{13,15,18} This has been previously observed in studies focussing on pilon fractures whereby a better-quality reduction is associated with better functional outcome scores at 2 years following injury.^{26,27} This systematic review also found that the quality of reduction is directly related to the risk of developing PTOA in the future, and was more frequent amongst patients sustaining Bosworth fractures, those with a posterior malleolar fracture, and those over the age of 35 years.^{16,19} This adds to the existing literature which demonstrates that the risk factors for PTOA include a high BMI, length of time since surgery, sustaining a Weber C fracture, and associated medial malleolar fracture.²⁸

Apart from being identified as a risk factor for PTOA, two studies identified increasing patient age^{11,18} and BMI¹⁸ as significant prognostic indicators for functional outcomes. This result mirrors a Swedish study which assessed the outcomes and QOL in patients over 65 years sustaining ankle fractures alone without dislocation, who underwent operative management. These patients experienced higher rates of difficulties with pain and reduced mobility when compared to younger patients at the end of a 2-year follow-up.²⁹

An ASA score of more than one was also found to be associated with a poor functional outcome in ankle fracture-dislocations.¹⁹ This trend has previously been observed in studies which illustrate that patients with a higher ASA had poorer functional outcomes at 1 year following ankle fracture injury.³⁰

At present, there are no qualitative studies exploring what effect ankle fracture-dislocations have on patients QOL and ADLs, with only ankle fractures being explored. One such qualitative study interviewed patients on their individual experiences of ankle fractures and essential factors personal to their recovery, and reported the impact on factors including ADLs, sleep disturbance, and social life. Participants described struggling with personal care resulting in the necessity of adapting to new routines. This was discussed in relation to individuals' weightbearing status and mobility aids. With regards to QOL, several participants reported negative impacts on social and family life, including psychological effects associated with depression and anxiety.⁵

Within this study, an analysis of bias was conducted according to Ramírez-Santana.¹⁰ Three studies had a greater risk of selection bias. The recruitment of the control group in the study by Cho et al.¹⁵ was not discussed at length, leading to potential bias in individual case selection.^{3,8,26} In the study by Lindsjö,¹⁶ it was not clear how much the patients' activities were affected by their injury within the follow-up period, having been assigned an outcome rated from poor to excellent by clinicians without a validated questionnaire being employed. The recruitment of the study group was not discussed at length in the study by Warner et al.¹² There was no evidence of confusion bias in any of the included studies. Hence, the effect of information bias in this study remains uncertain.¹⁴

The limitations of this review include the number of eligible studies, small patient cohorts, and the use of different functional outcome scores. These factors make it difficult to directly compare outcomes across the studies and therefore meaningful conclusions cannot be drawn due to low-level evidence. The results derived from this systematic review can be utilised in the development of future randomised control trials or prospective cohort studies with a similar focus.

Conclusion

Ankle fracture-dislocations are severe injuries which can be associated not only with poor clinical outcomes as assessed by validated foot and ankle scores, but also with an increased risk of PTOA. In addition to being identified as a risk factor for PTOA, increasing patient age and BMI were also concluded as prognostic indicators for poorer functional outcome in this systematic review.

Conflict of interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Ethical approval

Not applicable.

Informed consent

Not applicable.

Trial registration

Not applicable.

ORCID iDs

Ramsha Ahmed  <https://orcid.org/0000-0003-1758-2708>

Alex E Ward  <https://orcid.org/0000-0002-9609-830X>

References

- King CM, Hamilton GA, Cobb M, et al. Association between ankle fractures and obesity. *J Foot Ankle Surg* 2012; 51: 543–547.
- Therdyothin A, Phiphophatsanee N, Wajanavisit W, et al. Is ankle fracture related to low bone mineral density and subsequent fracture? A systematic review. *Osteoporosis Sarcopenia* 2020; 6: 151–159.
- Simske NM, Audet MA, Kim CY, et al. Open ankle fractures are associated with complications and reoperations. *OTA Int* 2019; 2: e042.
- The Management of Ankle Fractures, <https://www.boa.ac.uk/uploads/assets/f8b1c499-c38a-4805-8cb8d8eb3087bca7/8be763eb-5921-4cb2-b6802f3e65ce8e7f/the%20management%20of%20ankle%20fractures.pdf> (2016, accessed 7 July 2022).
- McKeown R, Kearney RS, Liew ZH, et al. Patient experiences of an ankle fracture and the most important factors in their recovery: a qualitative interview study. *BMJ Open* 2020; 10: e033539.
- Wang Y, Li Z, Wong DW, et al. Effects of ankle arthrodesis on biomechanical performance of the entire foot. *PLoS One* 2015; 10: e0134340.
- Pina G, Fonseca F, Vaz A, et al. Unstable malleolar ankle fractures: evaluation of prognostic factors and sports return. *Arch Orthop Trauma Surg* 2021; 141: 99–104.
- Lorente A, Gandía A, Mariscal G, et al. Quality of life and complications in elderly patients after pronation rotation type III ankle fractures treated with a cast and early weight-bearing. *BMC Musculoskelet Disord* 2021; 22: –8.
- The Centre for Evidence-Based Medicine (CEBM), <https://www.cebm.net/> (2019, accessed 7 July 2022).
- Ramirez-Santana M. *Limitations and biases in cohort studies*. London: IntechOpen, 2018.
- Stufkens SAS, van de Bekerom MPJ, Kerkhoffs GMMJ, et al. Long-term outcome after 1822 operatively treated ankle fractures: a systematic review of the literature. *Injury* 2010; 42: 119–127.
- Tantigate D, Ho G, Kirschenbaum J, et al. Functional outcomes after fracture-dislocation of the ankle. *Foot Ankle Specialist* 2020; 13: 18–26.
- Warner SJ, Schottel PC, Hinds RM, et al. Fracture-dislocations demonstrate poorer postoperative functional outcomes among pronation external rotation IV ankle fractures. *Foot Ankle Int* 2015; 36: 641–647.
- Wicks L, Faroug R and Mangwani J. Should pre-manipulation radiographs be obtained in ankle fracture-dislocations? *Foot* 2018; 1: 10–14.
- Cho BK, Choi SM and Shin YD. Prognostic factors for intermediate-term clinical outcomes following Bosworth fractures of the ankle joint. *Foot Ankle Surg* 2019; 25: 601–607.
- Lindsjö U. Operative treatment of ankle fracture-dislocations. A follow-up study of 306/321 consecutive cases. *Clin Orthop Relat Res* 1985; 199: 28–38.
- Sculco PK, Lazaro LE, Little MM, et al. Dislocation is a risk factor for poor outcome after supination external rotation type ankle fractures. *Arch Orthop Trauma Surg* 2016; 136: 9–15.
- Shou K, Adhikary R, Zou L, et al. The assessment of the reduction algorithm in the treatment for “Logsplitter” injury. *BioMed Res Int* 2020; 2020: 4139028.
- Testa G, Ganci M, Amico M, et al. Negative prognostic factors in surgical treatment for trimalleolar fractures. *Eur J Orthop Surg Traumatol* 2019; 29: 1325–1330.
- Wang Z, Tang X, Li S, et al. Treatment and outcome prognosis of patients with high-energy transsyndesmotic ankle fracture dislocation – the “Logsplitter” injury. *J Orthop Surg Res* 2017; 12: –0.
- Ceccarelli F, Calderazzi F and Pedrazzi G. Is there a relation between AOFAS ankle-hindfoot score and SF-36 in evaluation of Achilles ruptures treated by percutaneous technique? *J Foot Ankle Surg* 2014; 53: 16–21.
- Peterson ND, Shah F and Narayan B. An unusual ankle injury: the Bosworth-Pilon fracture. *J Foot Ankle Surg* 2015; 54: 751–753.
- Gaillard F, Qureshi P, Roberts D, et al. Weber classification of ankle fractures. Reference article, Radiopaedia.org (accessed 12 January 2023).
- Tartaglione JP, Rosenbaum AJ, Abousayed M, et al. Classifications in brief: Lauge–Hansen classification of ankle fractures. *Clin Orthop Relat Res* 2015; 473: 3323–3328.
- Burwell HN and Charnley AD. The treatment of displaced fractures at the ankle by rigid internal fixation and early joint movement. *J Bone Joint Surg Br* 1965; 47: 634–660.
- Harris AM, Patterson BM, Sontich JK, et al. Results and outcomes after operative treatment of high-energy tibial plafond fractures. *Foot Ankle Int* 2006; 27: 256–265.
- Lawson KA, Ayala AE, Morin ML, et al. Ankle fracture-dislocations: a review. *Foot Ankle Orthop* 2018; 3: 247301141876512.
- Lübbeke A, Salvo D, Stern R, et al. Risk factors for post-traumatic osteoarthritis of the ankle: an eighteen year follow-up study. *Int Orthop* 2012; 36: 1403–1410.
- Nilsson G, Jonsson K, Ekdahl C, et al. Outcome and quality of life after surgically treated ankle fractures in patients 65 years or older. *BMC Musculoskelet Disord* 2007; 8: –9.
- Egol KA, Tejwani NC, Walsh MG, et al. Predictors of short-term functional outcome following ankle fracture surgery. *J Bone Joint Surg* 2006; 88: 974–979.