

This is a repository copy of *Good functions, measures, and the Kleinbock-Tomanov conjecture*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/212388/>

Version: Accepted Version

Article:

Beresnevich, Victor orcid.org/0000-0002-1811-9697, Datta, Shreyasi and Ghosh, Anish (2024) Good functions, measures, and the Kleinbock-Tomanov conjecture. *Journal für die reine und angewandte Mathematik*. ISSN 1435-5345

<https://doi.org/10.1515/crelle-2024-0052>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

GOOD FUNCTIONS, MEASURES, AND THE KLEINBOCK-TOMANOV CONJECTURE

VICTOR BERESNEVICH, SHREYASI DATTA, AND ANISH GHOSH

ABSTRACT. In this paper we prove a conjecture of Kleinbock and Tomanov [13, Conjecture FP] on Diophantine properties of a large class of fractal measures on \mathbb{Q}_p^n . More generally, we establish the p -adic analogues of the influential results of Kleinbock, Lindenstrauss and Weiss [11] on Diophantine properties of friendly measures. We further prove the p -adic analogue of one of the main results in [10] concerning Diophantine inheritance of affine subspaces, which answers a question of Kleinbock. One of the key ingredients in the proofs of [11] is a result on (C, α) -good functions whose proof crucially uses the Mean Value Theorem. Our main technical innovation is an alternative approach to establishing that certain functions are (C, α) -good in the p -adic setting. We believe this result will be of independent interest.

CONTENTS

1. Introduction	1
2. Main results	2
3. Preliminaries	3
4. Proofs of Theorem 2.1, Theorem 2.2 and Theorem 2.3	8
5. Notes on the proof of Proposition 4.1	9
6. Good functions w.r.t. absolutely decaying measures	12
7. Polynomials satisfying the hypotheses in Theorem 6.1	17
8. Approximation by polynomials	21
9. Nondegeneracy and normalization	24
10. Completing the proof of Proposition 4.1	26
11. Appendix	27
References	32

1. INTRODUCTION

The problems considered in this paper go back to the landmark work of Kleinbock and Margulis [12] from 1998 which settled the Baker-Sprindžhuk conjecture and ushered in major sustained progress in the area of Diophantine approximation on manifolds. Several years later Kleinbock, Lindenstrauss and Weiss [11] transformed the area by introducing the general framework of Diophantine properties of measures and developing the relevant Diophantine theory for a large natural class of measures on \mathbb{R}^n which they named *friendly measures*. Specifically, they proved that almost every point in the support of a friendly measure is not very well approximable by rationals. Since the class of friendly measures includes smooth measures on

AG was supported by the Government of India, Department of Science and Technology, Swarnajayanti fellowship DSTSJFMSA-01/2016-17 and a grant from the Infosys Foundation. AG acknowledges support of the Department of Atomic Energy, Government of India [under project 12 - R&D - TFR - 5.01 - 0500].

Mathematics Subject Classification (2010). Primary 11J83; Secondary 11J54, 11J61, 37A45.

nondegenerate manifolds, [11] generalizes the results of Kleinbock and Margulis [12]. Moreover, there are many other interesting examples of friendly measures such as measures supported on self-similar sets satisfying the open set condition [11, 20], Patterson-Sullivan measures associated with convex cocompact Kleinian groups [18] and Patterson-Sullivan measures associated with some geometrically finite groups [3].

In another direction, Kleinbock and Tomanov [13] proved p -adic, and more generally S -arithmetic, versions of many of the results from [11]. In particular, they established the S -arithmetic analogue of the Baker-Sprindžhuk conjecture. In the same paper, Kleinbock and Tomanov conjectured that p -adic analogues of the main results in [11] should hold, see Conjecture FP in [13], where ‘FP’ stands for ‘Friendliness of Pushforwards’. Conjecture FP requires establishing certain properties of pushforwards of measures subject to natural constraints, and via the theory developed in [13], it leads to a coherent theory of Diophantine approximation over \mathbb{Q}_p . In this paper we settle this conjecture in full and prove several related results. Recently, there has been considerable progress in understanding p -adic Diophantine approximation on manifolds. We refer the reader to [4, 5, 6] for its account.

One of the main technical points of [11] is Proposition 7.3. This establishes that certain maps are “good” with respect to friendly measures. We provide all relevant definitions later in the paper. The proof of [11, Proposition 7.3] crucially uses the Mean Value Theorem which is unavailable in the p -adic setting and thus represents the main barrier to establishing the aforementioned conjectures in the p -adic case. In this paper we develop a new technique that bypasses this barrier and thus enables us to develop a coherent theory of Diophantine approximation for friendly measures on \mathbb{Q}_p^n .

2. MAIN RESULTS

While postponing the definitions that we use until Section 3, we now state our main results. In what follows, given a measure μ on an open subset U of \mathbb{Q}_ν^d and a map $\mathbf{f} : U \rightarrow \mathbb{Q}_\nu^n$, $\mathbf{f}_*\mu$ will denote the corresponding pushforward measure on \mathbb{Q}_ν^n , where ν is either a prime number p or ∞ . Thus \mathbb{Q}_ν is either \mathbb{Q}_p or \mathbb{R} . Our first main theorem reads as follows.

Theorem 2.1. *Let U be an open subset of \mathbb{Q}_ν^d , and $\mathbf{f} : U \rightarrow \mathbb{Q}_\nu^n$. Let μ be an absolutely decaying Federer measure on U . Suppose that \mathbf{f} is a C^{l+1} map that is nonsingular and l -nondegenerate at μ -almost every point. Then $\mathbf{f}_*\mu$ is friendly.*

One of the main results in [11] is the above theorem for $\nu = \infty$. Thus Theorem 2.1 is new when ν is a prime number p . As a corollary of Theorem 2.1 we prove the following theorem which resolves the conjecture of Kleinbock and Tomanov appearing as Conjecture FP in [13].

Theorem 2.2. *Let μ be a self-similar measure on the limit set of an irreducible family of contracting similitudes of \mathbb{Q}_ν^d satisfying the open set condition, and let $\mathbf{f} : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$ be a smooth map which is nonsingular and nondegenerate at μ -almost every point. Then $\mathbf{f}_*\mu$ is strongly extremal.*

We recall that a measure μ on \mathbb{Q}_ν^n is called strongly extremal if μ -almost every point is not “very well multiplicatively approximable”. This implies that μ is extremal, i.e. that μ -almost every point is not “very well approximable”. We refer the reader to Section 11 of [13] for the definitions of these Diophantine properties.

Another main theorem of this paper is the p -adic analogue of the results of Kleinbock [10] regarding Diophantine approximations on affine subspaces. For the definitions of Diophantine exponents $\omega(\cdot)$ of measures and subspaces, the reader is referred to Section 3.8 and [5].

Theorem 2.3. *Suppose \mathcal{L} is a d -dimensional affine subspace of \mathbb{Q}_ν^n . Let μ be an absolutely decaying and Federer measure on \mathbb{Q}_ν^d and let $\mathbf{f} : \mathbb{Q}_\nu^d \rightarrow \mathcal{L}$ be a smooth map which is nondegenerate in \mathcal{L} at μ -almost every point of \mathbb{Q}_ν^d . Then $\omega(\mathbf{f}_*\mu) = \omega(\mathcal{L})$.*

When $\nu = \infty$, the above theorem is one of the main results in [10]. When ν is a prime number, Theorem 2.3 is new.

2.1. Difficulties in p -adic fields. We want to point out the main difficulties in proving Theorem 2.1 in an ultrametric set-up.

- There is no analogue of the Mean Value Theorem for p -adic continuously differentiable maps, not even for C^1 maps defined in Section 3.4. But the proof in case of \mathbb{R} ([11]) uses the Mean Value Theorem in a crucial manner.
- In order to bypass the use of the Mean Value Theorem, one may use the definition of C^k maps. It is here that the difference quotients $\bar{\Phi}_\beta$ as in Section 3.4, kick into the picture. Note that $\bar{\Phi}_\beta$ is a multivariable function, even for maps from \mathbb{Q}_ν to \mathbb{Q}_ν . This makes the analogue of many crucial lemmata in [11] very different.
- The relation between $\bar{\Phi}^k f'$ with f^{k+1} for any C^{k+1} function from \mathbb{Q}_ν to \mathbb{Q}_ν , poses difficulties in various parts of the proof.
- For maps in higher dimensions, we use combinatorial arguments throughout the paper. For instance, we need to know the exact error terms in the higher dimensional “Taylor theorem” in \mathbb{Q}_ν^d . This seems to be unavailable in the literature and we provide the necessary arguments in Section 11.
- For the convenience of the reader, we have provided a detailed discussion of the main obstacles and our strategy in overcoming them in the simplest non-trivial case in Section 5.

We conclude this section with some remarks.

Remarks.

- (1) There has also been recent interest in the question of Diophantine approximation in positive characteristic, when one approximates Laurent series by ratios of polynomials. The positive characteristic version of the Baker-Sprindžhuk conjectures were settled in [7]. The methods of this paper can be adapted to the function field setting in a relatively straightforward manner as they work over any ultrametric field.
- (2) Similarly, it is possible to extend our main results to more than one place of \mathbb{Q} , namely to the S -arithmetic setting.
- (3) In a forthcoming work, we will use the results in this paper to investigate badly approximable points on manifolds in \mathbb{Q}_p . In short, we establish the p -adic version of the main results of [2].

3. PRELIMINARIES

In this section we recall some terminology and definitions from [11, 13] and [10]. Our exposition necessarily follows these papers quite closely.

3.1. Besicovitch spaces. A metric space X is called *Besicovitch* [13] if there exists a constant N_X such that the following holds: for any bounded subset A of X and for any family \mathcal{B} of nonempty open balls in X such that every $x \in A$ is a center of some ball in \mathcal{B} , there is a finite

or countable subfamily $\{B_i\}$ of \mathcal{B} with

$$1_A \leq \sum_i 1_{B_i} \leq N_X. \quad (3.1)$$

Let us recall [13, Lemma 2.3].

Lemma 3.1. *For a metric space X , let us define*

$$M_X := \sup \{k \mid \text{there are balls } B_i = B(x_i, r_i), 1 \leq i \leq k, \text{ s.t. } \cap_{i=1}^k B_i \neq \emptyset, x_i \notin \cup_{j \neq i} B_j\},$$

and for any $c > 1$,

$$D_X(c) := \sup \left\{ k \mid \begin{array}{l} \text{there are } x \in X, r > 0 \text{ and pairwise disjoint balls} \\ B_1, \dots, B_k \text{ of radius } r \text{ contained in } B(x, cr) \end{array} \right\}.$$

If both M_X and $D_X(8)$ are finite, then X is Besicovitch.

For any ultrametric space, any two balls either do not intersect, or one is contained inside another. Hence $M_X = 1$ for any ultrametric space.

Now when $X = \mathbb{Q}_p$, since the balls are all of radius p^n with $n \in \mathbb{Z}$, a ball of radius p^n can contain at most p many disjoint balls of radius p^{n-1} . Hence $D_X(8) < \infty$ for $X = \mathbb{Q}_p$ and applying Lemma 3.1 it follows that \mathbb{Q}_p is Besicovitch. One can also see that $\mathbb{R}^d, \mathbb{Q}_p^d$ are Besicovitch spaces; see [16, Theorem 2.7] and [1, Corollary 4.13].

In particular, we will use later that $N_{\mathbb{Q}_p^d} = 1$ since any two balls that intersect in this nonarchimedean space are contained in each other. Readers are also referred to [14, 15] for examples of several Besicovitch spaces.

3.2. Federer measures. Let μ be a locally finite Borel measure on a metric space X , and U be an open subset of X with $\mu(U) > 0$. Following [11] we say that μ is *D-Federer on U* if

$$\sup_{\substack{x \in \text{supp } \mu, r > 0 \\ B(x, 3r) \subset U}} \frac{\mu(B(x, 3r))}{\mu(B(x, r))} < D.$$

We say that μ as above is *Federer* if, for μ -almost every $x \in X$, there exists a neighbourhood U of x and $D > 0$ such that μ is *D-Federer on U* . We refer the reader to [11] and [13] for examples of Federer measures.

3.3. Nonplanar measures. Suppose μ is a measure on \mathbb{Q}_ν^d . We call μ *nonplanar* if $\mu(\mathcal{L}_1) = 0$ for any affine hyperplane \mathcal{L}_1 of \mathbb{Q}_ν^d . Let $\mathbf{f} = (f_1, \dots, f_n)$ be a map from $\mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$. The pair (\mathbf{f}, μ) will be called *nonplanar at $x_0 \in \mathbb{Q}_\nu^d$* if, for any neighbourhood B of x_0 , the restrictions of $1, f_1, \dots, f_n$ to $B \cap \text{supp } \mu$ are linearly independent over \mathbb{Q}_ν . This is equivalent to saying that $\mathbf{f}(B \cap \text{supp } \mu)$ is not contained in any proper affine subspace of \mathbb{Q}_ν^d . Note that for $d = n$, if μ on \mathbb{Q}_ν^n is nonplanar then (\mathbf{Id}, μ) is nonplanar at μ almost every point.

Suppose $\mathbf{f} : X \subset \mathbb{Q}_\nu^d \rightarrow \mathcal{L} \subset \mathbb{Q}_\nu^n$ be a map, μ be a measure on X , and \mathcal{L} be an affine subspace of \mathbb{Q}_ν^n . Then (\mathbf{f}, μ) is called nonplanar in \mathcal{L} if

$$\mathcal{L} = \langle \mathbf{f}(B \cap \text{supp } \mu) \rangle_a,$$

for every nonempty ball B such that $\mu(B) > 0$; see [10]. Here $\langle M \rangle_a$ means the intersection all subspaces in \mathbb{Q}_ν^n containing M .

3.4. **C^k functions in \mathbb{Q}_ν^d .** We recall the definition of p -adic C^k functions, following closely the exposition in [13]. We refer the reader to [17] for a detailed treatment of p -adic calculus. Let $f : U \subset \mathbb{Q}_\nu \rightarrow \mathbb{Q}_\nu$ be a function and assume that U does not have any isolated points. The first difference quotient $\Phi^1 f$ of f is the function of two variables given by

$$\Phi^1 f(x, y) := \frac{f(x) - f(y)}{x - y} \quad (x, y \in U, x \neq y)$$

defined on

$$\nabla^2 U := \{(x, y) \in U \times U \mid x \neq y\}.$$

We say that f is C^1 at a if the limit

$$\lim_{(x, y) \rightarrow (a, a)} \Phi^1 f(x, y)$$

exists, and that $f \in C^1(U)$ if f is C^1 at every point of U .

More generally, for $k \in \mathbb{N}$ set

$$\nabla^k U := \{(x_1, \dots, x_k) \in U^k \mid x_i \neq x_j \text{ for } i \neq j\},$$

and define the k -th order difference quotient $\Phi^k f : \nabla^{k+1} U \rightarrow \mathbb{Q}_\nu$ of f inductively by $\Phi^0 f := f$ and

$$\Phi^k f(x_1, x_2, \dots, x_{k+1}) := \frac{\Phi^{k-1} f(x_1, x_3, \dots, x_{k+1}) - \Phi^{k-1} f(x_2, x_3, \dots, x_{k+1})}{x_1 - x_2}.$$

We say that f is C^k if $\Phi^k f$ can be extended to a continuous function $\bar{\Phi}^k f : U^{k+1} \rightarrow \mathbb{Q}_\nu$.

Also, we say that f is C^k at a if the limit

$$\bar{\Phi}^k f(a, \dots, a) := \lim_{(x_1, \dots, x_{k+1}) \rightarrow (a, \dots, a)} \Phi^k f(x_1, \dots, x_{k+1})$$

exists. By [17, Theorem 29.9] $f \in C^k(U)$ if f is C^k at every point of U . It can be shown that C^k functions f are k times differentiable, and that

$$f^{(k)}(x) = k! \bar{\Phi}^k f(x, \dots, x).$$

The definition of C^k functions of several ultrametric variables follows along similar lines. If f is a \mathbb{Q}_ν -valued function on $U_1 \times \dots \times U_d$, where each U_i is a subset of \mathbb{Q}_ν without isolated points, denote by $\Phi_i^k f$ the k th order difference quotient of f with respect to the variable x_i , and, more generally, for a multi-index $\beta = (i_1, \dots, i_d)$ let

$$\Phi_\beta f := \Phi_1^{i_1} \circ \dots \circ \Phi_d^{i_d} f.$$

It is not hard to check that the composition can be taken in any order. For any fixed j permutation of coordinates in $U_j^{i_j+1}$ does not change the function; see [17, Lemma 29.2 (ii)]. This “difference quotient of order β ” is defined on $\nabla^{i_1} U_1 \times \dots \times \nabla^{i_d} U_d$, and as before we say that f belongs to $C^k(U_1 \times \dots \times U_d)$ if for any multi-index β with $|\beta| := \sum_{j=1}^d i_j \leq k$, $\Phi_\beta f$ is extendable to a continuous function $\bar{\Phi}_\beta f : U_1^{i_1+1} \times \dots \times U_d^{i_d+1} \rightarrow \mathbb{Q}_\nu$. As in the one-variable case, one can show that partial derivatives $\partial_\beta f := \partial_1^{i_1} \circ \dots \circ \partial_d^{i_d} f$ of a C^k function f exist and are continuous as long as $|\beta| \leq k$. Moreover, one has

$$\partial_\beta f(x_1, \dots, x_d) = \beta! \bar{\Phi}_\beta f(x_1, \dots, x_1, \dots, x_d, \dots, x_d). \quad (3.2)$$

where $\beta! := \prod_{j=1}^d i_j!$, and each of the variables x_j in the right hand side of Equation (3.2) is repeated $i_j + 1$ times. Lastly, we recall the definition of a function being nonsingular at a point.

Definition 3.1. A function C^1 function $\mathbf{f} = (f_1, \dots, f_n) : U \subset \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$ is called nonsingular at a point \mathbf{x}_0 if the matrix $\nabla \mathbf{f}(\mathbf{x}_0) = [\partial_{e_i} f_j(\mathbf{x}_0)]_{1 \leq i \leq d, 1 \leq j \leq n}$ has rank d .

3.5. Nondegeneracy in an affine subspace. Following [10], we say that a map $\mathbf{f} : U \rightarrow \mathbb{Q}_\nu^n$, where U is an open subset of \mathbb{Q}_ν^d , is l -nondegenerate in an affine subspace \mathcal{L} of \mathbb{Q}_ν^n at $\mathbf{x} \in U$ if $\mathbf{f}(U) \subset \mathcal{L}$, \mathbf{f} is C^l and the span of all the partial derivatives of \mathbf{f} at \mathbf{x} up to order l coincides with the linear part of \mathcal{L} . We say that \mathbf{f} is nondegenerate in an affine subspace \mathcal{L} of \mathbb{Q}_ν^n at $\mathbf{x} \in U$ if it is l -nondegenerate in an affine subspace \mathcal{L} of \mathbb{Q}_ν^n at \mathbf{x} for some $l \in \mathbb{N}$. We say that \mathbf{f} is (l) -nondegenerate at $\mathbf{x} \in U$ if the above hold with $\mathcal{L} = \mathbb{Q}_\nu^n$.

If \mathcal{M} is a d -dimensional submanifold of \mathcal{L} , we will say that \mathcal{M} is nondegenerate in \mathcal{L} at $y \in \mathcal{M}$ if any (equivalently, some) diffeomorphism \mathbf{f} between an open subset U of \mathbb{Q}_ν^d and a neighbourhood of y in \mathcal{M} is nondegenerate in \mathcal{L} at $\mathbf{f}^{-1}(y)$. We will say that $\mathbf{f} : U \rightarrow \mathcal{L}$ (resp. $\mathcal{M} \subset \mathcal{L}$) is nondegenerate in \mathcal{L} if it is nondegenerate in \mathcal{L} at λ -almost every point of U , where λ is the Lebesgue/Haar measure on U (resp. of \mathcal{M} , in the sense of the smooth measure class on \mathcal{M}).

3.6. Good maps and decaying measures. Let μ be a locally finite Borel measure on \mathbb{Q}_ν^d and U be an open subset of \mathbb{Q}_ν^d . Given $A \subset \mathbb{Q}_\nu^d$ with $\mu(A) > 0$ and a \mathbb{Q}_ν -valued function f on \mathbb{Q}_ν^d , let

$$\|f\|_{\mu,A} := \sup_{x \in A \cap \text{supp } \mu} |f(\mathbf{x})|_\nu,$$

where $|\cdot|_\nu$ is the standard ν -adic absolute value. A μ -measurable function $f : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$ is called (C, α) -good on U with respect to μ if for any open ball $B \subset U$ centered in $\text{supp}(\mu)$ and $\varepsilon > 0$ one has that

$$\mu(\{\mathbf{x} \in B \mid |f(\mathbf{x})|_\nu < \varepsilon\}) \leq C \left(\frac{\varepsilon}{\|f\|_{\mu,B}} \right)^\alpha \mu(B). \quad (3.3)$$

Similarly, a function $f : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$ is called absolutely (C, α) -good on U with respect to μ if for any open ball $B \subset U$ centered in $\text{supp}(\mu)$ and $\varepsilon > 0$ one has

$$\mu(\{\mathbf{x} \in B \mid |f(\mathbf{x})|_\nu < \varepsilon\}) \leq C \left(\frac{\varepsilon}{\|f\|_B} \right)^\alpha \mu(B), \quad (3.4)$$

where $\|f\|_B = \sup_{\mathbf{x} \in B} |f(\mathbf{x})|_\nu$. Since $\|f\|_{\mu,B} \leq \|f\|_B$, being absolutely (C, α) -good implies being (C, α) -good on U . The converse is true for measures having full support.

Remark 3.1. Note that if f is (absolutely) (C, α) -good w.r.t. μ , Equation (3.3) and Equation (3.4) will remain true if the strict inequality in the left hand side is replaced by a non-strict inequality. Indeed, we trivially have that $\{\mathbf{x} \in B \mid |f(\mathbf{x})|_\nu \leq \varepsilon\} \subset \{\mathbf{x} \in B \mid |f(\mathbf{x})|_\nu < \varepsilon + 1/N\}$ for $N > 1$. Then, for example in the case of absolutely (C, α) -good functions, we get that

$$\mu(\{\mathbf{x} \in B \mid |f(\mathbf{x})|_\nu \leq \varepsilon\}) \leq \mu(\{\mathbf{x} \in B \mid |f(\mathbf{x})|_\nu < \varepsilon + 1/N\}) \leq C \left(\frac{\varepsilon + 1/N}{\|f\|_B} \right)^\alpha \mu(B)$$

and it remains to let $N \rightarrow \infty$.

Next, let $\mathbf{f} = (f_1, \dots, f_n) : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$ be a map, and μ be a locally finite Borel measure on \mathbb{Q}_ν^d . We say that (\mathbf{f}, μ) is (C, α) -good at \mathbf{x} if there exists a neighborhood U of \mathbf{x} such that any linear combination of $1, f_1, \dots, f_n$ is (C, α) -good on U with respect to μ . We will simply say (\mathbf{f}, μ) is good at \mathbf{x} if it is (C, α) -good at \mathbf{x} for some positive C and α . We will say (\mathbf{f}, μ) is good (on U) if it is good at μ -almost every point \mathbf{x} (of U). When μ is Lebesgue/Haar measure on \mathbb{Q}_ν^d , we omit ‘with respect to μ ’ and just say that the map \mathbf{f} is good (at \mathbf{x}). As is well known, polynomials are (C, α) -good at every point for some C and α that depend only on d and the degree of the polynomial in question, see [19, Lemma 4.1]. In fact, there are many examples of good maps. We recall Lemma 2.5 from [12] and Proposition 4.2 from [13] which show that

nondegenerate maps are good. For the definition of C^l maps in the p -adic case, the reader is referred to Section 3.4.

Proposition 3.1 (See [12, Lemma 2.5] and [13, Proposition 5.1]). *Let $\mathbf{f} = (f_1, \dots, f_n)$ be a C^l map from an open subset $U \subset \mathbb{Q}_\nu^d$ to \mathbb{Q}_ν^n which is l -nondegenerate in \mathbb{Q}_ν^n at $\mathbf{x}_0 \in U$. Then there is a neighbourhood $V \subset U$ of \mathbf{x}_0 such that any linear combination of $1, f_1 \cdots f_n$ is (C', α) -good on V , where $C', \alpha > 0$ only depends on d, l and the field. In particular, the nondegeneracy of \mathbf{f} in \mathbb{Q}_ν^n at \mathbf{x}_0 implies that \mathbf{f} is good at \mathbf{x}_0 .*

We further have the following corollary whose proof is identical to that of [9, Corollary 3.2].

Corollary 3.1. *Let \mathcal{L} be an affine subspace of \mathbb{Q}_ν^n and let $\mathbf{f} = (f_1, \dots, f_n)$ be a map from an open subset U of \mathbb{Q}_ν^d to \mathcal{L} which is nondegenerate in \mathcal{L} at $\mathbf{x}_0 \in U$. Then \mathbf{f} is good at \mathbf{x}_0 .*

Given $C, \alpha > 0$ and an open subset U in \mathbb{Q}_ν^n , we say that μ is (absolutely) (C, α) -decaying on U if any affine map is (absolutely) (C, α) -good on U w.r.t. μ .

3.7. Friendly measures. We are now ready to define *friendly* measures following [11].

Definition 3.2 (Friendly measure). *A locally finite Borel measure μ on \mathbb{Q}_ν^n is called friendly if μ is nonplanar and for μ -almost every $\mathbf{x} \in \mathbb{Q}_\nu^n$, there exist a neighbourhood U of \mathbf{x} and positive $C, \alpha, D > 0$ such that μ is D -Federer on U and μ is (C, α) -decaying on U .*

Examples of friendly measures were constructed in [11] and their p -adic analogues were described in [13]. We briefly recall the discussion from [13, Section 11.6]. Let “dist” be a metric on \mathbb{Q}_ν^n induced by $|\cdot|_\nu$. A map $h : \mathbb{Q}_\nu^n \rightarrow \mathbb{Q}_\nu^n$ is a contracting similitude with contraction rate ρ if $0 < \rho < 1$ and

$$\text{dist}(h(\mathbf{x}), h(\mathbf{y})) = \rho \text{dist}(\mathbf{x}, \mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{Q}_\nu^n.$$

By a result of Hutchinson [8], for any finite family h_1, \dots, h_m of contracting similitudes there exists a unique nonempty compact set Q , called the limit set of the family, such that

$$Q = \bigcup_{i=1}^m h_i(Q).$$

The family h_1, \dots, h_m are said to satisfy the open set condition if there exists an open subset $U \subset \mathbb{Q}_\nu^n$ such that

$$h_i(U) \subset U \quad \text{for all } i = 1, \dots, m,$$

and

$$i \neq j \implies h_i(U) \cap h_j(U) = \emptyset.$$

By a result of Hutchinson [8], if $h_i, i = 1, \dots, m$, are contracting similitudes with contraction rates ρ_i satisfying the open set condition, and if $s > 0$ is the unique solution to the equation $\sum_i \rho_i^s = 1$ then the s -dimensional Hausdorff measure \mathcal{H}^s of Q is positive and finite. The number s is called the similarity dimension of the family $\{h_i\}$.

The family $\{h_i\}$ is irreducible if there does not exist a finite $\{h_i\}$ -invariant collection of proper affine subspaces of \mathbb{Q}_ν^n . The following proposition is slightly stronger than [13, Proposition 11.3], which in turn is a direct adaptation of [11, Theorem 2.3]. It provides a rich class of examples of friendly measures.

Proposition 3.2. *Let $\{h_1, \dots, h_m\}$ be an irreducible family of contracting similitudes on \mathbb{Q}_ν^n satisfying the open set condition, s its similarity dimension, and μ the restriction of \mathcal{H}^s to the limit set Q of the family $\{h_i\}$. Then μ is absolutely decaying and Federer.*

3.8. Diophantine exponents. For $\mathbf{y} \in \mathbb{Q}_\nu^n$, we define

$$\omega(\mathbf{y}) = \sup\{\omega \mid |\mathbf{q} \cdot \mathbf{y} + q_0|_\nu \leq \frac{1}{\|(\mathbf{q}, q_0)\|_\infty^\omega} \text{ for infinitely many } (\mathbf{q}, q_0) \in \mathbb{Z}^{n+1} \setminus 0\}.$$

Here $\|\cdot\|_\infty$ is the max norm in \mathbb{R}^{n+1} . For a Borel measure μ on \mathbb{Q}_ν^n , from [10] we recall the Diophantine exponent $\omega(\mu)$ of μ to be

$$\omega(\mu) = \sup\{v : \mu(\{y \mid \omega(y) > v\}) > 0\}.$$

If $\mathbf{f} : U \subset \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$ is a map such that, $\mathcal{M} = \mathbf{f}(U)$, where U is an open subset of \mathbb{Q}_ν^d , we define $\omega(\mathcal{M}) = \omega(\mathbf{f}_* \lambda)$. Here λ is the Haar measure on \mathbb{Q}_ν^d .

4. PROOFS OF THEOREM 2.1, THEOREM 2.2 AND THEOREM 2.3

We need the following proposition in order to prove Theorem 2.1 and Theorem 2.3.

Proposition 4.1. *Let $\mathbf{f} = (f_1, \dots, f_n) : U \subset \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$ be a C^{l+1} map, and let $\mathbf{x}_0 \in U$ be such that \mathbf{f} is l -nondegenerate at \mathbf{x}_0 . Let μ be a locally finite Borel measure which is D -Federer and absolutely (C, α) -decaying on U for some $D, C, \alpha > 0$. Then there exists a neighbourhood $V \subset U$ of \mathbf{x}_0 and a positive $C' > 0$ such that any*

$$g \in \mathcal{L}_{\mathbf{f}} := \left\{ c_0 + \sum_{i=1}^n c_i f_i : c_0, \dots, c_n \in \mathbb{Q}_\nu \right\} \quad (4.1)$$

is absolutely $(C', \frac{\alpha}{(2^{l+1}-2)})$ -good on V with respect to μ .

This proposition is the key to establishing the theorems stated in Section 2 and requires new technical ideas. We will prove it in the following sections. When $\nu = \infty$, Proposition 4.1 was proved in [11, Section 7]. One of the crucial facts used in [11] was the Mean Value Theorem. Since there is no Mean Value Theorem in the p -adic setting, we have come up with a different proof.

The following lemma gives a sufficient condition written in terms of μ , for $\mathbf{f}_* \mu$ to be friendly.

Lemma 4.1. *Let μ be a D -Federer measure on an open subset U of \mathbb{Q}_ν^d , let $\mathbf{f} : U \rightarrow \mathbb{Q}_\nu^n$ and $C, K, \alpha > 0$ be such that*

- *for any $\mathbf{x}_1, \mathbf{x}_2 \in U$ one has that*

$$\frac{1}{K} \text{dist}(\mathbf{x}_1, \mathbf{x}_2) \leq \text{dist}(\mathbf{f}(\mathbf{x}_1), \mathbf{f}(\mathbf{x}_2)) \leq K \text{dist}(\mathbf{x}_1, \mathbf{x}_2);$$

- *any $f \in \mathcal{L}_{\mathbf{f}}$ is absolutely (C, α) -good on U with respect to μ , where $\mathcal{L}_{\mathbf{f}}$ as in Equation (4.1).*

Then $\mathbf{f}_ \mu$ is friendly.*

When $\nu = \infty$ the above lemma was proved in of [11, Lemma 7.1] and the same proof applies verbatim when ν is a prime number.

4.1. Proof of Theorem 2.1 using Proposition 4.1 and Lemma 4.1. Since \mathbf{f} is a nonsingular C^{l+1} map at \mathbf{x}_0 , we can find a neighbourhood V of \mathbf{x}_0 such that $\mathbf{f}|_V$ is bi-Lipschitz; see Lemma 11.4. This satisfies the first condition of Lemma 4.1. By Proposition 4.1, we have that \mathbf{f} and μ satisfy the second condition of Lemma 4.1. Therefore, by Lemma 4.1 we have that $\mathbf{f}_* \mu$ is friendly.

4.2. Proof of Theorem 2.2 using Theorem 2.1. Theorem 2.2 follows by combining Proposition 3.2 and Theorem 2.1 and [13, Corollary 11.2]

4.3. Proof of Theorem 2.3 using Proposition 4.1. We first state the following theorem which is [5, Corollary 6.2].

Theorem 4.1 ([5, Corollary 6.2]). *Let μ be a Federer measure on a Besicovitch metric space X , \mathcal{L} an affine subspace of \mathbb{Q}_ν^n , and let $\mathbf{f} : X \rightarrow \mathcal{L}$ be a continuous map such that (\mathbf{f}, μ) is good and nonplanar in \mathcal{L} . Then*

$$\omega(\mathbf{f}_*\mu) = \omega(\mathcal{L}) = \inf\{\omega(\mathbf{y}) \mid \mathbf{y} \in \mathcal{L}\} = \inf\{\omega(\mathbf{f}(x)) \mid x \in \text{supp } \mu\}. \quad (4.2)$$

Remark 4.1. *Let μ be a friendly measure on \mathbb{Q}_ν^d , and let $\mathbf{f} : \mathbb{Q}_\nu^d \rightarrow \mathcal{L}$ be an affine isomorphism. Then by the above theorem, $\omega(\mathbf{f}_*\mu) = \omega(\mathcal{L})$.*

Since in Theorem 2.3, \mathbf{f} is nondegenerate in \mathcal{L} at μ almost all points, we get that (\mathbf{f}, μ) is nonplanar in \mathcal{L} . If not, then there exists a ball B such that $\mu(B) > 0$ and an affine subspace $\mathcal{L}', \mathcal{L}' \subset \mathcal{L}$ such that $\mathbf{f}(B \cap \text{supp } \mu) \subset \mathcal{L}'$. Let $\mathbf{x} \in B \cap \text{supp } \mu$ such that all partial derivatives of \mathbf{f} at \mathbf{x} up to order l spans the linear part of \mathcal{L} . But the fact $\mathbf{f}(B \cap \text{supp } \mu) \subset \mathcal{L}'$, implies all derivatives at \mathbf{x} must lie on the linear part of \mathcal{L}' which is proper in \mathcal{L} giving a contradiction.

Therefore, Theorem 2.3 follows from combining Proposition 4.1 and the above theorem. The rest of the paper is spent on proving Proposition 4.1.

5. NOTES ON THE PROOF OF PROPOSITION 4.1

In this section, we consider the simplest nontrivial situation in which we attempt to explain the obstacles we face and the ideas we use in the general case, where the obstacles and ideas are less transparent. Recall that to validate Proposition 4.1 we need to demonstrate that under certain conditions on the measure μ and the map $\mathbf{f} = (f_1, \dots, f_n) : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$, there exists a neighborhood V such that all linear combinations of $1, f_1, \dots, f_n$ are good w.r.t μ , a concept stated in Section 3.6.

5.1. Sufficient conditions. The task is that given any function $g : B \subset \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$, defined on a ball B of radius r , we want to find sufficient conditions on g , such that g is good in this ball B w.r.t μ ; see Theorem 6.1. It is too ambitious to hope that the same conditions as in the real case, [11, Theorem 7.6] would be enough. Next we discuss why this is so.

- Let us constrain ourselves to the $d = 1$ case. The lemmas [11, Lemma 7.4] and [11, Lemma 7.5], which are crucial in [11, Theorem 7.6] are not clear in the \mathbb{Q}_ν setting. For instance, in [11, Lemma 7.4] it was shown that for any C^1 function on ball $B \subset \mathbb{R}$ of radius r , we have that

$$r \inf_{x \in B} |g'(x)| \ll \sup_{x \in B} |g(x)|.$$

This is a direct implication of the Mean Value Theorem.

- However, the above is not true¹ in \mathbb{Q}_ν . For instance, we may take $B = \mathbb{Z}_\nu$, and $g_N(\sum_{i \geq 0} a_i \nu^i) = \sum_{i \geq N} a_i \nu^i$ for $N \in \mathbb{N}$. Then g_N differs from the identity map by a locally constant map. Hence for every $x \in \mathbb{Z}_\nu$, $|g'_N(x)|_\nu = 1$, whereas $\sup_{x \in B} |g_N(x)|_\nu \leq \nu^{-N}$, by definition. Hence for sufficiently large N , the above inequality is false. Nevertheless, we can use the definition of C^k functions; see Section 3.4 in order to prove a weaker statement; see Lemma 6.1. In particular, for any C^1 function defined on a ball $B \subset \mathbb{Q}_\nu$ of radius r , we show that

$$r \inf_{(x,y) \in B \times B} |\bar{\Phi}^1 g(x, y)|_\nu \ll \sup_{x \in B} |g(x)|_\nu.$$

¹We are thankful to an anonymous referee who suggested the example we provide here.

- For the same reason as above our Lemma 6.2 is weaker than [11, Lemma 7.5].
- Other than having two weaker lemmas, there is another issue that we face. In the real case, we have that $(g')' = g''$ for any C^2 function. But in case of any C^2 function g defined on \mathbb{Q}_ν , we have instead that

$$\bar{\Phi}^1 g'(x, y) = \bar{\Phi}^2 g(x, x, y) + \bar{\Phi}^2 g(x, y, y).$$

Hence it is not immediately clear what sort of relation we have between

$$\inf_{(x,y,z) \in B \times B \times B} |\bar{\Phi}^2 g(x, y, z)|_\nu \quad \text{and} \quad \inf_{(x,y) \in B \times B} |\bar{\Phi}^1 g'(x, y)|_\nu.$$

Since the first one is defined over a larger domain, one would expect that the first quantity should be less than the second quantity upto some constant multiple. But the second quantity is the infimum of a sum of two functions, hence there is a possibility that it becomes 0.

To tackle the aforementioned obstacles we impose three conditions on g in Theorem 6.1, namely Equation (6.9), Equation (6.10) and Equation (6.11). Since these conditions are a bit technical, we refrain from stating them here, except for the most important one namely Equation (6.9) which is discussed below. The latter two conditions showed up, albeit in a weaker sense, in the real setting. The first condition Equation (6.9) however, is entirely new and an important idea in our paper. No analogous version was needed in the proof of the real case. The second and third conditions are stronger than the respective conditions in the real case, [11, Theorem 7.6].

As we prove the main theorem Theorem 2.1 with no extra assumption than those are present in the real case, we need to show that these stronger sufficient conditions in Theorem 6.1 are satisfied by any linear combination of $1, f_1, \dots, f_n$.

5.2. The first condition. The first condition Equation (6.9) in Theorem 6.1 is the hardest to check among the three conditions in the theorem. In the case $d = 1$ and C^k function g , note that the first condition is to check a chain of inequalities as follows:

$$\inf |\bar{\Phi}^k g|_\nu \ll \inf |\bar{\Phi}^{k-1} g'|_\nu \ll \dots \ll \inf |\bar{\Phi}^1 g^{k-1}|_\nu \ll \inf |g^k|_\nu. \quad (5.1)$$

Here and elsewhere in this section we adopt the convention that

$$\inf |\bar{\Phi}^j g|_\nu := \inf_{(x_1, \dots, x_{j+1}) \in B \times \dots \times B} |\bar{\Phi}^j g(x_1, \dots, x_{j+1})|_\nu \quad \text{for } j = 0, \dots, k.$$

First, we show that polynomials satisfy all three conditions; see Theorem 7.1. We use the trick of ‘proof by maximality’, which is also used in [11, Lemma 7.7]. But the proof in [11, Lemma 7.7] uses the Mean Value Theorem and there were only two weaker conditions to check unlike our case, where we need to verify three stronger conditions.

5.3. A very simple case of polynomials. Let us sketch how we handle the simplest nontrivial case of polynomials, where three conditions in Theorem 6.1 are satisfied. Let P be a polynomial over \mathbb{Q}_ν in one variable such that $\sup_{x \in B(0,1)} |P(x)|_\nu = 1$. Suppose k is the largest integer such that there exists $(x_1, \dots, x_{k+1}) \in B(0, 1)^{k+1}$ with

$$|\bar{\Phi}^k P(x_1, \dots, x_{k+1})|_\nu > \frac{1}{\nu^{6\alpha+1}},$$

where $\alpha > 0$ is carefully chosen. Such a k will exist since we started with a ‘normalized’ polynomial. Now for simplicity, let $k = 2$. Hence there exists $(a, b, c) \in B(0, 1)^3$ such that

$$|\bar{\Phi}^2 P(a, b, c)|_\nu > \frac{1}{\nu^{6\alpha+1}}.$$

Then it is easy to see that for any $(x, y, z) \in B(0, 1)^3$,

$$\bar{\Phi}^2 P(x, y, z) - \bar{\Phi}^2 P(a, b, c) = \bar{\Phi}^3 P(x, y, z, a)(z - a) + \bar{\Phi}^3 P(x, y, a, b)(y - b) + \bar{\Phi}^3 P(x, a, b)(x - c).$$

Now we use ultrametric norm property to conclude that for any $(x, y, z) \in B(0, 1)^3$,

$$|\bar{\Phi}^2 P(x, y, z)|_\nu \leq \max\{|\bar{\Phi}^2 P(a, b, c)|_\nu, |\bar{\Phi}^3 P(x, y, z, a)|_\nu, |\bar{\Phi}^3 P(x, y, a, b)|_\nu, |\bar{\Phi}^3 P(x, a, b)|_\nu\}.$$

By maximality of $k = 2$, the max in the above should be attained by $|\bar{\Phi}^2 P(a, b, c)|_\nu$, giving

$$|\bar{\Phi}^2 P(x, y, z)|_\nu > \frac{1}{\nu^{6\alpha+2}}.$$

This verifies the second condition Equation (6.10). Next we claim,

$$\inf |\bar{\Phi}^2 P|_\nu \ll \inf |\bar{\Phi}^1 P'|_\nu.$$

The following simple observation is key for verifying the first inequality in Equation (5.1). For any $(x, y, z) \in B(0, 1)^3$, we have that

$$2\bar{\Phi}^2 P(x, y, z) = \bar{\Phi}^1 P'(x, y) + \bar{\Phi}^3 P(x, x, y, z)(z - x) + \bar{\Phi}^3 P(x, y, y, z)(z - y). \quad (5.2)$$

Now let $\inf |\bar{\Phi}^1 P'|_\nu = |\bar{\Phi}^1 P'(c, d)|_\nu$ for some $c, d \in B(0, 1)$. For any $z \in B(0, 1)$,

$$|2\bar{\Phi}^2 P(c, d, z)|_\nu \leq \max\{|\bar{\Phi}^1 P'(c, d)|_\nu, |\bar{\Phi}^3 P(c, c, d, z)|_\nu, |\bar{\Phi}^3 P(c, d, d, z)|_\nu\}.$$

If for some $z \in B(0, 1)$, $|\bar{\Phi}^1 P'(c, d)|_\nu \geq \max\{|\bar{\Phi}^3 P(c, c, d, z)|_\nu, |\bar{\Phi}^3 P(c, d, d, z)|_\nu\}$ then we are done. If not, then for every z ,

$$|2\bar{\Phi}^2 P(c, d, z)|_\nu \leq |\bar{\Phi}^3 P(c, c, d, z)|_\nu, \text{ or } |2\bar{\Phi}^2 P(c, d, z)|_\nu \leq |\bar{\Phi}^3 P(c, d, d, z)|_\nu.$$

We choose α such that either case will contradict the maximality.

5.4. Approximations by polynomials. Once we show that polynomials satisfy all three conditions in Theorem 6.1, we approximate difference quotients of functions of interest by difference quotients of polynomials in a uniform manner; see Theorem 8.1.

Consider the $d = 1$ case. We show the following for any C^{l+1} map f defined on $B(y, r) \subset B(0, 1)$, and for every $k \leq l$,

$$\begin{aligned} & \sup_{\substack{z_i \in B(y, r) \\ i=1, \dots, k+1}} |\bar{\Phi}^k f(z_1, z_2, \dots, z_{k+1}) - \bar{\Phi}^k P_{f, y, l}(z_1, z_2, \dots, z_{k+1})|_\nu \\ & \leq r^{l-k} \sup_{\substack{x_i, y_i \in B(y, r) \\ i=1, \dots, l+1}} |\bar{\Phi}^l f(x_1, x_2, \dots, x_{l+1}) - \bar{\Phi}^l f(y_1, y_2, \dots, y_{l+1})|_\nu. \end{aligned} \quad (5.3)$$

Here the l -th Taylor polynomial of f at $y \in B$ is denoted as

$$P_{f, y, l}(x) = f(y) + \bar{\Phi}^1 f(y, y)(x - y) + \dots + \bar{\Phi}^l f(y, \dots, y)(x - y)^l.$$

We prove Equation (5.3) by induction on l . The following is typically referred as “Taylor’s theorem” in \mathbb{Q}_ν ,

$$f(x) = P_{f, y, l}(x) + \bar{\Phi}^{l+1} f(x, y, \dots, y)(x - y)^{l+1}, \quad (5.4)$$

which one can find in a simple manner, [17, Theorem 29.4]. In order to prove Equation (5.3) by induction, we consider the function $f_z(x) := \bar{\Phi}^1 f(x, z)$ restricting one coordinate. We explore relations between $\bar{\Phi}^i P_{f, y, l}$ with $\bar{\Phi}^i P_{f_z, y, l}$; see Lemma 8.1, which we then apply to prove Theorem 8.1.

In higher dimension, one would expect to have an analogue of Equation (5.4), but we could not find any reference where the error term is explicitly calculated, which we require. So in an

appendix Section 11 we prove Taylor's theorem in higher dimension with explicitly given error term.

6. GOOD FUNCTIONS W.R.T. ABSOLUTELY DECAYING MEASURES

6.1. Notation. First of all, in what follows if there is no risk of confusion, we will omit the subscript ν from the notation of ν -adic absolute value, e.g. given $x \in \mathbb{Q}_\nu$, we will write $|x|$ for $|x|_\nu$. It will be clear from the context which absolute value we use. Throughout, the norm $\|\cdot\|$ on \mathbb{Q}_ν^i with $i \in \mathbb{N}$ induced by the ν -adic absolute value will be the supremum norm. For any multi-index $\beta = (i_1, \dots, i_d)$ we denote $|\beta| := i_1 + \dots + i_d$ by abuse of notation.

Next, given a ball B in \mathbb{Q}_ν , $B^i := B \times \dots \times B \subset \mathbb{Q}_\nu^i$ will denote the product of i copies of B and $\mathbf{y} \in B^i$ will be written as (y_j) , thus $\mathbf{y} = (y_j) \in B^i$. We set $e_i = (0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{Z}^d$, where 1 is at the i -th place. Given a multi-index $\beta = (i_1, \dots, i_d)$, we let $\beta(1) := \beta + \sum_{i=1}^d e_i$, thus $\beta(1) = (i_1+1, \dots, i_d+1)$. We denote by $B(\mathbf{y}, r) := \{\mathbf{x} \mid \|\mathbf{x} - \mathbf{y}\| \leq r\}$ a ball centred at \mathbf{y} of radius r . Given a multi-index $\alpha = (i_1, \dots, i_d)$, we also let $\mathbf{B}^\alpha = B(\mathbf{y}, r)^\alpha := B(y_1, r)^{i_1} \times \dots \times B(y_d, r)^{i_d}$ and denote by \mathbf{x}_α the elements of \mathbf{B}^α , thus $\mathbf{x}_\alpha = (\mathbf{x}_1, \dots, \mathbf{x}_d) \in \mathbf{B}^\alpha$, where $\mathbf{x}_j \in B(y_j, r)^{i_j}$. For a function $g : X \rightarrow \mathbb{Q}_\nu$, we set

$$\inf_X |g| := \inf_{x \in X} |g(x)|.$$

Similar notation will be used for supremum. Finally, given $y \in \mathbb{Q}_\nu$ and $k \in \mathbb{N}$, we let $y^k := (y, \dots, y) \in \mathbb{Q}_\nu^k$ and, given a multi-index $\beta = (i_1, \dots, i_d)$, we let $\bar{\mathbf{y}}_\beta := (y_1^{i_1+1}, \dots, y_d^{i_d+1})$, thus $\bar{\mathbf{y}}_\beta$ is an element of $\mathbb{Q}_\nu^{i_1+1} \times \dots \times \mathbb{Q}_\nu^{i_d+1}$. For a multi index $\beta = (i_1, \dots, i_d)$, let us recall $\beta! := \prod_{j=1}^d i_j!$.

Lemma 6.1. *Let B be a ball in \mathbb{Q}_ν of radius $r := \nu^{-t}$, where $t \in \mathbb{N}$. Suppose that $f : B^d \rightarrow \mathbb{Q}_\nu$ is a C^l map. Then for any multi-index $\beta = (i_1, \dots, i_d)$ with $|\beta| := i_1 + \dots + i_d \leq l$ we have that*

$$\inf_{\nabla^{i_1} B \times \dots \times \nabla^{i_d} B} |\Phi_\beta f| \leq |\beta|!^{-1} r^{-|\beta|} \|f\|_{B^d}, \quad (6.1)$$

where $|\beta|!$ is the ν -adic norm of $\beta!$, and $\|f\|_{B^d} = \sup_{\mathbf{x} \in B^d} |f(\mathbf{x})|$.

Proof. Let $\mathbf{x} = (x_1, \dots, x_d) \in B^d$ and for $j = 1, \dots, d$ define

$$D_j f(\mathbf{x}) := f(x_1, \dots, x_{j-1}, x_j + \nu^t, x_{j+1}, \dots, x_d) - f(\mathbf{x}).$$

Suppose that $x \in B$, then $x + k\nu^t \in B$ for any integer $k \geq 1$. Then for any $\mathbf{x} \in B^d$,

$$\Phi_j^k f(x_1, \dots, x_{j-1}, x_j, x_j + \nu^t, \dots, x_j + k\nu^t, x_{j+1}, \dots, x_d) = \frac{1}{k! \nu^{kt}} D_j^k f(\mathbf{x}). \quad (6.2)$$

For $k = 1$, the above equality follows from definition. Suppose, by the induction hypothesis, that the equality is true for $k - 1 \in \mathbb{N}$. This means that for any $\mathbf{x} \in B^d$,

$$\Phi_j^{k-1} f(x_1, \dots, x_{j-1}, x_j, x_j + \nu^t, \dots, x_j + (k-1)\nu^t, x_{j+1}, \dots, x_d) = \frac{1}{(k-1)! \nu^{(k-1)t}} D_j^{k-1} f(\mathbf{x}),$$

$$\begin{aligned} & \text{and } \Phi_j^{k-1} f(x_1, \dots, x_{j-1}, x_j + \nu^t, \dots, x_j + k\nu^t, x_{j+1}, \dots, x_d) \\ &= \frac{1}{(k-1)! \nu^{(k-1)t}} D_j^{k-1} f(x_1, \dots, x_{j-1}, x_j + \nu^t, x_{j+1}, \dots, x_d). \end{aligned}$$

Now using the two above equalities and definitions,

$$\begin{aligned}
& k\nu^t \Phi_j^k f(x_1, \dots, x_{j-1}, x_j, x_j + \nu^t, \dots, x_j + k\nu^t, x_{j+1}, \dots, x_d) \\
&= \Phi_j^{k-1} f(x_1, \dots, x_{j-1}, x_j + \nu^t, \dots, x_j + k\nu^t, x_{j+1}, \dots, x_d) - \\
& \Phi_j^{k-1} f(x_1, \dots, x_{j-1}, x_j, x_j + \nu^t, \dots, x_j + (k-1)\nu^t, x_{j+1}, \dots, x_d) \\
&= \frac{1}{(k-1)!\nu^{(k-1)t}} (D_j^{k-1} f(x_1, \dots, x_{j-1}, x_j + \nu^t, x_{j+1}, \dots, x_d) - D_j^{k-1} f(\mathbf{x})) \\
&= \frac{1}{(k-1)!\nu^{(k-1)t}} D_j^k f(\mathbf{x}).
\end{aligned}$$

Thus Equation (6.2) is valid.

Note that, by the ultrametric property, $|D_j^k f(\mathbf{x})| \leq |k!|^{-1} \|f\|_{B^d}$, where $|k!|$ is the ν -adic norm of $k! \in \mathbb{Z}$. Hence taking norms on both sides of Equation (6.2) gives

$$|k!| \nu^{-kt} |\Phi_j^k f(x_1, \dots, x_{j-1}, x_j, x_j + \nu^t, \dots, x_j + k\nu^t, x_{j+1}, \dots, x_d)| \leq \|f\|_{B^d}.$$

Given a multi-index $\beta = (i_1, \dots, i_d)$, let $D_\beta(\mathbf{x}) = D_1^{i_1} \circ \dots \circ D_d^{i_d}(\mathbf{x})$ and $\Phi_\beta = \Phi_1^{i_1} \circ \dots \circ \Phi_d^{i_d}$. Therefore when $|\beta| = m$ we get that

$$D_\beta(\mathbf{x}) = \beta! \nu^{mt} \Phi_\beta f(x_1, x_1 + \nu^t, \dots, x_1 + i_1 \nu^t, x_2, \dots, x_2 + i_2 \nu^t, \dots, x_d, x_d + \nu^t, \dots, x_d + i_d \nu^t). \quad (6.3)$$

Again, since $|D_\beta(\mathbf{x})| \leq \|f\|_{B^d}$, taking norms on both sides of Equation (6.3) gives

$$\inf_{\nabla^{i_1 B} \times \dots \times \nabla^{i_d B}} |\Phi_\beta f| \leq |\beta!|^{-1} r^{-|\beta|} \|f\|_{B^d}.$$

□

Lemma 6.2. *Let $C > 0$ and $\alpha > 0$ be given and let μ be an absolutely (C, α) -decaying measure on $U \subset \mathbb{Q}_\nu^d$. Let $c, r, \varepsilon > 0$, let $B(\mathbf{y}, r) \subset U$ be a ball centred at $\mathbf{y} = (y_j) \in \text{supp}(\mu)$ and let $f : B(\mathbf{y}, r) \rightarrow \mathbb{Q}_\nu$ be a C^l function with $l \geq 2$ such that*

$$\|\nabla f(\mathbf{y})\| \geq c \quad (6.4)$$

and

$$\|\bar{\Phi}_\beta f\|_{B(\mathbf{y}, r)^{\beta(1)}} \leq \frac{\varepsilon}{r^2} \quad \text{for all multi-indices } \beta \text{ with } |\beta| = 2, \quad (6.5)$$

where $\beta(1)$ is defined in Section 6.1. Then

$$\mu(\{\mathbf{x} \in B(\mathbf{y}, r) \mid |f(\mathbf{x})| < \varepsilon\}) \leq C \left(\frac{\varepsilon}{cr} \right)^\alpha \mu(B(\mathbf{y}, r)). \quad (6.6)$$

Proof. First, observe that for $\mathbf{x} \in B(\mathbf{y}, r)$, by Proposition 11.1 and (8.1), we have that

$$f(\mathbf{x}) = f(\mathbf{y}) + \sum_{i=1}^d (x_i - y_i) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i}) + \sum_{\substack{\beta=(i_1, \dots, i_d) \\ |\beta|=2}} \bar{\Phi}_\beta f(\mathbf{r}^\beta) \prod_{j=1}^d (x_j - y_j)^{i_j}.$$

Here the coordinates of \mathbf{r}^β depend on \mathbf{x} and \mathbf{y} . Therefore

$$\begin{aligned} & |f(\mathbf{x}) - f(\mathbf{y}) - \sum_{i=1}^d (x_i - y_i) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i})| \\ & \leq \max_{|\beta|=2} |\bar{\Phi}_\beta f(\mathbf{r}^\beta)| \prod_{j=1}^d |x_j - y_j|^{i_j} \\ & \stackrel{(6.5)}{\leq} \frac{\varepsilon}{r^2} r^2 = \varepsilon. \end{aligned}$$

Therefore, by the ultra-metric property, for any $\mathbf{x} = (x_1, \dots, x_d)$ and $\mathbf{x}_0 = (x_1^0, \dots, x_d^0) \in B(\mathbf{y}, r)$, we have that

$$|f(\mathbf{x}) - f(\mathbf{x}_0) - \sum_{i=1}^d (x_i - x_i^0) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i})| \leq \varepsilon. \quad (6.7)$$

Now take $\mathbf{x}_0 \in B(\mathbf{y}, r)$ for which $|f(\mathbf{x}_0)| < \varepsilon$, otherwise there is nothing to prove. Then, by Equation (6.7), for any $\mathbf{x} \in B(\mathbf{y}, r)$ with $|f(\mathbf{x})| < \varepsilon$, we have $\left| \sum_{i=1}^d (x_i - x_i^0) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i}) \right| \leq \varepsilon$. Hence,

$$\{\mathbf{x} \in B(\mathbf{y}, r) \mid |f(\mathbf{x})| < \varepsilon\} \subset \left\{ \mathbf{x} \in B(\mathbf{y}, r) \mid \left| \sum_{i=1}^d (x_i - x_i^0) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i}) \right| \leq \varepsilon \right\}. \quad (6.8)$$

Note that $\mathbf{x} \rightarrow \sum_{i=1}^d (x_i - x_i^0) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i})$ is an affine map. Therefore, since μ is absolutely (C, α) -decaying on U , by definition, this affine map is absolutely (C, α) -good on $B(\mathbf{y}, r) \subset U$ w.r.t. μ . Further note that,

$$\max_{\mathbf{x} \in B(\mathbf{y}, r)} \left| \sum_{i=1}^d (x_i - x_i^0) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i}) \right| = r \max_{1 \leq i \leq d} |\bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i})| \stackrel{(6.4)}{\geq} r c.$$

Hence, by the fact that the above affine map is absolutely (C, α) -good on $B(\mathbf{y}, r)$ w.r.t. μ , and in view of Remark 3.1, we get that

$$\mu \left\{ \mathbf{x} \in B(\mathbf{y}, r) \mid \left| \sum_{i=1}^d (x_i - x_i^0) \bar{\Phi}_{e_i} f(\bar{\mathbf{y}}_{e_i}) \right| \leq \varepsilon \right\} \leq C \left(\frac{\varepsilon}{cr} \right)^\alpha \mu(B(\mathbf{y}, r))$$

and Equation (6.8) now completes the proof. \square

Lemma 6.3. *Let $f : B^d \subset \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$ be a C^{k+1} map, where B is a ball of radius $r > 0$. Suppose that there exists $k \in \mathbb{N}$ and a multi-index $\beta = (i_1, \dots, i_d)$ with $|\beta| = k \in \mathbb{N}$ such that for any two pairs of multi-indices $(\alpha_1, \alpha_2), (\alpha'_1, \alpha'_2)$, with $\alpha_1 + \alpha_2 = \beta = \alpha'_1 + \alpha'_2$ and $|\alpha_1| > |\alpha'_1|$, we have that*

$$\inf_{B^{|\alpha_1|+d}} |\bar{\Phi}_{\alpha_1} \partial_{\alpha_2} f| \leq C_{\alpha_1, \alpha'_1} \inf_{B^{|\alpha'_1|+d}} |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} f| \quad (6.9)$$

and

$$\inf_{B^{|\beta|+d}} |\bar{\Phi}_\beta f| > \frac{s}{r^k} \|f\|_{B^d} \quad (6.10)$$

for some $C_{\alpha_1, \alpha'_1} \geq 1, s > 0$. Also, suppose that for some $S > 0$, for every multi-index α with $|\alpha| \leq k+1$,

$$\sup_{B^{|\alpha|+d}} |\bar{\Phi}_\alpha f| \leq \frac{S}{r^{|\alpha|}} \|f\|_{B^d}. \quad (6.11)$$

Then, if $i_j > 0$, the function $\partial_{e_j} f$ also satisfies Equation (6.9), Equation (6.10) and Equation (6.11) with multi-index $\beta - e_j$ with constants $\frac{s}{C_{\beta, \beta - e_j} S}$, $\frac{C_{\beta, \beta - e_j} S}{s}$ in place of s and S respectively. Consequently we have that

$$\frac{s|\beta!|}{C_{\beta, \beta - e_j} r} \|f\|_{B^d} < \|\partial_{e_j} f\|_{B^d} < \frac{S}{r} \|f\|_{B^d}. \quad (6.12)$$

Proof. By Lemma 11.1,

$$\bar{\Phi}^i g'(x_1, \dots, x_{i+1}) = \sum_{n=1}^{i+1} \bar{\Phi}^{i+1} g(x_1, \dots, x_{n-1}, x_n, x_n, x_{n+1}, \dots, x_{i+1}), \quad (6.13)$$

for a C^{j+1} map $g : \mathbb{Q}_\nu \rightarrow \mathbb{Q}_\nu$ and $i \leq j, j > 0$.

Using Equation (6.13) for $g = \partial_{e_j} f$ we have,

$$\begin{aligned} & \bar{\Phi}_{\beta - e_j} \partial_{e_j} f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{x}_j, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\ &= \sum_{n=1}^{i_j+1} \bar{\Phi}_\beta f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,n-1}, x_{j,n}, x_{j,n}, x_{j,n+1}, \dots, x_{j,i_j+1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d). \end{aligned}$$

Therefore Lemma 6.1 gives the following:

$$|\beta!|^{-1} \frac{1}{r^{k-1}} \|\partial_{e_j} f\|_{B^d} \stackrel{(6.1)}{\geq} \inf_{B^{(k-1)+d}} |\bar{\Phi}_{\beta - e_j} \partial_{e_j} f| \stackrel{(6.9)}{\geq} \frac{1}{C_{\beta, \beta - e_j}} \inf_{B^{k+d}} |\bar{\Phi}_\beta f| \stackrel{(6.10)}{\geq} \frac{s}{C_{\beta, \beta - e_j} r^k} \|f\|_{B^d}. \quad (6.14)$$

Above we have used the fact that $|\beta!| \leq |(\beta - e_j)!|$ if $i_j > 1$ an integer, which follows from the ultrametric property of the norm on \mathbb{Q}_ν . By Equation (6.11), $\|\partial_{e_j} f\|_{B^d} \leq \frac{S}{r} \|f\|_{B^d}$. Combining this with Equation (6.14) we have Equation (6.12). \square

Note that $s \leq S$, that follows by taking $\alpha = \beta$ in (6.11), and comparing it to (6.10). Define

$$\mathcal{A}_\alpha := \{(\alpha_1, \alpha'_1) \mid \exists \alpha_2, \alpha'_2 \text{ multi-indices such that } \alpha_1 + \alpha_2 = \alpha'_1 + \alpha'_2 = \alpha, |\alpha_1| > |\alpha'_1|\}.$$

Theorem 6.1. Suppose that $f : B^d \subset \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$ is a C^{k+1} map, $B \subset \mathbb{Q}_\nu$ is an open ball of radius $r > 0$ and that for some $k \in \mathbb{N}$ and multi-index β with $|\beta| = k$, f satisfies Equation (6.9), Equation (6.10) and Equation (6.11). Let μ be an absolutely (C, α) -decaying measure on B^d . Then for any $\varepsilon < \min(s, \frac{s}{S})$ one has

$$\mu(\{x \in B^d \mid |f(x)| < \varepsilon \|f\|_{B^d}\}) \leq C 2^k \left(\prod_{(\alpha_1, \alpha'_1) \in \mathcal{A}_\beta} C_{\alpha_1, \alpha'_1} \right)^{2^k \alpha} \frac{1}{|\beta!|^\alpha} \left(\frac{S}{s} \right)^{2^k \alpha} \varepsilon^{\eta_k \alpha} \mu(B^d), \quad (6.15)$$

where $\eta_k = \frac{1}{2^{k+1}-2}$ for $k \geq 1$, and $\eta_0 = \infty$ and $0! =: 1$

Proof. Let us define the following set for $\delta > 0$

$$E(f, \delta, B^d) := \{\mathbf{x} \in B^d \mid |f(\mathbf{x})| < \delta\}.$$

We will proceed by induction on k . If $k = 0$, then by (6.10) and $\varepsilon < s$, the left hand side set of (6.15) becomes empty, so the left hand side is 0. On the other hand, $\mathcal{A}_\beta = \{0\}$ and $C_0 \geq 1$, so the right hand side is infinite. Hence the conclusion of the theorem follows trivially in the

case $k = 0$. Suppose $\beta = (i_1, \dots, i_d)$ and $i_j > 0$. We know that $\partial_{e_j} f$ satisfies the conditions Equation (6.9), Equation (6.10) and Equation (6.11), by Lemma 6.3. Choose

$$\delta = \varepsilon^{\frac{1}{2(\eta_{k-1}+1)}}.$$

Applying the induction hypothesis, we get

$$\begin{aligned} \mu(E(\partial_{e_j} f, \delta \|\partial_{e_j} f\|_{B^d}, B^d)) &\leq \\ C 2^{k-1} \left(\prod_{(\alpha_1, \alpha'_1) \in \mathcal{A}_{\beta-e_j}} C_{\alpha_1, \alpha'_1} \right)^{2^{k-1}\alpha} &\frac{1}{|(\beta - e_j)!|^\alpha} \left(\frac{C_{\beta, \beta-e_j}^2 S^2}{s^2} \right)^{2^{k-1}\alpha} \delta^{\eta_{k-1}\alpha} \mu(B^d). \end{aligned} \quad (6.16)$$

Let us choose $t = r\sqrt{\frac{\varepsilon}{S}}$. For every $\mathbf{y} \in B^d \cap \text{supp } \mu \setminus E(\partial_{e_j} f, \delta \|\partial_{e_j} f\|_{B^d}, B^d)$, consider the ball $B(\mathbf{y}, t)$. By the choice of ε , and the fact that $s \leq S$ which we noted earlier, we have that $B(\mathbf{y}, t) \subset B^d$. Since $\mathbf{y} \notin E(\partial_{e_j} f, \delta \|\partial_{e_j} f\|_{B^d}, B^d)$,

$$|\partial_{e_j} f(\mathbf{y})| \geq \delta \|\partial_{e_j} f\|_{B^d}$$

and, by Equation (6.11) for every β_\star multi-index, with $|\beta_\star| = 2$, we have

$$\|\bar{\Phi}_{\beta_\star} f\|_{B(\mathbf{y}, t)^{\beta_\star(1)}} \leq \frac{S}{r^2} \|f\|_{B^d} = \frac{\varepsilon \|f\|_{B^d} S}{r^2 \varepsilon} = \frac{\varepsilon \|f\|_{B^d}}{t^2}.$$

Now we can use Lemma 6.2 and Equation (6.12) and conclude that

$$\begin{aligned} &\mu(B(\mathbf{y}, t) \cap E(f, \varepsilon \|f\|_{B^d}, B^d)) \\ &\stackrel{(6.6)}{\leq} C \left(\frac{\varepsilon \|f\|_{B^d}}{\delta t \|\partial_{e_j} f\|_{B^d}} \right)^\alpha \mu(B(\mathbf{y}, t)) \\ &\stackrel{(6.12)}{\leq} C \left(\frac{\varepsilon C_{\beta, \beta-e_j} r}{\delta t s |\beta|!} \right)^\alpha \mu(B(\mathbf{y}, t)) \\ &\leq C C_{\beta, \beta-e_j}^\alpha \frac{1}{|\beta|!^\alpha} \left(\frac{\sqrt{S}}{s} \right)^\alpha \frac{\varepsilon^{\frac{\alpha}{2}}}{\delta^\alpha} \mu(B(\mathbf{y}, t)). \end{aligned}$$

By the Besicovitch covering property of \mathbb{Q}_ν^d , one can cover $B^d \cap \text{supp } \mu \setminus E(\partial_{e_j} f, \delta \|\partial_{e_j} f\|_{B^d}, B^d)$ by balls $B(\mathbf{y}, t)$ with multiplicity 1. This means (3.1) holds with $N_X = 1, X = \mathbb{Q}_\nu^d$; see §3.1. Hence

$$\begin{aligned} &\mu \left((B^d \setminus E(\partial_{e_j} f, \delta \|\partial_{e_j} f\|_{B^d}, B^d)) \cap E(f, \varepsilon \|f\|_{B^d}, B^d) \right) \\ &\leq C C_{\beta, \beta-e_j}^\alpha \frac{1}{|\beta|!^\alpha} \left(\frac{\sqrt{S}}{s} \right)^\alpha \frac{\varepsilon^{\frac{\alpha}{2}}}{\delta^\alpha} \mu(B^d). \end{aligned} \quad (6.17)$$

Recall that $\delta = \varepsilon^{\frac{1}{2(\eta_{k-1}+1)}}$. Then combining Equation (6.16) and Equation (6.17), we have that

$$\begin{aligned}
& \mu(B^d \cap E(f, \varepsilon \|f\|_{B^d}, B^d)) \\
& \leq C 2^{k-1} \left(\prod_{(\alpha_1, \alpha'_1) \in \mathcal{A}_{\beta-e_j}} C_{\alpha_1, \alpha'_1} \right)^{2^{k-1}\alpha} \frac{1}{|(\beta - e_j)!|^\alpha} \left(\frac{C_{\beta, \beta-e_j}^2 S^2}{s^2} \right)^{2^{k-1}\alpha} \delta^{\eta_{k-1}\alpha} \mu(B^d) + \\
& \quad C C_{\beta, \beta-e_j}^\alpha \frac{1}{|\beta!|^\alpha} \left(\frac{\sqrt{S}}{s} \right)^\alpha \frac{\varepsilon^{\frac{\alpha}{2}}}{\delta^\alpha} \mu(B^d) \\
& \leq C 2^{k-1} \left(\prod_{(\alpha_1, \alpha'_1) \in \mathcal{A}_\beta} C_{\alpha_1, \alpha'_1} \right)^{2^k\alpha} \frac{1}{|\beta!|^\alpha} \left(\frac{S}{s} \right)^{2^k\alpha} \left(\delta^{\eta_{k-1}\alpha} + \frac{\varepsilon^{\frac{\alpha}{2}}}{\delta^\alpha} \right) \mu(B^d) \\
& \leq C 2^k \left(\prod_{(\alpha_1, \alpha'_1) \in \mathcal{A}_\beta} C_{\alpha_1, \alpha'_1} \right)^{2^k\alpha} \frac{1}{|\beta!|^\alpha} \left(\frac{S}{s} \right)^{2^k\alpha} \varepsilon^{\frac{\eta_{k-1}\alpha}{2(\eta_{k-1}+1)}} \mu(B^d).
\end{aligned}$$

Hence, the theorem is proved with $\eta_k = \frac{\eta_{k-1}}{2(\eta_{k-1}+1)}$, which gives $\eta_k = \frac{1}{2^{k+1}-2}$. \square

7. POLYNOMIALS SATISFYING THE HYPOTHESES IN THEOREM 6.1

Let us denote by $\mathcal{P}_{d,l}$ the family of polynomials of degree l in d variables.

Lemma 7.1. *Suppose $P \in \mathcal{P}_{d,l}$ be a nonzero element and $\eta : \mathbb{N} \cup \{0\} \rightarrow \mathbb{N}$ be a function such that $\eta(n+1) \geq \eta(n) + 1$. Suppose k is the largest integer such that there exists β , a multi-index with $|\beta| = k$, with*

$$|\bar{\Phi}_\beta P(\mathbf{x}_1, \dots, \mathbf{x}_d)| > \frac{1}{\nu^{\eta(k)}}$$

for some $(\mathbf{x}_1, \dots, \mathbf{x}_d) \in B(\mathbf{0}, 1)^{\beta(1)}$. Then for every $(\mathbf{y}_1, \dots, \mathbf{y}_d) \in B(\mathbf{0}, 1)^{\beta(1)}$,

$$|\bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{y}_d)| > \frac{1}{\nu^{\eta(k)+1}}. \quad (7.1)$$

Proof. Note

$$\begin{aligned}
& \bar{\Phi}_\beta P(\mathbf{x}_1, \dots, \mathbf{x}_d) - \bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{y}_d) \\
& = \sum_{j=0}^{d-1} \bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{y}_j, \mathbf{x}_{j+1}, \mathbf{x}_{j+2}, \dots, \mathbf{x}_d) - \bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{y}_j, \mathbf{y}_{j+1}, \mathbf{x}_{j+2}, \dots, \mathbf{x}_d) \\
& = \sum_{j=0}^{d-1} \sum_{q=0}^{i_{j+1}} \left(\bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{y}_j, x_{j+1,1}, \dots, x_{j+1, i_{j+1}+1-q}, y_{j+1,1}, \dots, y_{j+1,q}, \mathbf{x}_{j+2}, \dots, \mathbf{x}_d) \right. \\
& \quad \left. - \bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{y}_j, x_{j+1,1}, \dots, x_{j+1, i_{j+1}-q}, y_{j+1,1}, \dots, y_{j+1, q+1}, \mathbf{x}_{j+2}, \dots, \mathbf{x}_d) \right) \\
& = \sum_{j=0}^{d-1} \sum_{q=0}^{i_{j+1}} \bar{\Phi}_{\beta+e_j} P(\mathbf{y}_1, \dots, \mathbf{y}_j, x_{j+1,1}, \dots, x_{j+1, i_{j+1}+1-q}, y_{j+1,1}, \dots, y_{j+1, q+1}, \mathbf{x}_{j+2}, \dots, \mathbf{x}_d) \delta_{j,q}
\end{aligned}$$

where $\delta_{j,q} = (x_{j+1,i_{j+1}+1-q} - y_{j+1,q+1})$. Therefore, we have

$$\begin{aligned} & |\bar{\Phi}_\beta P(\mathbf{x}_1, \dots, \mathbf{x}_d)| - |\bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{x}_d)| \\ & \leq \max_{j=0}^{d-1} \max_{q=0}^{i_{j+1}} \left\{ |\bar{\Phi}_{\beta+e_j} P(\mathbf{y}_1, \dots, \mathbf{y}_j, x_{j+1,1}, \dots, x_{j+1,i_{j+1}+1-q}, y_{j+1,1}, \dots, y_{j+1,q+1}, \mathbf{x}_{j+2}, \dots, \mathbf{x}_d)| \right\}. \end{aligned}$$

We proceed by contradiction. Suppose there exists a $(\mathbf{y}_1, \dots, \mathbf{y}_d) \in B(\mathbf{0}, 1)^{\beta(1)}$ such that

$$|\bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{y}_d)| \leq \frac{1}{\nu^{\eta(k)+1}}. \quad (7.2)$$

Then we have

$$\frac{1}{\nu^{\eta(k+1)}} < \frac{1}{\nu^{\eta(k)}} - \frac{1}{\nu^{\eta(k)+1}} < |\bar{\Phi}_\beta P(\mathbf{x}_1, \dots, \mathbf{x}_d)| - |\bar{\Phi}_\beta P(\mathbf{y}_1, \dots, \mathbf{x}_d)|.$$

The above will imply that there exists $0 \leq j \leq d-1$, and $0 \leq q \leq i_{j+1}$ such that

$$|\bar{\Phi}_{\beta+e_j} P(\mathbf{y}_1, \dots, \mathbf{y}_j, x_{j+1,1}, \dots, x_{j+1,i_{j+1}+1-q}, y_{j+1,1}, \dots, y_{j+1,q+1}, \mathbf{x}_{j+2}, \dots, \mathbf{x}_d)| > \frac{1}{\nu^{\eta(k+1)}}.$$

Since $|\beta + e_j| = k+1$, the last inequality contradicts the maximality of k . \square

In the following theorem $\alpha, \alpha_1, \alpha'_1, \alpha_2, \alpha'_2$ and β are all multi-indices.

Theorem 7.1. *Fix d and l . There exist $s, S > 0$ and a collection $\bigcup_{\bar{\eta}, |\bar{\eta}| \leq l} \{C_{\alpha_1, \alpha'_1} : (\alpha_1, \alpha'_1) \in \mathcal{A}_{\bar{\eta}}\}$ of constants greater than 1 such that for any nonzero $P \in \mathcal{P}_{d,l}$ the following are satisfied.*

- *There exists $k \leq l$ and a multi-index $\beta = (i_1, \dots, i_d)$ with $|\beta| = k \in \mathbb{N}$ such that, for every $\alpha_1 + \alpha_2 = \beta = \alpha'_1 + \alpha'_2$, $|\alpha_1| > |\alpha'_1|$,*

$$\inf_{B(0,1)^{|\alpha_1|+d}} |\bar{\Phi}_{\alpha_1} \partial_{\alpha_2} P| \leq C_{\alpha_1, \alpha'_1} \inf_{B(0,1)^{|\alpha'_1|+d}} |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} P|. \quad (7.3)$$

and

$$\inf_{B(0,1)^{|\beta|+d}} |\bar{\Phi}_\beta P(\mathbf{x}_1, \dots, \mathbf{x}_d)| > s \|P\|_{B(0,1)^d}. \quad (7.4)$$

•

$$\sup_{B(0,1)^{|\alpha|+d}} |\bar{\Phi}_\alpha P(\mathbf{x}_1, \dots, \mathbf{x}_d)| < S \|P\|_{B(0,1)^d} \quad \forall |\alpha| \geq 0. \quad (7.5)$$

Proof. Without loss of generality we assume that $\|P\|_{B(0,1)^d} = 1$. Let $a > 0$ be an integer such that $\frac{1}{\nu^a} < |i| = |i|_\nu$ for $i = 1, \dots, l$. Consider the function $\eta : \mathbb{N} \cup \{0\} \rightarrow \mathbb{N}$, $\eta(n) := n^2 a + na + 1$. Let k be the largest integer such that there exists a multi-index $\beta = (i_1, \dots, i_d)$ with $|\beta| = k$ and $(\mathbf{x}_1, \dots, \mathbf{x}_d) \in B(\mathbf{0}, 1)^{\beta(1)}$ such that

$$|\bar{\Phi}_\beta P(\mathbf{x}_1, \dots, \mathbf{x}_d)| > \frac{1}{\nu^{\eta(k)}}.$$

Such a k exists because for $k = 0$ the above inequality holds by the fact that $\|P\|_{B(0,1)^d} = 1$ and $\eta(0) = 1$. Also note that $k \leq l$. By Lemma 7.1, we have that

$$\inf_{B(0,1)^{|\beta|+d}} |\bar{\Phi}_\beta P| > \frac{1}{\nu^{\eta(k)+1}}. \quad (7.6)$$

If $k = 0$ then Equation (7.3) will be true trivially. So, let us assume $k \neq 0$.

Claim 1: We claim that for any $j = 1, \dots, d$ if $i_j > 0$ then

$$\inf_{B(0,1)^{|\beta|+d}} |\bar{\Phi}_\beta P| \leq |i_j|^{-1} \inf_{B(0,1)^{|\beta|-1+d}} |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P|. \quad (7.7)$$

Let $(\mathbf{x}_1^0, \dots, \mathbf{x}_d^0) \in B(\mathbf{0}, 1)^{(\beta-e_j)(1)}$ be such that

$$\inf_{B(0,1)^{|\beta|-1+d}} |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P| = |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P(\mathbf{x}_1^0, \dots, \mathbf{x}_d^0)|.$$

Now by Lemma 11.2, for any $(\mathbf{x}_1, \dots, \mathbf{x}_d) \in B(\mathbf{0}, 1)^{\beta(1)}$,

$$\begin{aligned} & i_j \bar{\Phi}_{\beta} P(\mathbf{x}_1, \dots, \mathbf{x}_d) \\ &= \bar{\Phi}_{\beta-e_j} \partial_{e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\ &+ \sum_{i=1}^{i_j} \bar{\Phi}_{\beta+e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i-1}, x_{j,i}, x_{j,i}, x_{j,i+1}, \dots, x_{j,i_j+1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) (x_{j,i_j+1} - x_{j,i}). \end{aligned} \quad (7.8)$$

If for some $x_{j,i_j+1} \in B(0, 1)$,

$$\begin{aligned} & \max_{i=1}^{i_j} |\bar{\Phi}_{\beta+e_j} P(\mathbf{x}_1^0, \dots, \mathbf{x}_{j-1}^0, x_{j,1}^0, \dots, x_{j,i-1}^0, x_{j,i}^0, x_{j,i}^0, x_{j,i+1}^0, \dots, x_{j,i_j}^0, x_{j,i_j+1}, \mathbf{x}_{j+1}^0, \dots, \mathbf{x}_d^0)| \\ & \leq |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P(\mathbf{x}_1^0, \dots, \mathbf{x}_{j-1}^0, x_{j,1}^0, \dots, x_{j,i_j}^0, \mathbf{x}_{j+1}^0, \dots, \mathbf{x}_d^0)| \end{aligned}$$

then Equation (7.7) holds using Equation (11.5). If not, then for every $x_{j,i_j+1} \in B(0, 1)$ there exists some $1 \leq i \leq i_j$ such that

$$\begin{aligned} & |\bar{\Phi}_{\beta+e_j} P(\mathbf{x}_1^0, \dots, \mathbf{x}_{j-1}^0, x_{j,1}^0, \dots, x_{j,i-1}^0, x_{j,i}^0, x_{j,i}^0, x_{j,i+1}^0, \dots, x_{j,i_j}^0, x_{j,i_j+1}, \mathbf{x}_{j+1}^0, \dots, \mathbf{x}_d^0)| \\ & > |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P(\mathbf{x}_1^0, \dots, \mathbf{x}_{j-1}^0, x_{j,1}^0, \dots, x_{j,i_j}^0, \mathbf{x}_{j+1}^0, \dots, \mathbf{x}_d^0)|. \end{aligned}$$

Since $|i_j| > \frac{1}{\nu^a}$, we have that

$$\begin{aligned} \frac{1}{\nu^{\eta(k)+1+a}} & < \frac{|i_j|}{\nu^{\eta(k)+1}} < |i_j| |\bar{\Phi}_{\beta} P(\mathbf{x}_1, \dots, \mathbf{x}_d)| \\ & < |\bar{\Phi}_{\beta+e_j} P(\mathbf{x}_1^0, \dots, \mathbf{x}_{j-1}^0, x_{j,1}^0, \dots, x_{j,i-1}^0, x_{j,i}^0, x_{j,i}^0, x_{j,i+1}^0, \dots, x_{j,i_j}^0, x_{j,i_j+1}, \mathbf{x}_{j+1}^0, \dots, \mathbf{x}_d^0)|. \end{aligned}$$

Now note that

$$\frac{1}{\nu^{(\eta(k)+1+a)}} > \frac{1}{\nu^{((k+1)^2 a + (k+1)a+1)}} = \frac{1}{\nu^{\eta(k+1)}},$$

which will give a contradiction to the maximality of k . This yields Claim 1.

Claim 2: If $i_j > 1$ then we claim that

$$\inf_{B(0,1)^{|\beta|-1+d}} |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P| \leq |i_j - 1|^{-1} \inf_{B(0,1)^{|\beta|-2+d}} |\bar{\Phi}_{\beta-2e_j} \partial_{2e_j} P|. \quad (7.9)$$

Note that Claim 1 and Equation (7.6) imply that

$$\inf_{B(0,1)^{|\beta|-1+d}} |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P| > \frac{1}{\nu^{\eta(k)+1+a}}. \quad (7.10)$$

Now observe that by Lemma 11.2, for any $(\mathbf{x}_1, \dots, \mathbf{x}_d) \in B(\mathbf{0}, 1)^{(\beta-e_j)(1)}$,

$$\begin{aligned}
& (i_j - 1) \bar{\Phi}_{\beta-e_j} \partial_{e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_d) \\
&= \bar{\Phi}_{\beta-2e_j} \partial_{2e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i_j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\
&+ \sum_{i=1}^{i_j-1} \bar{\Phi}_{\beta} \partial_{e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i-1}, x_{j,i}, x_{j,i}, x_{j,i+1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) (x_{j,i_j} - x_{j,i}) \\
&\stackrel{\text{Lemma 11.1}}{=} \bar{\Phi}_{\beta-2e_j} \partial_{2e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i_j-1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\
&+ \sum_{i=1}^{i_j-1} \left(\sum_{b=1}^{i_j} \bar{\Phi}_{\beta+e_j} P(\mathbf{t}_{b,i}) \right) (x_{j,i_j} - x_{j,i}),
\end{aligned}$$

where $\mathbf{t}_{b,i} = (\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,i}, x_{j,1}, \dots, x_{j,b}, x_{j,b}, x_{j,b+1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d)$ has coordinates in terms of \mathbf{x} . In the last line of the above equality, in particular, we use Lemma 11.1 to get

$$\bar{\Phi}_{\beta} \partial_{e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i-1}, x_{j,i}, x_{j,i}, x_{j,i+1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) = \left(\sum_{b=1}^{i_j} \bar{\Phi}_{\beta+e_j} P(\mathbf{t}_{b,i}) \right).$$

Note that $\frac{1}{\nu^{\eta(k)+1+2a}} > \frac{1}{\nu^{\eta(k+1)}}$. Hence the same argument as in Claim 1 using Equation (7.10) yields Claim 2.

Claim c : Continuing this iterative method we get that for any $j = 1, \dots, d$, if $i_j > c + 1$ then,

$$\inf_{B(0,1)^{|\beta|-c+d}} |\bar{\Phi}_{\beta-ce_j} \partial_{ce_j} P| \leq |i_j - c|^{-1} \inf_{B(0,1)^{|\beta|-c-1+d}} |\bar{\Phi}_{\beta-(c+1)e_j} \partial_{(c+1)e_j} P|. \quad (7.11)$$

Claim $(\beta - e_j, \beta - e_j - e_m)$: We claim that

$$\inf_{B(0,1)^{|\beta|-1+d}} |\bar{\Phi}_{\beta-e_j} \partial_{e_j} P| \leq |i_m|^{-1} \inf_{B(0,1)^{|\beta|-2+d}} |\bar{\Phi}_{\beta-e_j-e_m} \partial_{e_j+e_m} P|. \quad (7.12)$$

If $i_m > 0$ and $i_j > 0$ then by Lemma 11.2 and Lemma 11.1 for any $(\mathbf{x}_1, \dots, \mathbf{x}_d) \in B(\mathbf{0}, 1)^{(\beta-e_j)(1)}$, one can write the following:

$$\begin{aligned}
& i_m \bar{\Phi}_{\beta-e_j} \partial_{e_j} P(\mathbf{x}_1, \dots, \mathbf{x}_d) \\
&= \bar{\Phi}_{\beta-e_j-e_m} \partial_{e_j+e_m} P(\mathbf{x}_1, \dots, \mathbf{x}_{m-1}, x_{m,1}, \dots, x_{m,i_m}, \mathbf{x}_{m+1}, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\
&+ \sum_{i=1}^{i_m} \left(\sum_{b=1}^{i_j} \bar{\Phi}_{\beta+e_m} P(\mathbf{t}'_{b,i}) \right) (x_{m,i_m+1} - x_{m,i}),
\end{aligned}$$

where

$$\mathbf{t}'_{b,i} = (\mathbf{x}_1, \dots, \mathbf{x}_{m-1}, \delta_{m,i}, \mathbf{x}_{m+1}, \dots, \mathbf{x}_{j-1}, \kappa_{j,b}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d), \text{ and}$$

$\delta_{m,i} = (x_{m,1}, \dots, x_{m,i}, x_{m,i}, x_{m,i+1}, \dots, x_{m,i_m+1})$ and $\kappa_{j,b} = (x_{j,1}, \dots, x_{j,b}, x_{j,b}, x_{j,b+1}, \dots, x_{j,i_j})$ have coordinates in terms of \mathbf{x} . The deduction of the above from Lemma 11.2 and Lemma 11.1 follows the same way as in (the first unnumbered equations) Claim 2. Now the same argument as in Claim 1 using Equation (7.10) yields Claim $(\beta - e_j, \beta - e_j - e_m)$. One can continue with the same arguments and get Equation (7.3).

The upper bound, Equation (7.5) follows from the observation that $P \rightarrow \|P\|_{B(0,1)^d}$ and $P \rightarrow \max_{\alpha} \|\bar{\Phi}_{\alpha} P\|_{B(0,1)^{|\alpha|+d}}$ are two norms on $\mathcal{P}_{d,l}$, a finite dimensional vector space over \mathbb{Q}_{ν} . Hence they must be equivalent.

It is clear from (7.6), that we may take $s = \frac{1}{\nu\eta(l)+1}$. Also note that C_{α_1, α'_1} , we find in the proof depends on the indices of $\alpha_1, \alpha'_1, \beta$, and since $|\beta| \leq l$, these constants can be made only dependent on l and d ; see (7.7), (7.9), (7.11), (7.12). \square

8. APPROXIMATION BY POLYNOMIALS

Let us denote $y^k := (y, \dots, y) \in \mathbb{Q}_\nu^k$ for $y \in \mathbb{Q}_\nu$, and $k \in \mathbb{N}$. We denote $\bar{\mathbf{y}}_\beta := (y_1^{i_1+1}, \dots, y_d^{i_d+1})$, where $\beta = (i_1, \dots, i_d)$. Suppose that $f : B^d \subset B(0, 1)^d \rightarrow \mathbb{Q}_\nu$ is a C^{l+1} map for some $l \in \mathbb{N}$. For a multi-index $\beta := (i_1, \dots, i_d)$ we set $L_{\beta, \mathbf{y}}(\mathbf{x}) := \prod_{j=1}^d (x_j - y_j)^{i_j}$, where $\mathbf{y} \in \mathbb{Q}_\nu^d$. When $\beta = 0$, we define $L_{\beta, \mathbf{y}} = 1$. Note that

$$\bar{\Phi}_{\beta'} L_{\beta, \mathbf{y}} = 1, \text{ when } \beta = \beta',$$

and

$$\bar{\Phi}_{\beta'} L_{\beta, \mathbf{y}} = 0, \text{ when } \beta \neq \beta', |\beta| = |\beta'|.$$

We denote the l -th Taylor polynomial of f at $\mathbf{y} \in B^d$ as

$$P_{f, \mathbf{y}, l}(\mathbf{x}) := f(\mathbf{y}) + \sum_{k=1}^l \sum_{|\beta|=k} \bar{\Phi}_\beta f(\bar{\mathbf{y}}_\beta) L_{\beta, \mathbf{y}}(\mathbf{x}). \quad (8.1)$$

Note that it follows from the definition that,

$$\bar{\Phi}_{\beta+e_1} f(z, \bar{\mathbf{y}}_\beta) = \bar{\Phi}_{\beta+e_1} f(\bar{\mathbf{y}}_{\beta+e_1}) + (z - y_1) \bar{\Phi}_{\beta+2e_1} f(z, \bar{\mathbf{y}}_{\beta+e_1}). \quad (8.2)$$

Let us denote $f_a^i(\mathbf{x}) := \bar{\Phi}_{e_i} f(a, \mathbf{x}^{(i)})$, where $(a, \mathbf{x}^{(i)}) := (x_1, \dots, x_{i-1}, a, x_i, x_{i+1}, \dots, x_d)$ for $\mathbf{x} \in B^d$, $a \in B$, and $i = 1, \dots, d$. Also, for $\mathbf{x}_\beta = (\mathbf{x}_1, \dots, \mathbf{x}_d) \in B^\beta$, let us denote $(a, \mathbf{x}_\beta^{(i)}) := (\mathbf{x}_1, \dots, \mathbf{x}_{i-1}, a, \mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_d)$, and $(a, b, \mathbf{x}_\beta^{(i)}) := (\mathbf{x}_1, \dots, \mathbf{x}_{i-1}, a, b, \mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_d)$.

Therefore note by definition,

$$\bar{\Phi}_{\alpha+e_i} f(y_i, \mathbf{x}_{\alpha(1)}^{(i)}) = \bar{\Phi}_\alpha f_{y_i}^i(\mathbf{x}_{\alpha(1)}).$$

Lemma 8.1. *Suppose that $z \in B, \mathbf{y} \in B^d$. Let $f : B^d \subset B(0, 1)^d \rightarrow \mathbb{Q}_\nu$ a C^{l+1} map. For any multi-index α with $|\alpha| \leq l-1$ and for any $\mathbf{x}_{\alpha(1)} \in B^{\alpha(1)}$ and $i = 1, \dots, d$ we have,*

$$\begin{aligned} & \bar{\Phi}_{\alpha+e_i} P_{f, \mathbf{y}, l}(z, \mathbf{x}_{\alpha(1)}^{(i)}) \\ &= (z - y_i) \bar{\Phi}_{\alpha+e_i} P_{f_{y_i}^i, \mathbf{y}, l-1}(z, \mathbf{x}_{\alpha(1)}^{(i)}) + \bar{\Phi}_\alpha P_{f_{y_i}^i, \mathbf{y}, l-1}(\mathbf{x}_{\alpha(1)}). \end{aligned} \quad (8.3)$$

Proof. Let us note that

$$\begin{aligned}
& \bar{\Phi}_{\alpha+e_i} P_{f,\mathbf{y},l}(z, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= \sum_{|\beta|=0} \bar{\Phi}_{\beta} f(\bar{\mathbf{y}}_{\beta}) \bar{\Phi}_{\alpha+e_i} L_{\beta,\mathbf{y}}(z, \mathbf{x}_{\alpha(1)}^{(i)}) + \cdots + \sum_{|\beta|=l} \bar{\Phi}_{\beta} f(\bar{\mathbf{y}}_{\beta}) \bar{\Phi}_{\alpha+e_i} L_{\beta,\mathbf{y}}(z, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= \sum_{|\beta|=0} \bar{\Phi}_{\beta+e_i} f(\bar{\mathbf{y}}_{\beta+e_i}) \bar{\Phi}_{\alpha+e_i} L_{\beta+e_i,\mathbf{y}}(z, \mathbf{x}_{\alpha(1)}^{(i)}) + \cdots + \sum_{|\beta|=l-1} \bar{\Phi}_{\beta+e_i} f(\bar{\mathbf{y}}_{\beta+e_i}) \bar{\Phi}_{\alpha+e_i} L_{\beta+e_i,\mathbf{y}}(z, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= \bar{\Phi}_{\alpha+e_i} \left(L_{e_i,\mathbf{y}} \times \left(\sum_{|\beta|=0} \bar{\Phi}_{\beta+e_i} f(\bar{\mathbf{y}}_{\beta+e_i}) L_{\beta,\mathbf{y}} + \cdots + \sum_{|\beta|=l-1} \bar{\Phi}_{\beta+e_i} f(\bar{\mathbf{y}}_{\beta+e_i}) L_{\beta,\mathbf{y}} \right) \right) (z, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= \bar{\Phi}_{\alpha+e_i} \left(L_{e_i,\mathbf{y}} \times \left(\sum_{|\beta|=0} \bar{\Phi}_{\beta} f_{y_i}^i(\bar{\mathbf{y}}_{\beta}) L_{\beta,\mathbf{y}} + \cdots + \sum_{|\beta|=l-1} \bar{\Phi}_{\beta} f_{y_i}^i(\bar{\mathbf{y}}_{\beta}) L_{\beta,\mathbf{y}} \right) \right) (z, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= \bar{\Phi}_{\alpha+e_i} \left(L_{e_i,\mathbf{y}} \times \left(P_{f_{y_i}^i,\mathbf{y},l-1} \right) \right) (z, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= L_{e_i,\mathbf{y}}(z) \bar{\Phi}_{\alpha+e_i} \left(P_{f_{y_i}^i,\mathbf{y},l-1} \right) (z, \mathbf{x}_{\alpha(1)}^{(i)}) + \bar{\Phi}_{\alpha} \left(P_{f_{y_i}^i,\mathbf{y},l-1} \right) (\mathbf{x}_{\alpha(1)}).
\end{aligned}$$

In the equations above, by \times we mean multiplication of functions. The last line follows from Lemma 11.3, which is a version of the chain rule. \square

Theorem 8.1. *Suppose that $f : B(\mathbf{y}, r) \subset B(0, 1)^d \rightarrow \mathbb{Q}_{\nu}$ be a C^{l+1} map, then for any $0 \leq |\eta| \leq l$,*

$$\begin{aligned}
& \sup_{\mathbf{x}_{\eta(1)} \in B(\mathbf{y}, r)^{\eta(1)}} |\bar{\Phi}_{\eta} f(\mathbf{x}_{\eta(1)}) - \bar{\Phi}_{\eta} P_{f,\mathbf{y},l}(\mathbf{x}_{\eta(1)})| \\
& \leq r^{l-|\eta|} \sup_{\substack{\mathbf{w}_1, \mathbf{w}_2 \in B(\mathbf{y}, r)^{\beta(1)}, \\ |\beta|=l}} |\bar{\Phi}_{\beta} f(\mathbf{w}_1) - \bar{\Phi}_{\beta} f(\mathbf{w}_2)|.
\end{aligned} \tag{8.4}$$

Proof. We proceed by induction on l . If $l = 0$, then $|\eta| = 0$ and $P_{f,\mathbf{y},0}(\mathbf{x}) = f(\mathbf{y})$, hence the statement of the theorem follows for $l = 0$. Assume the the statement is true for $l - 1$. We examine η a multi index with $|\eta| \leq l$ in three cases, $|\eta| = 0, l$ and $1 \leq |\eta| \leq l - 1$. Let $l - 1 \geq k \geq 1$ and fix any $1 \leq i \leq d$. Take α be a multi index such that $|\alpha| = k - 1$. Let

$$\mathbf{M} = \sup_{\substack{\mathbf{w}_1, \mathbf{w}_2 \in B(\mathbf{y}, r)^{\beta(1)}, \\ |\beta|=l}} |\bar{\Phi}_{\beta} f(\mathbf{w}_1) - \bar{\Phi}_{\beta} f(\mathbf{w}_2)|.$$

By the induction hypothesis, we have for any $\mathbf{x}_{\alpha(1)} \in B(\mathbf{y}, r)^{\alpha(1)}$,

$$|\bar{\Phi}_{\alpha} P_{f_{y_i}^i,\mathbf{y},l-1}(\mathbf{x}_{\alpha(1)}) - \bar{\Phi}_{\alpha} f_{y_i}^i(\mathbf{x}_{\alpha(1)})| \leq r^{l-k} \mathbf{M}. \tag{8.5}$$

Also using induction we have for any $(z, \mathbf{x}_{\alpha(1)}^{(i)}) \in B(\mathbf{y}, r)^{\alpha+e_i}$,

$$|\bar{\Phi}_{\alpha+e_i} P_{f_{y_i}^i,\mathbf{y},l-1}(z, \mathbf{x}_{\alpha(1)}^{(i)}) - \bar{\Phi}_{\alpha+e_i} f_{y_i}^i(z, \mathbf{x}_{\alpha(1)}^{(i)})| \leq r^{l-1-k} \mathbf{M}. \tag{8.6}$$

On the right hand side of Equation (8.5) and Equation (8.6) \mathbf{M} appears because for any multi index β with $|\beta| = l - 1$, $\bar{\Phi}_{\beta} f_{y_i}^i(\mathbf{w}) = \bar{\Phi}_{\beta+e_i} f(y_i, \mathbf{w}^{(i)})$.

Note that

$$\begin{aligned}
\bar{\Phi}_{\alpha+e_i} f(z, \mathbf{x}_{\alpha(1)}^{(i)}) &= (z - y_i) \bar{\Phi}_{\alpha+2e_i} f(z, y_i, \mathbf{x}_{\alpha(1)}^{(i)}) + \bar{\Phi}_{\alpha+e_i} f(y_i, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= (z - y_i) \bar{\Phi}_{\alpha+e_i} f_{y_i}^i(z, \mathbf{x}_{\alpha(1)}^{(i)}) + \bar{\Phi}_{\alpha} f_{y_i}^i(\mathbf{x}_{\alpha(1)}).
\end{aligned}$$

In the last line of the above equality, we use the definition of the function $f_{y_i}^i(\mathbf{x}) = \bar{\Phi}_{e_i} f(y_i, \mathbf{x}^{(i)})$. Therefore, by Lemma 8.1 we have the following:

$$|\bar{\Phi}_{\alpha+e_i} P_{f,\mathbf{y},l}(z, \mathbf{x}_{\alpha(1)}^{(i)}) - \bar{\Phi}_{\alpha+e_i} f(z, \mathbf{x}_{\alpha(1)}^{(i)})| \leq r^{l-k} \mathbf{M}. \quad (8.7)$$

Now let us take a multi-index α , such that $|\alpha| = l - 1$, then we have

$$\bar{\Phi}_{\alpha+e_i} P_{f,\mathbf{y},l}(z, \mathbf{x}_{\alpha(1)}^{(i)}) = \bar{\Phi}_{\alpha+e_i} f(\bar{\mathbf{y}}_{\alpha+e_i}).$$

Therefore for any $(z, \mathbf{x}_{\alpha(1)}^{(i)}) \in B(\mathbf{y}, r)^{\alpha+e_i}$,

$$|\bar{\Phi}_{\alpha+e_i} f(\bar{\mathbf{y}}_{\alpha+e_i}) - \bar{\Phi}_{\alpha+e_i} f(z, \mathbf{x}_{\alpha(1)}^{(i)})| \leq r^0 \mathbf{M}. \quad (8.8)$$

Now suppose $k = 0$, then by Proposition 11.1 we have,

$$|f(\mathbf{x}) - P_{f,\mathbf{y},l}(\mathbf{x})| \leq \max_{\substack{|\beta|=l+1, \\ \beta=(0,\dots,0,i_j,\dots,i_d), \\ i_j>0 \\ j=1,\dots,d}} |\bar{\Phi}_{\beta} f(x_1, \dots, x_j, y_j^{i_j}, y_{j+1}^{i_{j+1}+1}, \dots, y_d^{i_d+1}) L_{\beta,\mathbf{y}}(\mathbf{x})|$$

Note that for $\beta = (0, \dots, 0, i_j, \dots, i_d)$, $i_j > 0$ above,

$$L_{\beta,\mathbf{y}}(\mathbf{x}) = \prod_{s=j}^d (x_s - y_s)^{i_s} = (x_j - y_j) L_{\beta-e_j,\mathbf{y}}(\mathbf{x}).$$

Since $\mathbf{x} \in B(\mathbf{y}, r)$, and $|\beta - e_j| = l$, the right hand side of the above inequality can be bounded above by the following,

$$\leq r^l \max_{\substack{|\beta|=l+1, \\ \beta=(0,\dots,0,i_j,\dots,i_d), \\ i_j>0 \\ j=1,\dots,d}} |\bar{\Phi}_{\beta} f(x_1, \dots, x_j, y_j^{i_j}, y_{j+1}^{i_{j+1}+1}, \dots, y_d^{i_d+1}) (x_j - y_j)|.$$

Since for any multi-index β ,

$$\begin{aligned} & \bar{\Phi}_{\beta} f(x_1, \dots, x_j, y_j^{i_j}, y_{j+1}^{i_{j+1}+1}, \dots, y_d^{i_d+1}) (x_j - y_j) \\ &= \bar{\Phi}_{\beta-e_j} f(x_1, \dots, x_j, y_j^{i_j-1}, y_{j+1}^{i_{j+1}+1}, \dots, y_d^{i_d+1}) - \bar{\Phi}_{\beta-e_j} f(x_1, \dots, x_{j-1}, y_j^{i_j}, y_{j+1}^{i_{j+1}+1}, \dots, y_d^{i_d+1}), \end{aligned}$$

we conclude that for any $\mathbf{x} \in B(\mathbf{y}, r)$, we have that

$$|f(\mathbf{x}) - P_{f,\mathbf{y},l}(\mathbf{x})| \leq r^l \mathbf{M}. \quad (8.9)$$

Hence Equation (8.7), Equation (8.8) and Equation (8.9) yield the theorem. \square

For $\mathbf{f} = (f_1, \dots, f_n)$ we define

$$\mathcal{S}_{\mathbf{f}} = \{c_0 + c_1 f_1 + \dots + c_n f_n \mid \max_{i=0}^n |c_i| = 1\},$$

where $f_i : B \subset B(0, 1)^d \rightarrow \mathbb{Q}_{\nu}$ for all $i = 1, \dots, n$.

Lemma 8.2. *Suppose that $\mathbf{f} = (f_1, f_2, \dots, f_n) : B \subset B(0, 1)^d \rightarrow \mathbb{Q}_{\nu}^n$ is a C^{l+1} map, and $\mathbf{x}_0 \in B$. Then for any $\varepsilon > 0$ there exists a neighbourhood $V \subset B$ of \mathbf{x}_0 such that for any ball $B(\mathbf{y}, r) \subset V$, for any multi-index β , $0 \leq |\beta| \leq l$, and for any $g \in \mathcal{S}_{\mathbf{f}}$, we have*

$$\sup_{\mathbf{x}_{\beta(1)} \in B(\mathbf{y}, r)^{\beta(1)}} |\bar{\Phi}_{\beta} g(\mathbf{x}_{\beta(1)}) - \bar{\Phi}_{\beta} P_{g,\mathbf{y},l}(\mathbf{x}_{\beta(1)})| < \varepsilon r^{l-|\beta|}. \quad (8.10)$$

Proof. Using Theorem 8.1 it is enough to show that for any $\varepsilon > 0$, there exists a neighbourhood V such that for any $B(\mathbf{y}, r) \subset V$, we have

$$\sup_{\substack{\mathbf{w}_1, \mathbf{w}_2 \in B(\mathbf{y}, r)^{\beta(1)}, \\ |\beta|=l}} |\bar{\Phi}_{\beta} g(\mathbf{w}_1) - \bar{\Phi}_{\beta} g(\mathbf{w}_2)| < \varepsilon$$

for any $g \in \mathcal{S}_{\mathbf{f}}$. Since the family $\{\bar{\Phi}_{\beta} g \mid |\beta| = l, g \in \mathcal{S}_{\mathbf{f}}\}$ is equicontinuous, for any $\varepsilon > 0$, we can always guarantee a neighbourhood $V(\varepsilon)$ such that $|\bar{\Phi}_{\beta} g(\mathbf{w}_1) - \bar{\Phi}_{\beta} g(\mathbf{w}_2)| < \varepsilon$ for all β with $|\beta| = l$, $B(\mathbf{y}, r) \subset V(\varepsilon)$, $\mathbf{w}_1, \mathbf{w}_2 \in B(\mathbf{y}, r)^{\beta(1)}$ and $g \in \mathcal{S}_{\mathbf{f}}$. \square

9. NONDEGENERACY AND NORMALIZATION

Lemma 9.1. *Suppose that $\mathbf{f} = (f_1, f_2, \dots, f_n) : B \subset B(0, 1)^d \rightarrow \mathbb{Q}_{\nu}^n$ is a C^{l+1} map, which is l -nondegenerate at \mathbf{x}_0 . Then there is an open set $\mathbf{x}_0 \in V_0 \subset B$ and $\eta > 0$ such that for any $B(\mathbf{y}, r) \subset V_0$, and $g \in \mathcal{S}_{\mathbf{f}}$,*

$$\|g\|_{B(\mathbf{y}, r)} \geq \eta r^l \quad (9.1)$$

and

$$\frac{\|P_{g, \mathbf{y}, l}\|_{B(\mathbf{y}, r)}}{\|g\|_{B(\mathbf{y}, r)}} = 1. \quad (9.2)$$

Proof. Let V_0 be an open ball such that $\mathbf{f} = (f_1, \dots, f_n)$ is nondegenerate for any $\mathbf{y} \in V_0$. Let us rescale $P_{g, \mathbf{y}, l}$ as

$$Q_{g, \mathbf{y}}(\mathbf{x}) := P_{g, \mathbf{y}, l}(\mathbf{x} + \mathbf{y}), \mathbf{x} \in \mathbb{Q}_{\nu}^d,$$

where $g \in \mathcal{S}_{\mathbf{f}}$. Let us equip $\mathcal{P}_{d, l}$ with the norm $\|\cdot\|_{d, l}$ that is the maximum of all the ν -norms of all coefficients. By nondegeneracy of f and compactness of $\mathcal{S}_{\mathbf{f}}$ and V_0 , we have that the set $\{Q_{g, \mathbf{y}} : g \in \mathcal{S}_{\mathbf{f}}, \mathbf{y} \in V_0\}$ is bounded away from 0, which means for all $g \in \mathcal{S}_{\mathbf{f}}, \mathbf{y} \in V_0$, $\|Q_{g, \mathbf{y}}\|_{d, l} \geq \eta_1$ for some $\eta_1 > 0$. Moreover, we can rescale again, for each $\mathbf{y} \in V_0$ and $r = \nu^{-R} > 0$ such that $B(\mathbf{y}, r) \subset V_0$, we define

$$Q_{g, \mathbf{y}}^R(\mathbf{x}) := \nu^{-lR} Q_{g, \mathbf{y}}(\nu^R \mathbf{x}).$$

Since $\{Q_{g, \mathbf{y}} : g \in \mathcal{S}_{\mathbf{f}}, \mathbf{y} \in V_0\}$ is bounded away from 0, we will have $\{Q_{g, \mathbf{y}}^R : g \in \mathcal{S}_{\mathbf{f}}, B(\mathbf{y}, \nu^{-R}) \subset V_0\}$ is also bounded away from 0. The reason is $\|Q_{g, \mathbf{y}}^R\|_{d, l} \geq \|Q_{g, \mathbf{y}}\|_{d, l}$, since norm of β -th coefficient in $Q_{g, \mathbf{y}}^R$ is greater than norm of β -th coefficient in $Q_{g, \mathbf{y}}$ for every multi-index β , $|\beta| \leq l$. By comparing with the norm $P \rightarrow \|P\|_{B(0, 1)}$, the above implies that for all $g \in \mathcal{S}_{\mathbf{f}}, \|Q_{g, \mathbf{y}}^R\|_{B(0, 1)^d} \geq \eta$ for some $\eta > 0$. This gives us for all $B(\mathbf{y}, r) \subset V_0$ and $g \in \mathcal{S}_{\mathbf{f}}$.

$$\|P_{g, \mathbf{y}, l}\|_{B(\mathbf{y}, r)} = \nu^{-lR} \|Q_{g, \mathbf{y}}^R\|_{B(0, 1)^d} \geq r^l \eta.$$

Now using Lemma 8.2, for $|\beta| = 0$, for sufficiently small V_0 ,

$$\sup_{\mathbf{x} \in B(\mathbf{y}, r)} |g(\mathbf{x}) - P_{g, \mathbf{y}, l}(\mathbf{x})| \leq \frac{1}{\nu} \nu^{-lR} \eta = \frac{1}{\nu} r^l \eta.$$

Hence, we have

$$\|g\|_{B(\mathbf{y}, r)} \geq r^l \eta. \quad (9.3)$$

Note,

$$\|P_{g, \mathbf{y}, l}\|_{B(\mathbf{y}, r)} \leq \max\left(\frac{1}{\nu} r^l \eta, \|g\|_{B(\mathbf{y}, r)}\right).$$

Hence by Equation (9.3) we have,

$$\|P_{g, \mathbf{y}, l}\|_{B(\mathbf{y}, r)} \leq \|g\|_{B(\mathbf{y}, r)}.$$

Similarly we get,

$$\|g\|_{B(\mathbf{y},r)} \leq \|P_{g,\mathbf{y},l}\|_{B(\mathbf{y},r)}.$$

Therefore,

$$\frac{\|P_{g,\mathbf{y},l}\|_{B(\mathbf{y},r)}}{\|g\|_{B(\mathbf{y},r)}} = 1.$$

□

Similar to the way we have normalized polynomials to make them functions in the unit ball $B(0,1)^d$, we will normalize functions in $\mathcal{S}_{\mathbf{f}}$. So for V_0 a sufficiently small neighbourhood of \mathbf{x}_0 , $B(\mathbf{y},r) \subset V_0$ and $g \in \mathcal{S}_{\mathbf{f}}$, and $r > 0$ is a power of ν , we define the function

$$g_{r,\mathbf{y},g}(\mathbf{x}) := \|g\|_{B(\mathbf{y},r)} g(\mathbf{y} + r^{-1}\mathbf{x}),$$

which is defined on $B(0,1)^d$. We note that the above normalization and shift in the definition of $g_{r,\mathbf{y},g}$ are taken such that using ν -adic norm, $\|g_{r,\mathbf{y},g}\|_{B(0,1)^d} = 1$. We also consider the collection

$$\mathcal{G}(\mathbf{f}, V) := \{g_{r,\mathbf{y},g} \mid g \in \mathcal{S}_{\mathbf{f}}, B(\mathbf{y}, r) \subset V\}.$$

Lemma 9.2. *For any $\varepsilon > 0$ there exists a neighbourhood $V \subset B$ of \mathbf{x}_0 such that for any $\phi \in \mathcal{G}(\mathbf{f}, V)$ one has*

$$\max_{|\beta| \leq l+1} \sup_{\mathbf{x}_{\beta(1)} \in B(\mathbf{0},1)^{\beta(1)}} |(\bar{\Phi}_{\beta}\phi - \bar{\Phi}_{\beta}P_{\phi,\mathbf{0},l})(\mathbf{x}_{\beta(1)})| < \varepsilon. \quad (9.4)$$

Proof. Let us choose a small V that we get from Lemma 8.2 and Lemma 9.1. Let us denote $y_j + r^{-1}\mathbf{x}_j := y_j^{i_j} + r^{-1}(x_{j,1}, \dots, x_{j,i_j+1})$, and $\mathbf{y} + r^{-1}\mathbf{x}_{\beta(1)} = (y_1 + r^{-1}\mathbf{x}_1, \dots, y_d + r^{-1}\mathbf{x}_d)$. Note that if $\phi = g_{r,\mathbf{y},g} \in \mathcal{G}(\mathbf{f}, V)$, then

$$\bar{\Phi}_{\beta}\phi(\mathbf{x}_{\beta(1)}) = \frac{\|g\|_{B(\mathbf{y},r)}}{r^{|\beta|}} \bar{\Phi}_{\beta}g(\mathbf{y} + r^{-1}\mathbf{x}_{\beta(1)})$$

and

$$\bar{\Phi}_{\beta}P_{\phi,\mathbf{0},l}(\mathbf{x}_{\beta(1)}) = \frac{\|g\|_{B(\mathbf{y},r)}}{r^{|\beta|}} \bar{\Phi}_{\beta}P_{g,\mathbf{y},l}(\mathbf{y} + r^{-1}\mathbf{x}_{\beta(1)}).$$

Hence, we have

$$\begin{aligned} & |(\bar{\Phi}_{\beta}\phi - \bar{\Phi}_{\beta}P_{\phi,\mathbf{0},l})(\mathbf{x}_{\beta(1)})| \\ &= \frac{r^{|\beta|}}{\|g\|_{B(\mathbf{y},r)}} |\bar{\Phi}_{\beta}g(\mathbf{y} + r^{-1}\mathbf{x}_{\beta(1)}) - \bar{\Phi}_{\beta}P_{g,\mathbf{y},l}(\mathbf{y} + r^{-1}\mathbf{x}_{\beta(1)})|. \end{aligned}$$

Using Lemma 8.2 and Lemma 9.1 for $|\beta| \leq l$ we get,

$$|(\bar{\Phi}_{\beta}\phi - \bar{\Phi}_{\beta}P_{\phi,\mathbf{0},l})(\mathbf{x}_{\beta(1)})| \leq \frac{r^{|\beta|}}{\eta r^l} \varepsilon \eta r^{l-|\beta|} = \varepsilon.$$

For $|\beta| = l+1$ by compactness, we have a $K > 0$ such that $\|\bar{\Phi}_{\beta}g\|_V \leq K$ for all $g \in \mathcal{S}_{\mathbf{f}}$,

$$\begin{aligned} \|\bar{\Phi}_{\beta}\phi - \bar{\Phi}_{\beta}P_{\phi,\mathbf{0},l}\|_{B(0,1)^{l+1+d}} &= \|\bar{\Phi}_{\beta}\phi\|_{B(0,1)^{l+1+d}} \\ &= \frac{r^{l+1}}{\|g\|_{B(\mathbf{y},r)}} \sup_{\mathbf{x}_{\beta(1)} \in B(0,1)^{l+1+d}} |\bar{\Phi}_{\beta}g(\mathbf{y} + r^{-1}\mathbf{x}_{\beta(1)})| \leq \frac{r}{\eta} K < \varepsilon, \end{aligned}$$

if V is small enough.

□

10. COMPLETING THE PROOF OF PROPOSITION 4.1

Theorem 10.1. *Let $\mathbf{B} \subset B(0, 1)^d \subset \mathbb{Q}_\nu$ be an open ball and let $\mathbf{f} : \mathbf{B} \rightarrow \mathbb{Q}_\nu^n$ be a C^{l+1} map which is l nondegenerate at $\mathbf{x}_0 \in \mathbf{B}$. Let μ be a measure which is D -Federer and absolutely (C, α) -decaying on \mathbf{B} for some $D, C, \alpha > 0$. Then there exists a neighborhood $V \subset \mathbf{B}$ of \mathbf{x}_0 and a positive \tilde{C} such that for any $g \in \mathcal{S}_{\mathbf{f}}$ is absolutely (\tilde{C}, α') -good on V with respect to μ .*

Proof. By Theorem 6.1, it suffices to find $C_{\alpha_1, \alpha'_1}, s, S$ for multi-index α_1, α'_1 and $V \subset \mathbf{B}$ such that for any $\phi \in \mathcal{G}(\mathbf{f}, V)$ there is a multi-index β with $|\beta| = k \in \mathbb{N}$ such that, for every $\alpha_1 + \alpha_2 = \beta = \alpha'_1 + \alpha'_2$, $|\alpha_1| > |\alpha'_1|$,

$$\inf_{B(0,1)^{|\alpha_1|+d}} |\bar{\Phi}_{\alpha_1} \partial_{\alpha_2} \phi| \leq C_{\alpha_1, \alpha'_1} \inf_{B(0,1)^{|\alpha'_1|+d}} |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} \phi|$$

and

$$\inf_{B(0,1)^{|\beta|+d}} |\bar{\Phi}_{\beta} \phi| > s,$$

and for each $|\eta| \leq k+1$,

$$\|\bar{\Phi}_{\eta} \phi\|_{B(0,1)^{|\eta|+d}} < S.$$

Using Lemma 9.1, we can choose V_0 around \mathbf{x}_0 such that for $B(\mathbf{y}, r) \subset V_0$, $g \in \mathcal{S}_{\mathbf{f}}$,

$$\|P_{g, \mathbf{y}, l}\|_{B(\mathbf{y}, r)} = \|g\|_{B(\mathbf{y}, r)}.$$

This implies

$$\|P_{\phi, \mathbf{0}, l}\|_{B(\mathbf{0}, 1)} = \frac{1}{\|g\|_{B(\mathbf{y}, r)}} \sup_{\mathbf{x} \in B(\mathbf{0}, 1)} |P_{g, \mathbf{y}, l}(\mathbf{y} + r^{-1} \mathbf{x})| = 1.$$

Now using Theorem 7.1 we can find $|\beta| = k \leq l$ and $s, S, C_{\alpha_1, \alpha'_1} > 0$ such that, for every $\alpha_1 + \alpha_2 = \beta = \alpha'_1 + \alpha'_2$, $|\alpha_1| > |\alpha'_1|$,

$$\inf_{B(0,1)^{|\alpha_1|+d}} |\bar{\Phi}_{\alpha_1} \partial_{\alpha_2} P_{\phi, \mathbf{0}, l}| \leq C_{\alpha_1, \alpha'_1} \inf_{B(0,1)^{|\alpha'_1|+d}} |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} P_{\phi, \mathbf{0}, l}|, \quad (10.1)$$

and

$$\inf_{B(0,1)^{k+d}} |\bar{\Phi}_{\beta} P_{\phi, \mathbf{0}, l}| > s \quad (10.2)$$

and

$$\|\bar{\Phi}_{\eta} P_{\phi, \mathbf{0}, l}\|_{B(0,1)^{|\eta|+d}} < S \quad \forall |\eta| \geq 0. \quad (10.3)$$

Note that from Equation (10.1) for $\beta = \alpha'_1 + \alpha'_2$, $|\beta| > |\alpha'_1|$ we have,

$$\inf_{B(0,1)^{k+d}} |\bar{\Phi}_{\beta} P_{\phi, \mathbf{0}, l}| \leq C_{\beta, \alpha'_1} \inf_{B(0,1)^{|\alpha'_1|+d}} |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} P_{\phi, \mathbf{0}, l}|.$$

Let us take $a \in \mathbb{N}$ such that $\frac{1}{\nu^a} < \max_{\alpha_1 + \alpha_2 = \beta = \alpha'_1 + \alpha'_2} C_{\alpha_1, \alpha'_1}^{-1}$. Hence by Equation (10.2) we have that

$$\inf_{B(0,1)^{|\alpha'_1|+d}} |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} P_{\phi, \mathbf{0}, l}| > s C_{\beta, \alpha'_1}^{-1} > \frac{s}{\nu^a}. \quad (10.4)$$

Let us take $0 < \varepsilon < \frac{s}{\nu^a}$. If $\alpha'_1 + \alpha'_2 = \beta$, then for any $\mathbf{z}' \in B(0, 1)^{|\alpha'_1|+d}$, by Lemma 11.1,

$$\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} P_{\phi, \mathbf{0}, l}(\mathbf{z}') = \sum_{\mathbf{t}_{\beta}} \bar{\Phi}_{\beta} P_{\phi, \mathbf{0}, l}(\mathbf{t}_{\beta}), \text{ and } \bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} \phi(\mathbf{z}') = \sum_{\mathbf{t}_{\beta}} \bar{\Phi}_{\beta} \phi(\mathbf{t}_{\beta}),$$

where \mathbf{t}_β has coordinates in terms of \mathbf{z}' . It should be clear that applying Lemma 11.1 multiple times will give \mathbf{t}_β 's, whose coordinates will be a variation of coordinates of \mathbf{z}' . Therefore by Lemma 9.2 and the equality above, and property of ultrametric norm, we have

$$\|\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} \phi - \bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} P_{\phi, \mathbf{0}, l}\|_{B(0,1)^{|\alpha'_1|+d}} < \varepsilon. \quad (10.5)$$

Since $\varepsilon < \frac{s}{\nu^a}$, by Equation (10.4) and Equation (10.5) we have that for any $\mathbf{z}' \in B(0,1)^{|\alpha'_1|+d}$,

$$|\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} \phi(\mathbf{z}')| = |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} P_{\phi, \mathbf{0}, l}(\mathbf{z}')| \quad (10.6)$$

for any $\alpha'_1 + \alpha'_2 = \beta$. Hence by Equation (10.1) we have, for every $\alpha_1 + \alpha_2 = \beta = \alpha'_1 + \alpha'_2$, $|\alpha_1| > |\alpha'_1|$,

$$\inf_{B(0,1)^{|\alpha_1|+d}} |\bar{\Phi}_{\alpha_1} \partial_{\alpha_2} \phi| \leq C_{\alpha_1, \alpha'_1} \inf_{B(0,1)^{|\alpha'_1|+d}} |\bar{\Phi}_{\alpha'_1} \partial_{\alpha'_2} \phi|.$$

By Lemma 9.2 and Equation (10.2), we get that for any $\mathbf{x}_{\beta(1)} \in B(\mathbf{0}, 1)^{\beta(1)}$,

$$s < \max(|\bar{\Phi}_\beta \phi(\mathbf{x}_{\beta(1)})|, \varepsilon).$$

Since $\varepsilon < \frac{s}{\nu^a}$ we get,

$$\inf_{B(0,1)^{k+d}} |\bar{\Phi}_\beta \phi| > s.$$

Now for upper bound, we apply Lemma 9.2 and use Equation (10.3) to get

$$\|\bar{\Phi}_\eta \phi\|_{B(0,1)^{k+d}} \leq \max(\varepsilon, S) = S$$

for every $|\eta| \leq l$, since $\varepsilon < \frac{s}{\nu^a} < S$. □

11. APPENDIX

In this appendix, we will prove the following proposition which can be thought of as the analogue of Taylor's theorem for real variables. One of the main issues here is the careful computation of the 'error' term as we need it for Theorem 8.1. In the one dimensional case, the proof is much easier, and is well known [17, Theorem 29.4]. We could not find a reference in general and so we provide a proof. Let us recall that for a C^k function $f : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$, we denoted by $P_{f, \mathbf{y}, k}$ the Taylor polynomial of f at $\mathbf{y} \in \mathbb{Q}_\nu^d$; see (8.1).

Proposition 11.1. *For a C^{l+1} map $f : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$ and $\mathbf{y} \in \mathbb{Q}_\nu^d$, we have the following for $k \leq l$:*

$$\begin{aligned} f(\mathbf{x}) &= P_{f, \mathbf{y}, k}(\mathbf{x}) + \sum_{\substack{|\beta|=k+1 \\ \beta=(i_1, \dots, i_d) \\ i_1 > 0}} \bar{\Phi}_\beta f(x_1, y_1^{i_1}, y_2^{i_2+1}, \dots, y_d^{i_d+1}) L_{\beta, \mathbf{y}}(\mathbf{x}) \\ &+ \sum_{\substack{|\beta|=k+1 \\ \beta=(0, i_2, \dots, i_d) \\ i_2 > 0}} \bar{\Phi}_\beta f(x_1, x_2, y_2^{i_2}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) L_{\beta, \mathbf{y}}(\mathbf{x}) \\ &+ \dots + \bar{\Phi}_{(k+1)e_d} f(x_1, \dots, x_d, y_d^{k+1}) L_{(k+1)e_d, \mathbf{y}}(\mathbf{x}) \end{aligned} \quad (11.1)$$

Remark 11.1. *We remark that Proposition 11.1 differs from the Lagrange remainder formula and also from the Mean Value theorem. Let us explain the simplest meaningful case to point out the difference. Let $g : \mathbb{Q}_\nu \rightarrow \mathbb{Q}_\nu$ be a C^2 map. Proposition 11.1 shows*

$$g(x) = g(y) + \underbrace{\bar{\Phi}^1 g(y, y)}_{g'(y)} (x - y) + \bar{\Phi}^2 g(x, y, y) (x - y)^2. \quad (11.2)$$

Note that the remainder term contains the function $\bar{\Phi}^2 g$ evaluated at (x, y, y) and the function $\bar{\Phi}^2 g$ has a larger domain than that of g'' , and $\frac{1}{2}g''(z) = \bar{\Phi}^2 g(z, z, z)$ for all $z \in \mathbb{Q}_\nu$.

On the other hand for a function $\psi : \mathbb{R} \rightarrow \mathbb{R}$ a C^2 map, by Lagrange form, we have

$$\psi(x) = \psi(y) + \psi'(y)(x - y) + \frac{1}{2}\psi''(c)(x - y)^2,$$

where c is a point between x and y in \mathbb{R} . Proposition 11.1 is the generalization of (11.2) in higher dimensions.

Proof. We proceed by induction on k and accordingly, set $k = 0$. Then,

$$\begin{aligned} f(\mathbf{x}) - f(\mathbf{y}) &= \sum_{i=1}^d f(x_1, \dots, x_{i-1}, y_i, y_{i+1}, \dots, y_d) - f(x_1, \dots, x_{i-1}, x_i, y_{i+1}, \dots, y_d) \\ &= \sum_{i=1}^d \bar{\Phi}_{e_i} f(x_1, \dots, x_{i-1}, x_i, y_i, y_{i+1}, \dots, y_d) L_{e_i, \mathbf{y}}(\mathbf{x}). \end{aligned}$$

Since $P_{f, \mathbf{y}, 0}(\mathbf{x}) = f(\mathbf{y})$, we have the statement to be true for $k = 0$. By induction hypothesis let us assume that the statement is true for $k = m$:

$$f(\mathbf{x}) = P_{f, \mathbf{y}, m}(\mathbf{x}) + I_1 + \dots + I_d, \quad (11.3)$$

where

$$I_j := \sum_{\substack{|\beta|=m+1 \\ \beta=(0, \dots, 0, i_j, \dots, i_d) \\ i_j > 0}} \bar{\Phi}_\beta f(x_1, \dots, x_j, y_j^{i_j}, y_{j+1}^{i_{j+1}+1}, \dots, y_d^{i_d+1}) L_{\beta, \mathbf{y}}(\mathbf{x}).$$

We want to show that the statement is true for $k = m + 1 \leq l$.

We know by definition, that the coefficient of a term in the summand I_1 is,

$$\begin{aligned} \bar{\Phi}_\beta f(x_1, y_1^{i_1}, y_2^{i_2+1}, \dots, y_d^{i_d+1}) &= \bar{\Phi}_\beta f(y_1^{i_1+1}, y_2^{i_2+1}, \dots, y_d^{i_d+1}) \\ &\quad + \bar{\Phi}_{\beta+e_1} f(x_1, y_1^{i_1+1}, y_2^{i_2+1}, \dots, y_d^{i_d+1}) L_{e_1, \mathbf{y}}(\mathbf{x}). \end{aligned}$$

We also have, that the coefficient of a term in the summand I_2 is,

$$\begin{aligned} &\bar{\Phi}_\beta f(x_1, x_2, y_2^{i_2}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) \\ &= \bar{\Phi}_\beta f(x_1, y_2^{i_2+1}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) + \bar{\Phi}_{\beta+e_2} f(x_1, x_2, y_2^{i_2+1}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) L_{e_2, \mathbf{y}}(\mathbf{x}) \\ &= \bar{\Phi}_\beta f(y_1, y_2^{i_2+1}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) + \bar{\Phi}_{\beta+e_2} f(x_1, x_2, y_2^{i_2+1}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) L_{e_2, \mathbf{y}}(\mathbf{x}) \\ &\quad + \bar{\Phi}_{\beta+e_1} f(x_1, y_1, y_2^{i_2+1}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) L_{e_1, \mathbf{y}}(\mathbf{x}). \end{aligned}$$

Continuing like this we get that the coefficient of a term in I_d is,

$$\begin{aligned} &\bar{\Phi}_{(m+1)e_d} f(x_1, \dots, x_d, y_d^{m+1}) \\ &= \bar{\Phi}_{(m+1)e_d} f(x_1, \dots, x_{d-1}, y_d^{m+2}) + \bar{\Phi}_{(m+2)e_d} f(x_1, \dots, x_d, y_d^{m+2}) L_{e_d, \mathbf{y}}(\mathbf{x}) \\ &= \bar{\Phi}_{(m+1)e_d} f(x_1, \dots, y_{d-1}, y_d^{m+2}) + \bar{\Phi}_{(m+1)e_d+e_{d-1}} f(x_1, \dots, x_{d-1}, y_{d-1}, y_d^{m+2}) L_{e_{d-1}, \mathbf{y}}(\mathbf{x}) \\ &\quad + \bar{\Phi}_{(m+2)e_d} f(x_1, \dots, x_d, y_d^{m+2}) L_{e_d, \mathbf{y}}(\mathbf{x}) \\ &= \bar{\Phi}_{(m+1)e_d} f(y_1, \dots, y_{d-1}, y_d^{m+2}) + \sum_{j=1}^{d-1} \bar{\Phi}_{(m+1)e_d+e_j} f(x_1, \dots, x_j, y_j, \dots, y_d^{m+2}) L_{e_j, \mathbf{y}}(\mathbf{x}) \\ &\quad + \bar{\Phi}_{(m+2)e_d} f(x_1, \dots, x_d, y_d^{m+2}) L_{e_d, \mathbf{y}}(\mathbf{x}). \end{aligned}$$

One can observe that in all of the above, we can rewrite each coefficient in I_j as a sum of terms, where the first term is $\bar{\Phi}_\beta f(y_1, \dots, y_{j-1}, y_j^{i_j+1}, \dots, y_d^{i_d+1}) L_{\beta, \mathbf{y}}(\mathbf{x})$, and $\beta = (0, \dots, 0, i_j, \dots, i_d)$, $i_j > 0$, $|\beta| = m + 1$. Therefore, we can rewrite Equation (11.3) and use the previous equations as follows,

$$\begin{aligned}
f(\mathbf{x}) &= P_{f, \mathbf{y}, m}(\mathbf{x}) + \sum_{|\beta|=m+1} \bar{\Phi}_\beta f(\bar{\mathbf{y}}_\beta) L_{\beta, \mathbf{y}}(\mathbf{x}) + \\
&\quad \sum_{\substack{|\beta|=m+2 \\ \beta=(i_1, \dots, i_d) \\ i_1 > 0}} \bar{\Phi}_\beta f(x_1, y_1^{i_1}, y_2^{i_2+1}, \dots, y_d^{i_d+1}) L_{\beta, \mathbf{y}}(\mathbf{x}) \\
&\quad + \sum_{\substack{|\beta|=m+2 \\ \beta=(0, i_2, \dots, i_d) \\ i_2 > 0}} \bar{\Phi}_\beta f(x_1, x_2, y_2^{i_2}, y_3^{i_3+1}, \dots, y_d^{i_d+1}) L_{\beta, \mathbf{y}}(\mathbf{x}) \\
&\quad + \dots + \bar{\Phi}_{(m+2)e_d} f(x_1, \dots, x_d, y_d^{m+2}) L_{(m+2)e_d, \mathbf{y}}(\mathbf{x}).
\end{aligned} \tag{11.4}$$

Since $P_{f, \mathbf{y}, m+1}(\mathbf{x}) = P_{f, \mathbf{y}, m}(\mathbf{x}) + \sum_{|\beta|=m+1} \bar{\Phi}_\beta f(\bar{\mathbf{y}}_\beta) L_{\beta, \mathbf{y}}(\mathbf{x})$, we have the statement to be true for $k = m + 1$. This completes the proof. \square

We also give proof of the following lemmata whose statements is used in the paper several times.

Lemma 11.1. *Let $g : \mathbb{Q}_\nu \rightarrow \mathbb{Q}_\nu$ be a C^{k+1} map. Then for any $y_1, \dots, y_{k+1} \in \mathbb{Q}_\nu$,*

$$\bar{\Phi}^k g'(y_1, \dots, y_{k+1}) = \bar{\Phi}^{k+1} g(y_1, y_1, y_2, \dots, y_{k+1}) + \dots + \bar{\Phi}^{k+1} g(y_1, y_2, \dots, y_k, y_{k+1}, y_{k+1}).$$

Proof. We will prove this by induction on k . If $k = 0$, then $\bar{\Phi}^0 g'(y) = g'(y) = \bar{\Phi}^1 g(y, y)$. Suppose by the induction hypothesis, we know the conclusion for any k , $1 \leq k \leq i$. Now we wish to show that for $k = i + 1$ the conclusion of this lemma holds.

Note that for any $(y_1, \dots, y_{k+1}) \in \nabla^{k+1} \mathbb{Q}_\nu$

$$\bar{\Phi}^{i+1} g'(y_1, \dots, y_{i+2}) = \frac{\bar{\Phi}^i g'(y_1, y_3, \dots, y_{i+2}) - \bar{\Phi}^i g'(y_2, \dots, y_{i+2})}{(y_1 - y_2)}.$$

Using the induction hypothesis,

$$\begin{aligned}
&\bar{\Phi}^i g'(y_1, y_3, \dots, y_{i+2}) \\
&= \bar{\Phi}^{i+1} g(y_1, y_1, y_3, \dots, y_{i+2}) + \bar{\Phi}^{i+1} g(y_1, y_3, y_3, y_4, \dots, y_{i+2}) \dots + \bar{\Phi}^{i+1} g(y_1, y_3, \dots, y_{i+2}, y_{i+2})
\end{aligned}$$

and

$$\begin{aligned}
&\bar{\Phi}^i g'(y_2, \dots, y_{i+2}) \\
&= \bar{\Phi}^{i+1} g(y_2, y_2, \dots, y_{i+2}) + \bar{\Phi}^{i+1} g(y_2, y_3, y_3, y_4, \dots, y_{i+2}) + \dots + \bar{\Phi}^{i+1} g(y_2, \dots, y_{i+2}, y_{i+2}).
\end{aligned}$$

Now substituting the above two, we get

$$\begin{aligned}
& \bar{\Phi}^{i+1} g'(y_1, \dots, y_{i+2}) \\
&= \frac{\bar{\Phi}^{i+1} g(y_1, y_1, y_3, \dots, y_{i+2}) - \bar{\Phi}^{i+1} g(y_2, y_2, y_3, \dots, y_{i+2})}{y_1 - y_2} + \\
&\quad \bar{\Phi}^{i+2} g(y_1, y_2, y_3, y_3, \dots, y_{i+2}) + \dots + \bar{\Phi}^{i+2} g(y_1, y_2, \dots, y_{i+2}, y_{i+2}) \\
&= \frac{\bar{\Phi}^{i+1} g(y_1, y_1, y_3, \dots, y_{i+2}) - \bar{\Phi}^{i+1} g(y_1, y_2, \dots, y_{i+2})}{y_1 - y_2} + \\
&\quad \frac{\bar{\Phi}^{i+1} g(y_1, y_2, \dots, y_{i+2}) - \bar{\Phi}^{i+1} g(y_2, y_2, y_3, \dots, y_{i+2})}{y_1 - y_2} + \\
&\quad \bar{\Phi}^{i+2} g(y_1, y_2, y_3, y_3, \dots, y_{i+2}) + \dots + \bar{\Phi}^{i+2} g(y_1, y_2, \dots, y_{i+2}, y_{i+2}) \\
&= \bar{\Phi}^{i+2} g(y_1, y_1, y_2, y_3, \dots, y_{i+2}) + \bar{\Phi}^{i+2} g(y_1, y_2, y_2, y_3, \dots, y_{i+2}) + \\
&\quad \bar{\Phi}^{i+2} g(y_1, y_2, y_3, y_3, \dots, y_{i+2}) + \dots + \bar{\Phi}^{i+2} g(y_1, y_2, \dots, y_{i+2}, y_{i+2}).
\end{aligned}$$

Thus the proof is completed using the above and the definition of $\bar{\Phi}^{k+1} g'$. \square

Lemma 11.2. For $\beta = (i_1, \dots, i_d)$ any multi index with $|\beta| = l$ and any $(\mathbf{x}_1, \dots, \mathbf{x}_d) \in B(\mathbf{0}, 1)^{\beta(1)}$, for any $f : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$, that is C^{l+1} , $1 \leq j \leq d$, we get

$$\begin{aligned}
& i_j \bar{\Phi}_\beta f(\mathbf{x}_1, \dots, \mathbf{x}_d) \\
&= \bar{\Phi}_{\beta-e_j} \partial_{e_j} f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\
&+ \sum_{i=1}^{i_j} \bar{\Phi}_{\beta+e_j} f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i-1}, x_{j,i}, x_{j,i}, x_{j,i+1}, \dots, x_{j,i_j+1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) (x_{j,i_j+1} - x_{j,i}).
\end{aligned} \tag{11.5}$$

Proof. First note that by Lemma 11.1,

$$\begin{aligned}
& \bar{\Phi}_{\beta-e_j} \partial_{e_j} f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\
&= \bar{\Phi}_\beta f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, x_{j,1}, x_{j,2}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) + \\
&\quad \dots + \bar{\Phi}_\beta f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, x_{j,2}, \dots, x_{j,i_j-1}, x_{j,i_j}, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d).
\end{aligned}$$

Also, note there are i_j many summands in the above sum.

Then by definition, for any $1 \leq i \leq i_j$,

$$\begin{aligned}
& \bar{\Phi}_\beta f(\mathbf{x}_1, \dots, \mathbf{x}_d) - \bar{\Phi}_\beta f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i-1}, x_{j,i}, x_{j,i}, x_{j,i+1}, \dots, x_{j,i_j}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) \\
&= \bar{\Phi}_{\beta+e_j} f(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, x_{j,1}, \dots, x_{j,i-1}, x_{j,i}, x_{j,i}, x_{j,i+1}, \dots, x_{j,i_j}, x_{j,i_j+1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d) (x_{j,i_j+1} - x_{j,i}).
\end{aligned}$$

Combining these two observations, the lemma follows. \square

The following is a version of [17, Lemma 29.2, (v)]. We provide a proof to make this article self contained.

Lemma 11.3. Let $g : \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu$ be C^{l+1} map. For any α multi-index with $|\alpha| = l$, we have the following, for any $z \in \mathbb{Q}_\nu$, $\mathbf{x}_{\alpha(1)} \in \mathbb{Q}_\nu^{|\alpha|+d}$, $1 \leq i \leq d$,

$$\begin{aligned}
& \bar{\Phi}_{\alpha+e_i} (L_{e_i, \mathbf{y}} g)(z, \mathbf{x}_{\alpha(1)}^{(i)}) \\
&= L_{e_i, \mathbf{y}}(z) \bar{\Phi}_{\alpha+e_i} g(z, \mathbf{x}_{\alpha(1)}^{(i)}) + \bar{\Phi}_\alpha g(\mathbf{x}_{\alpha(1)}).
\end{aligned} \tag{11.6}$$

Proof. The proof is done by induction on l . When $l = 0$, then $\alpha = 0$, Suppose $(z, \mathbf{x}) \in \mathbb{Q}_\nu^{d+1}$, $z \neq x_i$ and $1 \leq i \leq d$. Then

$$\begin{aligned} & \Phi_{e_i}(L_{e_i, \mathbf{y}}g)(x_1, \dots, x_{i-1}, z, x_i, x_{i+1}, \dots, x_d) \\ &= \frac{(z - y_i)g(x_1, \dots, x_{i-1}, z, x_{i+1}, \dots, x_d) - (x_i - y_i)g(\mathbf{x})}{z - x_i} \\ &= (z - y_i)\Phi_{e_i}g(z, \mathbf{x}^{(i)}) + g(\mathbf{x}). \end{aligned}$$

Then taking limits gives us Equation (11.6) for $l = 0$. Now suppose by the induction hypothesis, that Equation (11.6) is true for when $|\alpha| \leq l - 1$. Now let us take α such that $|\alpha| = l$, and fix i , where $1 \leq i \leq d$. There exists some $1 \leq j \leq d$ such that $\alpha = \beta + e_j$, where $|\beta| = l - 1$. Then for $z \in \mathbb{Q}_\nu$, $\mathbf{x}_{\alpha(1)} \in \mathbb{Q}_\nu^{|\alpha|+d}$, $x_{j,1} \neq x_{j,2}$,

$$\begin{aligned} & \Phi_{\alpha+e_i}(L_{e_i, \mathbf{y}}g)(z, \mathbf{x}_{\alpha(1)}^{(i)}) \\ &= \frac{\Phi_{\beta+e_i}(L_{e_i, \mathbf{y}}g)(z, \hat{\mathbf{x}}_{\alpha(1),j,2}^{(i)}) - \Phi_{\beta+e_i}(L_{e_i, \mathbf{y}}g)(z, \hat{\mathbf{x}}_{\alpha(1),j,1}^{(i)})}{x_{j,1} - x_{j,2}}, \end{aligned}$$

where $\hat{\mathbf{x}}_{\alpha(1),j,1} = (\mathbf{x}_1, \mathbf{x}_{j-1}, x_{j,2}, \dots, x_{j,i_j+1}, \mathbf{x}_{j+1}, \dots, \mathbf{x}_d)$, and similarly for $\hat{\mathbf{x}}_{\alpha(1),j,2}$ with $x_{j,2}$ missing from $\mathbf{x}_{\alpha(1)}$ among the j -th coordinates. Now using the induction hypothesis, we get

$$\Phi_{\beta+e_i}(L_{e_i, \mathbf{y}}g)(z, \hat{\mathbf{x}}_{\alpha(1),j,2}^{(i)}) = L_{e_i, \mathbf{y}}(z)\Phi_{\beta+e_i}g(z, \hat{\mathbf{x}}_{\alpha(1),j,2}^{(i)}) + \Phi_{\beta}g(\hat{\mathbf{x}}_{\alpha(1),j,2}).$$

and

$$\Phi_{\beta+e_i}(L_{e_i, \mathbf{y}}g)(z, \hat{\mathbf{x}}_{\alpha(1),j,1}^{(i)}) = L_{e_i, \mathbf{y}}(z)\Phi_{\beta+e_i}g(z, \hat{\mathbf{x}}_{\alpha(1),j,1}^{(i)}) + \Phi_{\beta}g(\hat{\mathbf{x}}_{\alpha(1),j,1}).$$

Using the above two, and $\alpha = \beta + e_j$, we get

$$\begin{aligned} \Phi_{\alpha+e_i}(L_{e_i, \mathbf{y}}g)(z, \mathbf{x}_{\alpha(1)}) &= \frac{(z - y_i)\Phi_{\beta+e_i}g(z, \hat{\mathbf{x}}_{\alpha(1),j,2}^{(i)}) - (z - y_i)\Phi_{\beta+e_i}g(z, \hat{\mathbf{x}}_{\alpha(1),j,1}^{(i)})}{x_{j,1} - x_{j,2}} + \Phi_{\beta+e_j}g(\mathbf{x}_{\alpha(1)}) \\ &= (z - y_i)\Phi_{\alpha+e_i}g(z, \mathbf{x}_{\alpha(1)}^{(i)}) + \Phi_{\alpha}g(\mathbf{x}_{\alpha(1)}). \end{aligned}$$

□

Lemma 11.4. Suppose $\mathbf{f} = (f_1, \dots, f_n) : U \subset \mathbb{Q}_\nu^d \rightarrow \mathbb{Q}_\nu^n$ is a nonsingular C^{l+1} map at \mathbf{x}_0 , we can find a neighbourhood V of \mathbf{x}_0 such that $\mathbf{f}|_V$ is bi-Lipschitz.

Proof. For simplicity of notation and to make it more readable, we give a proof for $d = 2$. The proof for the other cases is exactly the same. Note that, for $\mathbf{x} = (x_1, x_2)$, $\mathbf{y} = (y_1, y_2)$, for every $1 \leq i \leq n$,

$$\begin{aligned} f_i(\mathbf{x}) - f_i(\mathbf{y}) &= f_i(x_1, x_2) - f_i(x_1, y_2) + f_i(x_1, y_2) - f_i(y_1, y_2) \\ &= \bar{\Phi}_{e_2}f_i(x_1, x_2, y_2)(x_2 - y_2) + \bar{\Phi}_{e_1}f_i(x_1, y_1, y_2)(x_1 - y_1). \end{aligned}$$

This implies that

$$\mathbf{f}(\mathbf{x}) - \mathbf{f}(\mathbf{y}) = A_{\mathbf{x}, \mathbf{y}}(\mathbf{x} - \mathbf{y})^T, \quad (11.7)$$

where

$$A_{\mathbf{x}, \mathbf{y}} = \begin{bmatrix} \bar{\Phi}_{e_1}f_1(x_1, y_1, y_2) & \bar{\Phi}_{e_2}f_1(x_1, x_2, y_2) \\ \vdots & \vdots \\ \bar{\Phi}_{e_1}f_n(x_1, y_1, y_2) & \bar{\Phi}_{e_2}f_n(x_1, x_2, y_2) \end{bmatrix}.$$

Let us also denote by $A_{\mathbf{x},\mathbf{y}}(r, s)$ the 2×2 matrix with r, s -th rows from the above matrix $A_{\mathbf{x},\mathbf{y}}$. Note that $A_{\mathbf{x}_0,\mathbf{x}_0} = \nabla \mathbf{f}(\mathbf{x}_0)$, by (3.2). Since \mathbf{f} is nonsingular at \mathbf{x}_0 , we know $\nabla \mathbf{f}(\mathbf{x}_0)$ has rank 2. This implies that there exist $1 \leq i_1 < i_2 \leq n$ such that

$$A_{\mathbf{x}_0,\mathbf{x}_0}(i_1, i_2) = \begin{bmatrix} \partial_{e_1} f_{i_1}(\mathbf{x}_0) & \partial_{e_2} f_{i_1}(\mathbf{x}_0) \\ \partial_{e_1} f_{i_2}(\mathbf{x}_0) & \partial_{e_2} f_{i_2}(\mathbf{x}_0) \end{bmatrix}$$

is invertible. By continuity of the determinant function, there exists a neighborhood ball V (which is both open and closed) of \mathbf{x}_0 , such that for any $\mathbf{x}, \mathbf{y} \in V$, we have that $A_{\mathbf{x},\mathbf{y}}(i_1, i_2)$ is also invertible. Moreover, using the ultrametric norm, one can also guarantee that $|\det(A_{\mathbf{x},\mathbf{y}}(i_1, i_2))|$ is nonzero and constant on $V \times V$. Thus $(\mathbf{x}, \mathbf{y}) \rightarrow A_{\mathbf{x},\mathbf{y}}^{-1}(i_1, i_2)$ is a continuous map defined on $V \times V$. Hence

$$\|\mathbf{x} - \mathbf{y}\| \leq \|A_{\mathbf{x},\mathbf{y}}^{-1}(i_1, i_2)\| \|A_{\mathbf{x},\mathbf{y}}(i_1, i_2)(\mathbf{x} - \mathbf{y})^T\|. \quad (11.8)$$

Also

$$\|A_{\mathbf{x},\mathbf{y}}(i_1, i_2)(\mathbf{x} - \mathbf{y})^T\| = \|(f_{i_1}, f_{i_2})(\mathbf{x}) - (f_{i_1}, f_{i_2})(\mathbf{y})\| \leq \|\mathbf{f}(\mathbf{x}) - \mathbf{f}(\mathbf{y})\|. \quad (11.9)$$

Since $(\mathbf{x}, \mathbf{y}) \rightarrow \|A_{\mathbf{x},\mathbf{y}}^{-1}(i_1, i_2)\|$ is continuous on $V \times V$, and V is a compact ball, there is a $K_1 > 0$ such that $\|A_{\mathbf{x},\mathbf{y}}^{-1}(i_1, i_2)\| \leq K_1$. This together with (11.8) and (11.9) completes the proof of the lower bound in the Lipschitz condition. The upper bound follows from (11.7) using the compactness of V and continuity of the map $(\mathbf{x}, \mathbf{y}) \rightarrow \|A_{\mathbf{x},\mathbf{y}}\|$. \square

Acknowledgements. The authors thank the anonymous referees for their valuable comments that helped to clarify many details of the first version of this paper. SD thanks Ralf Spatzier and Subhajit Jana for many helpful discussions and several helpful remarks which have improved the paper.

REFERENCES

- [1] J. M. Aldaz. Besicovitch type properties in metric spaces,. (in press), <https://arxiv.org/abs/1902.02184>.
- [2] V. Beresnevich, E. Nesharim, and L. Yang. Winning property of badly approximable points on curves. *Duke Math J.*, To appear, 2022.
- [3] T. Das, L. Fishman, D. Simmons, and M. Urbański. A variational principle in the parametric geometry of numbers. <https://arxiv.org/abs/1901.06602>, 2019.
- [4] S. Datta and A. Ghosh. Multiplicative p -adic metric Diophantine approximation on manifolds and dichotomy of exponents. *Bull. Soc. Math. France*, 148(4):733–747, 2020.
- [5] S. Datta and A. Ghosh. Diophantine inheritance for p -adic measures. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, XXIII:49–80, 2022.
- [6] S. Datta and A. Ghosh. S -arithmetic inhomogeneous Diophantine approximation on manifolds. *Adv. Math.*, 400:Paper No. 108239, 2022.
- [7] A. Ghosh. Metric Diophantine approximation over a local field of positive characteristic. *J. Number Theory*, 124(2):454–469, 2007.
- [8] J. E. Hutchinson. Fractals and self-similarity. *Indiana Univ. Math. J.*, 30(5):713–747, 1981.
- [9] D. Kleinbock. Extremal subspaces and their submanifolds. *Geometric and Functional Analysis (GAFA)*, 13(2):437–466, 2003.
- [10] D. Kleinbock. An extension of quantitative nondivergence and applications to Diophantine exponents. *Trans. Amer. Math. Soc.*, 360(12):6497–6523, 2008.
- [11] D. Kleinbock, E. Lindenstrauss, and B. Weiss. On fractal measures and Diophantine approximation. *Sel. math., New ser.*, 10(4):479 – 523, 2005.
- [12] D. Kleinbock and G. A. Margulis. Flows on homogeneous spaces and Diophantine approximation on manifolds. *Ann. Math.*, 148:339–360, 1998.
- [13] D. Kleinbock and G. Tomanov. Flows on S -arithmetic homogeneous spaces and applications to metric Diophantine approximation. *Comm. Math. Helv.*, 82:519–581, 2007.

- [14] E. Le Donne and S. Rigot. Besicovitch covering property for homogeneous distances on the Heisenberg groups. *J. Eur. Math. Soc. (JEMS)*, 19(5):1589–1617, 2017.
- [15] E. Le Donne and S. Rigot. Besicovitch covering property on graded groups and applications to measure differentiation. *J. Reine Angew. Math.*, 750:241–297, 2019.
- [16] P. Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [17] W. Schikhof. Ultrametric calculus. an introduction to p-adic analysis. In *Cambridge University Press*, pages 113–132. Academic Press, London, 1985.
- [18] B. O. Stratmann and M. Urbański. Diophantine extremality of the Patterson measure. *Math. Proc. Cambridge Philos. Soc.*, 140(2):297–304, 2006.
- [19] G. Tomanov. Orbits on homogeneous spaces of arithmetic origin and approximations. In *Analysis on homogeneous spaces and representation theory of Lie groups, Okayama–Kyoto (1997)*, volume 26 of *Adv. Stud. Pure Math.*, pages 265–297. Math. Soc. Japan, Tokyo, 2000.
- [20] M. Urbański. Diophantine approximation and self-conformal measures. *J. Number Theory*, 110(2):219–235, 2005.

Victor Beresnevich, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK, UK
Email address: victor.beresnevich@york.ac.uk

Shreyasi Datta, DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY, UPPSALA
Email address: shreyasi1992datta@gmail.com, shreyasi.datta@math.uu.se

Anish Ghosh, SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, MUMBAI, INDIA 400005
Email address: ghosh@math.tifr.res.in