



UNIVERSITY OF LEEDS

This is a repository copy of *Denture status is related to chewing difficulty and bite force, though not BMI in elderly Thai adults*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/213650/>

Version: Published Version

---

**Proceedings Paper:**

Nitsuwat, S., Marshall, L., Sranacharoenpong, K. et al. (2 more authors) (2024) Denture status is related to chewing difficulty and bite force, though not BMI in elderly Thai adults. In: Proceedings of the Nutrition Society. Summer Conference 2023, Nutrition at key stages of the lifecycle, 03-06 Jul 2023, Liverpool, UK. Cambridge University Press , E374.

<https://doi.org/10.1017/S0029665123004731>

---

© The Author(s), 2023. Published by Cambridge University Press on behalf of The Nutrition Society. This article has been published in a revised form in <https://doi.org/10.1017/S0029665123004731>. This version is free to view and download for private research and study only. Not for re-distribution or re-use.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## Denture status is related to chewing difficulty and bite force, though not BMI in elderly Thai adults

S. Nitsuwat<sup>1</sup>, L. Marshall<sup>1</sup>, K. Sranacharoenpong<sup>2</sup>, A. Sarkar<sup>1</sup> and J. Cade<sup>1</sup>

<sup>1</sup>*School of Food Science and Nutrition, University of Leeds, Leeds, UK and*

<sup>2</sup>*Institute of Nutrition, Mahidol University, Nakhon Pathom, Thailand*

Oral processing of food is often affected due to limiting oro-muscular capabilities associated with aging<sup>(1)</sup>. Particularly loss of teeth, reduced salivary flow rate, and declines in oro-masticatory muscle performance result in chewing difficulties and, consequently, malnutrition<sup>(2)</sup>. However, the quantitative relationship between chewing difficulty with foods of varying textures and instrumentally measured oro-muscular forces and denture status remains rarely investigated.

**Aims:** To explore the associations of denture status and objectively assessed biting force ability with self-perceived chewing difficulties and nutritional status of community-dwelling older adults in Thailand.

A total of 148 older adult participants aged 64 to 90 years old (mean:  $71.9 \pm 6.0$  years) within the Bangkok Metropolitan Region and Chonburi Province completed a self-assessment of chewing ability by rating the chewing difficulty of 20 common food items (5 difficulty grades of texture) using a 3-point Likert scale. The maximum biting force of posterior teeth, as the objective assessment, was measured from both sides, and denture status was recorded. Nutritional status was assessed based on the body mass index of the participants with their food intake obtained by the 24-hour dietary recall.

Results show that participants who did not wear dentures and had more than 11 teeth rated their chewing ability significantly higher than those without dentures but had less than 11 teeth ( $p < 0.001$ ) and those with complete dentures ( $p < 0.001$ ) with differences especially seen in food items with a difficulty grade higher than 3. Correspondingly, the maximum biting force was also significantly lower in these groups ( $p < 0.05$ ) than in those with more than 11 teeth, implants, and partial dentures. Nonetheless, nutritional status was not significantly associated with denture status, biting force, or chewing difficulty.

The findings suggest that dentures status is associated with self-perceived mastication ability, and reduced maximum biting force may limit the consumption of food groups with certain textures and affect the variety of the nutrient intake in the vulnerable group.

### References

1. Sarkar A (2019) *PNS* 78(3), 329–39.
2. Xu F, Laguna L, Sarkar A (2019) *J Texture Stud* 50, 27–35.