



This is a repository copy of *Gravitational waves in a cyclic Universe: resilience through cycles and vacuum state*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/214226/>

Version: Published Version

---

**Article:**

Angelis, M.D. [orcid.org/0000-0003-3432-8250](https://orcid.org/0000-0003-3432-8250), Smith, A. [orcid.org/0009-0000-2254-1551](https://orcid.org/0009-0000-2254-1551), Giarè, W. [orcid.org/0000-0002-4012-9285](https://orcid.org/0000-0002-4012-9285) et al. (1 more author) (2024) Gravitational waves in a cyclic Universe: resilience through cycles and vacuum state. *Journal of Cosmology and Astroparticle Physics*, 2024 (06). ISSN 1475-7516

<https://doi.org/10.1088/1475-7516/2024/06/036>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

PAPER • OPEN ACCESS

# Gravitational waves in a cyclic Universe: resilience through cycles and vacuum state

To cite this article: Mariaveronica De Angelis *et al* JCAP06(2024)036

View the [article online](#) for updates and enhancements.

You may also like

- [Does Planck actually “see” the Bunch-Davies state?](#)  
Rose Baunach, Nadia Bolis, R. Holman et al.
- [Trans-Planckian issues for inflationary cosmology](#)  
Robert H Brandenberger and Jérôme Martin
- [Design and optimization of a longitudinal feedback kicker cavity for the HLS-II storage ring](#)  
Wei Xu, , W. Z. Wu et al.

## Gravitational waves in a cyclic Universe: resilience through cycles and vacuum state

Mariaveronica De Angelis , Adam Smith , William Giarè   
and Carsten van de Bruck 

*School of Mathematics and Statistics, The University of Sheffield,  
Hounsfield Road, S3 7RH Sheffield, U.K.*

*E-mail: [mdeangelis1@sheffield.ac.uk](mailto:mdeangelis1@sheffield.ac.uk), [asmith69@sheffield.ac.uk](mailto:asmith69@sheffield.ac.uk),  
[w.giare@sheffield.ac.uk](mailto:w.giare@sheffield.ac.uk), [c.vandebruck@sheffield.ac.uk](mailto:c.vandebruck@sheffield.ac.uk)*

**ABSTRACT:** We present a generalised calculation for the spectrum of primordial tensor perturbations in a cyclic Universe, making no assumptions about the vacuum state of the theory and accounting for the contribution of tensor modes produced in the dark energy phase of the previous cycle. We show that these modes have minimal impact on the spectrum observed in the current cycle, except for corrections on scales as large as the comoving Hubble radius today. These corrections are due to sub-horizon modes produced towards the end of the dark energy phase, persisting into the *ekpyrotic* phase of the next cycle as additional quanta. In relation to the vacuum state, we argue that non-Bunch-Davies quanta can easily overwhelm the energy density driving the dark energy phase, potentially compromising the model. Therefore, avoiding backreaction effects sets restrictive constraints on deviations away from the Bunch-Davies vacuum during this phase, limiting the overall freedom to consider alternative vacua in the cyclic Universe.

**KEYWORDS:** alternatives to inflation, cosmological perturbation theory, physics of the early universe, gravitational waves / theory

**ARXIV EPRINT:** [2403.00533](https://arxiv.org/abs/2403.00533)



---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Cyclic model and background dynamics</b>	<b>4</b>
2.1	Evolution and continuity of the scale factor across stages	5
2.2	Parameter constraints	7
<b>3</b>	<b>General primordial tensor spectrum</b>	<b>8</b>
3.1	General solutions in the different phases	9
3.2	Matching phases	9
3.3	Strain spectrum today ( $\tau = \tau_0$ )	10
<b>4</b>	<b>Resilience through cycles and the vacuum state</b>	<b>12</b>
4.1	Gravitational waves from a non-Bunch Davies vacuum	12
4.2	Gravitational waves produced from different phases	14
<b>5</b>	<b>Conclusion</b>	<b>17</b>

---

## 1 Introduction

The most compelling observational evidence supporting cosmological inflation [1–4] as the leading theory of the early Universe is currently provided by the Planck satellite measurement of the spectral index of scalar perturbations,  $n_s = 0.9649 \pm 0.0042$  [5]. In the simplest single-field slow-roll inflationary models, the spectrum of scalar modes is expected to be almost but not exactly flat [6–9], with deviations from flatness are quantified in terms of how much  $n_s$  deviates from 1 [10–13]. As a result, the Planck data seem to be in excellent agreement with the theoretical predictions of inflationary models [5, 14], ruling out a Harrison-Zeldovich scale-invariant spectrum [15–17] (corresponding to  $n_s = 1$ ) at a statistical level exceeding 8.5 standard deviations and lending weight to the inflationary paradigm.

That being said, with no aim to downplay the significance of this result or its interpretation, it is crucial to emphasise that, on its own, it does not provide conclusive evidence for cosmological inflation. Even hinging on a certain level of optimism and setting aside the uncertainty surrounding constraints on  $n_s$  from CMB experiments other than Planck<sup>1</sup> — or the potential implications arising from the well-known tensions [27–31] characterising the

---

<sup>1</sup>Over the years, constraints on the spectral index have been released by a multitude of Planck-independent CMB experiments such as WMAP [18, 19], the Atacama Cosmology Telescope (ACT) [20, 21], and the South Pole Telescope (SPT) [22, 23]. When considering these data at face value, Planck is currently the only experiment excluding  $n_s = 1$  at a statistical significance much larger than  $3\sigma$ . Conversely, ACT shows a preference for  $n_s = 1$  [21, 24]. Different combinations of these data overall support the result  $n_s \neq 1$ , although sometimes they lead to discordant results in terms of the other inflationary parameters or the preferred inflationary models [25, 26].

recent debate<sup>2</sup> — alternative theoretical mechanisms have been put forth, yielding an almost scale-invariant spectrum of primordial density fluctuations without invoking inflation.

An illustrative example of such mechanisms is the cyclic Universe [43–49] that, in contrast to the conventional cosmological framework, suggests a periodic history for the Cosmos. The model has been extensively studied and discussed in relation to a broad range of topics, including quantum gravity, modified gravity, gravitational waves and dark energy, see e.g., refs. [50–84] or refs. [85, 86] for reviews. In broad terms, each cycle comprises a phase recasting the standard Hot Big Bang theory (during which large-scale structures take shape), followed by a phase of slow, accelerated expansion mirroring the present-day observational evidence for a Dark Energy dominated dynamics. In the cyclic Universe, this latter stage also serves to dilute inhomogeneities and flatten the spatial geometry. Subsequently, a contraction phase ensues, generating nearly scale-invariant density perturbations. Finally, the cycle concludes with a big-crunch/big-bang transition, during which matter and radiation are generated, setting the stage for the next cycle.

Notice that both inflation and the cyclic Universe provide physical mechanisms to produce an almost scale-invariant spectrum of density perturbations [87–89]. In addition, they can both explain observational facts such as the homogeneity in the cosmic microwave background (CMB) radiation [90] and the fact that the present-day spatial geometry of the Universe appears to be flat, or at the very least nearly flat.<sup>3</sup> Therefore, at first glance, one might wonder how to distinguish between the two models. Focusing solely on scalar modes, this is a challenging knot to unravel [112, 113]. However, the two scenarios yield significantly distinct predictions for the stochastic background of gravitational waves [50]. Similar to scalar modes, inflation predicts a nearly scale-invariant (red-tilted) spectrum of tensor modes [12, 13, 114]. Conversely, in the cyclic Universe, the tensor spectrum is typically blue-tilted, and its amplitude is many orders of magnitude lower than that predicted by inflation, remaining well below any observable threshold achievable in the near future. Consequently, any measurement of primordial gravitational waves (e.g., through the effects left in the CMB B-mode polarisation at large angular scales) would offer conclusive evidence for inflation, discounting the cyclic model.

Despite this fact being acknowledged as a strength for inflation and perhaps a limitation in predictive capacity for the cyclic model, it is worth emphasising a few caveats surrounding this conclusion. Firstly, despite the best efforts, the detection of primordial tensor modes remains elusive at present [115], making it impossible to discriminate between the two scenarios. Therefore, the cyclic Universe remains an alternative worth considering. Secondly, the inflationary predictions concerning the amplitude and tilt of the tensor spectrum depend significantly on the specific model. While well-known consistency relations among inflationary parameters can be derived within single-field slow-roll inflation minimally coupled to gravity [12, 13], these relations can be violated by various physical mechanisms. A long yet not exhaustive list of possibilities include considering modified gravity [116–119], multi-field inflation [120–124], additional (spectator) rolling axion fields [125–128], couplings to axion-gauge or spin-2 fields [129, 130], breaking spatial and/or temporal diffeomorphism invariance [131–

<sup>2</sup>For studies suggesting potential implications of cosmological tensions for inflation, see, e.g., refs. [32–42].

<sup>3</sup>For recent discussions surrounding the spatial geometry of the Universe, see, e.g., [91–111].

[134], higher curvature corrections to the effective gravitational action [135, 136], higher order operators in effective field theory [137, 138], violations of the null energy condition [139, 140], alternative vacuum state/initial conditions [141–143], sound speed resonances [144], inflation in an Universe filled with an elastic medium [145], and possible effects/models inspired by quantum gravity [146–149]. Many of these more elaborated scenarios yield completely different predictions, often resulting in a blue-tilted spectrum and possibly leaving signatures in different cosmological and astrophysical observables [150–159]. Furthermore, models with an arbitrarily small tensor amplitude can always be constructed (see, e.g., ref. [160]), making it virtually impossible to rule out inflation based solely on a lack of detection of primordial gravitational waves. This is a critique frequently raised against inflationary cosmology as it questions its actual predictive capability.

As concerns the cyclic Universe, since any difference with respect to inflation in terms of predictions is likely to be confined to the spectrum of tensor modes, it becomes interesting to test whether similar caveats apply or if the model demonstrates greater resilience.

In light of this, we review the production of primordial gravitational waves in a cyclic Universe, identifying (and eventually clarifying) some conceptual aspects related to its concrete predictivity. Specifically, prevailing calculations in the existing literature conventionally establish initial conditions for primordial scalar and tensor modes during the *ekpyrotic* contracting phase [50]. While for scalar perturbations the implications of setting the initial conditions in different phases have been examined [161], the calculation of the tensor spectrum has always been performed starting in the *ekpyrotic* phase, assuming a Bunch-Davies (BD) vacuum state and neglecting any potential contributions arising from tensor modes originating during the dark energy phase of the previous cycle. This leads us to question whether they exert any influence on the spectrum observed in the current cycle. Taking a broader perspective, one may wonder whether the predictions concerning tensor modes remain resilient throughout the diverse cycles of the model itself. Yet another aspect that is imperative to clarify is to what extent the predictions depend on the choice of the vacuum state, addressing the crucial question of what freedom exists in the cyclic Universe regarding the choice of the vacuum state and whether substantial alterations can arise in the tensor spectrum by assuming different vacua, akin to what is found in inflationary cosmology.

To address these points, we present a general model for the evolution of gravitational waves produced in a cyclic Universe, making no assumptions about the initial vacuum state and starting the calculation from the dark energy phase of the previous cycle. We find that the additional tensor models originated in the previous cycle have minimal impact on the tensor spectrum observed in the current cycle, except for corrections on scales as large as the comoving Hubble radius today that are due to sub-horizon modes produced towards the end of the previous dark energy phase. Most importantly, we find that non-BD quanta in the dark energy phase can easily overwhelm the energy density associated with the modulus field, potentially spoiling the model. Avoiding these backreaction effects sets restrictive constraints on deviations away from the BD vacuum during the dark energy phase, thereby limiting the overall freedom to consider alternative vacua in the cyclic Universe.

The paper is organised as follows. In section 2, we introduce the cyclic Universe model and review its background dynamics. In section 3, we consider the evolution of gravitational

waves in such a Universe, starting from the previous cycle's dark energy phase and deriving the evolution in full generality. In section 4, we discuss the implications for the model's predictions, deriving constraints on the choice of the vacuum state and analysing the contribution coming from tensor modes originated in the previous cycle. Finally, in section 5, we derive our main conclusions.

## 2 Cyclic model and background dynamics

We consider a simple scalar field setup in which the dynamics of the cyclic model in the Einstein frame are well described by the  $4D$  effective Lagrangian [161]

$$\mathcal{L} = \sqrt{-g} \left( \frac{M_{\text{Pl}}^2}{2} R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) \right), \quad (2.1)$$

where  $g$  is the determinant of the metric  $g_{\mu\nu}$ ,  $R$  is the Ricci scalar and we adopt units where  $c = 1$ . The scalar field  $\phi$  is a modulus field, driving the dark energy dominated phase and the subsequent *ekpyrotic* and contracting kinetic phases, which we discuss below. Assuming a spatially flat FRLW background, the scalar field satisfies the usual equation of motion

$$\ddot{\phi} + 3H\dot{\phi} + V_{,\phi} = 0, \quad (2.2)$$

where dots denote derivatives with respect to cosmological time  $t$ . On the other hand, ignoring any coupling between the scalar field and the other standard model species and neglecting any additional contributions from the latter to the total Universe energy density, the evolution of the scale factor is governed by the Friedmann equation that, in terms of the Hubble parameter  $H = \dot{a}/a$ , reads

$$H^2 = \frac{1}{3M_{\text{Pl}}^2} \left( \frac{1}{2} \dot{\phi}^2 + V(\phi) \right). \quad (2.3)$$

In what follows, to efficiently describe the dynamics of the cyclic Universe we focus on a phenomenological potential of the form

$$V = V_0 \left( 1 - e^{-c\phi/M_{\text{Pl}}} \right) Y(\phi), \quad (2.4)$$

where  $V_0$  is of the same order of the vacuum energy observed in today's Universe,  $c$  is a positive constant value and  $Y(\phi)$  is a step function. Notice that our choice concerning the specific potential employed in the work is, in part, motivated by the fact that the exponential form is convenient for analysis, and in part from the fact that the same potential has been widely adopted in similar studies, allowing a direct comparison between our findings and other results documented in the existing literature. However, it is important to emphasise that cyclic models can emerge from a broad spectrum of different potentials that should ultimately emerge from the higher-dimensional theory. Without loss of generality, the only constraint comes from requiring an acceptable spectrum of scalar perturbations that implies considering a steep, strongly negatively curved region across observational ranges of the field to reproduce.

Having that said, the potential (2.4) serves multiple purposes, including describing dark energy responsible for cosmic acceleration observed today. More importantly, it plays a

crucial role in transitioning the Universe from accelerated expansion to contraction. This is achieved by rolling from positive to negative values of the potential until reaching a time where  $H^2 = 0$  and consequently triggering a phase characterised by an equation of state  $\omega \gg 1$ . For instance, by solving eq. (2.3) it can be shown that when the negative potential dominates  $V \simeq -V_0 e^{-c\phi}$  (i.e. *ekpyrotic* phase), the scale factor behaves as [161]

$$a(t) \sim (-t)^{\tilde{\alpha}}, \quad (2.5)$$

where  $t$  has negative values and  $\tilde{\alpha} = 2/c^2$ . At this point it is also convenient to introduce the conformal time  $d\tau = dt/a(t)$ , which we will frequently use later. In terms of the conformal time, the scale factor during the *ekpyrotic* phase evolves as:

$$a(\tau) \sim \left[ (-1)^{\tilde{\alpha}} (\tau - \tilde{\alpha}\tau)^{\tilde{\alpha}/(1-\tilde{\alpha})} \right]_{\tau_i}^{\tau_f}, \quad (2.6)$$

underscoring that the Universe is gradually contracting while the scalar field slowly descends along its sharply decreasing negative potential, to produce an acceptable spectrum of cosmological scalar perturbations.

In the literature, the *ekpyrotic* phase is typically assumed as the starting point of the cycle where initial conditions of primordial perturbations are imposed, and the calculations of the relative scalar and tensor spectra begin. However, in this study, we want to extend the model to include the contribution of tensor perturbations produced during the dark energy phase of the previous cycle to investigate whether they could have any impact on the spectrum we observe in the current cycle and eventually clarify why (not). To do so, the overall strategy will be to start the calculation in the dark energy phase of the previous cycle (making no assumptions on the vacuum state) and evolve the system through four regimes. For this reason, before dealing with the explicit calculation of the tensor spectrum, given that in our case we consider one more phase than in previous studies, it is useful to dedicate the following two subsections to reviewing the background dynamics of the model. In the same spirit of the discussion outlined so far, we start from the dark energy phase of the previous cycle and evolve the scale factor, ensuring its continuity across the boundaries of each phase. Additionally, we derive constraints on the model's parameters based on minimum theoretical requirements, such as the continuity of  $H(t)$  and the consistency of the theory across cycles.

## 2.1 Evolution and continuity of the scale factor across stages

### 2.1.1 Dark Energy phase

We start from the dark energy phase, in which the expansion rate  $H$  is roughly constant and the scale factor  $a(t)$  behaves as

$$a(t) = a(t_{\text{tr}}) e^{H(t-t_{\text{tr}})} \quad t < t_{\text{tr}}, \quad (2.7)$$

with  $t_{\text{tr}}$  transition time between dark energy and *ekpyrotic* phase. The equation above can be translated in terms of conformal time as

$$a(\tau) = \frac{1}{H(B - \tau)}, \quad (2.8)$$

where, for the continuity across  $\tau = \tau_{\text{tr}}$ ,  $B$  is fixed to<sup>4</sup>

$$B = \frac{1}{a(\tau_{\text{tr}})H} + \tau_{\text{tr}}. \quad (2.9)$$

Additionally, by means of eq. (2.6), we can infer

$$\frac{a(\tau_{\text{tr}})}{a(\tau_{\text{end}})} = \left( \frac{\tau_{\text{tr}} - \tau_{\text{ek}}}{\tau_{\text{end}} - \tau_{\text{ek}}} \right)^\alpha, \quad (2.10)$$

where  $\alpha \equiv \tilde{\alpha}/(1 - \tilde{\alpha})$  and  $\tau_{\text{ek}} \equiv (1 - 2\alpha)\tau_{\text{end}}$  is the conformal time corresponding to when the potential diverges to minus infinity.

### 2.1.2 Ekpyrotic phase

As a next step, we transition to the *ekpyrotic* phase. In this phase the potential becomes negative and the Einstein frame expansion forces the scale factor to contract

$$\frac{a(\tau)}{a(\tau_{\text{end}})} = \left( \frac{\tau - \tau_{\text{ek}}}{\tau_{\text{end}} - \tau_{\text{ek}}} \right)^\alpha \quad \tau_{\text{tr}} < \tau < \tau_{\text{end}}. \quad (2.11)$$

We note again that, being  $\alpha \ll 1$ , the contraction is very slow.

### 2.1.3 Contracting kinetic phase

Once  $\tau > \tau_{\text{end}}$  we enter the region where the effects of the potential are negligible, namely  $\phi < \phi_{\text{end}}$ . During this period — known as contracting kinetic phase — we have:

$$\frac{a(\tau)}{a(\tau_{\text{r}})} = \left( \frac{-\tau}{(1 + \chi)\tau_{\text{r}}} \right)^{\frac{1}{2}} \quad \tau_{\text{end}} < \tau < 0, \quad (2.12)$$

where  $\chi$  is a small positive constant that measures the amount of radiation created at the bounce ( $\tau = 0$ ).

### 2.1.4 Expanding kinetic phase

Finally, for the last phase of the cycle, the so-called expanding kinetic phase, we get

$$a(\tau) = \left( \frac{\tau}{\tau_{\text{r}}} \right)^{\frac{1}{2}}, \quad 0 < \tau < \tau_{\text{r}}, \quad (2.13)$$

Notice that, for convenience, we work in a coordinate frame where the scale factor is set to  $a(\tau_{\text{r}}) = 1$  for the time  $t_{\text{r}}$  corresponding to the beginning of the radiation-dominated era. The corresponding conformal time  $\tau_{\text{r}} = (2H_{\text{r}})^{-1}$  is constrained by the radiation temperature  $T_{\text{r}}$ , being  $H_{\text{r}} \propto T_{\text{r}}^2/M_{\text{Pl}}$ . Following ref. [46], we choose a quite conservative value  $T_{\text{r}} \sim 10^7$  GeV, akin to that obtained in the more familiar standard cosmology at the end of the reheating phase following inflation. This choice also ensures that we can safely recover predictions of primordial Big Bang Nucleosynthesis (BBN).

---

<sup>4</sup>The value of  $B$  is achieved by considering  $H_0, a(\tau_{\text{r}})$  and  $\tau_{\text{r}}$  given in the next sections.

## 2.2 Parameter constraints

### 2.2.1 Continuity of the Hubble parameter

In order to constrain the length of the *ekpyrotic* contracting phase, we require that the Hubble parameter returns to its original value after every cycle. Following ref. [161], this implies that

$$H_{\text{end}}/H_0 \approx \sqrt{\frac{-V_{\text{end}}}{V_0}} \quad (2.14)$$

where  $V_{\text{end}}$  is the depth of the potential well and  $V_0$  is the height of the potential plateau. The spectral range of perturbations produced when the field rolls from  $V \approx 0$  to  $V \approx -V_{\text{end}}$ , satisfies

$$\frac{k_{\text{max}}}{k_{\text{min}}} \approx \sqrt{\frac{-V_{\text{end}}}{V_0}}, \quad (2.15)$$

and it needs to span at least  $N = 60$  e-folds for the *ekpyrotic* phase to produce a scale-invariant spectrum over a broad enough range of scales for us to observe today [5, 14]. Hence, the transition time between dark energy and *ekpyrotic* phase is constrained to

$$\frac{H_{\text{tr}}}{H_{\text{end}}} = 2\alpha\tau_{\text{end}} \frac{a_{\text{end}} a_{\text{tr}}}{(\tau_{\text{tr}} - \tau_{\text{ek}})} < e^{-60}. \quad (2.16)$$

Moreover, as  $a(\tau) \approx \text{const}$  during the *ekpyrotic* contracting phase [161], from eq. (2.16) it follows that

$$|\tau_{\text{tr}} - \tau_{\text{ek}}| > 2\alpha\tau_{\text{end}} a_{\text{end}}^2 e^{60}. \quad (2.17)$$

### 2.2.2 Cycling constraint

To place constraints on the duration of the kinetic evolution phases, we require that they must last enough time for the scalar field to have started at the potential minimum (i.e.  $\phi = \phi_{\text{end}}$ ), moved off to the Bounce (i.e.  $\phi \rightarrow -\infty$ ), and returned all the way back past  $\phi_{\text{end}}$  and made it up to the potential plateau to begin a radiation-dominated Universe.

Barring some brief  $\omega \gg 1$  period (which divides the expanding kinetic phase into two parts) as the field moves back up past  $\phi_{\text{end}}$  to the potential plateau, from eq. (2.2) and eq. (2.4) we find

$$\phi - \phi_{\text{end}} = c_1 \ln\left(\frac{t}{t_{\text{end}}}\right), \quad (2.18)$$

where the factor  $c_1^2 = 2/3$  comes from eq. (2.3) during kinetic domination,  $t_{\text{end}}$  is the time taken to reach  $\phi_{\text{end}}$  starting at  $\phi \rightarrow -\infty$ .

Notice that in the region where  $\omega \gg 1$ , contributions from  $Y(\phi)$  becomes relevant and  $V \approx V_0(1 - e^{-c\phi})$ . Therefore, in this case, the time  $t_{\text{r}}$  required to climb the potential well and reach the plateau at  $V \approx V_0$  can be bounded to

$$\frac{t_{\text{r}}}{t_{\text{end}}} > \left(\frac{V_{\text{end}}}{V_0}\right) \sqrt{\frac{3}{2c^2}}. \quad (2.19)$$

Since the time taken for the field to cross the negative region of the potential before radiation domination begins is given by

$$\frac{t_r}{t_{\text{end}}} \approx \frac{\sqrt{V_{\text{end}}}}{H_r}, \quad (2.20)$$

from eq. (2.19) we can infer an upper limit on the Hubble parameter at  $t_r$  which reads

$$H_r \lesssim \frac{\sqrt{V_{\text{end}}}}{M_{\text{Pl}}} \left( \frac{V_0}{V_{\text{end}}} \right)^{\frac{3}{2c^2}}. \quad (2.21)$$

This upper limit constrains the ratio between  $\tau_r$  and  $\tau_{\text{end}}$  to

$$\Gamma = \left| \frac{\tau_r}{\tau_{\text{end}}} \right| \gtrsim \left( \frac{V_{\text{end}}}{V_0} \right)^{\sqrt{\frac{2}{3c^2}}} \simeq 10^8, \quad (2.22)$$

where we used that the vacuum energy density  $\rho_\Lambda = V_0 \sim 10^{-120} M_{\text{Pl}}^4$  and  $V_{\text{end}} \sim 10^{-20} M_{\text{Pl}}^4$ . Following ref. [50], throughout this paper, we always consider the dimensionless parameter  $\Gamma \sim 10^8$ .

In conclusion, the final constraints we derive for the cyclic model at the background level (and that are important to bear in mind for the following discussion on tensor perturbations) are:

$$\left| \frac{\tau_r}{\tau_{\text{end}}} \right| \gtrsim 10^8, \quad (\tau_{\text{tr}} - \tau_{\text{ek}}) \gtrsim 10^9, \quad \tau_r = \frac{1}{2H_r}. \quad (2.23)$$

### 3 General primordial tensor spectrum

Considering a spatially flat FLRW metric, in the synchronous gauge the perturbed line element reads:

$$ds^2 = a^2(\tau) \left[ d\tau^2 - (\delta_{ij} + h_{ij}) dx^i dx^j \right]. \quad (3.1)$$

Tensor modes (i.e., metric perturbations) are described in terms of the transverse and traceless part of the symmetric  $3 \times 3$  matrix  $h_{ij}$ . To characterise the contribution of each wavenumber  $k$  to  $h_{ij}(t, \mathbf{x})$ , we consider a Fourier representation  $\tilde{h}_{ij}(t, \mathbf{k})$ . Moving to the Fourier space, focusing on one particular polarisation state, and assuming isotropy, the gravitational wave field  $h_k$  satisfies the following equation:

$$h_k'' + 2\frac{a'}{a}h_k' + k^2 h_k = 0, \quad (3.2)$$

where  $(\dots)'$  denotes the derivative with respect to conformal time  $\tau$ . However, it is more convenient to use a new variable  $f_k(\tau) \equiv a(\tau)h_k(\tau)$  satisfying

$$f_k'' + \left( k^2 + \frac{a''}{a} \right) f_k = 0. \quad (3.3)$$

After redefining  $f_k = i\sqrt{\tau} u_k$ , eq. (3.3) assumes the more familiar form of a Bessel equation and, for each phase of the cyclic model, the general solution involves a linear combination of the Hankel functions of the first and second kind, that we denote as  $H^{(1,2)}$ .

In this section, we present a generalised calculation for the primordial tensor spectrum in the cyclic Universe. In section 3.1, we derive the general solution of eq. (3.3) making no assumptions about the vacuum state of the theory and considering the contribution of tensor modes produced in the dark energy phase of the previous cycle rather than starting directly from the *ekpyrotic* phase of the present cycle. In section 3.2, we require internal consistency in the evolution of the tensor mode amplitudes throughout the four phases of the model, ensuring that both  $h_k(\tau)$  and  $h'_k(\tau)$  remain continuous functions and matching the general solutions across the different phases. Finally, in section 3.3, we derive the expression of the primordial tensor spectrum and briefly discuss its strain today.

### 3.1 General solutions in the different phases

#### 3.1.1 General solution in the Dark Energy phase

During the dark energy phase,  $a(\tau)$  is given by eq. (2.8), and the general solution of eq. (3.3) reads

$$f_k(\eta) = \sqrt{-k\eta} \left( D_1(k) H_{\frac{3}{2}}^{(1)}(-k\eta) + D_2(k) H_{\frac{3}{2}}^{(2)}(-k\eta) \right) \quad \tau < \tau_{\text{tr}}, \quad (3.4)$$

where  $H_n^{(1)}$  and  $H_n^{(2)}$  denote the Hankel functions of first and second kind respectively,  $\eta = \tau - B$ , where  $B$  is given by eq. (2.9) and, for each wave-number  $k$ ,  $D_{1,2}(k)$  are arbitrary constants.

#### 3.1.2 General solution in the *ekpyrotic* contraction

In the stage of the *ekpyrotic* contraction,  $a(\tau)$  is given by eq. (2.11) and the general solution reads

$$f_k(\tau) = \sqrt{y} \left( A_1(k) H_n^{(1)}(y) + A_2(k) H_n^{(2)}(y) \right) \quad \tau_{\text{tr}} < \tau < \tau_{\text{end}}, \quad (3.5)$$

where  $y \equiv -k(\tau - \tau_{\text{ek}})$ .

#### 3.1.3 General solution in the kinetic contracting phase

Moving to the kinetic contracting phase and making use of eq. (2.12), we achieve

$$f_k(\tau) = \sqrt{-k\tau} \left( B_1(k) H_0^{(1)}(-k\tau) + B_2(k) H_0^{(2)}(-k\tau) \right) \quad \tau_{\text{end}} < \tau < 0, \quad (3.6)$$

#### 3.1.4 General solution in the kinetic expanding phase

Considering eq. (2.13) in the kinetic expanding phase the general solution is

$$f_k(\tau) = \sqrt{k\tau} \left( C_1(k) H_0^{(1)}(k\tau) + C_2(k) H_0^{(2)}(k\tau) \right) \quad 0 < \tau < \tau_{\text{r}}. \quad (3.7)$$

### 3.2 Matching phases

In what follows we obtain expressions for the coefficients  $D_{1,2}(k)$ ,  $A_{1,2}(k)$ ,  $B_{1,2}(k)$  and  $C_{1,2}(k)$  by matching  $h_k(\tau)$  and  $h'_k(\tau)$  at the boundaries of each phase.

### 3.2.1 Dark energy — *ekpyrotic*

At the boundary between dark energy and *ekpyrotic* stage, we require continuity of  $f_k(\tau)$  namely

$$\sqrt{x_{\text{tr}}} \left( D_1(k) H_{\frac{3}{2}}^{(1)}(x_{\text{tr}}) + D_2(k) H_{\frac{3}{2}}^{(2)}(x_{\text{tr}}) \right) = \sqrt{y_{\text{tr}}} \left( A_1(k) H_n^{(1)}(y_{\text{tr}}) + A_2(k) H_n^{(2)}(y_{\text{tr}}) \right), \quad (3.8)$$

where  $x_{\text{tr}} = -k\eta(\tau_{\text{tr}})$ , and continuity of  $f'(\tau)$  which we write in the matrix form  $\underline{\mathbf{y}} \mathbf{A} = \underline{\mathbf{x}}_1 \mathbf{D}$  where

$$\begin{aligned} \underline{\mathbf{y}} &= \begin{pmatrix} \sqrt{y_{\text{tr}}} H_n^{(1)}(y_{\text{tr}}) & \sqrt{y_{\text{tr}}} H_n^{(2)}(y_{\text{tr}}) \\ \sqrt{y_{\text{tr}}} H_{n-1}^{(1)}(y_{\text{tr}}) - \frac{(n-\frac{1}{2})}{\sqrt{y_{\text{tr}}}} H_n^{(1)}(y_{\text{tr}}) & \sqrt{y_{\text{tr}}} H_{n-1}^{(2)}(y_{\text{tr}}) - \frac{(n-\frac{1}{2})}{\sqrt{y_{\text{tr}}}} H_n^{(2)}(y_{\text{tr}}) \end{pmatrix}, \\ \underline{\mathbf{x}}_1 &= \begin{pmatrix} \sqrt{x_{\text{tr}}} H_{\frac{3}{2}}^{(1)}(x_{\text{tr}}) & \sqrt{x_{\text{tr}}} H_{\frac{3}{2}}^{(2)}(x_{\text{tr}}) \\ \sqrt{x_{\text{tr}}} H_{\frac{1}{2}}^{(1)}(x_{\text{tr}}) - \frac{1}{\sqrt{x_{\text{tr}}}} H_{\frac{3}{2}}^{(1)}(x_{\text{tr}}) & \sqrt{x_{\text{tr}}} H_{\frac{1}{2}}^{(2)}(x_{\text{tr}}) - \frac{1}{\sqrt{x_{\text{tr}}}} H_{\frac{3}{2}}^{(2)}(x_{\text{tr}}) \end{pmatrix}, \end{aligned} \quad (3.9)$$

and

$$\mathbf{A} = \begin{pmatrix} A_1 \\ A_2 \end{pmatrix}, \quad \mathbf{D} = \begin{pmatrix} D_1 \\ D_2 \end{pmatrix}, \quad \text{with } x_{\text{tr}} \equiv -k\tau_{\text{tr}}. \quad (3.10)$$

### 3.2.2 *Ekpyrotic* — kinetic contraction

Similarly here, at  $\tau = \tau_{\text{end}}$  namely the end of the *ekpyrotic* phase, we require  $\underline{\mathbf{y}}_2 \mathbf{B} = \underline{\mathbf{x}}_2 \mathbf{A}$  where

$$\begin{aligned} \underline{\mathbf{y}}_2 &= \begin{pmatrix} H_0^{(1)}(x_e) & H_0^{(2)}(x_e) \\ \sqrt{x_e} H_{-1}^{(1)}(x_e) + \frac{H_0^{(1)}(x_e)}{2\sqrt{x_e}} & \sqrt{x_e} H_{-1}^{(2)}(x_e) + \frac{H_0^{(2)}(x_e)}{2\sqrt{x_e}} \end{pmatrix}, \\ \underline{\mathbf{x}}_2 &= \begin{pmatrix} \sqrt{2\alpha} H_n^{(1)}(2\alpha x_e) & \sqrt{2\alpha} H_n^{(2)}(2\alpha x_e) \\ \sqrt{2\alpha x_e} H_{n-1}^{(1)}(2\alpha x_e) - \frac{(n-\frac{1}{2})}{\sqrt{2\alpha x_e}} H_n^{(1)}(2\alpha x_e) & \sqrt{2\alpha x_e} H_{n-1}^{(2)}(2\alpha x_e) - \frac{(n-\frac{1}{2})}{\sqrt{2\alpha x_e}} H_n^{(2)}(2\alpha x_e) \end{pmatrix}, \end{aligned} \quad (3.11)$$

and

$$\mathbf{B} = \begin{pmatrix} B_1 \\ B_2 \end{pmatrix}, \quad \text{with } x_e \equiv -k\tau_{\text{end}}. \quad (3.12)$$

### 3.2.3 Kinetic contraction — kinetic expansion

The matching for these two final stages arises at  $\tau = 0$ , which is trivial as we have

$$C_{1,2} = -\sqrt{1 + \chi} B_{2,1}. \quad (3.13)$$

### 3.3 Strain spectrum today ( $\tau = \tau_0$ )

The quantity in terms of which we assess the production of primordial gravitational waves in the cyclic Universe is the dimensionless strain spectrum

$$\Delta h = k^{\frac{3}{2}} \frac{|h_k(\tau)|}{\pi}. \quad (3.14)$$

It is convenient to evaluate the dimensionless strain spectrum in the radiation-dominated epoch. Using the general solutions we derived and matched across the different phases of the model, it reads:

$$\Delta h(k, \tau_r) = \frac{k^2 \sqrt{2\tau_r}}{a(\tau_r) \pi M_{\text{Pl}}} \left| C_1(k) H_0^{(1)}(x_r) + C_2(k) H_0^{(2)}(x_r) \right|, \quad (3.15)$$

where  $x_r \equiv k \tau_r$ . The strain tensor spectrum today  $\Delta h(k, \tau_0)$  can be easily related to the spectrum in the radiation dominated epoch by means of the transfer function formalism as:

$$\Delta h(k, \tau_0) \equiv \mathcal{T}(k) \Delta h(k, \tau_r), \quad (3.16)$$

where  $\mathcal{T}(k)$  is the transfer function, responsible for propagating the spectrum forwards to today's observed spectrum. Following ref. [50], we use a transfer function of the form

$$\mathcal{T}(k) \approx \left( \frac{k_0}{k} \right)^2 \left( 1 + \frac{k}{k_{\text{eq}}} + \frac{k^2}{k_{\text{eq}} k_r} \right), \quad (3.17)$$

where  $k_r = a_r H_r \approx T_r^2 / M_{\text{Pl}}$  is the wave number of modes crossing the horizon at the start of radiation domination.

So far, the whole calculation is done by setting  $a(\tau_r) = 1$ . However, for ease of comparison with data and the other results documented in the literature, it is convenient to return to the usual coordinate system where  $a_0 = 1$ . This can be easily done by means of the well-known inverse scaling between the scale factor and the CMB temperature (i.e.  $a \propto 1/T$ ). This relation fixes the ratio  $a(\tau_0)/a(\tau_r) \propto T_r/T_0 \sim 10^{20}$ . As a result, we can re-scale the comoving frequency accordingly, which now takes the more familiar value  $\hat{k}_r = k_r/a(\tau_0) = 10^{-1}$  Hz. From now on, we will name the re-scaled comoving frequency  $\hat{k}_r$  as  $k_r$ . Additionally, always following from the inverse proportionality between the scale factor and the CMB temperature, we expect a spectral range of modes entering between the start of radiation domination and today of the order of

$$\frac{k_0}{k_r} \propto \frac{T_0}{T_r} \sim 6.6 \times 10^{-20}. \quad (3.18)$$

Taking in mind that during matter domination  $H \sim a^{-3/2}$ , we get

$$\frac{k_{\text{eq}}}{k_0} \approx \sqrt{1 + z_{\text{eq}}} \sim 10^2, \quad (3.19)$$

where  $z_{\text{eq}}$  is the redshift at the equivalence.

We can also place a bound on the wave number of modes on the horizon at the previous dark energy-*ekpyrotic* transition,  $k_{\text{tr}}$ , which will be useful later. The wavelength of such modes can be estimated using the duration of the *ekpyrotic* phase, which we require to last at least 60 e-folds to effectively homogenise and flatten the Universe for the subsequent cycle. However, a much stronger constraint comes from eq. (2.15), as the number of e-folds during the *ekpyrotic* phase is given by  $N \sim \ln(H_{\text{end}}/H_0) \approx \ln(\sqrt{-V_{\text{end}}/V_0})$ . Taking our previous constraints on  $V_0$  at today's dark energy density and  $V_{\text{end}}$  at the GUT scale, we arrive at  $N \sim 115$ . This forces our horizon to shrink by a factor of  $\sim 10^{50}$  during the dark *ekpyrotic* phase, and hence we would expect

$$\frac{k_{\text{end}}}{k_{\text{tr}}} \approx 10^{50}. \quad (3.20)$$

## 4 Resilience through cycles and the vacuum state

The calculation of the spectrum of gravitational waves introduced in the previous section, in addition to considering the contribution of tensor modes produced in the dark energy phase of the previous cycle, is entirely general regarding the vacuum state and applies to any particular choice. This allows us to test several conceptual aspects of the theory (partially highlighted in the introduction), such as its resilience throughout cycles and the implications of the choice of the vacuum states. In this section, we examine both these issues in more detail. In section 4.1, we focus on the choice of the vacuum state of the theory, deriving novel constraints on the latter based on internal consistency through cycles and epochs. In section 4.2, we quantify the extent to which including the contribution from tensor modes originating during the dark energy phase of the previous cycle changes the spectrum of tensor perturbations observed in the current cycle.

### 4.1 Gravitational waves from a non-Bunch Davies vacuum

As often speculated both in quantum field theory and effective field theory, the choice of the vacuum state represents an important topic of discussion and significance as it plays a crucial role in determining the properties of the theory and the physical predictions it makes. On the one hand, in quantum field theory, the vacuum state is the lowest-energy state of a quantum field, which usually corresponds to the state with no physical particles. On the other hand, effective field theories often deal with specific energy scales or regimes of a more fundamental theory (such as in our effective representation of the cyclic Universe), and the choice of the vacuum state may involve integrating out high-energy degrees of freedom and focusing on the low-energy behavior. This makes the choice of the vacuum state far from trivial when considering the possibility of new physics at sufficiently high energies.

The selection of the vacuum state has been the subject of intense study and attention in the context of inflationary models where considering more exotic (though physically motivated) vacuum states other than BD) can lead to markedly different predictions for the spectrum of scalar and tensor modes, see e.g.[141–143] and references therein. Just to mention one example among many, even sticking to the framework of single-field inflation minimally coupled to gravity, considering an exotic vacuum can result in the violation of the usual slow-roll consistency relations allowing for a blue-tilted spectrum of gravitational waves, with an enhanced amplitude on small scales that can eventually produce observable signatures visible by CMB and Gravitational Waves experiments.

However, the same issue has not been investigated in the cyclic Universe. Previous analyses of the spectrum of the gravitational wave assume a BD vacuum for perturbations produced in the *ekpyrotic* phase, enforcing the solution to converge to that of a plane wave in flat space at sufficiently short distances [50]. As argued in ref. [44], this choice mostly relies on the classical treatment of the dynamics of the evolving modulus field at arbitrarily small length scales. However, just like in inflation, the choice may no longer be trivial if the dynamics of the modulus field is influenced by new physics at some characteristic energy scale  $M$ . As a result, one may wonder whether in the cyclic Universe, the choice of the vacuum state is somehow affected by the same level of “arbitrariness” as inflation, or if additional constraints can be derived based on the strong interconnection between the different phases and/or from

the need to maintain consistency through cycles. A different, yet related, question is whether, also in the cyclic Universe, this potential degree of arbitrariness impacts the predictions for the tensor spectrum or if they remain robust under the choice of the vacuum state.

Here, we take a first step forward in the discussion and, considering a non-BD vacuum state in the dark energy phase of the previous cycle, we argue that this initial state imposes a more fundamental constraint on the cyclic model, primarily due to the backreaction effects that a non-BD state could have on the background evolution of the modulus field.

To prove this point, following refs. [141, 162, 163], we allow the second Bogoliubov coefficient  $\beta_k$  to be non-zero. Notice that, since the Bogoliubov coefficient is related to  $D_2$  by

$$\beta_k = 2D_2\sqrt{(k/\pi)}, \quad (4.1)$$

and parameterizes deviations from the BD vacuum. As the effective theory becomes invalid at energy scales beyond those of new physics, we ensure that no modes are excited past this point, therefore requiring  $\beta_k \rightarrow 0$  for  $k > Ma(\tau_c)$ . In addition, we choose  $\beta_k$  of the form

$$\beta_k \sim \beta_0 e^{-\frac{k^2}{M^2 a(\tau_c)^2}}, \quad (4.2)$$

where the energy scale of the effective theory is  $M \sim 10^{-4}M_{\text{Pl}}$  and  $\beta_0$  is a proportionality factor. The effects produced by the backreaction in the dark energy-dominated phase set stringent limits on how large the non-vanishing  $\beta_k$  can be. In particular, requiring that the energy density of the non-BD quanta does not overwhelm the energy density associated with the modulus field — hence spoiling the model — implies

$$\rho_{\text{non-BD}} \sim \frac{1}{a_i^4} \int \frac{d^3k}{(2\pi)^3} |\beta_0|^2 k \sim \left(\frac{a(\tau_c)}{a_i(\tau)}\right)^4 |\beta_0|^2 M^4 \ll M_{\text{Pl}}^2 H^2, \quad (4.3)$$

where  $a_i$  refers to the value of the scale factor when we set our initial vacuum conditions, and we have used  $M_{\text{Pl}}^2 H^2$  as the energy density associated with the background evolution. We take the cutoff time  $\tau_c$ , beyond which we cannot be certain of the modulus field dynamics to be the start of radiation domination regime in the previous cycle  $\tau_{r(pc)}$ . As long as  $(a(\tau_{r(pc)})/a(\tau))^4 \ll 1$ , eq. (4.3) sets an upper bound for the parameter  $\beta_0$ :

$$\beta_0 < \frac{HM_{\text{Pl}}}{M^2} \sim 10^{-53} \quad (4.4)$$

where we used that, during the dark energy phase, the Hubble parameter is approximately constant  $H \simeq H_0 = 100 h \simeq 2.1h \times 10^{-42}$  GeV. Notice that this bound is independent of the cutoff time  $\tau_c$  when starting the calculation in the dark energy phase. This is because of the three expansion phases preceding this epoch and the large net expansion of the scale factor from cycle to cycle, ensuring that  $(a(\tau_c)/a(\tau_i))^4 \ll 1$  for any  $\tau_c$ .

We emphasise once again that avoiding problems with backreaction during the dark energy-dominated phase leads to an extremely restrictive constraint on deviations away from the BD vacuum. To gain a rough idea of how restrictive our bound is, we can compare constraints on the same cutoff scale obtained in inflationary cosmology. In that case, as highlighted in ref. [163], the restriction reads  $\beta_0 < 10^{-6}$ ; i.e., about 47 orders of magnitude

weaker than our bound. The reason for such a large difference lies in the fact that, although inflation and dark energy share several common aspects in terms of background dynamics, these two phases span energy scales that differ by over 100 orders of magnitude in characteristic energy density. The energy scale of inflation is much higher, making it way more challenging for perturbations to overwhelm the dynamics and allowing larger freedom for deviations away from BD.

To further appreciate the strength of our constraint, it is also worth briefly discussing what happens when setting the vacuum state in a different phase of the model rather than in the dark energy-dominated one. In particular, we focus on the *ekpyrotic* phase, which, as highlighted multiple times in this work, is the phase where initial conditions are typically fixed, and the calculation of the spectrum of tensor perturbations is initiated. In this phase, the Hubble parameter drastically grows, see eq. (2.15), while the scale factor remains approximately constant. As a consequence, the constraint on  $\beta_0$  is relaxed up to  $\beta_0 < 10^{45}$  for  $M \sim 10^{-4} M_{\text{Pl}}$ , allowing us complete freedom to choose a wide range of vacuum states in the *ekpyrotic* phase without compromising the model during this stage. That being said, it is necessary to ensure that significant deviations away from a BD vacuum during the *ekpyrotic* phase do not lead to other issues during the subsequent evolutionary states of the cycle. However, given that the evolution of the dark energy phase is governed by the background dynamics of the lowest energy scale and considering the cyclic nature of the model and its resilience through cycles (meaning that starting from the dark energy phase of the previous cycle does not produce differences in observable quantities as we prove in the next subsection), we argue that selecting the vacuum during the dark energy phase is the most restrictive and conservative choice to circumvent this issue entirely. This makes relying on the BD vacuum a safer assumption from a model-building perspective.

## 4.2 Gravitational waves produced from different phases

Our general calculation for the strain spectrum of gravitational waves enables us to investigate how the predictions change when incorporating the contribution from tensor modes originating in the dark energy phase of the previous cycle. We can then test the robustness of these predictions by comparing our results with those already documented in the literature, derived from starting in the *ekpyrotic* contracting phase.


In this section, we compare the predictions for the strain spectrum of gravitational waves obtained in the following two cases:

- (a) Using our general calculation and starting in the dark energy phase of the previous cycle. In this case we impose BD vacuum conditions on the coefficients of the dark energy stage  $D_1$  and  $D_2$  given by:

$$D_1 = \frac{1}{2} \sqrt{\frac{\pi}{k}}, \quad D_2 = 0. \quad (4.5)$$

- (b) Starting the calculation in the *ekpyrotic* phase (disregarding the matching at  $\tau = \tau_{tr}$ ) and imposing BD initial conditions as done in Res. [50]

$$A_1 = \frac{1}{2} \sqrt{\frac{\pi}{k}}, \quad A_2 = 0. \quad (4.6)$$




**Figure 1.** Strain spectrum plotted against comoving frequency  $k$ , of gravitational waves produced starting the cycle with the dark energy phase in blue, and with the *ekpyrotic* phase in orange. BD initial conditions are assumed for both spectra.  $k_0, k_{eq}, k_r$  and  $k_{end}$  are the comoving frequencies on the horizon today, at matter-radiation equivalence, at the start of radiation domination, and at the *ekpyrotic*-kinetic transition respectively.

Notice that, although our calculation is fully generic concerning the choice of the vacuum state, as we proved in the previous subsection, deviations away from the BD vacuum state in the dark energy phase are strongly constrained, providing us with a valid physical reason to impose this vacuum state in the dark energy phase for the case (a). Instead, for the case (b), we impose the BD initial conditions in the *ekpyrotic* phase to work in the same framework as ref. [50] and allow direct comparison.

After matching the relevant phases, the strain spectra of tensor modes predicted in the two cases can be derived using eq. (3.15) and eq. (3.17) and are shown in figure 1 in blue for the case (a) and in orange for the case (b). As evident from the figure, a difference of up to an order of magnitude in the strain  $\Delta h$  is observed for modes  $k_0$  at the present-day horizon.

This difference can be understood by considering the evolution of modes produced during the dark energy phase of the previous cycle. To further clarify this point, we refer to figure 2 where we show the evolution of the comoving Hubble horizon,  $1/aH$ , throughout the different phases of the model. From the figure we note that the horizon at the end of the dark energy phase (i.e.  $k_{tr}^{-1}$ ) is several orders of magnitude greater than the present-day horizon (i.e.  $k_0^{-1}$ ) ensuring that none of these super horizon modes can have re-entered by today or in any subsequent cycle. Modes produced during this dark energy phase that exit the horizon become frozen and subsequently experience further stretching throughout this epoch.

On the other hand, sub-horizon modes produced in the same phase oscillate with decaying amplitude  $h \propto a^{-1}$ . This feature can be shown by solving eq. (3.3) in the dark energy background dynamics described by eq. (2.7). In view of that, we expect sub-horizon



**Figure 2.** Illustrative plot of the comoving Hubble horizon,  $1/aH$ , throughout the *ekpyrotic*, kinetic contracting, kinetic expanding, radiation domination, and finally matter domination phases in magenta, orange, dark blue, green and cyan respectively. Key modes on the horizon are illustrated as horizontal dashed lines, along with a label of their comoving wavenumber, and some less important modes are labelled.

modes produced deep within the dark energy phase (as well as in any previous phase of the previous cycle) to decay away to negligible amplitudes when compared to modes produced at the end of the same phase or during the subsequent *ekpyrotic* phase. Instead, sub-horizon modes produced *near* the end of the dark energy phase can survive into the *ekpyrotic* phase of the next cycle, acting as extra quanta in the vacuum initial conditions for the subsequent *ekpyrotic* phase. This contribution is encoded in the coefficient  $A_2$  in eq. (3.5) and can lead to observable effects on scales corresponding to the long wavelength portion of the strain spectrum. Referring back to figure 1, we can appreciate how the differences produced by amplitude oscillations expected from an under-damped simple harmonic oscillator solution affect only the range of frequencies between  $k_0$  and  $k_{eq}$  before becoming frozen in during the *ekpyrotic* phase.

Moving forward, as outlined in ref. [161], the subsequent phases will provide a red-tilt to the scale-invariant dark energy spectrum. Notice that for an *ekpyrotic* phase lasting  $N \sim 115$  e-folds, the magnitude of the coefficient  $A_2$  becomes  $\mathcal{O}(10^{10})$  between  $k_0$  and  $k_{eq}$  then rapidly decays to order  $\mathcal{O}(10^{-15})$  for modes around  $k_r$ . This explains the discrepancy between imposing BD initial conditions in the dark energy phase given by eq. (4.5), and in the *ekpyrotic* phase, given by eq. (4.6).

We conclude with a last important remark: as argued in section 4.1, to preserve the background evolution of the modulus field, it is important to have a BD-like vacuum state during the dark energy phase. This requirement implies that perturbations existing in the

current cycle must decay to negligible levels in amplitude compared to the energy density present in the BD vacuum, preventing them from acting as additional quanta on top of the vacuum. We emphasise that this condition is satisfied by the evolution of tensor perturbations after the bounce, particularly during the subsequent radiation and matter-dominated phases. In particular, the transfer function, eq. (3.17), ensures that the amplitudes of short-wavelength modes re-entering the horizon first (that are potentially the most problematic), are decreased by a factor  $(k_0/k)^2$ . Consequently, the amplitude of the shortest wavelength modes in the observable spectrum ( $k \sim k_{\text{end}}$ ), is suppressed by more than 20 orders of magnitude during the radiation and matter-dominated phases. As a result, the evolution through these phases guarantees the decaying amplitude of all modes in every cycle preceding each dark energy phase and restoring the vacuum to a BD state in the latter. This shields the dark energy phase from possible backreaction effects, underscoring the resilience of the model and demonstrating once again that, from a theoretical standpoint, fixing initial conditions in this phase is much safer from a model-building perspective.

## 5 Conclusion

In this study, we investigate the production of gravitational waves in a cyclic Universe, focusing on certain conceptual aspects of the theory, such as the resilience of observable predictions against the phase of the cycle in which initial conditions are set and the choice of the vacuum state.

In most of the analyses carried out in the literature, the *ekpyrotic* phase is typically assumed as the starting point of the cycle where initial conditions of primordial perturbations are imposed, and the calculations of the relative spectra begin. While for scalar perturbations the implications of setting the initial conditions in different phases of the theory have been examined in a few studies surrounding this topic, to the best of our knowledge, the calculation of the tensor spectrum has always been performed starting in the *ekpyrotic* phase, assuming a BD vacuum state and neglecting any potential contributions arising from tensor modes originating during the dark energy phase of the previous cycle.

In light of this, a few (we believe) interesting questions remained somewhat pending. For instance, one might wonder whether setting initial conditions in the dark energy phase of the previous cycle and considering the potential contribution of tensor modes generated in this phase could lead to any observable differences in the gravitational wave strain spectrum observed in the current cycle and eventually clarify why (not). Most importantly, in effective field theory descriptions of the cyclic Universe, the choice of the vacuum state may involve integrating out high-energy degrees of freedom, focusing on low-energy behaviors of the modulus field. This makes the choice far from trivial when considering the possibility of new physics acting at sufficiently high characteristic energy scales. Consequently, one might wonder about the implications of assuming a BD vacuum state in the *ekpyrotic* phase and, more broadly, what freedom exists in the cyclic Universe regarding the choice of the vacuum state and how such freedom affects the predictions for the spectrum of gravitational waves.

Fuelled by these questions, we consider a cyclic Universe described by the effective 4D Lagrangian (2.1) with a potential given by eq. (2.4). After reviewing the background dynamics of the model (section 2), we focus on the production and evolution of tensor modes. In

section 3, we analytically solve the equation of motion of the gravitational wave field through all the different phases of the cycle by starting from the dark energy phase of the previous cycle and making no assumptions about the vacuum state. The results presented in this section generalise the predictions for the tensor spectrum to include the contribution of tensor perturbations produced during the dark energy phase of the previous cycle and apply to any choice of the vacuum state of the theory. Therefore, they extend the treatment presented so far in the literature, allowing us to gain important insights into the issues posed earlier.

As argued in section 4, our findings reveal a significant resilience of the cyclic Universe model concerning predictions for the spectrum of primordial gravitational waves, with the most relevant results reading as follows.

- *Initial Conditions:* to quantify the impact of the contribution arising from tensor modes produced in the previous dark energy phase of the cycle, in section 4.2 we compared the spectra obtained by starting from the *ekpyrotic* contracting phase (in orange in figure 1) and the dark energy phase of the previous cycle (in blue in figure 1), assuming in both cases a BD vacuum. The two spectra are essentially identical except for a difference up to an order of magnitude in the strain  $\Delta h$  for modes on the scale  $k_0$  at the present-day horizon. The reason is that the horizon at the end of the dark energy phase is several orders of magnitude greater than the present-day horizon (see also figure 2), implying that none of these super-horizon modes can have re-entered by today or in any subsequent cycle. On the other hand, sub-horizon modes produced in the same phase oscillate with decaying amplitude  $h \propto a^{-1}$ . Therefore, sub-horizon modes produced deep within the dark energy phase (as well as in any previous phase of the previous cycle) decay away to negligible amplitudes, while sub-horizon modes produced near the end of the dark energy phase can survive into the *ekpyrotic* phase of the next cycle, being responsible for the small differences in the strain spectrum observed around  $k_0$ .
- *Vacuum state:* although including tensor modes originating from the previous dark energy phase does not lead to significant differences in the spectrum of primordial perturbations, we argue that starting the calculation in this phase imposes a more fundamental constraint on the choice of the vacuum state of the model. To highlight this aspect, in section 4.1 we consider a non-BD vacuum state in the dark energy phase parametrized in terms of the second Bogoliubov coefficient given by eq. (4.2). We show that requiring the energy density of the non-BD quanta not to overwhelm the energy density associated with the modulus field — hence, potentially spoiling the model — implies an extremely restrictive constraint on deviations away from the BD vacuum. On the other hand, for the same non-BD vacuum state, the constraint is completely lost in the *ekpyrotic* phase where we are left with complete freedom to choose a wide range of vacuum states without compromising the model during this stage (but possibly spoiling the subsequent evolution). As a result, selecting the vacuum during the dark energy phase (where we are basically forced to BD) seems to be the most restrictive and conservative choice, as well as a safer assumption from a model-building perspective. Given the cyclic nature of the model and the resilience of its predictions against the phase where we set initial conditions (as proven in the previous point), this strongly reduces our freedom to consider exotic vacuum states in the cyclic Universe.

## Acknowledgments

WG and CvdB are supported by the Lancaster-Sheffield Consortium for Fundamental Physics under STFC grant: ST/X000621/1.

## References

- [1] A.H. Guth, *The inflationary universe: a possible solution to the horizon and flatness problems*, *Phys. Rev. D* **23** (1981) 347 [INSPIRE].
- [2] A.D. Linde, *A new inflationary universe scenario: a possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems*, *Phys. Lett. B* **108** (1982) 389 [INSPIRE].
- [3] A. Albrecht and P.J. Steinhardt, *Cosmology for grand unified theories with radiatively induced symmetry breaking*, *Phys. Rev. Lett.* **48** (1982) 1220 [INSPIRE].
- [4] A. Vilenkin, *The birth of inflationary universes*, *Phys. Rev. D* **27** (1983) 2848 [INSPIRE].
- [5] PLANCK collaboration, *Planck 2018 results. X. Constraints on inflation*, *Astron. Astrophys.* **641** (2020) A10 [arXiv:1807.06211] [INSPIRE].
- [6] V.F. Mukhanov and G.V. Chibisov, *Quantum fluctuations and a nonsingular universe*, *JETP Lett.* **33** (1981) 532 [INSPIRE].
- [7] J.M. Bardeen, P.J. Steinhardt and M.S. Turner, *Spontaneous creation of almost scale-free density perturbations in an inflationary universe*, *Phys. Rev. D* **28** (1983) 679 [INSPIRE].
- [8] S.W. Hawking, *The development of irregularities in a single bubble inflationary universe*, *Phys. Lett. B* **115** (1982) 295 [INSPIRE].
- [9] A.H. Guth and S.Y. Pi, *Fluctuations in the new inflationary universe*, *Phys. Rev. Lett.* **49** (1982) 1110 [INSPIRE].
- [10] J.E. Lidsey et al., *Reconstructing the inflation potential: an overview*, *Rev. Mod. Phys.* **69** (1997) 373 [astro-ph/9508078] [INSPIRE].
- [11] D.H. Lyth and A. Riotto, *Particle physics models of inflation and the cosmological density perturbation*, *Phys. Rept.* **314** (1999) 1 [hep-ph/9807278] [INSPIRE].
- [12] D. Baumann, *Inflation*, in the proceedings of the *Theoretical Advanced Study Institute in Elementary Particle Physics: physics of the large and the small*, (2011) [DOI:10.1142/9789814327183\_0010] [arXiv:0907.5424] [INSPIRE].
- [13] J. Martin, C. Ringeval and V. Vennin, *Encyclopædia inflationaris*, *Phys. Dark Univ.* **5-6** (2014) 75 [arXiv:1303.3787] [INSPIRE].
- [14] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [Erratum *ibid.* **652** (2021) C4] [arXiv:1807.06209] [INSPIRE].
- [15] E.R. Harrison, *Fluctuations at the threshold of classical cosmology*, *Phys. Rev. D* **1** (1970) 2726 [INSPIRE].
- [16] Y.B. Zeldovich, *A hypothesis, unifying the structure and the entropy of the universe*, *Mon. Not. Roy. Astron. Soc.* **160** (1972) 1P [INSPIRE].
- [17] P.J.E. Peebles and J.T. Yu, *Primeval adiabatic perturbation in an expanding universe*, *Astrophys. J.* **162** (1970) 815 [INSPIRE].

- [18] WMAP collaboration, *Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: final maps and results*, *Astrophys. J. Suppl.* **208** (2013) 20 [[arXiv:1212.5225](#)] [[INSPIRE](#)].
- [19] WMAP collaboration, *Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: cosmological parameter results*, *Astrophys. J. Suppl.* **208** (2013) 19 [[arXiv:1212.5226](#)] [[INSPIRE](#)].
- [20] ACT collaboration, *The Atacama Cosmology Telescope: a measurement of the cosmic microwave background power spectra at 98 and 150 GHz*, *JCAP* **12** (2020) 045 [[arXiv:2007.07289](#)] [[INSPIRE](#)].
- [21] ACT collaboration, *The Atacama Cosmology Telescope: DR4 maps and cosmological parameters*, *JCAP* **12** (2020) 047 [[arXiv:2007.07288](#)] [[INSPIRE](#)].
- [22] SPT-3G collaboration, *SPT-3G: a next-generation cosmic microwave background polarization experiment on the South Pole Telescope*, *Proc. SPIE Int. Soc. Opt. Eng.* **9153** (2014) 91531P [[arXiv:1407.2973](#)] [[INSPIRE](#)].
- [23] SPT-3G collaboration, *Measurements of the E-mode polarization and temperature-E-mode correlation of the CMB from SPT-3G 2018 data*, *Phys. Rev. D* **104** (2021) 022003 [[arXiv:2101.01684](#)] [[INSPIRE](#)].
- [24] W. Giarè et al., *Is the Harrison-Zel'dovich spectrum coming back? ACT preference for  $n_s \sim 1$  and its discordance with Planck*, *Mon. Not. Roy. Astron. Soc.* **521** (2023) 2911 [[arXiv:2210.09018](#)] [[INSPIRE](#)].
- [25] M. Forconi, W. Giarè, E. Di Valentino and A. Melchiorri, *Cosmological constraints on slow roll inflation: an update*, *Phys. Rev. D* **104** (2021) 103528 [[arXiv:2110.01695](#)] [[INSPIRE](#)].
- [26] W. Giarè et al., *Inflationary potential as seen from different angles: model compatibility from multiple CMB missions*, *JCAP* **09** (2023) 019 [[arXiv:2305.15378](#)] [[INSPIRE](#)].
- [27] J.L. Bernal, L. Verde and A.G. Riess, *The trouble with  $H_0$* , *JCAP* **10** (2016) 019 [[arXiv:1607.05617](#)] [[INSPIRE](#)].
- [28] L. Verde, T. Treu and A.G. Riess, *Tensions between the early and the late universe*, *Nature Astron.* **3** (2019) 891 [[arXiv:1907.10625](#)] [[INSPIRE](#)].
- [29] E. Di Valentino et al., *Snowmass2021 — letter of interest cosmology intertwined II: the Hubble constant tension*, *Astropart. Phys.* **131** (2021) 102605 [[arXiv:2008.11284](#)] [[INSPIRE](#)].
- [30] E. Di Valentino et al., *In the realm of the Hubble tension — a review of solutions*, *Class. Quant. Grav.* **38** (2021) 153001 [[arXiv:2103.01183](#)] [[INSPIRE](#)].
- [31] E. Abdalla et al., *Cosmology intertwined: a review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies*, *JHEAp* **34** (2022) 49 [[arXiv:2203.06142](#)] [[INSPIRE](#)].
- [32] E. Di Valentino, A. Melchiorri, Y. Fantaye and A. Heavens, *Bayesian evidence against the Harrison-Zel'dovich spectrum in tensions with cosmological data sets*, *Phys. Rev. D* **98** (2018) 063508 [[arXiv:1808.09201](#)] [[INSPIRE](#)].
- [33] G. Ye, B. Hu and Y.-S. Piao, *Implication of the Hubble tension for the primordial universe in light of recent cosmological data*, *Phys. Rev. D* **104** (2021) 063510 [[arXiv:2103.09729](#)] [[INSPIRE](#)].

- [34] G. Ye, J.-Q. Jiang and Y.-S. Piao, *Toward inflation with  $n_s = 1$  in light of the Hubble tension and implications for primordial gravitational waves*, *Phys. Rev. D* **106** (2022) 103528 [[arXiv:2205.02478](#)] [[INSPIRE](#)].
- [35] J.-Q. Jiang and Y.-S. Piao, *Toward early dark energy and  $n_s = 1$  with Planck, ACT, and SPT observations*, *Phys. Rev. D* **105** (2022) 103514 [[arXiv:2202.13379](#)] [[INSPIRE](#)].
- [36] J.-Q. Jiang, G. Ye and Y.-S. Piao, *Return of Harrison-Zel'dovich spectrum in light of recent cosmological tensions*, *Mon. Not. Roy. Astron. Soc.* **527** (2023) L54 [[arXiv:2210.06125](#)] [[INSPIRE](#)].
- [37] F. Takahashi and W. Yin, *Cosmological implications of  $n_s \approx 1$  in light of the Hubble tension*, *Phys. Lett. B* **830** (2022) 137143 [[arXiv:2112.06710](#)] [[INSPIRE](#)].
- [38] C.-M. Lin, *D-term inflation in braneworld models: consistency with cosmic-string bounds and early-time Hubble tension resolving models*, *Phys. Rev. D* **106** (2022) 103511 [[arXiv:2204.10475](#)] [[INSPIRE](#)].
- [39] D.K. Hazra, A. Antony and A. Shafieloo, *One spectrum to cure them all: signature from early universe solves major anomalies and tensions in cosmology*, *JCAP* **08** (2022) 063 [[arXiv:2201.12000](#)] [[INSPIRE](#)].
- [40] M. Braglia, X. Chen and D.K. Hazra, *Uncovering the history of cosmic inflation from anomalies in cosmic microwave background spectra*, *Eur. Phys. J. C* **82** (2022) 498 [[arXiv:2106.07546](#)] [[INSPIRE](#)].
- [41] R.E. Keeley, A. Shafieloo, D.K. Hazra and T. Souradeep, *Inflation wars: a new hope*, *JCAP* **09** (2020) 055 [[arXiv:2006.12710](#)] [[INSPIRE](#)].
- [42] J.-Q. Jiang, G. Ye and Y.-S. Piao, *Impact of the Hubble tension on the  $r$ - $n_s$  contour*, *Phys. Lett. B* **851** (2024) 138588 [[arXiv:2303.12345](#)] [[INSPIRE](#)].
- [43] P.J. Steinhardt and N. Turok, *Cosmic evolution in a cyclic universe*, *Phys. Rev. D* **65** (2002) 126003 [[hep-th/0111098](#)] [[INSPIRE](#)].
- [44] P.J. Steinhardt, N. Turok and N. Turok, *A cyclic model of the universe*, *Science* **296** (2002) 1436 [[hep-th/0111030](#)] [[INSPIRE](#)].
- [45] P.J. Steinhardt and N. Turok, *The cyclic universe: an informal introduction*, *Nucl. Phys. B Proc. Suppl.* **124** (2003) 38 [[astro-ph/0204479](#)] [[INSPIRE](#)].
- [46] J. Khoury, P.J. Steinhardt and N. Turok, *Designing cyclic universe models*, *Phys. Rev. Lett.* **92** (2004) 031302 [[hep-th/0307132](#)] [[INSPIRE](#)].
- [47] N. Turok and P.J. Steinhardt, *Beyond inflation: a cyclic universe scenario*, *Phys. Scripta T* **117** (2005) 76 [[hep-th/0403020](#)] [[INSPIRE](#)].
- [48] J. Khoury, *A briefing on the ekpyrotic/cyclic universe*, in the proceedings of the 6<sup>th</sup> RESCEU international symposium on frontier in astroparticle physics and cosmology, (2004) [[astro-ph/0401579](#)] [[INSPIRE](#)].
- [49] J.-L. Lehners, *Ekpyrotic and cyclic cosmology*, *Phys. Rept.* **465** (2008) 223 [[arXiv:0806.1245](#)] [[INSPIRE](#)].
- [50] L.A. Boyle, P.J. Steinhardt and N. Turok, *The cosmic gravitational wave background in a cyclic universe*, *Phys. Rev. D* **69** (2004) 127302 [[hep-th/0307170](#)] [[INSPIRE](#)].
- [51] A. Ashtekar, M. Bojowald and J. Lewandowski, *Mathematical structure of loop quantum cosmology*, *Adv. Theor. Math. Phys.* **7** (2003) 233 [[gr-qc/0304074](#)] [[INSPIRE](#)].

- [52] M. Bojowald, R. Maartens and P. Singh, *Loop quantum gravity and the cyclic universe*, *Phys. Rev. D* **70** (2004) 083517 [[hep-th/0407115](#)] [[INSPIRE](#)].
- [53] H.-H. Xiong, T. Qiu, Y.-F. Cai and X. Zhang, *Cyclic universe with quintom matter in loop quantum cosmology*, *Mod. Phys. Lett. A* **24** (2009) 1237 [[arXiv:0711.4469](#)] [[INSPIRE](#)].
- [54] P.H. Frampton, *Cyclic universe and infinite past*, *Mod. Phys. Lett. A* **22** (2007) 2587 [[arXiv:0705.2730](#)] [[INSPIRE](#)].
- [55] J.V. Narlikar, G. Burbidge and R.G. Vishwakarma, *Cosmology and cosmogony in a cyclic universe*, *J. Astrophys. Astron.* **28** (2007) 67 [[arXiv:0801.2965](#)] [[INSPIRE](#)].
- [56] L. Baum and P.H. Frampton, *Entropy of contracting universe in cyclic cosmology*, *Mod. Phys. Lett. A* **23** (2008) 33 [[hep-th/0703162](#)] [[INSPIRE](#)].
- [57] T. Biswas, *The Hagedorn soup and an emergent cyclic universe*, [arXiv:0801.1315](#) [[INSPIRE](#)].
- [58] T. Cailleteau, P. Singh and K. Vandersloot, *Non-singular ekpyrotic/cyclic model in loop quantum cosmology*, *Phys. Rev. D* **80** (2009) 124013 [[arXiv:0907.5591](#)] [[INSPIRE](#)].
- [59] R.H. Brandenberger, *Processing of cosmological perturbations in a cyclic cosmology*, *Phys. Rev. D* **80** (2009) 023535 [[arXiv:0905.1514](#)] [[INSPIRE](#)].
- [60] A.R. El-Nabulsi, *Accelerated dilatonic-Brans-Dicke cyclic and non-singular universe from string theory*, *Res. Astron. Astrophys.* **11** (2011) 1249 [[INSPIRE](#)].
- [61] Y.-F. Cai, C. Gao and E.N. Saridakis, *Bounce and cyclic cosmology in extended nonlinear massive gravity*, *JCAP* **10** (2012) 048 [[arXiv:1207.3786](#)] [[INSPIRE](#)].
- [62] Y.-F. Cai and E.N. Saridakis, *Cyclic cosmology from Lagrange-multiplier modified gravity*, *Class. Quant. Grav.* **28** (2011) 035010 [[arXiv:1007.3204](#)] [[INSPIRE](#)].
- [63] S. Nojiri, S.D. Odintsov and D. Saez-Gomez, *Cyclic, ekpyrotic and little rip universe in modified gravity*, *AIP Conf. Proc.* **1458** (2012) 207 [[arXiv:1108.0767](#)] [[INSPIRE](#)].
- [64] H.-Y. Chang and R.J. Scherrer, *Coincidence problem in cyclic phantom models of the universe*, *Phys. Rev. D* **86** (2012) 027303 [[arXiv:1204.6329](#)] [[INSPIRE](#)].
- [65] R.I. Ivanov and E.M. Prodanov, *Cyclic universe with an inflationary phase from a cosmological model with real gas quintessence*, *Phys. Rev. D* **86** (2012) 083536 [[arXiv:1210.0186](#)] [[INSPIRE](#)].
- [66] K. Saaidi, H. Sheikahmadi and A.H. Mohammadi, *Interacting new agegraphic dark energy in a cyclic universe*, *Astrophys. Space Sci.* **338** (2012) 355 [[arXiv:1201.0275](#)] [[INSPIRE](#)].
- [67] I. Bars, P.J. Steinhardt and N. Turok, *Cyclic cosmology, conformal symmetry and the metastability of the Higgs*, *Phys. Lett. B* **726** (2013) 50 [[arXiv:1307.8106](#)] [[INSPIRE](#)].
- [68] Y. Tavakoli and J.C. Fabris, *Creation of particles in a cyclic universe driven by loop quantum cosmology*, *Int. J. Mod. Phys. D* **24** (2015) 1550062 [[arXiv:1412.0775](#)] [[INSPIRE](#)].
- [69] D. Oriti, L. Sindoni and E. Wilson-Ewing, *Emergent Friedmann dynamics with a quantum bounce from quantum gravity condensates*, *Class. Quant. Grav.* **33** (2016) 224001 [[arXiv:1602.05881](#)] [[INSPIRE](#)].
- [70] M. de Cesare, A.G.A. Pithis and M. Sakellariadou, *Cosmological implications of interacting group field theory models: cyclic universe and accelerated expansion*, *Phys. Rev. D* **94** (2016) 064051 [[arXiv:1606.00352](#)] [[INSPIRE](#)].
- [71] P. Pavlovic and M. Sossich, *Cyclic cosmology in modified gravity*, *Phys. Rev. D* **95** (2017) 103519 [[arXiv:1701.03657](#)] [[INSPIRE](#)].

- [72] E.N. Saridakis, S. Banerjee and R. Myrzakulov, *Bounce and cyclic cosmology in new gravitational scalar-tensor theories*, *Phys. Rev. D* **98** (2018) 063513 [[arXiv:1807.00346](#)] [[INSPIRE](#)].
- [73] P. Das, S. Pan, S. Ghosh and P. Pal, *Cosmological time crystal: cyclic universe with a small cosmological constant in a toy model approach*, *Phys. Rev. D* **98** (2018) 024004 [[arXiv:1801.07970](#)] [[INSPIRE](#)].
- [74] A. Ijjas, *Cyclic completion of the anamorphic universe*, *Class. Quant. Grav.* **35** (2018) 075010 [[arXiv:1610.02752](#)] [[INSPIRE](#)].
- [75] N. Ahmed and S.Z. Alamri, *A cyclic universe with varying cosmological constant in  $f(R,T)$  gravity*, *Can. J. Phys.* **97** (2019) 1075 [[arXiv:1902.03104](#)] [[INSPIRE](#)].
- [76] S.-L. Li et al., *Emergent universe scenario, bouncing universes, and cyclic universes in degenerate massive gravity*, *Phys. Rev. D* **99** (2019) 104057 [[arXiv:1903.03940](#)] [[INSPIRE](#)].
- [77] R.J. Scherrer, *The coincidence problem and the swampland conjectures in the Ijjas-Steinhardt cyclic model of the universe*, *Phys. Lett. B* **798** (2019) 134981 [[arXiv:1907.11293](#)] [[INSPIRE](#)].
- [78] G. Barca, P. Di Antonio, G. Montani and A. Patti, *Semiclassical and quantum polymer effects in a flat isotropic universe*, *Phys. Rev. D* **99** (2019) 123509 [[arXiv:1902.02128](#)] [[INSPIRE](#)].
- [79] A. Ijjas and P.J. Steinhardt, *Entropy, black holes, and the new cyclic universe*, *Phys. Lett. B* **824** (2022) 136823 [[arXiv:2108.07101](#)] [[INSPIRE](#)].
- [80] N. Gorkavyi, *Gravitational wave background discovered by NANOGrav as evidence of a cyclic universe*, *New Astron.* **91** (2022) 101698 [[arXiv:2110.10218](#)] [[INSPIRE](#)].
- [81] M. Martín-Benito, R.B. Neves and J. Olmedo, *States of low energy in bouncing inflationary scenarios in loop quantum cosmology*, *Phys. Rev. D* **103** (2021) 123524 [[arXiv:2104.03035](#)] [[INSPIRE](#)].
- [82] A. Calcinari and S. Gielen, *Towards anisotropic cosmology in group field theory*, *Class. Quant. Grav.* **40** (2023) 085004 [[arXiv:2210.03149](#)] [[INSPIRE](#)].
- [83] E. Giovannetti and G. Montani, *Is Bianchi I a bouncing cosmology in the Wheeler-DeWitt picture?*, *Phys. Rev. D* **106** (2022) 044053 [[arXiv:2203.01062](#)] [[INSPIRE](#)].
- [84] E. Giovannetti, F. Maione and G. Montani, *Quantum big bounce of the isotropic universe using relational time*, *Universe* **9** (2023) 373 [[arXiv:2301.06149](#)] [[INSPIRE](#)].
- [85] D. Battfeld and P. Peter, *A critical review of classical bouncing cosmologies*, *Phys. Rept.* **571** (2015) 1 [[arXiv:1406.2790](#)] [[INSPIRE](#)].
- [86] R. Brandenberger and P. Peter, *Bouncing cosmologies: progress and problems*, *Found. Phys.* **47** (2017) 797 [[arXiv:1603.05834](#)] [[INSPIRE](#)].
- [87] J. Khoury, B.A. Ovrut, P.J. Steinhardt and N. Turok, *Density perturbations in the ekpyrotic scenario*, *Phys. Rev. D* **66** (2002) 046005 [[hep-th/0109050](#)] [[INSPIRE](#)].
- [88] J.-L. Lehners, P. McFadden, N. Turok and P.J. Steinhardt, *Generating ekpyrotic curvature perturbations before the big bang*, *Phys. Rev. D* **76** (2007) 103501 [[hep-th/0702153](#)] [[INSPIRE](#)].
- [89] E.I. Buchbinder, J. Khoury and B.A. Ovrut, *On the initial conditions in new ekpyrotic cosmology*, *JHEP* **11** (2007) 076 [[arXiv:0706.3903](#)] [[INSPIRE](#)].
- [90] J.-L. Lehners and P.J. Steinhardt, *Planck 2013 results support the cyclic universe*, *Phys. Rev. D* **87** (2013) 123533 [[arXiv:1304.3122](#)] [[INSPIRE](#)].

- [91] C.-G. Park and B. Ratra, *Using the tilted flat- $\Lambda$ CDM and the untilted non-flat  $\Lambda$ CDM inflation models to measure cosmological parameters from a compilation of observational data*, *Astrophys. J.* **882** (2019) 158 [[arXiv:1801.00213](#)] [[INSPIRE](#)].
- [92] W. Handley, *Curvature tension: evidence for a closed universe*, *Phys. Rev. D* **103** (2021) L041301 [[arXiv:1908.09139](#)] [[INSPIRE](#)].
- [93] E. Di Valentino, A. Melchiorri and J. Silk, *Planck evidence for a closed universe and a possible crisis for cosmology*, *Nature Astron.* **4** (2019) 196 [[arXiv:1911.02087](#)] [[INSPIRE](#)].
- [94] G. Efstathiou and S. Gratton, *The evidence for a spatially flat universe*, *Mon. Not. Roy. Astron. Soc.* **496** (2020) L91 [[arXiv:2002.06892](#)] [[INSPIRE](#)].
- [95] E. Di Valentino, A. Melchiorri and J. Silk, *Investigating cosmic discordance*, *Astrophys. J. Lett.* **908** (2021) L9 [[arXiv:2003.04935](#)] [[INSPIRE](#)].
- [96] D. Benisty and D. Staicova, *Testing late-time cosmic acceleration with uncorrelated baryon acoustic oscillation dataset*, *Astron. Astrophys.* **647** (2021) A38 [[arXiv:2009.10701](#)] [[INSPIRE](#)].
- [97] S. Vagnozzi et al., *The galaxy power spectrum take on spatial curvature and cosmic concordance*, *Phys. Dark Univ.* **33** (2021) 100851 [[arXiv:2010.02230](#)] [[INSPIRE](#)].
- [98] S. Vagnozzi, A. Loeb and M. Moresco, *Eppur è piatto? The cosmic chronometers take on spatial curvature and cosmic concordance*, *Astrophys. J.* **908** (2021) 84 [[arXiv:2011.11645](#)] [[INSPIRE](#)].
- [99] E. Di Valentino et al., *Interacting dark energy in a closed universe*, *Mon. Not. Roy. Astron. Soc.* **502** (2021) L23 [[arXiv:2011.00283](#)] [[INSPIRE](#)].
- [100] W. Yang et al., *2021- $H_0$  odyssey: closed, phantom and interacting dark energy cosmologies*, *JCAP* **10** (2021) 008 [[arXiv:2101.03129](#)] [[INSPIRE](#)].
- [101] S. Cao, J. Ryan and B. Ratra, *Using Pantheon and DES supernova, baryon acoustic oscillation, and Hubble parameter data to constrain the Hubble constant, dark energy dynamics, and spatial curvature*, *Mon. Not. Roy. Astron. Soc.* **504** (2021) 300 [[arXiv:2101.08817](#)] [[INSPIRE](#)].
- [102] S. Dhawan, J. Alsing and S. Vagnozzi, *Non-parametric spatial curvature inference using late-universe cosmological probes*, *Mon. Not. Roy. Astron. Soc.* **506** (2021) L1 [[arXiv:2104.02485](#)] [[INSPIRE](#)].
- [103] B.R. Dinda, *Cosmic expansion parametrization: implication for curvature and  $H_0$  tension*, *Phys. Rev. D* **105** (2022) 063524 [[arXiv:2106.02963](#)] [[INSPIRE](#)].
- [104] J.E. Gonzalez, M. Benetti, R. von Marttens and J. Alcaniz, *Testing the consistency between cosmological data: the impact of spatial curvature and the dark energy EoS*, *JCAP* **11** (2021) 060 [[arXiv:2104.13455](#)] [[INSPIRE](#)].
- [105] O. Akarsu et al., *Testing spatial curvature and anisotropic expansion on top of the  $\Lambda$ CDM model*, *Phys. Dark Univ.* **39** (2023) 101162 [[arXiv:2112.07807](#)] [[INSPIRE](#)].
- [106] S. Cao and B. Ratra, *Using lower redshift, non-CMB, data to constrain the Hubble constant and other cosmological parameters*, *Mon. Not. Roy. Astron. Soc.* **513** (2022) 5686 [[arXiv:2203.10825](#)] [[INSPIRE](#)].
- [107] A. Glanville, C. Howlett and T.M. Davis, *Full-shape galaxy power spectra and the curvature tension*, *Mon. Not. Roy. Astron. Soc.* **517** (2022) 3087 [[arXiv:2205.05892](#)] [[INSPIRE](#)].
- [108] J. Bel et al., *Constraining spatial curvature with large-scale structure*, *JCAP* **09** (2022) 076 [[arXiv:2206.03059](#)] [[INSPIRE](#)].
- [109] W. Yang et al., *Revealing the effects of curvature on the cosmological models*, *Phys. Rev. D* **107** (2023) 063509 [[arXiv:2210.09865](#)] [[INSPIRE](#)].

- [110] J. Stevens, H. Khoraminezhad and S. Saito, *Constraining the spatial curvature with cosmic expansion history in a cosmological model with a non-standard sound horizon*, *JCAP* **07** (2023) 046 [[arXiv:2212.09804](#)] [[INSPIRE](#)].
- [111] A. Favale, A. Gómez-Valent and M. Migliaccio, *Cosmic chronometers to calibrate the ladders and measure the curvature of the universe. A model-independent study*, *Mon. Not. Roy. Astron. Soc.* **523** (2023) 3406 [[arXiv:2301.09591](#)] [[INSPIRE](#)].
- [112] J. Khoury, P.J. Steinhardt and N. Turok, *Inflation versus cyclic predictions for spectral tilt*, *Phys. Rev. Lett.* **91** (2003) 161301 [[astro-ph/0302012](#)] [[INSPIRE](#)].
- [113] S. Gratton, J. Khoury, P.J. Steinhardt and N. Turok, *Conditions for generating scale-invariant density perturbations*, *Phys. Rev. D* **69** (2004) 103505 [[astro-ph/0301395](#)] [[INSPIRE](#)].
- [114] C. Caprini and D.G. Figueroa, *Cosmological backgrounds of gravitational waves*, *Class. Quant. Grav.* **35** (2018) 163001 [[arXiv:1801.04268](#)] [[INSPIRE](#)].
- [115] BICEP and KECK collaborations, *Improved constraints on primordial gravitational waves using Planck, WMAP, and BICEP/Keck observations through the 2018 observing season*, *Phys. Rev. Lett.* **127** (2021) 151301 [[arXiv:2110.00483](#)] [[INSPIRE](#)].
- [116] T. Kobayashi, M. Yamaguchi and J. Yokoyama, *G-inflation: inflation driven by the Galileon field*, *Phys. Rev. Lett.* **105** (2010) 231302 [[arXiv:1008.0603](#)] [[INSPIRE](#)].
- [117] M. Kawasaki, N. Kitajima and S. Yokoyama, *Gravitational waves from a curvaton model with blue spectrum*, *JCAP* **08** (2013) 042 [[arXiv:1305.4464](#)] [[INSPIRE](#)].
- [118] K. Nozari and S. Shafizadeh, *Realization of blue spectrum in generalized Galileon super-inflation models*, *Int. J. Mod. Phys. D* **26** (2016) 1750016 [[arXiv:1712.09530](#)] [[INSPIRE](#)].
- [119] W. Giarè and F. Renzi, *Propagating speed of primordial gravitational waves*, *Phys. Rev. D* **102** (2020) 083530 [[arXiv:2007.04256](#)] [[INSPIRE](#)].
- [120] D.I. Kaiser and E.I. Sfakianakis, *Multifield inflation after Planck: the case for nonminimal couplings*, *Phys. Rev. Lett.* **112** (2014) 011302 [[arXiv:1304.0363](#)] [[INSPIRE](#)].
- [121] L.C. Price, H.V. Peiris, J. Frazer and R. Easther, *Gravitational wave consistency relations for multifield inflation*, *Phys. Rev. Lett.* **114** (2015) 031301 [[arXiv:1409.2498](#)] [[INSPIRE](#)].
- [122] A. Achúcarro et al., *Features of heavy physics in the CMB power spectrum*, *JCAP* **01** (2011) 030 [[arXiv:1010.3693](#)] [[INSPIRE](#)].
- [123] M. De Angelis and C. van de Bruck, *Adiabatic and isocurvature perturbations in extended theories with kinetic couplings*, *JCAP* **10** (2023) 023 [[arXiv:2304.12364](#)] [[INSPIRE](#)].
- [124] W. Giarè, M. De Angelis, C. van de Bruck and E. Di Valentino, *Tracking the multifield dynamics with cosmological data: a Monte Carlo approach*, *JCAP* **12** (2023) 014 [[arXiv:2306.12414](#)] [[INSPIRE](#)].
- [125] S. Mukohyama, R. Namba, M. Peloso and G. Shiu, *Blue tensor spectrum from particle production during inflation*, *JCAP* **08** (2014) 036 [[arXiv:1405.0346](#)] [[INSPIRE](#)].
- [126] R. Namba et al., *Scale-dependent gravitational waves from a rolling axion*, *JCAP* **01** (2016) 041 [[arXiv:1509.07521](#)] [[INSPIRE](#)].
- [127] M. Peloso, L. Sorbo and C. Unal, *Rolling axions during inflation: perturbativity and signatures*, *JCAP* **09** (2016) 001 [[arXiv:1606.00459](#)] [[INSPIRE](#)].
- [128] O. Özsoy, *Synthetic gravitational waves from a rolling axion monodromy*, *JCAP* **04** (2021) 040 [[arXiv:2005.10280](#)] [[INSPIRE](#)].

- [129] E. Dimastrogiovanni, M. Fasiello and T. Fujita, *Primordial gravitational waves from axion-gauge fields dynamics*, *JCAP* **01** (2017) 019 [[arXiv:1608.04216](#)] [[INSPIRE](#)].
- [130] L. Iacconi et al., *Interferometer constraints on the inflationary field content*, *JCAP* **03** (2020) 031 [[arXiv:1910.12921](#)] [[INSPIRE](#)].
- [131] S. Endlich, A. Nicolis and J. Wang, *Solid inflation*, *JCAP* **10** (2013) 011 [[arXiv:1210.0569](#)] [[INSPIRE](#)].
- [132] D. Cannone, G. Tasinato and D. Wands, *Generalised tensor fluctuations and inflation*, *JCAP* **01** (2015) 029 [[arXiv:1409.6568](#)] [[INSPIRE](#)].
- [133] L. Graef and R. Brandenberger, *Breaking of spatial diffeomorphism invariance, inflation and the spectrum of cosmological perturbations*, *JCAP* **10** (2015) 009 [[arXiv:1506.00896](#)] [[INSPIRE](#)].
- [134] A. Ricciardone and G. Tasinato, *Primordial gravitational waves in supersolid inflation*, *Phys. Rev. D* **96** (2017) 023508 [[arXiv:1611.04516](#)] [[INSPIRE](#)].
- [135] D. Baumann, H. Lee and G.L. Pimentel, *High-scale inflation and the tensor tilt*, *JHEP* **01** (2016) 101 [[arXiv:1507.07250](#)] [[INSPIRE](#)].
- [136] W. Giarè, F. Renzi and A. Melchiorri, *Higher-curvature corrections and tensor modes*, *Phys. Rev. D* **103** (2021) 043515 [[arXiv:2012.00527](#)] [[INSPIRE](#)].
- [137] G. Capurri, N. Bartolo, D. Maino and S. Matarrese, *Let effective field theory of inflation flow: stochastic generation of models with red/blue tensor tilt*, *JCAP* **11** (2020) 037 [[arXiv:2006.10781](#)] [[INSPIRE](#)].
- [138] W. Giarè, M. Forconi, E. Di Valentino and A. Melchiorri, *Towards a reliable calculation of relic radiation from primordial gravitational waves*, *Mon. Not. Roy. Astron. Soc.* **520** (2023) 2 [[arXiv:2210.14159](#)] [[INSPIRE](#)].
- [139] Y. Cai, *Generating enhanced parity-violating gravitational waves during inflation with violation of the null energy condition*, *Phys. Rev. D* **107** (2023) 063512 [[arXiv:2212.10893](#)] [[INSPIRE](#)].
- [140] G. Ye, M. Zhu and Y. Cai, *Null energy condition violation during inflation and pulsar timing array observations*, *JHEP* **02** (2024) 008 [[arXiv:2312.10685](#)] [[INSPIRE](#)].
- [141] A. Ashoorioon, K. Dimopoulos, M.M. Sheikh-Jabbari and G. Shiu, *Reconciliation of high energy scale models of inflation with Planck*, *JCAP* **02** (2014) 025 [[arXiv:1306.4914](#)] [[INSPIRE](#)].
- [142] A. Ashoorioon, K. Dimopoulos, M.M. Sheikh-Jabbari and G. Shiu, *Non-Bunch-Davis initial state reconciles chaotic models with BICEP and Planck*, *Phys. Lett. B* **737** (2014) 98 [[arXiv:1403.6099](#)] [[INSPIRE](#)].
- [143] S. Choudhury, *Single field inflation in the light of pulsar timing array data: quintessential interpretation of blue tilted tensor spectrum through non-Bunch Davies initial condition*, *Eur. Phys. J. C* **84** (2024) 278 [[arXiv:2307.03249](#)] [[INSPIRE](#)].
- [144] Y.-F. Cai, C. Lin, B. Wang and S.-F. Yan, *Sound speed resonance of the stochastic gravitational wave background*, *Phys. Rev. Lett.* **126** (2021) 071303 [[arXiv:2009.09833](#)] [[INSPIRE](#)].
- [145] A. Gruzinov, *Elastic inflation*, *Phys. Rev. D* **70** (2004) 063518 [[astro-ph/0404548](#)] [[INSPIRE](#)].
- [146] A. Ashoorioon, J.L. Hovdebo and R.B. Mann, *Running of the spectral index and violation of the consistency relation between tensor and scalar spectra from trans-Planckian physics*, *Nucl. Phys. B* **727** (2005) 63 [[gr-qc/0504135](#)] [[INSPIRE](#)].
- [147] R.H. Brandenberger, A. Nayeri, S.P. Patil and C. Vafa, *Tensor modes from a primordial Hagedorn phase of string cosmology*, *Phys. Rev. Lett.* **98** (2007) 231302 [[hep-th/0604126](#)] [[INSPIRE](#)].

- [148] R.H. Brandenberger, A. Nayeri and S.P. Patil, *Closed string thermodynamics and a blue tensor spectrum*, *Phys. Rev. D* **90** (2014) 067301 [[arXiv:1403.4927](#)] [[INSPIRE](#)].
- [149] M. Baumgart, J.J. Heckman and L. Thomas, *CFTs blueshift tensor fluctuations universally*, *JCAP* **07** (2022) 034 [[arXiv:2109.08166](#)] [[INSPIRE](#)].
- [150] A. Stewart and R. Brandenberger, *Observational constraints on theories with a blue spectrum of tensor modes*, *JCAP* **08** (2008) 012 [[arXiv:0711.4602](#)] [[INSPIRE](#)].
- [151] Y.-F. Cai et al., *On the possibility of blue tensor spectrum within single field inflation*, *Nucl. Phys. B* **900** (2015) 517 [[arXiv:1412.7241](#)] [[INSPIRE](#)].
- [152] Y. Wang and W. Xue, *Inflation and alternatives with blue tensor spectra*, *JCAP* **10** (2014) 075 [[arXiv:1403.5817](#)] [[INSPIRE](#)].
- [153] S. Kuroyanagi, T. Takahashi and S. Yokoyama, *Blue-tilted tensor spectrum and thermal history of the universe*, *JCAP* **02** (2015) 003 [[arXiv:1407.4785](#)] [[INSPIRE](#)].
- [154] S. Kuroyanagi, T. Takahashi and S. Yokoyama, *Blue-tilted inflationary tensor spectrum and reheating in the light of NANOGraV results*, *JCAP* **01** (2021) 071 [[arXiv:2011.03323](#)] [[INSPIRE](#)].
- [155] W. Giarè and A. Melchiorri, *Probing the inflationary background of gravitational waves from large to small scales*, *Phys. Lett. B* **815** (2021) 136137 [[arXiv:2003.04783](#)] [[INSPIRE](#)].
- [156] S. Vagnozzi, *Implications of the NANOGraV results for inflation*, *Mon. Not. Roy. Astron. Soc.* **502** (2021) L11 [[arXiv:2009.13432](#)] [[INSPIRE](#)].
- [157] S. Vagnozzi, *Inflationary interpretation of the stochastic gravitational wave background signal detected by pulsar timing array experiments*, *JHEAp* **39** (2023) 81 [[arXiv:2306.16912](#)] [[INSPIRE](#)].
- [158] J.-Q. Jiang, Y. Cai, G. Ye and Y.-S. Piao, *Broken blue-tilted inflationary gravitational waves: a joint analysis of NANOGraV 15-year and BICEP/Keck 2018 data*, *JCAP* **05** (2024) 004 [[arXiv:2307.15547](#)] [[INSPIRE](#)].
- [159] V.K. Oikonomou, P. Tsyba and O. Razina, *Red or blue tensor spectrum from GW170817-compatible Einstein-Gauss-Bonnet theory: a detailed analysis*, *EPL* **145** (2024) 49001 [[arXiv:2402.02049](#)] [[INSPIRE](#)].
- [160] N.K. Stein and W.H. Kinney, *Simple single-field inflation models with arbitrarily small tensor/scalar ratio*, *JCAP* **03** (2023) 027 [[arXiv:2210.05757](#)] [[INSPIRE](#)].
- [161] J.K. Erickson, S. Gratton, P.J. Steinhardt and N. Turok, *Cosmic perturbations through the cyclic ages*, *Phys. Rev. D* **75** (2007) 123507 [[hep-th/0607164](#)] [[INSPIRE](#)].
- [162] A. Aravind, D. Lorshbough and S. Paban, *Non-Gaussianity from excited initial inflationary states*, *JHEP* **07** (2013) 076 [[arXiv:1303.1440](#)] [[INSPIRE](#)].
- [163] R. Holman and A.J. Tolley, *Enhanced non-Gaussianity from excited initial states*, *JCAP* **05** (2008) 001 [[arXiv:0710.1302](#)] [[INSPIRE](#)].