



This is a repository copy of *Informed decision making for the elderly patient with newly diagnosed coeliac disease*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/216218/>

Version: Published Version

Article:

Shiha, M.G. orcid.org/0000-0002-2713-8355, Marks, L.J. and Sanders, D.S. (2020) Informed decision making for the elderly patient with newly diagnosed coeliac disease. *Gastroenterology and Hepatology from Bed to Bench*, 13 (3). p. 273. ISSN 2008-2258

<https://doi.org/10.22037/ghfbb.v13i3.1989>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Informed decision making for the elderly patient with newly diagnosed coeliac disease

Mohamed G Shiha, Lauren J Marks, David S. Sanders

Academic Unit of Gastroenterology, Royal Hallamshire Hospital, Sheffield, United Kingdom

(Please cite as: **Shiha MG, Marks LJ, Sanders DS. Informed decision making for the elderly patient with newly diagnosed coeliac disease. Gastroenterol Hepatol Bed Bench 2020;13(3):273).**

To The Editor

We thank Dr. W. Shanahan and Dr. J. Shanahan for their interest in our study (1). In their letter, they have raised a few important points that we wish to clarify.

Firstly, they implied that we had a lack of appreciation for the recognition and prevention of bone-related morbidity in elderly Coeliac Disease (CD) patients. We respectfully disagree with their statement; throughout our paper, we repeatedly highlighted the importance of recognising such complications in this group of patients. Moreover, we reported that elderly CD patients had a significantly higher risk of osteopenia/osteoporosis compared to younger patients. In our practice, we routinely measure serum vitamin D, alkaline phosphatase and calcium levels for all CD patients and refer those older than 55 years or with known osteoporotic risk factors for bone density scans following the British Society of Gastroenterology guidelines. (2) A single centre audit, while valuable in improving local practices, should not be used to generalise poor adherence to national guidelines.

Secondly, we agree that CD is associated with an increased risk of osteoporosis and fractures. However, GFD adherence rarely reverses bone loss in adults. This is particularly evident in elderly patients with long-term exposure to gluten and established bone disease at the time of diagnosis (3, 4). In our study, 57% of the osteopenic/osteoporotic elderly patients with CD were completely asymptomatic. Therefore, without strong evidence from randomised controlled trials or large prospective studies showing a clear benefit from adhering to a strict GFD over nutritional supplementation, only in this age group, we have to

raise the question whether it is worthwhile breaking lifetime dietary habits in asymptomatic elderly patients. Finally, should we be diagnosing Coeliac Disease in the elderly? The answer is probably yes. However, the majority of these elderly patients show little to no gastrointestinal symptoms and are often referred due to cancer or metabolic pathways with unexplained anaemia and bone loss. Hence, the diagnosis of CD may come as a shock to them. We feel that empowering elderly patients to adopt a GFD on an individual level based on the presence or absence of symptoms might be appropriate. Nonetheless, future research is needed to evaluate the alternatives to GFD and the other long-term risks of CD, such as malignancy and anaemia, before accepting such an approach.

References

1. Shiha MG, Marks LJ, Sanders DS. Diagnosing coeliac disease in the elderly: a United Kingdom cohort study. *Gastroenterol Hepatol Bed Bench* 2020;13:37-43
2. Ludvigsson JF, Bai JC, Biagi F, Card TR, Ciacci C, Ciclitira PJ, et al. diagnosis and management of adult coeliac disease: Guidelines from the British society of gastroenterology. *Gut* 2014;63:1210-28.
3. Bianchi M, Bardella M. Bone in celiac disease. *Osteoporos Int* 2008;19:1705-16.
4. Capriles VD, Martini LA, Arêas JAG. Metabolic osteopathy in celiac disease: Importance of a gluten-free diet. *Nutr Rev* 2009;67:599-606