



This is a repository copy of *Reduced fractionation in lung cancer patients treated with curative-intent radiotherapy during the COVID-19 pandemic*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/216461/>

Version: Accepted Version

Article:

Faivre-Finn, C., Fenwick, J.D., Franks, K.N. et al. (11 more authors) (2020) Reduced fractionation in lung cancer patients treated with curative-intent radiotherapy during the COVID-19 pandemic. *Clinical Oncology*, 32 (8). pp. 481-489. ISSN 0936-6555

<https://doi.org/10.1016/j.clon.2020.05.001>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Reduced fractionation in lung cancer patients treated with curative-intent radiotherapy during the COVID-19 pandemic

Corinne Faivre-Finn^{a,b*}, John D Fenwick^{c,d}, Kevin N Franks^{e,f}, Stephen Harrow^{g,h}, Matthew QF Hattonⁱ, Crispin Hiley^{j,k}, Jonathan J McAleese^l, Fiona McDonald^m, Jolyne O'Hare^l, Clive Peedellⁿ, Ceri Powell^{o,p}, Tony Pope^q, Robert Rulach^{g,h}, Elizabeth Toy^r

* To whom correspondence should be addressed

^aThe Christie NHS Foundation Trust, Wilmslow Road, Manchester, M20 4BX

^bThe University of Manchester, Oxford Road, Manchester, M13 9PT

^cDepartment of Molecular and Clinical Cancer Medicine, Institute of Translational Medicine, University of Liverpool, Crown Street, Liverpool, L69 3BX

^dDepartment of Physics, Clatterbridge Cancer Centre, Bebington, Wirral, CH63 4JY

^eLeeds Cancer Centre, St James's University Hospital, Beckett Street, Harehills, Leeds, LS9 7LP

^fUniversity of Leeds, Woodhouse, Leeds, LS2 9JT

^gBeatson West of Scotland Cancer Centre, 1053 Great Western Road, Glasgow, G12 0YN

^hUniversity of Glasgow, University Avenue, Glasgow, G12 8QQ

ⁱWeston Park Hospital, Whitham Road, Broomhall, Sheffield, S10 2SJ

^jCRUK Lung Cancer Centre of Excellence, University College London

^kDepartment of Clinical Oncology, University College London Hospitals NHS Foundation Trust, 235 Euston Road, Bloomsbury, London, NW1 2BU

^lNorthern Ireland Cancer Centre, Belfast City Hospital, 10 Jubilee Road, Belfast, BT9 7JL

^mThe Royal Marsden NHS Foundation Trust, Fulham Road, London. SW3 6JJ

ⁿJames Cook University Hospital, Marton Road, Middlesbrough, TS4 3BW

^oSouth West Wales Cancer Centre, Singleton Hospital, Sketty Lane, Swansea, SA2 8QA

^pVelindre Cancer Centre, Velindre Road, Cardiff, CF14 2TL

^qClatterbridge Cancer Centre, Bebington, Wirral, CH63 4JY

^rRoyal Devon and Exeter NHS Foundation Trust, Exeter Hospital, EX2 5DW

Email addresses:

Corinne.Finn@christie.nhs.uk; John.Fenwick@liverpool.ac.uk; Kevin.Franks@nhs.net; Stephen.Harrow@ggc.scot.nhs.uk; MatthewHatton@nhs.net; Crispin.Hiley@ucl.ac.uk; Jonathan.McAleese@belfasttrust.hscni.net; Fiona.McDonald@rmh.nhs.uk; Jolyne.OHare@belfasttrust.hscni.net; ClivePeedell@nhs.net; Ceri.Powell@wales.nhs.uk; Tony.Pope@nhs.net; Robert.Rulach@ggc.scot.nhs.uk; Liz.Toy@nhs.net

Acknowledgements

Corinne Faivre-Finn was supported by the NIHR Manchester Biomedical Research Centre. The authors would like to thank Dr Kate Wicks for her assistance in preparing this manuscript.

Reduced fractionation in lung cancer patients treated with curative-intent radiotherapy during the COVID-19 pandemic

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

The World Health Organisation (WHO) declared COVID-19, the disease caused by the 2019 novel coronavirus SARS-CoV-2, a pandemic on the 11th of March 2020. This situation is resulting in unprecedented demands on the NHS as a whole posing a major burden on cancer services in the UK.

Approximately 49,000 new patients are diagnosed with lung cancer each year in the UK and >50% require radiotherapy treatment. The lung cancer population requiring active treatment has been classified as 'extremely vulnerable' with a significant proportion of previously treated lung cancer patients included in this category due to co-existing severe co-morbidities [1, 2]. There is therefore a need to mitigate the risks of their anti-cancer treatments by addressing risks associated with multiple visits to hospital, treatment-induced immune suppression, and radiation-associated lung injury. This means adapting our current treatment protocols rapidly to reflect the shifting risk-benefit ratio and diminished resources. Furthermore, the impact of this pandemic is likely to last for a significant length of time beyond resumption of normal services. This is due to the anticipated backlog of patients diagnosed with lung cancer, and the increased demands on the radiotherapy departments due to the deferral of radiotherapy for disease sites such as breast and prostate.

General guidance on delivery of radiotherapy during the COVID-19 pandemic has been provided by NICE [3]. One recommendation is to consider alternative dose-fractionation schedules or radiotherapy techniques.

The objectives of this paper are: to identify reduced-fractionation and curative-intent radiotherapy regimes in lung cancer, assess their evidence base, and provide organs-at-risk (OAR) dose constraints. We also discuss limitations and practical considerations associated with the implementation of these reduced-fractionation regimes. The anticipated impact of this work is firstly, to reduce hospital visits and limit exposure to COVID-19 in patients having curative-intent radiotherapy for lung cancer, and secondly, to increase radiotherapy service capacity for operable patients with stage I-III lung cancer, who may not be able to have surgery during the pandemic.

Methods

Systematic reviews and relevant papers were identified by a group of UK clinical oncologists through a PubMed search between 20/3/2020 and 30/3/2020. We also included published and unpublished audits of hypofractionated regimes from UK centres.

Early stage NSCLC

UK practice is based on the recommendations from the UK SABR consortium (18). Here we outline the evidence for reduction in SABR fraction number and provide OAR dose constraints from existing international protocols. We also outline the evidence for hypofractionation (beyond 55Gy in 20 fractions) for central/ultra-central early-stage NSCLC not suitable for SABR due to OAR constraints being exceeded.

1. Single-fraction SABR

Advice

- Consider 30Gy to 34Gy in a single fraction in patients with tumours that are ≤ 2 cm, > 1 cm from the chest wall, and are outside of the no-fly zone. This is in keeping with the current NCCN guidelines [4].

Evidence

Single-fraction schedules of 30-34Gy have been compared to multi-fraction SABR in two phase 2 studies (RTOG 0915, Roswell Park) [5-7]. Local control rate, progression-free survival (PFS), and overall survival (OS), as well as late toxicity and quality of life, were comparable between single-fraction and multi-fraction SABR regimens. Chest wall toxicity did not exceed grade 2 in either arm of both studies. A retrospective study including 146 lesions showed that grade 2-4 chest wall toxicity was 30.6% for lesions abutting the chest wall, 8.2% for tumours ≤ 1 cm from the chest wall, and 3.8% for tumours 1 to 2 cm from the chest wall [8]. Overall grade ≥ 3 chest wall toxicity was 1.4%.

Limitations

- A range of SABR dose/fractionation schedules have been described, but no single regimen has been established as the standard of care.
- Evidence is based on phase 2 data only where the numbers treated within 2 cm of the chest wall is very small.

Practical Considerations

- Only centres with prior experience of delivering lung SABR should offer single-fraction SABR.
- Patients considered for single-fraction SABR are those typically treated with 54Gy in 3 fractions, rather than 55Gy in 5 fractions.
- It is advised only to consider tumours that move less than 1cm after appropriate motion management on 4DCT imaging.
- The dose constraints recommended are those set out in the RTOG 0915 study (see Tables 1 and 2; online appendix).

2. SABR for tumours within 2.5 cm of the chest wall

Advice

- Consider 3-fraction regimes (e.g. 54Gy/3 fractions).
- Where the planning target volume (PTV) abuts or overlaps the chest wall, consider 54Gy/3 fractions or a reduced dose to minimise toxicity (e.g. 48Gy/3 fractions).

Evidence

1 The rate of grade 3 chest wall toxicity with SABR from a large meta-analysis (combining several
2 different dose and fractionations) is 1.2% [9]. Individual papers have found that the tumour to chest
3 wall distance is a significant factor, as well as the maximum dose (Dmax) and volume of chest wall
4 receiving 30Gy (V30) [10-13]. Multi-fraction retrospective data specifically looking at patients with
5 tumours near the chest wall are shown in Table 3 (online appendix). Where the gross tumour
6 volume (GTV) is within 2.5cm of the chest wall, no increased risk was seen with 3 fractions compared
7 to 5 fractions (1.6% compared to 3.2% respectively) [12]. Where the PTV is abutting the chest wall,
8 data from Andolino et al suggests 48Gy/3 fractions has a lower toxicity than 54Gy/3 fractions [10].
9

10 **Limitations**

- 11 • The effect of fractionation schedules on chest wall toxicity has not been investigated in
12 prospective trials.

13 **Practical Considerations**

- 14 • Suggested chest wall dose constraints for 3 fraction schedules are D0.5cc<60Gy, D5cc<40Gy
15 and V30<30cc (Tables 4.1 and 4.2; online appendix).

16 **3. SABR for moderately central tumours**

17 **Advice**

- 18 • Consider 50Gy/5 fractions in moderately central tumours.

19 **Evidence**

20 Moderately central early-stage NSCLC is defined as a lesion within 2cm of the bronchial tree,
21 trachea, major vessels, oesophagus, heart, pericardium, or brachial plexus, or a PTV abutting
22 mediastinal pleura or pericardium, excluding ultra-central disease. An ultra-central lesion is where
23 the PTV abuts either the main bronchi or trachea.

24 Two fractionations are commonly used:

- 25 • 4-5 fractions as per ASTRO guidelines (based largely on studies using a total dose of 45-
26 50Gy) [14].
- 27 • 8 fractions as per UK SABR consortium (total dose 60Gy) [15].

28 Retrospective studies show similar grade 3 or above toxicity rates between 0 and 7.7%, and local
29 control rates between 77.6 - 95%. There is a lack of prospective evidence to suggest which regime is
30 superior. The safest arm in the prospective RTOG 0813 trial was the 50Gy/5 fractions cohort with no
31 \geq grade 3 toxic events. 50Gy in 5 fractions has been used in Glasgow based on the RTOG 0813 dose
32 constraints [16]. In a study of 50 patients, there was a 4% grade 3 toxicity rate and a median OS of 27
33 months, which is consistent with other published literature (Table 5; online appendix). 50Gy/4
34 fractions has also been used in North America but lacks prospective trial data and dose constraints.

35 **Limitations**

- 36 • There is no evidence to support one dose fractionation regime being superior in terms of
37 efficacy or safety.

38 **Practical Considerations**

- 39 • The dose constraints set out in RTOG 0813 are recommended (Tables 6-8; online appendix).

4. SABR for tumours >5cm

Advice

- Tumours >5cm in diameter can be treated with caution, provided that the OAR constraints for tumours <5cm can be met.

Evidence

SABR is currently recommended for T1-2 tumours (or T3 tumours by virtue of invading chest wall) with a maximum size of 5cm [15]. Clinical trials have predominantly excluded lesions larger than 5cm, and therefore conventional fractionation schedules have been favoured in this group. Woody et al reported on 40 patients with a median tumour size of 5.6cm (range: 5.1-10cm) treated with a median dose of 50Gy in 5 fractions [17]. The 18-month local control, OS and grade 3 toxicity rates were 91.2%, 59.7% and 7.5% respectively. A Dutch series reported on 63 patients with a median diameter of 5.8cm (range: 5.1-10.1) with a longer median follow up of 54.7 months [18]. The median OS, 2-year local control and out-of-field distant recurrence rates were 28.3 months, 95.8% and 10% respectively. 30% developed grade \geq 3 toxicity (radiation pneumonitis was the most common toxicity) and 19% of deaths were treatment-related.

Limitations

- There is no prospective data to support SABR for tumours >5cm.

Practical Considerations

- Dose constraints to OARs must be met as when treating lesions \leq 5cm.
- Following treatment, patients should closely followed-up to detect and manage toxicity and expected higher distant relapse rates.

5. Hypofractionation for central/ultra-central early-stage tumours not suitable for SABR

Advice

- Consider 50-60Gy in 15 fractions in patients with central/ultra-central early stage NSCLC not suitable for SABR based on OAR constraints.

Evidence

A prospective phase 1 dose-escalation trial for patients of performance status (PS \geq 2) with stage \geq II NSCLC not suitable for surgery, SABR or chemoradiation used increasing doses in 15 fractions (50Gy, 55Gy or 60Gy) to validate OAR constraints for a 15 fraction schedule in the IMRT/IGRT era. They reported acceptable toxicities and no dose-limiting toxicity was documented [19]. The subsequent randomised phase 3 study comparing 60Gy in either 15 or 30 fractions in patients with PS \geq 2 stage II-III NSCLC has published interim results in abstract form [20]. 60 patients had been enrolled (88% stage III). Chemotherapy was given to some patients sequentially (pre or post RT) but not concurrently. Less toxicity was reported in the 15 fraction arm. Cho et al [21] retrospectively reviewed hypofractionated RT for medically inoperable T1-T3 N0 NSCLC using a risk-adaptive dose schedule (60Gy in 4, 15 or 20 fractions depending on location, size and geometry of the tumour in relation to the oesophagus). 124 patients were included in the study; 72.6% had T1-2 N0 tumours; 65.3% had centrally located disease; 44.1% had PS 2-3; and 20.2% received 60Gy/15 fractions. In patients treated with 15 fractions, the rate of grade 3 pneumonitis was 4% with no grade 4-5 pneumonitis and no grade 2-5 oesophagitis reported.

Limitations

- OAR constraints for 15-fraction schedules were mostly derived from studies including patients with PS \geq 2 and stage II-III disease.
- There is no prospective data to support 50-60Gy in 15 fractions specifically in central or ultra-central early stage NSCLC.

Practical Considerations

- Dose constraints to OARs for the 15-fraction schedule must be met with particular attention to the oesophageal constraint (see Table 9; online appendix).

Stage III NSCLC

1. Concurrent Chemoradiotherapy

Advice

- Consider for selected patients (see practical considerations below).
- Consider accelerated fractionation (i.e. 55Gy/20 fractions).
- Limit chemotherapy dose (see practical considerations below). Consider limiting chemotherapy to two cycles only and starting radiotherapy with cycle one.

Evidence

The randomised phase 2 SOCCAR trial [22] compared sequential versus concurrent chemotherapy combined with 55Gy in 20 fractions. The median number of cycles delivered was 2.8 in the concurrent arm. Toxicity was similar across both arms, with a median survival of 24 months (concurrent arm) in a UK population of patients with stage III NSCLC using 3D planning and treatment techniques. Following the study, a number of the participating centres adopted the schedule, fine-tuning chemotherapy regimens, and evolving treatment techniques by applying PET-CT staging, 4D planning, IMRT and VMAT. With these adaptations, UK centres are reporting encouraging 58% 2-year survival and acceptable rates of acute toxicity [23], which compares favourably to more recent trials, e.g. PACIFIC [24].

Limitations

The SOCCAR study only included 70 patients in the concurrent arm. It was published before many of the more modern staging and treatment techniques were in routine use. The evidence base for contemporary concurrent chemoradiotherapy using a hypofractionated accelerated fractionation schedule is therefore limited (particularly concerning acute and late toxicity) and of a retrospective nature [23].

Practical Considerations

- The inclusion criteria for the SOCCAR study can guide patient selection [22]. OARs constraints as per SOCCAR protocol are detailed in Table 9 (online appendix).
- Chemotherapy as per SOCCAR protocol [22] can be adapted during the COVID-19 epidemic. Consideration should be given to omitting the adjuvant cycles and delivering the concurrent chemotherapy cycles only (cisplatin 60mg/m² IV or carboplatin AUC5 D1 and oral vinorelbine 40mg/m² D1 and 8).

2. Radical radiotherapy +/- sequential chemotherapy

Advice

- Consider for selected patients (see practical considerations below).
- Offer accelerated fractionation (55Gy/20 fractions).
- Consider further hypofractionation (50-58Gy in 15 fractions).
- If offered, limit chemotherapy to 2 cycles and consider delivering it following radiotherapy (see practical considerations below).

Evidence

The hypofractionated regimen of 55Gy/20 fractions has been widely used in the UK [25], with audit data showing similar outcomes to CHART, 99% of patients completing treatment, and a 7% grade ≥3 toxicity rate [26]. Retrospective data on 45Gy in 15 fractions over 3 weeks (BED₁₀ 58.5Gy) showed comparable outcomes to doses ≥60Gy given with conventional fractionation [27]. However,

1 radiobiological calculation suggests this schedule would not be isoeffective in comparison to
2 55Gy/20 fractions (BED_{10} 70.1Gy). A higher dose hypofractionated regime of 60Gy/15 fractions
3 (BED_{10} 90Gy) has been reported by Sunnybrook in patients with stage I-III NSCLC [28]. 47 patients
4 (52.8%) had stage II-III disease and the 2-year survival was 68% for this group. Importantly, the dose
5 constraints derived for this study correspond well to those generated by Fenwick et al using
6 conversion from the I-START 20 fraction schedule [29] (Table 9; online appendix).

7 **Limitations**

8
9 15 fraction schedules have generally been used to treat central early-stage disease, with the
10 treatment of stage III patients limited to selected patients [28]. It should be noted that the toxicity of
11 this regime has not been reported specifically for patients with stage II-III.

12 **Practical considerations**

- 13 • Concerns over hypofractionated dose-escalated radiotherapy in NSCLC are dominated by late
14 radiation toxicity involving central and perihilar structures [30]. The experience of accelerated
15 schedules led to a UK research strategy that tested 4 separate escalation protocols in phase 1/2
16 studies. Two of these protocols used once daily hypofractionated schedules (IDEAL-CRT, I-START)
17 with reassuring toxicity profiles [31, 32]. Applying the principles that Fenwick et al [29] used to
18 develop these schedules to a 15-fraction schedule delivered over 19 – 21 days:
 - 19 ○ Using an α/β of 10, 52Gy/15 fractions is the isoeffective dose for tumour control and
20 using an α/β of 3, 50Gy/15 fractions is isotoxic to 55Gy/20 fractions for late
21 complications.
 - 22 ○ 58Gy/15 fractions would be the equivalent of the highest dose cohorts in these two
23 studies (IDEAL-CRT 73Gy/30 fractions over 6 weeks, I-START 65Gy/20 fractions over 4
24 weeks).
- 25 • The use of IMRT/VMAT is strongly recommended. The radiotherapy planning guidelines for
26 current stage III studies [33] are a resource that can help guide patient selection, outlining and
27 planning using the modified dose constraints in Table 9 (online appendix).
- 28 • The addition of chemotherapy in the sequential setting will need careful consideration,
29 balancing a 4% absolute OS benefit over RT alone [34] against the additional infective risk posed by
30 COVID-19. Consideration should be given to RT first, with deferred chemotherapy given later when
31 the risks related to COVID-19 start decreasing.

Small cell lung cancer

1. Early-stage SCLC

Advice

- Consider SABR (with or without chemotherapy) in T1-2 N0 M0 patients as an alternative to surgery or fractionated radiotherapy. Dose/fractionation and OAR constraints should be the same as those used for early-stage NSCLC.

Evidence

SABR is standard of care in medically inoperable early-stage NSCLC and is increasingly being delivered for early-stage SCLC [35-38]. SABR for early-stage SCLC is a treatment option in the ASTRO 2020 guidelines [39] and in the 2020 NCCN guidelines [40].

The largest series of SABR for LS-SCLC is a retrospective multicentre study including 74 patients [38], of which only 59% of the patients received chemotherapy, 23% received PCI and >30% of patients had a PS ECOG 2-3. Toxicity was mild with 5.2% grade ≥ 2 pneumonitis. Local progression-free survival was 96.1% and overall survival was 34% at 3 years.

Limitations

- Evidence base for SABR is limited to the peripheral early-stage SCLC setting. The risk of toxicity and development of lymph node metastases for central/ultra-central tumours is higher compared to peripheral tumours [41,42]. As data is lacking in ultra-central early-stage SCLC, conventionally fractionated RT is more appropriate for these patients.
- Given the risk of distant metastases, chemotherapy is generally considered in this setting for those patients who are suitable [35, 38].

Practical considerations

- In the context of the COVID-19 pandemic, the risk-benefit ratio of giving chemotherapy should be considered carefully. In patients who are suitable for chemotherapy, it is advisable to give SABR first as the tumour volume may decrease significantly after the first or second cycle of chemotherapy and become difficult to visualize on image-guidance.

2. Radiotherapy Fractionation in Good Performance Status Limited-Stage (LS) SCLC Patients

Advice

- Consider 40Gy in 15 daily fractions given with 1st or 2nd cycle of chemotherapy in patients with good PS LS-SCLC.
- Consider 40Gy in 15 daily fractions after induction chemotherapy in patients who are not suitable for concurrent treatment.
- Limit chemotherapy to a maximum of four cycles

Evidence

The current standard of care is twice-daily radiotherapy (45Gy in 30 fractions) delivered concurrently with cycle 1 or 2 chemotherapy [43-45]. However hypofractionated regimes are also used in UK centres and include: 40Gy in 15 fractions and 50-55Gy in 20 fractions. A randomised study by NCIC demonstrated a survival benefit with early concurrent radiotherapy (week 1) versus late (week 15) using 40Gy in 15 daily fractions [46]. Toxicity in both arms was acceptable. Grade 4 neutropenia was

1 common and pneumonitis was <3%. Grønberg et al [47] reported a randomised phase 2 trial of 157
2 patients with LS-SCLC treated with 42Gy in 15 fractions once daily (OD) or 45Gy in 30 fractions twice
3 daily (BD). There was no difference in one-year or median progression-free survival. There were no
4 differences in \geq grade 3 oesophagitis (OD: 31%, BD: 33%, $p=0.80$) or pneumonitis (OD: 2%, BD: 3%,
5 $p=1.0$) [47]. Videtic et al [48] retrospectively reviewed 122 LS-SCLC patients who received concurrent
6 chemotherapy with 50Gy in 25 fractions over 5 weeks (92pts) or 40Gy in 15 fractions over 3 weeks.
7 There was no difference in treatment related toxicity, overall survival and thoracic local control. Xia
8 et al [49] reported results on 59 LS-SCLC patients treated with 55Gy in 22 fractions over 30 days and
9 concurrent chemotherapy. 25% of patients developed \geq grade 3 oesophagitis and 10% of patients
10 developed \geq grade 3 pneumonitis. 40Gy in 15 fractions has been used concurrently and sequentially
11 in Leeds for limited stage SCLC for >10 years. Institutional dose constraints are listed in Table 10
12 (online appendix) and a recent unpublished audit of 43 LS-SCLC patients treated with concurrent
13 chemoradiotherapy (40Gy in 15 fractions) showed a 1-year OS of 88% and a median OS of 26.9
14 months [15.6-50.4].
15
16
17

18 **Limitations**

- 19 • The initial data on 40Gy in 15 fractions is from 1993 [46], and therefore radiotherapy planning
20 and delivery would be considered sub-optimal.
- 21 • Most data on hypofractionated regimes are from retrospective single-institution studies.
- 22 • A variety of different hypofractionated regimes are used in the published literature and in
23 routine UK practice.
- 24
- 25
- 26
- 27
- 28
- 29

30 **Practical considerations**

- 31 • When treating LS-SCLC with hypofractionated radiotherapy, IV contrast (if not contraindicated),
32 and 3DCT/IMRT planning with an offline IGRT protocol with volumetric imaging are considered
33 the standard of care. 4DCT planning and daily online CBCT is highly recommended, particularly if
34 OAR doses are close to tolerance.
- 35
- 36
- 37
- 38 • Leeds OAR constraints for 40Gy/15 fractions regime are listed in Table 10 (online appendix).
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

Discussion

This guidance document on reduced fractionation for lung cancer being treated with curative intent during the COVID-19 pandemic builds on a long tradition of hypofractionated radiotherapy in the UK. It reflects the current published literature and the combined experience of the authors and their colleagues in the UK and globally. However, it is acknowledged that for many centres, the fractionation regimens outlined will represent a significant change to current practice and standard of care. The extent of adoption of this guidance may reflect geographical pressures, although it is likely that all radiotherapy departments will need to adapt during this global pandemic.

This guidance document should be discussed with other specialist lung MDT members as access to adequate nodal staging procedures (e.g. EBUS-TBNA) and respiratory function testing is likely to be compromised during the peak of the virus pandemic. That discussion will disseminate the potential changes to radiotherapy practice that could be made in order to alleviate pressure on other departments such as thoracic surgery.

Adequate discussion with the patient about the risk and benefits of treatment during the COVID-19 pandemic and uncertainties about toxicity from reduced fractionation where there is limited experience in a department are an essential component of the consent process.

Centres should document deviations from standard pre-treatment work-up as well as deviations from standard of care treatments. We consider prospective and multi-centre documentation of outcome (including toxicity) from these reduced fractionation regimens as essential. We also urge colleagues to join national/international data collection initiatives on the impact of the COVID-19 pandemic.

References

- [1] Global Initiative for Chronic Obstructive Lung Disease. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease (2020 Report). 2020.
- [2] May R, Powis S. Caring for people at highest risk during COVID-19 incident. 2020.
- [3] The National Institute for Health and Care Excellence. COVID-19 rapid guideline: delivery of radiotherapy. NICE guideline [NG162]. 2020.
- [4] National Comprehensive Cancer Network. NCCN Guidelines Non-Small Cell Lung Cancer. https://www.nccn.org/professionals/physician_gls/pdf/nscl_blocks.pdf
- [5] Singh AK, Gomez-Suescun JA, Stephans KL, Bogart JA, Hermann GM, Tian L, et al. One Versus Three Fractions of Stereotactic Body Radiation Therapy for Peripheral Stage I to II Non-Small Cell Lung Cancer: A Randomized, Multi-Institution, Phase 2 Trial. *Int J Radiat Oncol Biol Phys.* 2019;105:752-9.
- [6] Videtic GM, Hu C, Singh AK, Chang JY, Parker W, Olivier KR, et al. A Randomized Phase 2 Study Comparing 2 Stereotactic Body Radiation Therapy Schedules for Medically Inoperable Patients With Stage I Peripheral Non-Small Cell Lung Cancer: NRG Oncology RTOG 0915 (NCCTG N0927). *Int J Radiat Oncol Biol Phys.* 2015;93:757-64.
- [7] Videtic GM, Paulus R, Singh AK, Chang JY, Parker W, Olivier KR, et al. Long-term Follow-up on NRG Oncology RTOG 0915 (NCCTG N0927): A Randomized Phase 2 Study Comparing 2 Stereotactic Body Radiation Therapy Schedules for Medically Inoperable Patients With Stage I Peripheral Non-Small Cell Lung Cancer. *Int J Radiat Oncol Biol Phys.* 2019;103:1077-84.

- 1 [8] Manyam BV, Videtic GMM, Verdecchia K, Reddy CA, Woody NM, Stephans KL. Effect of Tumor
2 Location and Dosimetric Predictors for Chest Wall Toxicity in Single-Fraction Stereotactic Body
3 Radiation Therapy for Stage I Non-Small Cell Lung Cancer. *Pract Radiat Oncol*. 2019;9:e187-e95.
- 4 [9] Ma JT, Liu Y, Sun L, Milano MT, Zhang SL, Huang LT, et al. Chest Wall Toxicity After Stereotactic
5 Body Radiation Therapy: A Pooled Analysis of 57 Studies. *Int J Radiat Oncol Biol Phys*. 2019;103:843-
6 50.
- 7 [10] Andolino DL, Forquer JA, Henderson MA, Barriger RB, Shapiro RH, Brabham JG, et al. Chest wall
8 toxicity after stereotactic body radiotherapy for malignant lesions of the lung and liver. *Int J Radiat
9 Oncol Biol Phys*. 2011;80:692-7.
- 10 [11] Asai K, Shioyama Y, Nakamura K, Sasaki T, Ohga S, Nonoshita T, et al. Radiation-induced rib
11 fractures after hypofractionated stereotactic body radiation therapy: risk factors and dose-volume
12 relationship. *Int J Radiat Oncol Biol Phys*. 2012;84:768-73.
- 13 [12] Bongers EM, Haasbeek CJ, Lagerwaard FJ, Slotman BJ, Senan S. Incidence and risk factors for
14 chest wall toxicity after risk-adapted stereotactic radiotherapy for early-stage lung cancer. *J Thorac
15 Oncol*. 2011;6:2052-7.
- 16 [13] Nambu A, Onishi H, Aoki S, Tominaga L, Kuriyama K, Araya M, et al. Rib fracture after
17 stereotactic radiotherapy for primary lung cancer: prevalence, degree of clinical symptoms, and risk
18 factors. *BMC Cancer*. 2013;13:68.
- 19 [14] Videtic GMM, Donington J, Giuliani M, Heinzerling J, Karas TZ, Kelsey CR, et al. Stereotactic body
20 radiation therapy for early-stage non-small cell lung cancer: Executive Summary of an ASTRO
21 Evidence-Based Guideline. *Pract Radiat Oncol*. 2017;7:295-301.
- 22 [15] UK SABR Consortium. UK SABR Consortium Guidelines v6.1. 2019.
- 23 [16] Rulach R, McLoone P, Lumsden G, McKay S, MacLaren V, Macphee J, et al. Toxicity and Efficacy
24 of Stereotactic Ablative Body Radiotherapy for Moderately Central Non-small Cell Lung Cancers
25 Using 50 Gy in Five Fractions. *Clin Oncol (R Coll Radiol)*. 2020;32:250-8.
- 26 [17] Woody NM, Stephans KL, Marwaha G, Djemil T, Videtic GM. Stereotactic Body Radiation
27 Therapy for Non-Small Cell Lung Cancer Tumors Greater Than 5 cm: Safety and Efficacy. *Int J Radiat
28 Oncol Biol Phys*. 2015;92:325-31.
- 29 [18] Tekatli H, van 't Hof S, Nossent EJ, Dahele M, Verbakel W, Slotman BJ, et al. Use of Stereotactic
30 Ablative Radiotherapy (SABR) in Non-Small Cell Lung Cancer Measuring More Than 5 cm. *J Thorac
31 Oncol*. 2017;12:974-82.
- 32 [19] Westover KD, Loo BW, Jr., Gerber DE, Iyengar P, Choy H, Diehn M, et al. Precision
33 Hypofractionated Radiation Therapy in Poor Performing Patients With Non-Small Cell Lung Cancer:
34 Phase 1 Dose Escalation Trial. *Int J Radiat Oncol Biol Phys*. 2015;93:72-81.
- 35 [20] Iyengar P, Westover KD, Court LE, Patel MK, Shivnani AT, Saunders MW, et al. A Phase III
36 Randomized Study of Image Guided Conventional (60 Gy/30 fx) Versus Accelerated,
37 Hypofractionated (60 Gy/15 fx) Radiation for Poor Performance Status Stage II and III NSCLC
38 Patients—An Interim Analysis. *International Journal of Radiation Oncology, Biology, Physics*.
39 2016;96:E451.
- 40 [21] Cho WK, Noh JM, Ahn YC, Oh D, Pyo H. Radiation Therapy Alone in cT1-3N0 Non-small Cell Lung
41 Cancer Patients Who Are Unfit for Surgical Resection or Stereotactic Radiation Therapy: Comparison
42 of Risk-Adaptive Dose Schedules. *Cancer Res Treat*. 2016;48:1187-95.
- 43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [22] Maguire J, Khan I, McMenemin R, O'Rourke N, McNee S, Kelly V, et al. SOCCAR: A randomised
2 phase II trial comparing sequential versus concurrent chemotherapy and radical hypofractionated
3 radiotherapy in patients with inoperable stage III Non-Small Cell Lung Cancer and good performance
4 status. *Eur J Cancer*. 2014;50:2939-49.
- 5 [23] Iqbal MS, Vashisht G, McMenemin R, Atherton P, McDonald F, Simmons T, et al.
6 Hypofractionated Concomitant Chemoradiation in Inoperable Locally Advanced Non-small Cell Lung
7 Cancer: A Report on 100 Patients and a Systematic Review. *Clin Oncol (R Coll Radiol)*. 2019;31:e1-
8 e10.
- 9 [24] Antonia SJ, Villegas A, Daniel D, Vicente D, Murakami S, Hui R, et al. Overall Survival with
10 Durvalumab after Chemoradiotherapy in Stage III NSCLC. *N Engl J Med*. 2018;379:2342-50.
- 11 [25] Prewett SL, Aslam S, Williams MV, Gilligan D. The management of lung cancer: a UK survey of
12 oncologists. *Clin Oncol (R Coll Radiol)*. 2012;24:402-9.
- 13 [26] Robinson SD, Tahir BA, Absalom KAR, Lankathilake A, Das T, Lee C, et al. Radical accelerated
14 radiotherapy for non-small cell lung cancer (NSCLC): A 5-year retrospective review of two dose
15 fractionation schedules. *Radiother Oncol*. 2020;143:37-43.
- 16 [27] Amini A, Lin SH, Wei C, Allen P, Cox JD, Komaki R. Accelerated hypofractionated radiation
17 therapy compared to conventionally fractionated radiation therapy for the treatment of inoperable
18 non-small cell lung cancer. *Radiat Oncol*. 2012;7:33.
- 19 [28] Zeng K, Poon I, Ung Y, Zhang L, Cheung P. Accelerated Hypofractionated Radiation Therapy for
20 Centrally Located Lung Tumors Not Suitable for Stereotactic Body Radiation Therapy (SBRT) or
21 Concurrent Chemoradiotherapy (CRT). *International Journal of Radiation Oncology, Biology, Physics*
22 2018. p. e719-e20.
- 23 [29] Fenwick JD, Nahum AE, Malik ZI, Eswar CV, Hatton MQ, Laurence VM, et al. Escalation and
24 intensification of radiotherapy for stage III non-small cell lung cancer: opportunities for treatment
25 improvement. *Clin Oncol (R Coll Radiol)*. 2009;21:343-60.
- 26 [30] Cannon DM, Mehta MP, Adkison JB, Khuntia D, Traynor AM, Tome WA, et al. Dose-limiting
27 toxicity after hypofractionated dose-escalated radiotherapy in non-small-cell lung cancer. *J Clin*
28 *Oncol*. 2013;31:4343-8.
- 29 [31] Landau DB, Hughes L, Baker A, Bates AT, Bayne MC, Counsell N, et al. IDEAL-CRT: A Phase 1/2
30 Trial of Isotoxic Dose-Escalated Radiation Therapy and Concurrent Chemotherapy in Patients With
31 Stage II/III Non-Small Cell Lung Cancer. *Int J Radiat Oncol Biol Phys*. 2016;95:1367-77.
- 32 [32] Lester J, Courtier N, Eswar C, Mohammed N, Fenwick J, Griffiths G, et al. Initial results of the
33 phase Ib/II, I-START trial: Isotoxic accelerated radiotherapy for the treatment of stage II-IIIb NSCLC.
34 *Journal of Clinical Oncology*. 2018;36:e20551-e.
- 35 [33] Hatton MQF, Lawless CA, Faivre-Finn C, Landau D, Lester JF, Fenwick J, et al. Accelerated, Dose
36 escalated, Sequential Chemoradiotherapy in Non-small-cell lung cancer (ADSCaN): a protocol for a
37 randomised phase II study. *BMJ Open*. 2019;9:e019903.
- 38 [34] Auperin A, Le Pechoux C, Rolland E, Curran WJ, Furuse K, Fournel P, et al. Meta-analysis of
39 concomitant versus sequential radiochemotherapy in locally advanced non-small-cell lung cancer. *J*
40 *Clin Oncol*. 2010;28:2181-90.
- 41 [35] Paximadis P, Beebe-Dimmer JL, George J, Schwartz AG, Wozniak A, Gadgeel S. Comparing
42 Treatment Strategies for Stage I Small-cell lung Cancer. *Clin Lung Cancer*. 2018;19:e559-e65.
- 43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [36] Stahl JM, Corso CD, Verma V, Park HS, Nath SK, Husain ZA, et al. Trends in stereotactic body
2 radiation therapy for stage I small cell lung cancer. *Lung Cancer*. 2017;103:11-6.
- 3 [37] Verma V, Hasan S, Wegner RE, Abel S, Colonias A. Stereotactic ablative radiation therapy versus
4 conventionally fractionated radiation therapy for stage I small cell lung cancer. *Radiother Oncol*.
5 2019;131:145-9.
- 6
7 [38] Verma V, Simone CB, 2nd, Allen PK, Gajjar SR, Shah C, Zhen W, et al. Multi-Institutional
8 Experience of Stereotactic Ablative Radiation Therapy for Stage I Small Cell Lung Cancer. *Int J Radiat*
9 *Oncol Biol Phys*. 2017;97:362-71.
- 10
11 [39] Simone CB, 2nd, Bogart JA, Cabrera AR, Daly ME, DeNunzio NJ, Detterbeck F, et al. Radiation
12 Therapy for Small Cell Lung Cancer: An ASTRO Clinical Practice Guideline. *Pract Radiat Oncol*. 2020.
- 13
14 [40] National Comprehensive Cancer Network. NCCN Guidelines Small Cell Lung Cancer.
15 https://www.nccn.org/professionals/physician_gls/pdf/scl_blocks.pdf
- 16
17 [41] Tekatli H, Haasbeek N, Dahele M, De Haan P, Verbakel W, Bongers E, et al. Outcomes of
18 Hypofractionated High-Dose Radiotherapy in Poor-Risk Patients with "Ultracentral" Non-Small Cell
19 Lung Cancer. *J Thorac Oncol*. 2016;11:1081-9.
- 20
21 [42] Chen H, Laba JM, Zayed S, Boldt RG, Palma DA, Louie AV. Safety and Effectiveness of
22 Stereotactic Ablative Radiotherapy for Ultra-Central Lung Lesions: A Systematic Review. *J Thorac*
23 *Oncol*. 2019;14:1332-42.
- 24
25 [43] The National Institute for Health and Care Excellence. Lung cancer: diagnosis and management.
26 NICE guideline [NG122]. 2019.
- 27
28 [44] Faivre-Finn C, Snee M, Ashcroft L, Appel W, Barlesi F, Bhatnagar A, et al. Concurrent once-daily
29 versus twice-daily chemoradiotherapy in patients with limited-stage small-cell lung cancer
30 (CONVERT): an open-label, phase 3, randomised, superiority trial. *Lancet Oncol*. 2017;18:1116-25.
- 31
32 [45] Turrisi AT, 3rd, Kim K, Blum R, Sause WT, Livingston RB, Komaki R, et al. Twice-daily compared
33 with once-daily thoracic radiotherapy in limited small-cell lung cancer treated concurrently with
34 cisplatin and etoposide. *N Engl J Med*. 1999;340:265-71.
- 35
36 [46] Murray N, Coy P, Pater JL, Hodson I, Arnold A, Zee BC, et al. Importance of timing for thoracic
37 irradiation in the combined modality treatment of limited-stage small-cell lung cancer. The National
38 Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol*. 1993;11:336-44.
- 39
40 [47] Grønberg BH, Halvorsen TO, Flotten O, Brustugun OT, Brunsvig PF, Aasebo U, et al. Randomized
41 phase II trial comparing twice daily hyperfractionated with once daily hypofractionated thoracic
42 radiotherapy in limited disease small cell lung cancer. *Acta Oncol*. 2016;55:591-7.
- 43
44 [48] Videtic GM, Truong PT, Dar AR, Yu EW, Stitt LW. Shifting from hypofractionated to
45 "conventionally" fractionated thoracic radiotherapy: a single institution's 10-year experience in the
46 management of limited-stage small-cell lung cancer using concurrent chemoradiation. *Int J Radiat*
47 *Oncol Biol Phys*. 2003;57:709-16.
- 48
49 [49] Xia B, Hong LZ, Cai XW, Zhu ZF, Liu Q, Zhao KL, et al. Phase 2 study of accelerated
50 hypofractionated thoracic radiation therapy and concurrent chemotherapy in patients with limited-
51 stage small-cell lung cancer. *Int J Radiat Oncol Biol Phys*. 2015;91:517-23.
- 52
53
54
55
56
57
58
59
60
61
62
63
64
65

Supplementary Materials

Table 1. Dose Gradient Requirements Based on Target Volume (from NRG Oncology RTOG 0915)

PTV Volume (cc)	Ratio of Prescription Isodose Volume to the PTV Volume		Ratio of 50% Prescription Isodose Volume to the PTV Volume, R _{50%}		Maximum Dose (in % of dose prescribed) @ 2 cm from PTV in Any Direction, D _{2cm} (%)		Percentage of Lung Receiving 20Gy Total or More, V ₂₀ (%)	
	Deviation		Deviation		Deviation		Deviation	
	None	Minor	None	Minor	None	Minor	None	Minor
1.8	<1.2	<1.5	<5.9	<7.5	<50.0	<57.0	<10	<15
3.8	<1.2	<1.5	<5.5	<6.5	<50.0	<57.0	<10	<15
7.4	<1.2	<1.5	<5.1	<6.0	<50.0	<58.0	<10	<15
13.2	<1.2	<1.5	<4.7	<5.8	<50.0	<58.0	<10	<15
22.0	<1.2	<1.5	<4.5	<5.5	<54.0	<63.0	<10	<15
34.0	<1.2	<1.5	<4.3	<5.3	<58.0	<68.0	<10	<15
50.0	<1.2	<1.5	<4.0	<5.0	<62.0	<77.0	<10	<15
70.0	<1.2	<1.5	<3.5	<4.8	<66.0	<86.0	<10	<15
95.0	<1.2	<1.5	<3.3	<4.4	<70.0	<89.0	<10	<15
126.0	<1.2	<1.5	<3.1	<4.0	<73.0	<91.0	<10	<15
163.0	<1.2	<1.5	<2.9	<3.7	<77.0	<94.0	<10	<15

PTV: planning target volume

Table 2. Organ dose-volume limits for 30-34Gy single fraction (from NRG Oncology RTOG 0915)

Serial Tissue	Volume (cc)	Volume Max (Gy)	Max Point Dose (Gy)
Spinal Cord	<0.35	10	14
	<1.2	7	
Oesophagus	<5	11.9	15.4
Brachial Plexus	<3	14	17.5
Heart/Pericardium	<15	16	22
Great vessels	<10	31	37
Trachea and Large Bronchus	<4	10.5	20.2
Rib	<1	22	30
Skin	<10	23	26
Stomach	<10	11.2	12.4
Parallel Tissue	Critical Volume (cc)	Critical Volume Dose Max (Gy)	
Lung (Right & Left)	1500	7	
Lung (Right & Left)	1000	7.4	

Table 3. Dose/fractionation, biological effective dose, tumour to chest wall distance and rate of toxicity

Paper	Number (n)	Dose/fx	BED ₃ Gy	BED ₁₀ Gy	GTV to CWD (cm)	Rate of toxicity
Andolino [1]	18	54/3 (median)	378	151	0.1	100% any grade
Andolino [1]	61	48/3	304	125	0.2	0% any grade
Asai[2]	116	48/4	240	106	2 (0.3 – 6.2)	24.1% rib fracture, 0.86% G3
Bongers [3]	183	60/3	460	180	<2.5 85.5%*	Any grade CWP: 10.4% G3 CWP: 1.6%
Bongers [3]	187	60/5	300	132	<2.5 91%*	Any grade CWP: 14.4% G3 CWP: 3.2%
Bongers [3]	73	60/8	210	105	<2.5 71.4%*	Any grade CWP: 15% G3 CWP: 1.4%
Nambu [4]	95	48/4	240	106	0.6 (0 - 5.3)	G3 CWP 0%
Nambu [4]	45	60/10	180	96	0.6 (0 - 5.3)	G3 CWP 0%
Nambu [4]	37	70/10	233.3	119	0.6 (0 - 5.3)	G3 CWP 0%

fx: fractions, BED: biological effective dose, CWD: chest wall distance, CWP: chest wall pain, GTV: gross tumour volume , G: grade

* Percentage of patients with tumours within 2.5cm of the chest wall

Table 4.1. Biological effective dose, Dmax to chest wall and ribs

Paper	Number (n)	Dose/fx	BED ₃ Gy	BED ₁₀ Gy	Dmax CW (Gy)	Dmax rib (Gy)	Rate of toxicity
Andolino [1]	18	54/3	378	151	64	64	100% any grade, worst possible G3 rate 16.6%
Andolino [1]	61	48/3	304	125	57	52	0% any grade
Taremi [5]	29	54/3	378	151	-	50.2	No rib fracture
		60/3*	460	180			
	17	54/3	378	151	-	63.7	Rib fracture
		60/3*	460	180			
	21	54/3	378	151	-	62.8	CW pain
60/3*		460	180				
25	54/3	378	151	-	47.2	No CW pain	
	60/3*	460	180				

CW: chest wall, fx: fractions, BED: biological effective dose

*unable to separate number of patients by fractionation as data not available in paper

Table 4.2. Volumetric constraints to the chest wall

Paper	Number (n)	Dose(Gy)/fx (median)	BED ₃ Gy	BED ₁₀ Gy	Dose constraint	Toxicity endpoint
Andolino [7]	347 lesions	18–72/2–5 (54/3)	378	151	D15Gy <240cc	Limits CW toxicity (any grade) to 30%
					D20Gy <130cc	
					D30Gy <40cc	
					D40Gy <15cc	
					D5cc 40Gy	Predicts 10% CW tox
					D15cc 40Gy	Predicts 30% CW tox
					Dmax >50Gy	Significantly increases risk of CW pain and rib fracture
Pettersson [6]	33	45/3	270	112.5	D2cc < 21 Gy	0% rib fracture
					D2cc < 27.2 Gy	5% rib fracture
					D2cc < 49.8 Gy	50% rib fracture
Taremi [5]	46	54/3	378	151	D0.5cc 60 Gy	50% rib fracture
Dunlap [7]	60	60/3*	460	180		
		21-60/3-5 (60/3)	460	180	V30 (30cc)	G2 CWP 30% if V30>35cc
Mutter [8]	126	40-60/3-5 (54/3)	378	151	V30 (70cc)	G2 CWP 27.8% correlated with V30 >70cc
Stephans [9]	45	60/3	460	180	V30 <30cc	G2 CWP 10-15% if V30<30cc
Welsh [10]	265	50/4	258.3	112.5	V30 <30cc	If V30<30cc G2 CWP rate 2.7%

CW: chest wall, fx: fraction

*unable to separate number of patients by fractionation as data not available in paper

Table 5. Dose fractionation for moderately central early-stage NSCLC

Fractionation	Tumour BED ₁₀ Gy	OARs BED ₃ Gy	Risk of ≥G3 toxicity	Tumour control	Number (n)	References
60/8	105	210	6.3%	mOS 47 months, 3 yr LCR 92.6%	63	Haasbeek [11]
			Unknown G3 rate, but 0% G4 toxicity	mOS, n/a, 4 yr LCR 77.8%*	9	Taremi [12]
			6.4%	mOS 38 months, LCR n/a	80	Tekatli [13]
50/5	100	216.67	4% (10% risk of chest infection 90 days post SABR)	mOS 27 months, 2 yr LCR 77.6%	50	Rulach [14]
			0%	mOS NR, LCR 100%	10	Olsen [15]
			0%	mOS 41.6, 2 yr LCR 87.5	8	Bezjak [16]
			2.9%	2 yr LCR 90%, 2 yr OS 63.2%	24	*Chaudhuri [17]
			7.7% late toxicity	mOS 42.1, 3 yr LCR 95%	65	§Arnett [18]
50/4	112.5	258.3	2.9%	2 yr LCR 90%, 2 yr OS 63.2%	10	*Chaudhuri [17]
			11%	2 yr LCR 100%	47	#Rowe [19]
			1.2%	mOS 55.6 months, 3 yr LCR 96.5%	82	Chang [20]
48/4	105.6	240	<14.7%	mOS 42.1, 3 yr LCR 95%	34	§Arnett [18]
60/4	150	360	41% acute toxicity	Crude LCR 5.8%, 2year OS 52%	17	Bral [21]
60/3	180	460	27.3%	mOS 24.4 months	22	Fakiris [22]

* Includes 7 ultracentral patients

Includes metastases, mixed cohort with median dose and fractionation 50 Gy/4 fx

§ Treated on consecutive days

mOS: median overall survival; LCR: local control rate

Table 6. Conformality of Prescribed Dose for Calculations Based on Deposition of Photon Beam Energy in Heterogeneous Tissue for 50Gy in 5 fraction regime (from RTOG 0813)

PTV Volume (cc)	Ratio of Prescription Isodose Volume to PTV		Ratio of 50% Prescription Isodose Volume to PTV, R50%		Maximum Dose (% of dose prescribed) 2 cm from PTV in any direction, D2cm (Gy)		Percentage of Lung Receiving ≥ 20 Gy, V20 (%)	
	Deviation None	Deviation Minor	Deviation None	Deviation Minor	Deviation None	Deviation Minor	Deviation None	Deviation Minor
1.8	<1.2	<1.5	<5.9	<7.5	<50.0	<57.0	<10	<15
3.8	<1.2	<1.5	<5.5	<6.5	<50.0	<57.0	<10	<15
7.4	<1.2	<1.5	<5.1	<6.0	<50.0	<58.0	<10	<15
13.2	<1.2	<1.5	<4.7	<5.8	<50.0	<58.0	<10	<15
22.0	<1.2	<1.5	<4.5	<5.5	<54.0	<63.0	<10	<15
34.0	<1.2	<1.5	<4.3	<5.3	<58.0	<68.0	<10	<15
50.0	<1.2	<1.5	<4.0	<5.0	<62.0	<77.0	<10	<15
70.0	<1.2	<1.5	<3.5	<4.8	<66.0	<86.0	<10	<15
95.0	<1.2	<1.5	<3.3	<4.4	<70.0	<89.0	<10	<15
126.0	<1.2	<1.5	<3.1	<4.0	<73.0	>91.0	<10	<15
163.0	<1.2	<1.5	<2.9	<3.7	<77.0	>94.0	<10	<15

PTV: planning target volume

Table 7. Maximum dose limits to a point or volume within several critical organs. These are absolute limits, and treatment delivery that exceeds these limits will constitute a major protocol violation (from RTOG 0813)

Serial Tissue	Volume (cc)	Volume Max (Gy)	Max Point Dose (Gy)	Avoidance Endpoint
Spinal Cord	<0.25	22.5 (4.5 Gy/fx)	30 (6 Gy/fx)	Myelitis
	<0.5	13.5 (2.7 Gy/fx)		
Ipsilateral Brachial Plexus	<3	30 (6 Gy/fx)	32 (6.4 Gy/fx)	Neuropathy
Skin	<10	30 (6 Gy/fx)	32 (6.4 Gy/fx)	Ulceration
Parallel Tissue	Critical Volume	Critical Volume Dose Max (Gy)		Avoidance Endpoint
Lung (Right & Left)	1500	12.5 (2.5 Gy/fx)		Basic Lung Function
Lung (Right & Left)	1000	13.5 (2.7 Gy/fx)		Pneumonitis

Fx: fractions

Table 8. Suggested volume limits are listed for these organs to be used for treatment planning purposes. Since the tumour and normal tissue may not allow strict avoidance, the volume limits (columns 2 and 3) will not be scored as protocol violations if exceeded. However, the maximum point dose limits (column 4) must be respected (from RTOG 0813)

Serial Tissue*	Volume	Volume Max (Gy)	Max Point Dose (Gy)	Avoidance Endpoint
Esophagus, non-adjacent wall	<5 cc	27.5 Gy (5.5 Gy/fx)	105% of PTV prescription	Stenosis/fistula
Heart/Pericardium	<15 cc	32 Gy (6.4 Gy/fx)	105% of PTV prescription	Pericarditis
Great vessels, non-adjacent wall	<10 cc	47 Gy (9.4 Gy/fx)	105% of PTV prescription	Aneurysm
Trachea and ipsilateral bronchus, non-adjacent wall	<4 cc	18 Gy (3.6 Gy/fx)	105% of PTV prescription	Stenosis/fistula

Fx: fractions, PTV: Planning Target Volume

Table 9. Dose constraints for hypofractionated radiotherapy in stage 3 NSCLC

Dose (Gy)	Volume	Concurrent CRTT 55Gy/20fx	RT only UK * 50 – 58Gy/15fx	RT only Canadian ** 50 – 60Gy/15fx
Spinal Cord	Max D 0.1cc	44Gy	<42Gy	38Gy
Oesophagus*	Max Vol	D 1cc <55Gy	D1cc <52Gy	50Gy V45 <10cc
Brachial Plexus	Max Vol	55Gy	<50Gy 0.5cc <42Gy	<50Gy
Heart/Pericardium	D100% D67% D33%	V ₃₀ <36%	<33Gy <40Gy <52Gy	Max 63Gy V57 <10cc
Mediastinal envelope	Max Vol		58Gy	(Great Vessels) 63Gy V57 <10cc
Trachea and Large Bronchus	Max Vol		58Gy	63Gy V57 <10cc
Rib	Max Vol			63Gy V30 <30cc
Skin	Max			0Gy
Stomach	Max Vol			50Gy V45 <10cc
Lung – GTV		V20 <35% MLD <18Gy	V19 <35% MLD <16Gy	V20 <30% V5 <60% MLD <20Gy
Contralateral lung	V5		<60%	

*15 fraction conversion from the I-START 20 fraction schedule (23)

** Constraints based on Sunnybrook study (24) and clinical update via personal communication with Dr Patrick Cheung

MLD-mean lung dose; GTV: Gross Tumour Volume, CRTT: chemo-radiotherapy; fx: fractions

Table 10. Leeds organs at risk constraints in LS-SCLC

Lung-GTV	Controlateral lung (not mandatory)	Spinal canal PRV	Heart	Oesophagus	Brachial plexus
V20 <30% (ideally); up to 35% (accepted); MLD <15Gy (ideally); up to 18Gy (accepted)*	V20 <10% V10 < 50% V5 <70% MLD <8Gy	Max 35Gy D0.5cc <36Gy	D100%<33 %	Ideally, <12 cm should receive total dose	D0.5cc <42Gy

Constraints based on practice in Leeds, via personal communication with Dr Kevin Franks and Dr Mike Snee

* A MLD (mean lung dose) of 18-20Gy and V20 of 35-40% can be considered in very selected cases

** A margin of 5mm should be used to create a spinal cord PRV. A smaller margin may be used (e.g. 3mm) if the tumour is close to cord provided daily on-line imaging is requested and the cone beam CT is matched to bone

References

- [1] Andolino DL, Forquer JA, Henderson MA, Barriger RB, Shapiro RH, Brabham JG, et al. Chest wall toxicity after stereotactic body radiotherapy for malignant lesions of the lung and liver. *Int J Radiat Oncol Biol Phys.* 2011;80:692-7.
- [2] Asai K, Shioyama Y, Nakamura K, Sasaki T, Ohga S, Nonoshita T, et al. Radiation-induced rib fractures after hypofractionated stereotactic body radiation therapy: risk factors and dose-volume relationship. *Int J Radiat Oncol Biol Phys.* 2012;84:768-73.
- [3] Bongers EM, Haasbeek CJ, Lagerwaard FJ, Slotman BJ, Senan S. Incidence and risk factors for chest wall toxicity after risk-adapted stereotactic radiotherapy for early-stage lung cancer. *J Thorac Oncol.* 2011;6:2052-7.
- [4] Nambu A, Onishi H, Aoki S, Tominaga L, Kuriyama K, Araya M, et al. Rib fracture after stereotactic radiotherapy for primary lung cancer: prevalence, degree of clinical symptoms, and risk factors. *BMC Cancer.* 2013;13:68.
- [5] Taremi M, Hope A, Lindsay P, Dahele M, Fung S, Purdie TG, et al. Predictors of radiotherapy induced bone injury (RIBI) after stereotactic lung radiotherapy. *Radiat Oncol.* 2012;7:159.
- [6] Pettersson N, Nyman J, Johansson KA. Radiation-induced rib fractures after hypofractionated stereotactic body radiation therapy of non-small cell lung cancer: a dose- and volume-response analysis. *Radiother Oncol.* 2009;91:360-8.
- [7] Dunlap NE, Cai J, Biedermann GB, Yang W, Benedict SH, Sheng K, et al. Chest wall volume receiving >30 Gy predicts risk of severe pain and/or rib fracture after lung stereotactic body radiotherapy. *Int J Radiat Oncol Biol Phys.* 2010;76:796-801.
- [8] Mutter RW, Liu F, Abreu A, Yorke E, Jackson A, Rosenzweig KE. Dose-volume parameters predict for the development of chest wall pain after stereotactic body radiation for lung cancer. *Int J Radiat Oncol Biol Phys.* 2012;82:1783-90.
- [9] Stephans KL, Djemil T, Tendulkar RD, Robinson CG, Reddy CA, Videtic GM. Prediction of chest wall toxicity from lung stereotactic body radiotherapy (SBRT). *Int J Radiat Oncol Biol Phys.* 2012;82:974-80.
- [10] Welsh J, Thomas J, Shah D, Allen PK, Wei X, Mitchell K, et al. Obesity increases the risk of chest wall pain from thoracic stereotactic body radiation therapy. *Int J Radiat Oncol Biol Phys.* 2011;81:91-6.
- [11] Haasbeek CJ, Lagerwaard FJ, Slotman BJ, Senan S. Outcomes of stereotactic ablative radiotherapy for centrally located early-stage lung cancer. *J Thorac Oncol.* 2011;6:2036-43.
- [12] Taremi M, Hope A, Dahele M, Pearson S, Fung S, Purdie T, et al. Stereotactic body radiotherapy for medically inoperable lung cancer: prospective, single-center study of 108 consecutive patients. *Int J Radiat Oncol Biol Phys.* 2012;82:967-73.
- [13] Tekatli H, Senan S, Dahele M, Slotman BJ, Verbakel WF. Stereotactic ablative radiotherapy (SABR) for central lung tumors: Plan quality and long-term clinical outcomes. *Radiother Oncol.* 2015;117:64-70.
- [14] Rulach R, McLoone P, Lumsden G, McKay S, MacLaren V, Macphee J, et al. Toxicity and Efficacy of Stereotactic Ablative Body Radiotherapy for Moderately Central Non-small Cell Lung Cancers Using 50 Gy in Five Fractions. *Clin Oncol (R Coll Radiol).* 2020;32:250-8.

- [15] Olsen JR, Robinson CG, El Naqa I, Creach KM, Drzymala RE, Bloch C, et al. Dose-response for stereotactic body radiotherapy in early-stage non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys.* 2011;81:e299-303.
- [16] Bezjak A, Paulus R, Gaspar LE, Timmerman RD, Straube WL, Ryan WF, et al. Safety and Efficacy of a Five-Fraction Stereotactic Body Radiotherapy Schedule for Centrally Located Non-Small-Cell Lung Cancer: NRG Oncology/RTOG 0813 Trial. *J Clin Oncol.* 2019;37:1316-25.
- [17] Chaudhuri AA, Tang C, Binkley MS, Jin M, Wynne JF, von Eyben R, et al. Stereotactic ablative radiotherapy (SABR) for treatment of central and ultra-central lung tumors. *Lung Cancer.* 2015;89:50-6.
- [18] Arnett ALH, Mou B, Owen D, Park SS, Nelson K, Hallemeier CL, et al. Long-term Clinical Outcomes and Safety Profile of SBRT for Centrally Located NSCLC. *Adv Radiat Oncol.* 2019;4:422-8.
- [19] Rowe BP, Boffa DJ, Wilson LD, Kim AW, Detterbeck FC, Decker RH. Stereotactic body radiotherapy for central lung tumors. *J Thorac Oncol.* 2012;7:1394-9.
- [20] Chang JY, Li QQ, Xu QY, Allen PK, Rebuena N, Gomez DR, et al. Stereotactic ablative radiation therapy for centrally located early stage or isolated parenchymal recurrences of non-small cell lung cancer: how to fly in a "no fly zone". *Int J Radiat Oncol Biol Phys.* 2014;88:1120-8.
- [21] Bral S, Gevaert T, Linthout N, Versmessen H, Collen C, Engels B, et al. Prospective, risk-adapted strategy of stereotactic body radiotherapy for early-stage non-small-cell lung cancer: results of a Phase II trial. *Int J Radiat Oncol Biol Phys.* 2011;80:1343-9.
- [22] Fakiris AJ, McGarry RC, Yiannoutsos CT, Papiez L, Williams M, Henderson MA, et al. Stereotactic body radiation therapy for early-stage non-small-cell lung carcinoma: four-year results of a prospective phase II study. *Int J Radiat Oncol Biol Phys.* 2009;75:677-82.
- [23] Lester J, Courtier N, Eswar C, Mohammed N, Fenwick J, Griffiths G, et al. Initial results of the phase Ib/II, I-START trial: Isotoxic accelerated radiotherapy for the treatment of stage II-IIIb NSCLC. *Journal of Clinical Oncology.* 2018;36:e20551-e.
- [24] Zeng K, Poon I, Ung Y, Zhang L, Cheung P. Accelerated Hypofractionated Radiation Therapy for Centrally Located Lung Tumors Not Suitable for Stereotactic Body Radiation Therapy (SBRT) or Concurrent Chemoradiotherapy (CRT). *International Journal of Radiation Oncology, Biology, Physics* 2018. p. e719-e20.

Author contributions

- 1 guarantor of integrity of the entire study - CFF
- 2 study concepts and design - CFF
- 3 literature research – all authors
- 4 clinical studies – N/A
- 5 experimental studies / data analysis – N/A
- 6 statistical analysis – N/A
- 7 manuscript preparation – all authors
- 8 manuscript editing – all authors

*Declaration of Interest Statement

Corinne Faivre-Finn reports grants from AstraZeneca, and grants from Elektra during the conduct of the study. Fiona McDonald reports speaker fees and consulting fees from AstraZeneca, speaker fees from Elektra, and consulting fees from Accuray outside the study. The other authors have nothing to disclose.