



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/216579/>

Version: Accepted Version

Article:

Gould, Victoria, Yang, Dandan, Ruskuc, Nik et al. (2025) Graph products of residually finite monoids are residually finite. *Journal of Algebra*. 662. pp. 514-527. ISSN: 1090-266X

<https://doi.org/10.1016/j.jalgebra.2024.07.057>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

GRAPH PRODUCTS OF RESIDUALLY FINITE MONOIDS ARE RESIDUALLY FINITE

JUNG WON CHO, VICTORIA GOULD, NIK RUŠKUC, AND DANDAN YANG¹

ABSTRACT. We show that any graph product of residually finite monoids is residually finite. As a special case we obtain that any free product of residually finite monoids is residually finite. The corresponding results for graph products of semigroups follow.

1. INTRODUCTION

An algebra A is *residually finite* if for any $a, b \in A$ with $a \neq b$ there is a finite algebra B and a morphism $\theta: A \rightarrow B$ such that $a\theta \neq b\theta$. Residual finiteness is a finitary property for algebras, in that clearly any finite algebra is residually finite. We are concerned here with residual finiteness for monoids, in particular, the preservation of that property under taking free products and, more generally, graph products.

The free product of a set of disjoint monoids $\mathcal{M} = \{M_\alpha : \alpha \in V\}$ is the monoid freely generated by $\bigcup_{\alpha \in V} M_\alpha$. It contains a copy of each M_α , and these copies do not have any non-identity elements in common. For a simple graph Γ with set of vertices V the graph product of \mathcal{M} with respect to Γ is the monoid freely generated by $\bigcup_{\alpha \in V} M_\alpha$, subject to the additional constraints that elements of M_α and M_β commute whenever there is an edge joining α and β in Γ [12, 2]. Again, the graph product contains a copy of each M_α , and these copies do not have any non-identity elements in common. We usually refer to the monoids in \mathcal{M} as vertex monoids. If the graph has no edges, then the graph product construction returns the free product. For finite complete graphs, the graph product simply yields the direct product. We give a formal definition in terms of presentations in Section 2.

For general reasons, it is true that the direct product $A \times B$ of two residually finite algebras is residually finite; in fact the residually finite algebras of the same type form a prevariety, the point of view adopted in [15]. The converse is not true in general, but is for monoids, and indeed whenever A and B are contained in $A \times B$; the converse is also

2020 *Mathematics Subject Classification.* 20M10, 20M05.

Key words and phrases. Monoid, free products, graph products, residual finiteness.

The second author is supported by the Engineering and Physical Sciences Research Council Grant EP/V002953/1. The third author is supported by the Engineering and Physical Sciences Research Council Grant EP/V003224/1. The fourth author is supported by the National Natural Science Foundation of China Grant No. 12171380, the Natural Science Basic Research Program of Shaanxi Province Grant No. 2023-JC-JQ-04, and the Shaanxi Fundamental Science Research Project for Mathematics and Physics Grant No. 22JSQ034.

¹ The corresponding author.

true for semigroups for somewhat non-obvious reasons [8]. The situation for residual finiteness of free products is exactly reverse: if the free product $A * B$ is residually finite, then so are A and B , because they are contained in $A * B$. The converse is not true in general, is true for monoids, but we have not been able to find it in literature. It is a special case of our main theorem in this paper. Green [9] showed that graph products of groups are residually finite if and only if each vertex group is residually finite; the proof relies on group-theoretic arguments, not available for arbitrary monoids.

The thrust of this paper is to prove the following result (see Theorem 5.1).

Theorem. *Any graph product of monoids is residually finite if and only if each vertex monoid is residually finite.*

We present a number of corollaries, including the corresponding result for semigroups.

The structure of this paper is as follows. In Section 2 we set out the background notions needed for this article, including the notion of left Foata normal form for elements of graph products. Section 3 introduces the notion of block length of words and elements of graph products, and examines the behaviour of block length under composition. In Section 4 we turn our attention to a special case of our main theorem and show that graph products of finite monoids are residually finite. We then are able to complete our main theorem, and the desired corollaries, in Section 5.

2. PRELIMINARIES

To keep this article as self-contained as possible, here we run through the main concepts required. For more details, we recommend [1, 10].

2.1. Monoid presentations. Let X be a set. The *free monoid* X^* on X consists of all words over X with operation of juxtaposition. We denote a non-empty word by $x_1 \circ \cdots \circ x_n$ where $x_i \in X$ for $1 \leq i \leq n$; here n is the *length* of the word. We also use \circ for juxtaposition of words. The empty word is denoted by ϵ and is the identity of X^* . Throughout, our convention is that if we say $x_1 \circ \cdots \circ x_n \in X^*$, then we mean that $x_i \in X$ for all $1 \leq i \leq n$, unless we explicitly say otherwise.

A *monoid presentation* $\langle X \mid R \rangle$, where X is a set and $R \subseteq X^* \times X^*$, determines the monoid X^*/R^\sharp , where R^\sharp is the congruence on X^* generated by R . In the usual way, we identify $(u, v) \in R$ with the formal equality $u = v$ in a presentation $\langle X \mid R \rangle$. We denote elements of the quotient by $[w]$, where $w \in X^*$.

2.2. Graph products of monoids. In this article, all graphs are simple, that is, they are undirected with no multiple edges or loops. Let $\Gamma = (V, E)$ be a graph; here V is the non-empty set of *vertices* and E is the set of *edges* of Γ . We denote an edge joining vertices α and β by (α, β) or (since our graph is undirected) by (β, α) .

Let $\mathcal{M} = \{M_\alpha : \alpha \in V\}$ be a set of monoids. The *direct product* $\prod_{\alpha \in V} M_\alpha$ of \mathcal{M} is the monoid where the underlying set consists of all functions $f: V \rightarrow \bigcup_{\alpha \in V} M_\alpha$ such that $\alpha f \in M_\alpha$ for all $\alpha \in V$ and the multiplication is defined to be $fg: V \rightarrow \bigcup_{\alpha \in V} M_\alpha$

such that $\alpha(fg) = (\alpha f)(\alpha g)$ for all $\alpha \in V$. The *restricted direct product* of \mathcal{M} is the submonoid of $\prod_{\alpha \in V} M_\alpha$ consisting of those elements $f: V \rightarrow \bigcup_{\alpha \in V} M_\alpha$ such that $|\{\alpha \in V: \alpha f \neq 1_\alpha\}| < \infty$. We note that if $|V| < \infty$, then the two products coincide. The definition of the free product of the monoids in the set $\mathcal{M} = \{M_\alpha: \alpha \in V\}$, and its universal properties, may be found in [1] and [10]. We do not give details here since free products are immediately obtained as a special case of the following.

Definition 2.1 ([9, 2]). Let $\Gamma = (V, E)$ be graph and let $\mathcal{M} = \{M_\alpha: \alpha \in V\}$ be a set of mutually disjoint monoids. For each $\alpha \in V$, we write 1_α for the identity of M_α and put $I = \{1_\alpha: \alpha \in V\}$. The *graph product* $\mathcal{G}\mathcal{P} = \mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$ of \mathcal{M} with respect to Γ is defined by the monoid presentation

$$\mathcal{G}\mathcal{P} = \langle X \mid R \rangle$$

where $X = \bigcup_{\alpha \in V} M_\alpha$ and the relations in $R = R_{\text{id}} \cup R_v \cup R_e$ are given by:

$$\begin{aligned} R_{\text{id}} &= \{1_\alpha = \epsilon: \alpha \in V\}, \\ R_v &= \{x \circ y = xy: x, y \in M_\alpha, \alpha \in V\}, \\ R_e &= \{x \circ y = y \circ x: x \in M_\alpha, y \in M_\beta, (\alpha, \beta) \in E\}. \end{aligned}$$

The monoids M_α in Definition 2.1 are referred to as *vertex monoids*. We may also say for brevity that $\mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$ is the *graph product of the monoids* M_α , $\alpha \in V$. We denote the R^\sharp -class of $u \in X^*$ by $[u]$.

Remark 2.2. Let $\mathcal{M} = \{M_\alpha: \alpha \in V\}$ be a set of mutually disjoint monoids.

- If $E = \emptyset$, then the graph product $\mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$ is the free product of the monoids in \mathcal{M} .
- If $E = V \times V$, then the graph product $\mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$ is isomorphic to the restricted direct product of the monoids in \mathcal{M} , and so to the direct product in the case where $|V| < \infty$.
- If the monoids in \mathcal{M} are all free monoids, then $\mathcal{G}\mathcal{P}$ is a *trace monoid* (see, for example, [4]).
- If the monoids in \mathcal{M} are all free groups, then $\mathcal{G}\mathcal{P}$ is a *free partially commutative group*, or, in the case where Γ is finite and each vertex group is finitely generated, a *right angled Artin group* (see, for example, [5]).

For the remainder of this section we consider the monoid $\mathcal{G}\mathcal{P} = \mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$, and give some of its properties that will be needed in the proof of our main result. We begin with a couple of universal properties.

Lemma 2.3. ([6, Proposition 1.7]) *Suppose $\{N_\alpha: \alpha \in V\}$ is another set of mutually disjoint monoids, and $\phi_\alpha: M_\alpha \rightarrow N_\alpha$, $\alpha \in V$, are morphisms. Then there is a unique morphism $\phi: \mathcal{G}\mathcal{P}(\Gamma, \mathcal{M}) \rightarrow \mathcal{G}\mathcal{P}(\Gamma, \mathcal{N})$ such that $m_\alpha \phi = m_\alpha \phi_\alpha$ for all $\alpha \in V$ and $m_\alpha \in M_\alpha$. \square*

Lemma 2.4. ([3, Proposition 2.3]) *Let $V' \subseteq V$, let $\Gamma' = (V', E')$ be the induced subgraph of Γ on V' , and let $\mathcal{M}' = \{M_\alpha: \alpha \in V'\}$. Then $\mathcal{G}\mathcal{P}(\Gamma', \mathcal{M}')$ is a retract of $\mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$. \square*

2.3. Reduced words.

Definition 2.5. We refer to an application of a relation in R_e to a word $w \in X^*$ as a *shuffle*. Two words in X^* are *shuffle equivalent* if one can be obtained from the other by shuffles.

Definition 2.6. We refer to an application of a relation in R to a word $w \in X^*$ that reduces the length of w as a *reduction*. A word $w \in X^*$ is *reduced* if it is not possible to apply a reduction step to any word shuffle equivalent to w .

Note that a reduction is either an application of a relation from R_{id} replacing 1_α by ϵ , or an application of a relation from R_v replacing $x \circ y$ by xy .

The following is key. It appears for groups in [9]; several subsequent papers have noted the result for monoids e.g. [2, 6].

Theorem 2.7. ([2, Theorem 5.5]) *Every element of X^* is equivalent to a reduced word. If w, v are reduced words, then w is equivalent to v if and only if w and v are shuffle equivalent.* \square

Definition 2.8. If $u, w \in X^*$ with $[u] = [w]$ and such that w is reduced, then we say that w is a *reduced form* of u .

It follows from Theorem 2.7 that all reduced forms of $u \in X^*$ are shuffle equivalent and hence have the same length as words in X^* . We now proceed to show that amongst the reduced forms of u we can pick words with useful properties.

2.4. Foata normal form.

Definition 2.9. Let $w = x_1 \circ \dots \circ x_n \in X^*$ with $x_i \in M_{\alpha_i}$ for all $1 \leq i \leq n$. We call the set $\{\alpha_1, \dots, \alpha_n\}$ the *support* of w and denote it by $\text{supp}(w)$. If $\text{supp}(w)$ is a singleton, i.e. $\text{supp}(w) = \{\alpha\}$, then for the sake of notational convenience we may identify $\text{supp}(w)$ with the vertex α .

Definition 2.10. A reduced word $w \in X^*$ is called a *complete block* if the subgraph of Γ induced by $\text{supp}(w)$ is complete.

Remark 2.11. A word $w \in X^*$ is a complete block if and only if $\text{supp}(w)$ is a complete subgraph, no letters in w are from I and for each $\alpha \in \text{supp}(w)$ there exists a unique letter contained in w with support α .

Definition 2.12. A word $w \in X^*$ is a *left Foata normal form* with blocks $w_i \in X^*$, $1 \leq i \leq k$, if the following three conditions are satisfied:

- (i) $w = w_1 \circ \dots \circ w_k$ is a reduced word;
- (ii) w_i is a complete block for all $1 \leq i \leq k$;
- (iii) for any $1 \leq i < k$ and $\alpha \in \text{supp}(w_{i+1})$, there is some $\beta \in \text{supp}(w_i)$ such that $(\alpha, \beta) \notin E$.

If $[u] = [w]$ where w is a left Foata normal form, then we may say w is a *left Foata normal form* of u . Sometimes we will abbreviate the term left Foata normal form to *LFNF*.

Note that the empty word ϵ is a LFNF by definition.

Theorem 2.13 ([3]). *Every element w of X^* has a LFNF $w_1 \circ w_2 \circ \cdots \circ w_k$, with blocks w_i for $1 \leq i \leq k$.*

If $w'_1 \circ w'_2 \circ \cdots \circ w'_h$ is any LFNF of w with blocks w'_j for $1 \leq j \leq h$, then $k = h$ and $[w_i] = [w'_i]$ for $1 \leq i \leq k$. \square

Theorem 2.13 allows us to define the *left block length* $\ell_\Gamma^l(u)$ of $u \in X^*$ as the number of blocks of any LFNF of u ; where the graph Γ is understood, we write more simply $\ell^l(u)$. The notion of *right Foata normal form* is defined in a dual way and we may define the *right block length* $\ell_\Gamma^r(u)$, or $\ell^r(u)$ of $u \in X^*$. In fact, these are equal, and equal to a further useful parameter, as we will show in the next section.

3. BLOCK LENGTH

Throughout this section we consider a graph product $\mathcal{G}\mathcal{P} = \mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$.

Proposition 3.1. *Let u be a reduced word written as a product $u = u_1 \circ \cdots \circ u_n$ such that each u_i is a complete block. Then $\ell^l(u) \leq n$.*

Proof. We suppose that the assertion is not true, and pick a counter-example $u = u_1 \circ \cdots \circ u_n$ where n is as small as possible, and, for that n , the length of u_n is as small as possible. Let $p_1 \circ \cdots \circ p_m$ be a LFNF of $u_1 \circ \cdots \circ u_{n-1}$ with blocks p_1, \dots, p_m . Then $m \leq n - 1$. Consider $p_1 \circ \cdots \circ p_m \circ u_n$. If $m < n - 1$ we obtain a contradiction by the minimality of n . If $m = n - 1$ and there is a letter x in u_n such that $\text{supp}(x)$ is adjacent to all vertices in $\text{supp}(p_m)$, then we have a contradiction to the minimality of u_n . In the remaining case, $p_1 \circ \cdots \circ p_m \circ u_n$ is a LFNF of u and so $\ell^l(u) = n$, which contradicts the choice of counter-example. The result follows. \square

The dual to Proposition 3.1 holds for right block length.

From Proposition 3.1 and its dual we deduce the main result of this section.

Theorem 3.2. *Let $w \in X^*$ be a reduced word. Then $\ell^l(w) = \ell^r(w)$ and is the least number of complete blocks whose product is equivalent to w .*

Definition 3.3. For any reduced word $w \in X^*$ we write $\ell(w)$ for $\ell^l(w) = \ell^r(w)$ and refer to $\ell(w)$ as the *block length* of w . More generally, for any $u \in X^*$ we define $\ell(u) = \ell(w)$ where w is any reduced form of u , and again refer to $\ell(u)$ as the *block length* of u .

We note in passing that for the case of graph products of semigroups (which we will consider in Section 5) the fact that $\ell^l(w) = \ell^r(w)$ appears in [16, Lemma 3.8].

We now want to examine what happens to the block length of a word when multiplied by a single letter, for which we need a bit of preparation.

Lemma 3.4. *Let $x_1 \circ \cdots \circ x_n \in X^+$ be a reduced word. If there exists $1 \leq k \leq n$ such that $(\text{supp}(x_k), \text{supp}(x_j)) \in E$ for all $k < j \leq n$, then $x_1 \circ \cdots \circ x_{k-1} \circ x_{k+1} \circ \cdots \circ x_n$ is also reduced.*

Proof. Clearly, $x_1 \circ \cdots \circ x_n$ is shuffle equivalent to $x_1 \circ \cdots \circ x_{k-1} \circ x_{k+1} \circ \cdots \circ x_n \circ x_k$, implying that $x_1 \circ \cdots \circ x_{k-1} \circ x_{k+1} \circ \cdots \circ x_n \circ x_k$ is reduced. Hence so is $x_1 \circ \cdots \circ x_{k-1} \circ x_{k+1} \circ \cdots \circ x_n$. \square

Lemma 3.5. *Let $w = w_1 \circ \cdots \circ w_k$ be a LFNF with blocks w_i , $1 \leq i \leq k$. Let $1 \leq j \leq k$ and let $t \in M_\alpha$ be a letter appearing in w_j .*

(i) *Suppose that $(\alpha, \beta) \in E$ for all $\beta \in \text{supp}(w_{j+1} \circ \cdots \circ w_k)$, and let w'_j be obtained from w_j by deleting t . If $w'_j \neq \epsilon$, then $w_1 \circ \cdots \circ w_{j-1} \circ w'_j \circ w_{j+1} \circ \cdots \circ w_k$ is a LFNF with blocks $w_1, \dots, w_{j-1}, w'_j, w_{j+1}, \dots, w_k$. Further, $w'_j = \epsilon$ can only occur if $j = k$.*

(ii) *Let w'_j be obtained from w_j by replacing t by any $z \in M_\alpha \setminus \{1_\alpha\}$. Then $w_1 \circ \cdots \circ w_{j-1} \circ w'_j \circ w_{j+1} \circ \cdots \circ w_k$ is a LFNF with blocks $w_1, \dots, w_{j-1}, w'_j, w_{j+1}, \dots, w_k$.*

Proof. We begin by remarking that t is the unique letter of w_j with support α .

(i) We first verify the second claim. Suppose that $j < k$ and $w'_j = \epsilon$, that is, $w_j = t$. It follows that

$$w = w_1 \circ \cdots \circ w_{j-1} \circ (t \circ w_{j+1}) \circ \cdots \circ w_k$$

is a decomposition of w into $k - 1$ complete blocks, contradicting Theorem 3.2. Hence, if $w'_j = \epsilon$, then $j = k$.

Now suppose that $w'_j \neq \epsilon$. By Lemma 3.4, deleting the letter t , and bearing in mind that w_j is a complete block, we obtain that $w_1 \circ \cdots \circ w_{j-1} \circ w'_j \circ w_{j+1} \circ \cdots \circ w_k$ is reduced. Since w is a LFNF, for each $\beta \in \text{supp}(w_{j+1})$, there exists $\delta \in \text{supp}(w_j)$ such that $(\beta, \delta) \notin E$. Further, as $(\text{supp}(t), \gamma) \in E$ for all $\gamma \in \text{supp}(w_{j+1} \circ \cdots \circ w_k)$, we deduce that $\delta \neq \text{supp}(t)$, so that $\delta \in \text{supp}(w'_j)$, implying that $w_1 \circ \cdots \circ w_{j-1} \circ w'_j \circ w_{j+1} \circ \cdots \circ w_k$ is a LFNF.

(ii) The proof is immediate. \square

We can now state and prove three results which describe the effect on a LFNF and block length of multiplication by a single letter.

Lemma 3.6. *Let $w = w_1 \circ \cdots \circ w_k$ be a LFNF with blocks w_i , $1 \leq i \leq k$, and let $x \in X \setminus I$. Then $\ell(w \circ x) = k + 1$ if and only if $\text{supp}(w_k \circ x)$ is an incomplete subgraph, in which case*

$$w_1 \circ \cdots \circ w_k \circ x$$

is a LFNF for $w \circ x$ with blocks w_1, \dots, w_k, x .

Proof. Suppose $\text{supp}(x) = \{\alpha\}$, i.e. $x \in M_\alpha$. If $\text{supp}(w_k \circ x)$ is incomplete, then $\alpha \notin \text{supp}(w_k)$ and there exists $\beta \in \text{supp}(w_k)$ such that $(\alpha, \beta) \notin E$, implying that $w \circ x$ is reduced. Further, $w_1 \circ \cdots \circ w_k \circ x$ is a LFNF with blocks w_1, \dots, w_k, x . Conversely, if $\text{supp}(w_k \circ x)$ is complete, then from Theorem 3.2, $\ell(w \circ x) \leq k$. \square

Lemma 3.7. *Let $w = w_1 \circ \cdots \circ w_k$ be a LFNF with blocks w_i , $1 \leq i \leq k$, and let $x \in M_\alpha \setminus \{1_\alpha\}$. Suppose that $\text{supp}(w_k \circ x)$ is a complete subgraph.*

(i) *We have that $\ell(w \circ x) = k - 1$ if and only if $\text{supp}(w_k) = \{\alpha\}$ and $w_k x = 1_\alpha$ in M_α , in which case $w_1 \circ \cdots \circ w_{k-1}$ is a LFNF for $w \circ x$ with blocks w_i , $1 \leq i \leq k - 1$.*

(ii) Otherwise, $\ell(w \circ x) = k$ and $w \circ x$ has a LFNF $w_1 \circ \cdots \circ w_{i-1} \circ w'_i \circ w_{i+1} \circ \cdots \circ w_k$ with blocks $w_1, \dots, w_{i-1}, w'_i, w_{i+1}, \dots, w_k$, where $1 \leq i \leq k$ and $w'_i \neq \epsilon$ is a reduced form of $w_i \circ x$.

Proof. If $w \circ x$ is not reduced, then there exists $1 \leq i \leq k$ and a letter y in w_i such that $y \in M_\alpha$ and $(\alpha, \beta) \in E$ for all $\beta \in \text{supp}(w_{i+1} \circ \cdots \circ w_k)$. Let w'_i be a reduced form of $w_i \circ x$. If $yx = 1_\alpha$, then w'_i is obtained by deleting y from w_i . If $yx \neq 1_\alpha$, then w'_i is obtained by replacing y by yx in w_i .

By Lemma 3.5, the case $w'_i = \epsilon$ can only occur if $i = k$. In this case, $w_1 \circ \cdots \circ w_{k-1}$ is a LFNF for $w \circ x$ with blocks w_1, \dots, w_{k-1} . Otherwise, $w'_i \neq \epsilon$, and from Lemma 3.5 $w \circ x$ has a LFNF

$$w_1 \circ \cdots \circ w_{i-1} \circ w'_i \circ w_{i+1} \circ \cdots \circ w_k$$

with blocks $w_1, \dots, w_{i-1}, w'_i, w_{i+1}, \dots, w_k$.

If $w \circ x$ is reduced, then $\alpha \notin \text{supp}(w_k)$. Further, as $\text{supp}(w_k \circ x)$ is complete, we have that $(\alpha, \beta) \in E$ for all $\beta \in \text{supp}(w_k)$. Let $1 \leq i \leq k$ be the smallest such that $(\alpha, \beta) \in E$ for all $\beta \in \text{supp}(w_i \circ \cdots \circ w_k)$. Then

$$w_1 \circ \cdots \circ w_{i-1} \circ (w_i \circ x) \circ w_{i+1} \circ \cdots \circ w_k$$

is a LFNF of $w \circ x$ with blocks $w_1, \dots, w_{i-1}, w_i \circ x, w_{i+1}, \dots, w_k$. □

Corollary 3.8. *Let w be a LFNF of block length k and let $x, y \in M_\alpha \setminus \{1_\alpha\}$. Then $\ell(w \circ x) \leq k$ if and only if $\ell(w \circ y) \leq k$.* □

4. THE FINITE CASE

The main step in the proof of our main theorem is to deal with the special case of a graph product $\mathcal{G}\mathcal{P} = \mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$ where both the graph Γ and all the vertex monoids in \mathcal{M} are finite. This is the subject of this section, and so these assumptions will apply throughout.

In outline, we will define actions of the vertex monoids of $\mathcal{G}\mathcal{P}$ on a finite set F_k . These actions will extend to an action of $\mathcal{G}\mathcal{P}$ itself, and hence they will induce a morphism from $\mathcal{G}\mathcal{P}$ to the finite full transformation monoid on F_k . We will then show how to pick a suitable k so that the morphism separates pairs of distinct elements of $\mathcal{G}\mathcal{P}$ having block length no greater than k .

Before we embark on this, we briefly review the basic definitions and notation for actions.

Definition 4.1. Let M be a monoid and let A be a non-empty set. An *action* of M on A is a map $* : A \times M \rightarrow A$, where $(a, m) \mapsto a * m$, such that for all $a \in A$ and $s, t \in M$ we have $a * 1 = a$ and $(a * s) * t = a * st$.

An action of M on A corresponds precisely to a monoid morphism $M \rightarrow \mathcal{T}_A$, where \mathcal{T}_A is the full transformation monoid of all maps from A to A . Further details may be found in the standard text [11].

Lemma 4.2. *Let $k \in \mathbb{N}$ and let*

$$F_k = \{[w] \in \mathcal{GP} : \ell(w) \leq k\}.$$

Each vertex monoid M_α acts on F_k via:

$$[w] * x = \begin{cases} [w \circ x] & \text{if } \ell(w \circ x) \leq k, \\ [w] & \text{if } \ell(w \circ x) = k + 1. \end{cases}$$

Proof. Let $m \leq k$ and let $w = w_1 \circ \cdots \circ w_m$ be a LFNF with blocks w_i , $1 \leq i \leq m$.

Clearly, $[w \circ 1_\alpha] = [w]$, so that $\ell(w \circ x) = \ell(w) \leq k$, and hence

$$[w] * 1_\alpha = [w \circ 1_\alpha] = [w].$$

Let $x, y \in M_\alpha$. We must show that $([w] * x) * y = [w] * xy$.

If $x = 1_\alpha$ or $y = 1_\alpha$, then the claim is clearly true. Assume therefore that $x, y \neq 1_\alpha$.

By Lemmas 3.6 and 3.7 we have that $\ell(w \circ x) \leq m + 1$. Hence, if $m \leq k - 1$, then $\ell(w \circ x), \ell(w \circ y)$ and $\ell(w \circ x \circ y) = \ell(w \circ xy)$ are all no greater than $m + 1 \leq k$, so that in this case,

$$([w] * x) * y = [w \circ x] * y = [w \circ x \circ y] = [w \circ (xy)] = [w] * (xy).$$

Suppose that $m = k$. If $\ell(w \circ x) = k + 1$, then by Corollary 3.8 we have that $\ell(w \circ y) = k + 1$, and if $xy \neq 1_\alpha$, then also $\ell(w \circ xy) = k + 1$. Hence, whether or not we have $xy \neq 1_\alpha$, we deduce

$$([w] * x) * y = [w] * y = [w] = [w] * (xy). \quad \square$$

The above action of each M_α induces a morphism from M_α to \mathcal{T}_{F_k} . We now aim to show that this induces a morphism from the entire \mathcal{GP} to \mathcal{T}_{F_k} . The key step is the following lemma, which shows that the defining relations R_e are respected by the actions of the M_α .

With some abuse of notation we use $*$ for the action of any M_α on F_k . It will always be clear from the context to which specific action $*$ refers.

Lemma 4.3. *If $x \in M_\alpha$ and $y \in M_\beta$ and $(\alpha, \beta) \in E$, then*

$$([w] * x) * y = ([w] * y) * x$$

for all $[w] \in F_k$.

Proof. Without loss of generality we may assume that $w = w_1 \circ \cdots \circ w_m$ is a LFNF with blocks w_i , $1 \leq i \leq m \leq k$. Clearly, the result is true if either $x = 1_\alpha$ or $y = 1_\beta$.

Suppose that $x, y \notin I$. We split our considerations into cases depending on $\ell(w \circ x)$ and $\ell(w \circ y)$.

Case 1: $\ell(w \circ x) = k + 1$ and $\ell(w \circ y) = k + 1$. We have $[w] * x = [w]$ and $[w] * y = [w]$, and so

$$([w] * x) * y = [w] * y = [w] = [w] * x = ([w] * y) * x.$$

Case 2: $\ell(w \circ x) \leq k$ and $\ell(w \circ y) = k + 1$. This time $[w] * x = [w \circ x]$ and $[w] * y = [w]$. It follows from Lemmas 3.6 and 3.7 that $m = \ell(w) = k$ and that $\text{supp}(w_k \circ y)$ is incomplete. If $\ell(w \circ x) = k - 1$, then, by Lemma 3.7, we have $\text{supp}(w_k) = \{\alpha\}$ and $w_k x = 1_\alpha$. On the other hand, as $\ell(w \circ y) = k + 1$, we have that $\text{supp}(w_k \circ y)$ is incomplete, again, by Lemma 3.7. But this contradicts the assumption that $(\alpha, \beta) \in E$. Therefore, $\ell(w \circ x) = k$. It follows from Lemma 3.7 that $w \circ x$ has a LFNF

$$w_1 \circ \cdots \circ w_{i-1} \circ (w_i \circ x)' \circ w_{i+1} \circ \cdots \circ w_k$$

$(w_i \circ x)' \neq \epsilon$ is a reduced form of $w_i \circ x$ for some $1 \leq i \leq k$. If $i < k$, then we know $\text{supp}(w_k \circ y)$ is incomplete. If $i = k$, then $\text{supp}(w'_k)$ is equal to one of $\text{supp}(w_k)$, $\text{supp}(w_k) \cup \{\alpha\}$ or $\text{supp}(w_k) \setminus \{\alpha\}$; but as $(\alpha, \beta) \in E$ we have in any of these cases that $\text{supp}(w'_k \circ y)$ is incomplete. It follows that $\ell(w \circ x \circ y) = k + 1$. Then

$$([w] * x) * y = [w \circ x] * y = [w \circ x] = [w] * x = ([w] * y) * x.$$

Case 3: $\ell(w \circ x) = k + 1$ and $\ell(w \circ y) \leq k$. This is dual to Case 2.

Case 4: $\ell(w \circ x) \leq k$ and $\ell(w \circ y) \leq k$. We have $[w] * x = [w \circ x]$ and $[w] * y = [w \circ y]$. We claim that

$$\ell(w \circ x \circ y) = \ell(w \circ y \circ x) \leq m + 1 \leq k. \quad (1)$$

To see this, first suppose that $m < k$. Then $[w \circ x \circ y] = [w_1 \circ \cdots \circ w_m \circ (x \circ y)] = [w_1 \circ \cdots \circ w_m \circ (y \circ x)] = [w \circ y \circ x]$. Certainly $\text{supp}(x \circ y)$ is a complete block since $(\alpha, \beta) \in E$, and (1) follows by Theorem 3.2. Next, suppose that $m = k$. By Lemma 3.7 we know that $\text{supp}(w_k \circ x)$ and $\text{supp}(w_k \circ y)$ are complete graphs. Then, again since $(\alpha, \beta) \in E$, we have that $\text{supp}(w_k \circ x \circ y)$ and $\text{supp}(w_k \circ y \circ x)$ are complete graphs, so their (common) reduced form is a complete block. Hence (1) follows using Theorem 3.1.

Having established (1), we have

$$([w] * x) * y = [w \circ x] * y = [w \circ x \circ y] = [w \circ y \circ x] = [w \circ y] * x = ([w] * y) * x,$$

completing the proof of this case and the lemma. \square

Let us denote by θ_α the morphism $M_\alpha \rightarrow \mathcal{T}$ induced by the action of M_α on F_k ; thus

$$[w](x\theta_\alpha) = [w] * x \quad ([w] \in F_k, x \in M_\alpha).$$

We let $\theta : X \rightarrow \mathcal{T}_{F_k}$ be $\bigcup_{\alpha \in V} \theta_\alpha$.

Proposition 4.4. *Let $k \in \mathbb{N}$. Then there is a morphism $\Theta : \mathcal{G}\mathcal{P} \rightarrow \mathcal{T}_{F_k}$ given by*

$$[w]([u]\Theta) = [w](x_1\theta)(x_2\theta) \dots (x_m\theta) \quad \text{where } u = x_1 \circ x_2 \circ \cdots \circ x_m \in X^*.$$

Proof. Since X^* is free on X , the map θ lifts to a morphism $\theta' : X^* \rightarrow F_k$, where $(x_1 \circ x_2 \circ \cdots \circ x_m)\theta' = (x_1\theta)(x_2\theta) \dots (x_m\theta)$. To show that θ' induces Θ as given, we need only show that the relations $R = R_{\text{id}} \cup R_\vee \cup R_e \subseteq \ker \theta'$. The fact that $R_{\text{id}} \cup R_\vee \subseteq \ker \theta'$ follows from Lemma 4.2 and the fact that $R_e \subseteq \ker \theta'$ from Lemma 4.3. \square

Note that in the language of actions, Proposition 4.4 says that $\mathcal{G}\mathcal{P}$ acts on F_k by

$$[w] * [u] = [w] * x_1 * x_2 * \cdots * x_n, \quad \text{where } u = x_1 \circ x_2 \circ \cdots \circ x_m \in X^*.$$

Proposition 4.5. *Any graph product of finite monoids with respect to a finite graph is residually finite.*

Proof. Let $\mathcal{G}\mathcal{P}$ be a finite graph product of finite monoids. Let $[u]$ and $[v]$ be distinct elements of $\mathcal{G}\mathcal{P}$; we may assume that u and v are reduced. Choose k to be the greater of $\ell(u)$ and $\ell(v)$ and consider the morphism $\Theta : \mathcal{G}\mathcal{P} \rightarrow \mathcal{T}_{F_k}$ from Proposition 4.4. It follows from Theorem 3.2 that for any $w \in X^*$ where $u = w \circ t$ or $v = w \circ t$, we have that $\ell(w) \leq k$. Consequently, writing $u = u_1 \circ \cdots \circ u_h \in X^*$, we see that

$$\begin{aligned} [\epsilon]([u]\Theta) &= [\epsilon](u_1\theta \circ \cdots \circ u_h\theta) = [\epsilon] * u_1 * u_2 * \cdots * u_h \\ &= [u_1] * u_2 * \cdots * u_h = [u_1 \circ u_2] * \cdots * u_h = \cdots = [u_1 \circ \cdots \circ u_h] = [u]. \end{aligned}$$

Similarly, $[\epsilon]([v]\Theta) = [v]$ and since $[u] \neq [v]$ we obtain that $[u]\Theta \neq [v]\Theta$, as required. \square

5. THE MAIN RESULTS

We can now prove our main result.

Theorem 5.1. *Any graph product of monoids is residually finite if and only if each vertex monoid is residually finite.*

Proof. Let $\mathcal{G}\mathcal{P} = \mathcal{G}\mathcal{P}(\Gamma, \mathcal{M})$ be a graph product of monoids, with $\Gamma = (V, E)$ and $\mathcal{M} = \{M_\alpha : \alpha \in V\}$. As already mentioned in the Introduction, the ‘only if’ part follows from the fact that all M_α embed into $\mathcal{G}\mathcal{P}$. For the ‘if’ part, suppose that all M_α are residually finite. Let $[v]$ and $[w]$ be distinct elements of $\mathcal{G}\mathcal{P}$. We may assume that $v = x_1 \circ \cdots \circ x_n, w = y_1 \circ \cdots \circ y_m \in X^*$ are reduced. Let $V' = \text{supp}(v) \cup \text{supp}(w)$ and let $Y = \{x_1, \dots, x_n, y_1, \dots, y_m\}$. For each $\alpha \in V'$ the residual finiteness of M_α allows us to choose a finite monoid M'_α and a morphism $\theta_\alpha : M_\alpha \rightarrow M'_\alpha$ which separates the distinct elements of $(M_\alpha \cap Y) \cup \{1_\alpha\}$. For $\alpha \notin V'$ we let $M'_\alpha = M_\alpha$ and let $\theta_\alpha : M_\alpha \rightarrow M_\alpha$ be the identity morphism. Let $\mathcal{G}\mathcal{P}' = \mathcal{G}\mathcal{P}(\Gamma, \mathcal{M}')$ where $\mathcal{M}' = \{M'_\alpha : \alpha \in V\}$. By Lemma 2.3, $\mathcal{G}\mathcal{P}'$ is a morphic image of $\mathcal{G}\mathcal{P}$ via the morphism θ which extends each θ_α .

We claim that $[v]\theta \neq [w]\theta$. Clearly, $[v]\theta = [x_1\theta \circ \cdots \circ x_n\theta]$, $[w]\theta = [y_1\theta \circ \cdots \circ y_m\theta]$ and $x_1\theta \circ \cdots \circ x_n\theta, y_1\theta \circ \cdots \circ y_m\theta$ are reduced words. If $[v]\theta = [w]\theta$, then we would be able to shuffle $x_1\theta \circ \cdots \circ x_n\theta$ to $y_1\theta \circ \cdots \circ y_m\theta$ by Theorem 2.7. Hence, by a corresponding sequence of moves, we could shuffle $x_1 \circ \cdots \circ x_n$ to $y_1 \circ \cdots \circ y_m$, a contradiction. Therefore, $[v]\theta \neq [w]\theta$, as claimed.

Let $\Gamma' = (V', E')$ be the induced subgraph on V' . By Lemma 2.4 there exists a retraction $\phi : \mathcal{G}\mathcal{P}' \rightarrow \mathcal{G}\mathcal{P}''$, where $\mathcal{M}'' = \{M'_\alpha : \alpha \in V'\}$ and $\mathcal{G}\mathcal{P}'' = \mathcal{G}\mathcal{P}(\Gamma', \mathcal{M}'')$. By construction, we have $[v]\theta\phi \neq [w]\theta\phi$.

Note that $\mathcal{G}\mathcal{P}''$ is a graph product of finite monoids with respect to a finite graph. Hence it is residually finite by Proposition 4.5. Thus there exist a finite monoid M and a morphism $\psi : \mathcal{G}\mathcal{P}'' \rightarrow M$ such that $[v]\theta\phi\psi \neq [w]\theta\phi\psi$. But then the composition $\theta\phi\psi$ is a morphism from our original graph product $\mathcal{G}\mathcal{P}$ into the finite monoid M separating $[v]$ and $[w]$, and the theorem is proved. \square

We record the following immediate consequence:

Corollary 5.2. *A free product or a restricted direct product of monoids is residually finite if and only if each vertex monoid is residually finite.* \square

As commented in the Introduction, the result for finite direct products follows from generic considerations. The result for free products does not seem to have appeared in literature. The result for residual finiteness of graph products of groups appears in [9]; in contrast to the argument there, our proof for monoids does not require any tools specific for groups.

We now briefly consider the notion of a graph product of semigroups (see [3]). Here we start with a *semigroup presentation*, which is a quotient of the free *semigroup* $X^+ = X^* \setminus \{\epsilon\}$ on a set X .

Definition 5.3. Let $\Gamma = (V, E)$ be graph and let $\mathcal{S} = \{S_\alpha : \alpha \in V\}$ be a set of mutually disjoint semigroups. The *graph product* $\mathcal{GP} = \mathcal{GP}(\Gamma, \mathcal{S})$ of \mathcal{S} with respect to Γ is defined by the semigroup presentation

$$\mathcal{GP} = \langle X \mid R \rangle$$

where $X = \bigcup_{\alpha \in V} S_\alpha$ and the relations in $R = R_v \cup R_e$ are given as in Definition 2.1.

Corollary 5.4. *Any graph product of semigroups is residually finite if and only if each vertex semigroup is residually finite.*

Proof. Consider a graph product $\mathcal{GP} = \mathcal{GP}(\Gamma, \mathcal{S})$ of semigroups $\mathcal{S} = \{S_\alpha : \alpha \in V\}$. For each $\alpha \in V$ we may embed S_α into a monoid S_α^{\perp} where \perp_α is an adjoined identity. From [3, Proposition 7.3], there is a semigroup embedding of \mathcal{GP} into $\mathcal{GP}' = \mathcal{GP}(\Gamma, \mathcal{M})$ where $\mathcal{M} = \{S_\alpha^{\perp} : \alpha \in V\}$.

If each semigroup S_α is residually finite, then so is each monoid S_α^{\perp} and hence from Theorem 5.1, so is \mathcal{GP}' . But, \mathcal{GP} embeds into \mathcal{GP}' , so that \mathcal{GP} is residually finite.

Conversely, if \mathcal{GP} is residually finite, then as each S_α embeds into \mathcal{GP} , so is each S_α . \square

Again, we obtain the following immediate consequence:

Corollary 5.5 (Golubov [7]). *A free product of semigroups is residually finite if and only if all the vertex semigroups are residually finite.* \square

We also remark that a finite direct product of semigroups is not realised by a graph product, and the notion of restricted direct product cannot be formulated, since there are no given identities. We observe, however, that it is shown in [13] that a finite direct product of semigroups is residually finite if and only if each constituent semigroup is residually finite.

ACKNOWLEDGEMENT

The authors are grateful to Prof. Mikhail Volkov for alerting them to the existence of [15] and for providing them with references [7] and [15].

REFERENCES

- [1] A.H. Clifford, G.B. Preston, *The algebraic theory of semigroups Vol. II*, Amer. Math. Soc. (1967).
- [2] A. Veloso da Costa, On graph products of monoids, *Semigroup Forum* 63 (2001), 247–277.
- [3] Y. Dandan, V. Gould, On graph products of monoids, *J. Algebra* 620 (2023), 113–156.
- [4] V. Diekert, *Combinatorics on traces*, Lecture Notes in Computer Science 454, Springer (1990).
- [5] A.J. Duncan, V. N. Remeslennikov, A.V. Treier, A survey of free partially commutative groups, *J. Phys.: Conf. Ser.* 1441 (2020), 012136.
- [6] J. Fountain, M. Kambites, Graph products of right cancellative monoids, *J. Austral. Math. Soc.* 87 (2009), 227–252.
- [7] É.A. Golubov, Free product and wreath product of finitely approximable semigroups (Russian), *Ural. Gos. Univ. Mat. Zap.* 9 (1971), 3–15.
- [8] R. Gray, N. Ruškuc, On residual finiteness of direct products of algebraic systems, *Monatsh. Math.* 158 (2009), 63–69.
- [9] E.R. Green, *Graph products of groups*, PhD Thesis, University of Leeds (1990).
- [10] J.M. Howie, *Fundamentals of semigroup theory*, Oxford University Press (1995).
- [11] M. Kilp, U. Knauer, A. Mikhalev, *Monoids, Acts, and Categories*, Walter de Gruyter (2000).
- [12] S. Hermiller, J. Meier, Algorithms and geometry for graph products of groups, *J. Algebra* 171 (1995), 230–257.
- [13] P. Mayr, N. Ruškuc, Finiteness properties of direct products of algebraic structures, *J. Algebra* 494 (2018), 167–187.
- [14] R.N. McKenzie, G.F. McNulty, W.F. Taylor, *Algebras, Lattices, Varieties Vol. I*, Wadsworth (1987).
- [15] M. Sapir, An implicative characterisation of prevarieties of semigroups and rings (Russian), *Ural. Gos. Univ. Mat. Zap.* 13 (1982), 121–132.
- [16] D. Yang, H. Li, Centralizers in graph products of semigroups, *Semigroup Forum* 106 (2023), 285–326.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ST ANDREWS, ST ANDREWS, FIFE KY16 9SS, UK.

Email address: `jwc21@st-andrews.ac.uk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK YO10 5DD, UK

Email address: `victoria.gould@york.ac.uk`

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ST ANDREWS, ST ANDREWS, FIFE KY16 9SS, UK.

Email address: `nik.ruskuc@st-andrews.ac.uk`

SCHOOL OF MATHEMATICS AND STATISTICS, XIDIAN UNIVERSITY, XI'AN 710071, P. R. CHINA

Email address: `ddyang@xidian.edu.cn`