

## REVIEW

# Outcomes and care priorities for older people living with frailty and advanced chronic kidney disease: a multi-professional scoping review

HELEN HURST<sup>1</sup>, HANNAH M.L. YOUNG<sup>2,3</sup>, ANDREW C. NIXON<sup>4</sup>, PAULA ORMANDY<sup>5</sup>, ALISON BRETTLER<sup>5</sup>,  
The scoping review author collaborative for outcomes and care priorities for older people with  
ACKD<sup>†</sup>

<sup>1</sup>The University of Manchester, Manchester Academic Health Science Centre, Manchester University NHS Foundation Trust, Manchester, UK

<sup>2</sup>Department of Respiratory Sciences, University of Leicester, Leicester, UK

<sup>3</sup>Department of Research and Innovation, University Hospitals of Leicester NHS Trust, Leicester, UK

<sup>4</sup>Department of Renal Medicine, Lancashire Teaching Hospitals NHS Foundation Trust, Preston, UK

<sup>5</sup>School of Health and Society, University of Salford, Salford, UK

Address correspondence to: Helen Hurst. Tel: 07903259485. Email: [helen.hurst@mft.nhs.uk](mailto:helen.hurst@mft.nhs.uk)

<sup>†</sup>Acknowledgement of Protocol Collaborative Authorship: In addition to the named authors above, the scoping review collaborative for outcomes and care priorities for older people with ACKD members are: Andrew Mooney, Anna Winterbottom, Hilary Bekker, Edwina A. Brown, Fliss E.M. Murtagh, Maria Da Silva-Gane, David Coyle and Rob Finnigan.

---

## Abstract

The growing older population with advanced chronic kidney disease (ACKD stages 4–5) poses a challenge for healthcare worldwide. The high prevalence of frailty and associated adverse health outcomes highlights concerns for management and interventions specific to this population. The aim was to objectively review the evidence relating to older people ( $\geq 65$  years) living with frailty and ACKD. More specifically how frailty is identified, what interventions have been studied and what outcomes have been reported including outcomes important to patients, families and carers. A scoping review was undertaken following the PRISMA-Scr guidelines. Nine databases were searched and a review team of five people followed a process using defined inclusion and exclusion criteria. Data were then analysed to answer the specific questions of the review. The World Health Organization's International Classification of Functioning Disability and Health was used to map outcomes across the domains. A total of 90 studies were included. The most reported frailty measure was the frailty phenotype. The most reported outcomes were mortality, hospitalisation and healthcare utilisation. Health-related quality of life was the most common patient-reported outcome measure. There were few intervention studies and limited evidence of patient and carer perspectives. This scoping review highlights important areas for further research in older people living with frailty and ACKD. This includes a 'gold standard' measure for identifying frailty, interventions and improvements in outcome measures that matter to patients (including studies that focus on carers and carer burden) and priority setting for future research.

**Keywords:** chronic kidney disease (CKD), chronic renal failure, renal replacement therapy, dialysis, transplantation, older people, aged, frailty, comprehensive geriatric assessment

## Key Points

- The prevalence of frailty in advanced chronic kidney disease (ACKD) is greater than in the general population.
  - Understanding care priorities and outcomes relevant to individuals living with frailty and ACKD is important.
  - Providing interventions that reflect improvements in outcome measures that matter to patients is essential.
  - Poor health outcomes in individuals living with frailty and ACKD are of concern.
-

## Introduction

By 2066, there will be a predicted 20.4 million people over the age of 65 years living in the UK [1]. Generally, someone over the age of 65 might be considered an older person accepting also that people can biologically age at different rates [2]. With advancing years come the increased likelihood of living with multimorbidity, frailty and associated risk of adverse outcomes, including falls, hospitalisation, institutionalisation and death [3]. Evidence-based health care services are essential to mitigate these risks. The advanced chronic kidney disease (ACKD) population is ageing rapidly. In all, 17–45% of people accepted for renal replacement therapy (RRT) are now >75 years old [4]. Nephrology services are therefore not exempted from the challenge, with the prevalence of frailty being greater in ACKD than in the general population [5–7]. Action is especially important within nephrology given that interventions such as RRT, which have the potential to prolong life, may not promise improved well-being [8–12].

To inform and direct the development of appropriate and effective management strategies for older people living with frailty and ACKD, we must first understand the care priorities of this group. Moreover, to effectively evaluate proposed interventions, we need to identify outcomes that are important and meaningful to these groups. The James Lind Alliance [13] and the Standardised Outcomes in Nephrology initiative [14] highlight important research priorities and outcomes in ACKD generally; however, neither have explored priorities and outcomes specifically pertinent for older people living with frailty and ACKD.

Using scoping review methodology, our aims were to: (i) identify and map the available evidence relating to older adults living with ACKD and (ii) identify knowledge gaps [15]. Following discussion with a stakeholder group, comprising older people living with ACKD, their caregivers and relatives, we identified four research questions:

1. Is frailty formally identified in studies involving older people living with ACKD and if so what methods are used?
2. What interventions have been reported and how might they enhance both care and outcomes for older people with ACKD?

3. What outcomes have been reported in studies involving older people living with frailty and ACKD?
4. What outcomes matter most to older people living with frailty and ACKD?

## Methods

The review was guided by Arksey and O'Malley's framework [16] and the PRISMA-ScR guidelines (Appendix 1 checklist), as per protocol [17]. The research team provided expertise in frailty, kidney care and scoping review methodology.

Table 1 outlines the inclusion and exclusion criteria.

### Literature searching

Nine databases were searched covering multiple disciplines and grey literature; full list included in the protocol [15]. Search results were managed using Endnote web reference management software (online). Each search strategy was recorded in full (see Appendix 2 that provides an example record of one of the searches). This search approach was replicated across all databases, changing thesaurus terms and search operators as appropriate. Searches were completed in 2019 and updated in 2021.

### Study selection

Five members of the review team used the eligibility criteria to identify potentially relevant studies by title. Three members (H.H., H.M.L.Y. and A.C.N.) then independently sifted by abstract, and subsequently by full text to create the final list of included studies.

### Data charting

The final set of papers included in the review are summarised in Appendix 4. Initial charting was piloted and completed by two reviewers (H.H. and H.M.L.Y.). No authors were contacted for information or clarity. Outcomes were charted using the World Health Organization (WHO)'s International Classification of Functioning Disability and Health (ICF). The ICF provides a comprehensive conceptual framework for understanding and organising health functioning


**Table 1.** Study eligibility criteria

#### Inclusion criteria

- Population: older people (aged  $\geq 65$  years) with ACKD stages 4 and 5 living with frailty, who may be receiving RRT, conservative care or have received a renal transplant, and carers/families of these individuals.
- Studies: all study designs that include older people with ACKD, carers/families in line with the population inclusion criteria.
- Interventions: where interventions are described, all types of interventions will be included.
- Outcomes: examples such as HRQOL, symptom burden, patient experience, functional and cognitive ability, life participation.

#### Exclusion criteria

- Studies not including  $\geq 65$ -year-old population and studies where subgroup analysis was not performed in this age group.
- Studies not published in English.
- Studies prior to 1995.



**Figure 1.** International Classification of Functioning Disability and Health.

and disability (Figure 1) [18], which facilitated the comparison of outcome used between studies and provided a structure for reporting findings [17, 18].

For all studies reporting information on outcomes, details of the measures used and what the selected measure aimed to capture were extracted. For studies that used measures with multiple components, each component was charted. Studies that included questionnaires, each question item was included.

### Synthesis of results

Results were synthesised using descriptive statistics and/or narrative summaries where appropriate. For outcome data, two reviewers familiar with the ICF (H.M.L.Y. and A.C.N.) used published linking rules to map outcomes to the ICF [19, 20]. Meaningful concepts for each measure, component (for composite measure) or question item and responses (for questionnaires) were identified and linked to the most precise category, using the ICF browser [21]. Where this was unclear, an interpretative approach was used, based on the study aims and using supporting literature that described the tests in detail [19]. Agreement on these interpretations was reached via discussion, including decisions regarding whether a measure was related to activity or participation, using operational definitions by Whitehead and Dejeckers [22].

### Results

Ninety papers, from 69 studies, were included in this review (Figure 2). The manuscripts were categorised as: observational ( $n = 61$ , 88%), cohort ( $n = 37$ , 53%), cross-sectional ( $n = 24$ , 35%), protocols ( $n = 11$ , 16%), systematic reviews ( $n = 7$ , 10%), qualitative ( $n = 4$ , 6%), quality improvement and programme evaluations ( $n = 3$ , 3%), randomised controlled trials (RCT) ( $n = 2$ , 3%), convenience series ( $n = 1$ , 1%) or where described as a ‘comparative’ design ( $n = 1$ ,


1%). The populations most studied were people receiving haemodialysis (HD) ( $n = 45$ , 65%), ACKD ( $n = 34$ , 50%), peritoneal dialysis (PD) ( $n = 15$ , 22%), transplant ( $n = 5$ , 7%) and conservative management ( $n = 4$ , 6%).

### Is frailty formally identified in this population and if so, what methods are used to identify frailty?

Across included studies, frailty was identified using 37 different assessments, with a median of three per study (IQR 1–6). Appendix 3 describes all assessments. The use of frailty measures varied according to the objectives of included studies. Twelve studies specifically compared self-reported frailty with other frailty measures to examine correlations and comparability.

The most frequently used assessment was the frailty phenotype, employed in 40 (44%) studies. In identifying frailty, 36 (43%) studies used one objective measure of frailty: frailty phenotype ( $n = 17$ , 16%), the clinical frailty scale (CFS;  $n = 10$ , 14%), Edmonton frailty scale ( $n = 4$ , 4%) and the Frail scale ( $n = 5$ , 4%). All other studies used multiple assessments, either directly or by proxy measure. For example, functional and cognitive assessments were used as a proxy measure of frailty. The majority of the assessments have all been used in the general population, specific CKD measures were the CKD frailty index ( $n = 6$ , 5%) and the Dialysis Morbidity Mortality Study ( $n = 1$ , 1%).

Five self-assessments of frailty were identified in 11 (16%) studies (Appendix 3). Information gathered was used in conjunction with assessments undertaken by the researcher/healthcare professional. The perception of frailty by an individual did not always correlate with the healthcare professional [23]. Eighteen studies (19%) included a cognitive assessment, using four different assessments. Eleven studies (13%) included in-depth assessment of related geriatric domains including cognition, depression, falls and nutrition. Six types of functional assessments were used across 18 studies. Nutritional assessments were completed in eight studies (9%). Only two studies (2%) included social



**Figure 2.** Scoping review consort diagram.


isolation as part of the assessment process. Caregiver burden was included in two studies (2%), with a further study including caregiver perspective using the Frailty Advanced Care Tool [24].

Seven systematic reviews highlighted issues with the consistency of the identification of frailty across CKD studies. Van Loon *et al.* (2016) examined the prevalence of frailty and geriatric impairments in people >70 years old commencing dialysis [25]. Twenty-seven studies were included. Three studies focused on frailty specifically, others focused on domains of frailty, including impairment of cognitive function, mood, performance status or (instrumental) activities of daily living (ADL), mobility (including falls), social environment and nutritional status. Malnutrition and frailty were the most systematically assessed [25]. Four reviews included studies that did not use a measure of frailty [4, 26–28] and most included all age ranges. Corroborating the findings of the current review, Mei *et al.* [7] found that the frailty phenotype was the most used frailty assessment across studies included within their review.

### What interventions have been reported and how might they enhance both care and outcomes specifically for older people with ACKD?

Few studies reported interventions ( $n=5$ , 6%). Those that did broadly evaluated two types of interventions: comprehensive geriatric assessment (CGA) and exercise. CGA is a model of health care based upon comprehensive multidisciplinary assessment taking account of medical, psychological, functional, social and environmental problems

[29]. Four studies (6%) reported on the implementation of CGA in practice, with variability in the populations included (HD, PD and pre-dialysis ACKD), the domains assessed and methodology used (qualitative, quality improvement, comparative and cross-sectional) to evaluate implementation [30–33]. CGA appeared to enhance education, training and awareness of the holistic and palliative needs of this population, facilitating appropriate onward referrals and advance care planning [31, 33, 34]. Barriers to CGA included lack of communication about goals, interpretation of geriatric assessments by clinicians, assessment burden for patients, poor health literacy and organisational challenges [33]. The latter related to the time needed to complete CGA and additional costs of involving geriatricians and additional multidisciplinary team members [33]. Although not formally measured, some studies observed an improvement in discussions about treatment options and adjustments in addition to increased advice seeking relating to social support [30, 33, 34]. These findings suggest that CGA may enhance decision-making and facilitate informed person-centred interventions for older people living with ACKD and frailty [33]. Other interventions included exercise. Two included papers (the protocol and completed study) referred to the same intervention [35, 36]. This was a pilot RCT evaluating a multicomponent home-based exercise programme for pre-frail and frail older adults with CKD. Seventy-three percent completed the programme of two exercise sessions per week, leading to potential improvements in both function and symptoms [36]. The other included intervention studies were all protocols of ongoing trials of exercise, focusing on hospitalised individuals with ACKD [37], and the effects of



**Figure 3.** Reported outcome measures. ICU, intensive care unit. 'Bad outcome' refers to delayed graft functioning of >14 days, never functioning kidney, readmission before day 90, surgical complications, discharge to a care facility or death.

muscle stimulation during HD [38]. Finally, a protocol by Chang *et al.* [39] described an RCT evaluating intervention individualised according to the findings of CGA.

### What outcomes have been reported in older people living with frailty and ACKD?

Fifty-two papers, from 41 studies, reported outcome measures. A total of 121 separate outcomes were identified, with a median of two outcomes per study (IQR 1–7) (Figure 3). Three (2%) measures were not reported in sufficient detail to enable classification. Thirty-two (26%) outcomes could not be classified according to the ICF domains. Of these, most reported mortality ( $n = 24$ , 58%), and hospitalisation and healthcare utilisation ( $n = 13$ , 32%).


Most outcomes ( $n = 86$ , 71%) were classified according to the ICF. Most ( $n = 47$ , 55%) were objective tests or appraisals from the perspective of the healthcare professional/researcher; the remainder were patient-reported outcomes (PROMS;  $n = 21$ , 24%) and biomarkers ( $n = 18$ , 21%). The most frequently reported objective measures examined capacity, performance or dependency related to ADL ( $n = 21$ , 22%), physical function ( $n = 19$ , 20%), cognition ( $n = 11$ , 12%) and body composition ( $n = 10$ , 11%). No measures clearly predominated; however, grip strength and gait speed (both  $n = 6$ , 6%) were most frequently used measures of function, and the Lawton and Barthel most frequently used ADL measures ( $n = 9$ , 10%;  $n = 7$ , 7%, respectively). The mini-mental state examination was the most frequently used measure of cognition ( $n = 5$ , 5%). Frailty was measured as an outcome within 11 (27%) studies, primarily using the frailty phenotype ( $n = 8$ , 50%) or the CFS ( $n = 4$ , 25%).

Of the PROMS, ten (30%) related to health-related quality of life (HRQOL), eight (24%) to mental health outcomes, particularly anxiety and depression, and three (9%) to physical activity. None examined the impact of ACKD and frailty upon carers or significant others. Again, no preferred PROMs emerged, but the SF-36 appeared to be the most frequently used HRQOL measure ( $n = 4$ , 12%). Most biomarkers were related to the management of kidney disease ( $n = 24$ , 73%), but 12 (36%) were markers of nutrition. The tests most commonly reported were albumin ( $n = 8$ , 22%) and haemoglobin ( $n = 5$ , 13%).

Overall, outcomes mapped to 172 categories within the ICF (summarised in Figure 4). Categories of body function most frequently examined related to mental functions ( $n = 99$ , 46%), digestive and metabolic functions ( $n = 41$ , 19%), cardiovascular, haematological, immunological and respiratory functions ( $n = 30$ , 14%) and genitourinary and reproductive functions ( $n = 14$ , 6%). Categories of activity and participation most frequently examined related to mobility ( $n = 80$ , 40%), self-care ( $n = 37$ , 19%) and domestic life ( $n = 29$ , 15%). Categories of body structures examined related only to movement, specifically muscle ( $n = 7$ , 78%) and bone ( $n = 2$ , 22%). Categories of environment related only to products and technology, specifically financial assets and medications ( $n = 10$ , 91%) and support and relationships ( $n = 1$ , 9%).

### What outcomes matter most to older people (living with frailty and ACKD), their relatives and carers?

Two studies explored HRQOL outcomes important to people living with frailty and receiving HD [40, 41]. Hall *et al.*



**Figure 4.** ICF categorisation of outcome measures.

[40] collected qualitative data to compare with domains measured in validated HRQOL questionnaires. Two themes were identified from the qualitative data: physical well-being and social support. Physical well-being related to symptom control, being alive and maintaining health status, whereas social support related to practical and emotional support and socialisation [41]. In Young *et al.*'s [41] study, maintaining mobility, the ability to undertake a range of ADLs and participation in social roles were viewed as key outcomes. The qualitative findings from Hall *et al.*'s [40] study did not correspond with domains in the Kidney Disease Quality of Life Instrument and WHO Quality of Life for Older Persons Questionnaire instruments, suggesting that a bespoke HRQOL measure is needed for this population. Interestingly in Young *et al.*'s study, only 13 (52%) participants agreed to complete a falls diary, many preferred falls information to be collected during HD treatment. The majority who had fallen rarely reported them to healthcare professionals, believing that they were an expected consequence of HD or having experienced their concerns about falls being overlooked. Consequently, falls prevention was not viewed as a key outcome from the patient perspective [41].

Reporting of patient and public involvement was noticeably absent from the studies reviewed; the four qualitative studies explored patient and carer perspectives. The scoping review findings, in particular outcome measurements, were presented to a patient focus group with five participants. The group raised concerns regarding the use of HRQOL tools as outcome measures because of the language used, limited options available for selection and lack of inclusion of aspects group members felt to be important. Overall, members of the focus group stressed that they felt interviews were the most appropriate method to gain an in-depth patient perspective regarding the impact of interventions, though they acknowledged that this is not always feasible for studies involving large numbers of participants.

## Discussion

The aim of this review was to identify and map available evidence relating to older adults living with ACKD and frailty, specifically addressing frailty assessments used, interventions reported and outcomes. Of the studies included, the majority were observational, reporting on incidence, prevalence and associations of frailty with outcomes such as mortality and hospitalisation. The measurement of frailty was inconsistent and varied across studies. The small number of intervention studies adds little to our understanding of which outcomes are a priority and are meaningful to this population. There was an underrepresentation of carers' perspectives and measures of carer burden, both are important considering carers often have great involvement with this population.

The high prevalence of frailty across many chronic diseases and associations with the frailty phenotype is now being recognised [42–44]. The importance of frailty and how it may change the prognosis and therapeutic approaches to several conditions have also been documented [42, 43]. A wide range of measures were used to identify frailty in the ACKD population. Whilst the purpose of this review was not to examine the validity of assessments used, several studies have demonstrated the diagnostic and prognostic accuracy of assessments in ACKD including the frailty phenotype and the CFS [45–48]. The frailty phenotype was most common across the studies, and is also a popular frailty assessment used in the general population [49]. Some studies described modifications to the frailty phenotype, adding a further complexity. A lack of standardised approach makes the comparison and synthesis of studies difficult. Arguably, some of the assessments used are not validated measures of frailty, or serve only as proxy measures, with moderate specificity for the identification of frailty [50]. This challenge has been identified in previous reviews of older people with ACKD and within the general population [4, 7, 25, 49, 51]. To provide guidance on the use of assessments in research

and practice, existing recommendations suggest considering the context and purpose when selecting a frailty measure; for example, if risk prediction is the goal, then frailty assessments validated for this purpose should be selected [51]. Different assessments are recommended for research and practice; for example, the frailty phenotype is commonly used in research and but less so in practice [49, 51], and the CFS is now gaining traction in clinical practice. In support of this, a consensus process by Voorend *et al.* [52] recommended the CFS be used within nephrology.

This scoping review highlights a lack of interventional studies within this population. The limited number primarily reported on the implementation of CGA, the gold standard of care for older people living with frailty in the general population [29]. Although CGA has been recommended within the European Best Practice guidelines [53], existing evidence only suggests that CGA is feasible in a nephrology setting, and robust evidence that CGA improves outcomes relevant to older adults with ACKD is lacking. However, preliminary findings suggest that CGA supports enhanced decision-making and provides opportunities to embed shared decision-making in practice [33]. This can improve the sharing of prognostic information, facilitate discussions about advance care planning and assist in the planning of more supportive, palliative approaches to treatment [33, 53–55].

Other interventions described related to exercise, which has been shown to be important for falls prevention and rehabilitation of older people living with frailty in the general population [56, 57]. Exercise has been shown to improve physical functioning, fatigue and HRQOL in the CKD population [58]. Whilst exercise appears to be feasible for older people with ACKD and frailty [36], further exploration of the effectiveness of tailored approaches for this group is required. This review identifies those multicomponent interventions that target all the relevant parts of the CGA process that are lacking, although may prove more beneficial than single interventions in this population.

Across all intervention studies, very little research has measured the impact of interventions on carer burden. A recent systematic review highlighted the high levels of burden in those caring for people with ACKD [59]. This appeared to be the result of a wide range of factors, including gender, caregiver and patient income, time providing daily care, duration of caregiving, the relationship to person, and cohabiting arrangements, reinforcing the need for further research into interventions to support this group [59]. The need to assess caregiver burden with a consistent approach within standard care was highlighted. Doing so may prompt healthcare professionals to offer interventions that aim to improve caregiver well-being [59].

This review underlines the broad range, but limited scope, of outcomes currently utilised within existing studies. Mortality and hospitalisation were the most common reported outcomes. There was little consensus amongst the PROMs and objective measures used. Outcomes were most frequently mapped to the ICF category of activities and

participation, but the majority focused on activities, rather than participation. Whilst this review exposes how little is currently known about what outcomes matter to people living with ACKD and frailty, existing evidence indicates that social support (including emotional and practical), participation in activities, maintaining mobility and being supported in decision-making are all important [33, 40, 41].

The number of core outcome sets, designed to improve the consistency, relevance and reporting of outcome measures, has grown in recent years [60]. The International Consortium for Health Outcomes Measurement has published core outcome sets for people with CKD and older people separately, both including different measures [61]. Matching outcomes important to individuals with measures used in assessment tools has been highlighted previously [40].

### Implications for future research

This review highlights several priorities for future research. Two priority setting exercises have been undertaken in the general population and within the dialysis population, but neither specifically addresses the priorities of older people living with frailty and ACKD [11, 12]. The National Institute for Health Research also published a report of important priorities for older people living with multiple conditions, including carers, highlighting that carer burden, support and social isolation were considered important [62]. Using the results of this review as a basis for discussion, our group will undertake a consensus process using nominal group technique to better understand the key outcomes and care priorities for patients and carers. This will enable us to design studies and make recommendations, which focus on areas that are meaningful and important to this group.

### Limitations

This is the first scoping review to specifically examine outcomes and care priorities for older adults living with frailty and ACKD. Despite these strengths, there are limitations with this review. Scoping review methodology does not include quality assessments of included studies. This limitation was mitigated by following clear eligibility guidelines and utilising all members of the team. Mapping to the ICF was made following assessment, interpretation and discussion between authors. Consequently, the classifications may have been assessed differently by other researchers. To mitigate this, the mapping process followed established linking rules and the procedures used described in detail to increase transparency [19]. Where the common generic PROMS (e.g. the SF-36) have been mapped within other published works, the mapping decisions have been compared to increase confidence and accuracy of classification decisions. Finally, this review only included studies of people aged >65 years old, therefore cannot be used to infer outcomes and care priorities for younger people living with frailty and ACKD.

## Conclusion

In summary, this scoping review highlights important areas for further research in older people living with frailty and ACKD that will contribute to improvements in services and care of this population. Researchers and clinicians should use a standardised approach to measure and assess frailty whilst recognising the accumulation and interaction of geriatric domains and in particular the complicated treatment decisions required in this vulnerable population [25]. Future research should build on the evidence for the implementation of CGA into practice, and to develop and test robust holistic interventions that improve outcomes that matter to people living with frailty and ACKD. This should include studies that focus on carers and carer burden.

**Supplementary Data:** Supplementary data mentioned in the text are available to subscribers in *Age and Ageing* online.

**Acknowledgements:** With thanks to the multi-professional collaborative for outcomes and care priorities for older people with ACKD-protocol team for their work on the protocol.

**Declaration of Conflicts of Interest:** None.

**Declaration of Sources of Funding:** British Renal Society and Kidney Care UK.

## References

1. Storey A. Living longer: How Our Population Is Changing and Why It Matters. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/ageing/articles/livinglongerhowourpopulationischangingandwhyitmatters/2018-08-13> (2019, date last accessed).
2. NHS England. 2021. <https://www.england.nhs.uk/ourwork/clinical-policy/older-people/improving-care-for-older-people/>.
3. Fried LP, Tangen CM, Walston J *et al.* Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001; 56: M146–57. <https://doi.org/10.1093/gerona/56.3.M146>.
4. Walker SR, Gill K, Macdonald K *et al.* Association of frailty and physical function in patients with non-dialysis CKD: a systematic review. *BMC Nephrol* 2013; 14. <https://doi.org/10.1186/1471-2369-14-228>.
5. Zhao Y, Liu Q, Ji J. The prevalence of frailty in patients on hemodialysis: a systematic review and meta-analysis. *Int Urol Nephrol* 2020; 52: 115–20.
6. Zhang Q, Ma Y, Lin F, Zhao J, Xiong J. Frailty and mortality among patients with chronic kidney disease and end-stage renal disease: a systematic review and meta-analysis. *Int Urol Nephrol* 2020; 52: 363–70.
7. Mei F, Gao Q, Chen F *et al.* Frailty as a predictor of negative health outcomes in chronic kidney disease: a systematic review and meta-analysis. *J Am Med Dir Assoc* 2021; 22: 535–43.e7.
8. Goto NA, Van Loon IN, Boereboom FTJ *et al.* Association of initiation of maintenance dialysis with functional status and caregiver burden. *Clin J Am Soc Nephrol* 2019; 14: 1039–47.
9. Iyasere O, Brown EA, Johansson L *et al.* Quality of life with conservative care compared with assisted peritoneal dialysis and haemodialysis. *Clin Kidney J* 2019; 12: 262–8.
10. Van Loon I, Hamaker ME, Boereboom FTJ *et al.* A closer look at the trajectory of physical functioning in chronic hemodialysis. *Age Ageing* 2017; 46: 594–9.
11. Van Loon IN, Goto NA, Boereboom FTJ *et al.* Geriatric assessment and the relation with mortality and hospitalizations in older patients starting dialysis. *Nephron* 2019; 143: 108–19.
12. Van Loon IN, Goto NA, Boereboom FTJ, Verhaar MC, Bots ML, Hamaker ME. Quality of life after the initiation of dialysis or maximal conservative management in elderly patients: a longitudinal analysis of the geriatric assessment in OLder patients starting dialysis (GOLD) study. *BMC Nephrol* 2019; 20: 108.
13. Parker SG, Corner L, Laing K *et al.* Priorities for research in multiple conditions in later life (multi-morbidity): findings from a James Lind Alliance Priority Setting Partnership. *Age and Ageing* 2019; 48: 401–06.
14. Matus Gonzalez A, Gutman T, Lopez-Vargas P *et al.* Patient and caregiver priorities for outcomes in chronic kidney disease: multinational nominal group technique study. *American Journal of Kidney Diseases* [accepted 5th April 2020].
15. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* 2018; 18: 143. <https://doi.org/10.1186/s12874-018-0611-x>.
16. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005; 8: 19–32.
17. Hurst H, Jones E, Ormandy P *et al.* Outcomes and care priorities for older people living with frailty and advanced chronic kidney disease: a multiprofessional scoping review protocol. *BMJ Open* 2021; 11: e040715. <https://doi.org/10.1136/bmjopen-2020-040715>.
18. World Health Organization (WHO). International Classification of Functioning, Disability and Health: ICF. Geneva: World Health Organization, 2001.
19. Cieza A, Fayed N, Bickenbach J, Prodinger B. Refinements of the ICF linking rules to strengthen their potential for establishing comparability of health information. *Disabil Rehabil* 2019; 41: 574–83.
20. Cieza A, Geyh S, Chatterji S, Kostanjsek N, Ustün B, Stucki G. ICF linking rules: an update based on lessons learned. *J Rehabil Med* 2005; 37: 212–8.
21. ICF. <https://www.who.int/standards/classifications/international-classification-of-functioning-disability-and-health>.
22. Whiteneck G, Dijkers MP. Difficult to measure constructs: conceptual and methodological issues concerning participation and environmental factors. *Arch Phys Med Rehabil* 2009; 1: S22–35.
23. Salter ML, Gupta N, Massie AB *et al.* Perceived frailty and measured frailty among adults undergoing hemodialysis: a cross-sectional analysis. *BMC Geriatr* 2015; 15: 52. <https://doi.org/10.1186/s12877-015-0051-y>.
24. Clark DA, Khan U, Kiberd BA *et al.* Frailty in end-stage renal disease: comparing patient, caregiver, and clinician perspectives. *BMC Nephrol* 2017; 18: 148. <https://doi.org/10.1186/s12882-017-0558-x>.
25. Van Loon IN, Wouters TR, Boereboom FT, Bots ML, Verhaar MC, Hamaker ME. The relevance of geriatric



- impairments in patients starting dialysis: a systematic review. *Clin J Am Soc Nephrol* 2016; 11: 1245–59.
26. Kallenberg MH, Kleinveld HA, Dekker FW *et al.* Functional and cognitive impairment, frailty, and adverse health outcomes in older patients reaching ESRD—a systematic review. *Clin J Am Soc Nephrol* 2016; 11: 1624–39.
  27. Shen Z, Ruan Q, Yu Z, Sun Z. Chronic kidney disease-related physical frailty and cognitive impairment: a systemic review. *Geriatr Gerontol Int* 2017; 17: 529–44.
  28. Song Y-H, Cai G-Y, Xiao Y-F, Chen X-M. Risk factors for mortality in elderly haemodialysis patients: a systematic review and meta-analysis. *BMC Nephrol* 2020; 21: 377. <https://doi.org/10.1186/s12882-020-02026-x>.
  29. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet* 2013; 381: 752–62.
  30. Hall RK, Haines C, Gorbalkin SM *et al.* Incorporating geriatric assessment into a nephrology clinic: preliminary data from two models of care. *J Am Geriatr Soc* 2016; 64: 2154–8.
  31. Nixon AC, Brown J, Brotherton A *et al.* Implementation of a frailty screening programme and geriatric assessment service in a nephrology Centre: a quality improvement project. *J Nephrol* 2021; 34: 1215–24.
  32. Parlevliet JL, Buurman BM, Pannekeet MM *et al.* Systematic comprehensive geriatric assessment in elderly patients on chronic dialysis: a cross-sectional comparative and feasibility study. *BMC Nephrol* 2012; 13: 30. <https://doi.org/10.1186/1471-2369-13-30>.
  33. Voorend CGN, Berkhout-Byrne NC, Meuleman Y, Mooijaart SP, Bos WJW, van Buren M. Perspectives and experiences of patients and healthcare professionals with geriatric assessment in chronic kidney disease: a qualitative study. *BMC Nephrol* 2021; 22: 9. <https://doi.org/10.1186/s12882-020-02206-9>.
  34. Moffatt H, Moorhouse P, Mallery L, Landry D, Tennankore K. Using the frailty assessment for care planning tool (fact) to screen elderly chronic kidney disease patients for frailty: the nurse experience. *Clin Interv Aging* 2018; 13: 843–52.
  35. Nixon AC, Bampouras TM, Gooch HJ *et al.* The EX-FRIL CKD trial: a study protocol for a pilot randomised controlled trial of a home-based EXercise programme for pre-frail and FRIL, older adults with chronic kidney disease. *BMJ Open* 2020; 10: e035344. <https://doi.org/10.1136/bmjopen-2019-035344>.
  36. Nixon AC, Bampouras TM, Gooch HJ *et al.* Home-based exercise for people living with frailty and chronic kidney disease: a mixed-methods pilot randomised controlled trial. *PLoS One* 2021; 16: e0251652. <https://doi.org/10.1371/journal.pone.0251652>.
  37. Wytmsa-Fisher K, Mustata S, Cowan T, Ester M, Culos-Reed SN. A physical activity intervention feasibility study for kidney inpatients: a basic research protocol. *Can J Kidney Health Dis* 2021; 8: 205435812098705. <https://doi.org/10.1177/2054358120987052>.
  38. Suzuki Y, Kamiya K, Tanaka S *et al.* Effects of electrical muscle stimulation in frail elderly patients during haemodialysis (DIAL): rationale and protocol for a crossover randomised controlled trial. *BMJ Open* 2019; 9: e025389. <https://doi.org/10.1136/bmjopen-2018-025389>.
  39. Chang J, Gao Y, Fang X-Y, Zhao S-M, Hou Y-P, Sun Q-M. Individualized intervention for frail non-dialysis elderly patients with chronic kidney disease: protocol for a randomized controlled trial. *BMC Geriatr* 2020; 20: 159. <https://doi.org/10.1186/s12877-020-1491-6>.
  40. Hall RK, Cary MP, Washington TR, Colon-Emeric CS. Quality of life in older adults receiving hemodialysis: a qualitative study. *Qual Life Res* 2020; 29: 655–63.
  41. Young HML, Eborall HC, Conroy S *et al.* Living with frailty and falls: a qualitative study exploring the experiences of patients with end-stage renal disease receiving haemodialysis. *Physiotherapy* 2019; 105: e28–9.
  42. Marengoni A, Vetrano DL, Manes-Gravina E, Bernabei R, Onder G, Palmer K. The relationship between COPD and frailty: a systematic review and meta-analysis of observational studies. *Chest* 2018; 154: 21–40.
  43. Marengoni A, Zucchelli A, Vetrano DL *et al.* Heart failure, frailty, and pre-frailty: a systematic review and meta-analysis of observational studies. *Int J Cardiol* 2020; 316: 161–71.
  44. Musso CG, Jauregui JR, Macias Nunez JF. Frailty phenotype and chronic kidney disease: a review of the literature. *Int Urol Nephrol* 2015; 47: 1801–7.
  45. Anderson BM, Qasim M, Correa G *et al.* Correlations, agreement and utility of frailty instruments in prevalent haemodialysis patients: baseline cohort data from the FITNESS study. *Clin Kidney J* 2022; 15: 145–52.
  46. Chao C-T, Chiang C-K, Huang J-W, Hung K-Y, COGENT study group. Self-reported frailty among end-stage renal disease patients: a potential predictor of dialysis access outcomes. *Nephrology* 2017; 22: 333–4.
  47. Gopinathan JC, Hafeeq B, Aziz F, Narayanan S, Aboobacker IN, Uvais NA. The prevalence of frailty and its association with cognitive dysfunction among elderly patients on maintenance hemodialysis: a cross-sectional study from South India. *Saudi J Kidney Dis Transpl* 2020; 31: 767–74.
  48. Nixon AC, Bampouras TM, Pendleton N, Mitra S, Dhaygude AP. Diagnostic accuracy of frailty screening methods in advanced chronic kidney disease. *Nephron* 2019; 141: 147–55.
  49. Dent E, Kowal P, Hoogendijk EO. Frailty measurement in research and clinical practice: a review. *Eur J Intern Med* 2016; 31: 3–10. <https://doi.org/10.1016/j.ejim.2016.03.007>.
  50. BGS. Fit for Frailty. British Geriatric Society, 2014. Rebranded edition 2017, British Geriatrics Society Marjory, London.
  51. Buta BJ, Walston JD, Godino JG *et al.* Frailty assessment instruments: systematic characterization of the uses and contexts of highly-cited instruments. *Ageing Res Rev* 2016; 26: 53–61.
  52. Voorend CA-O, Joosten H, Berkhout-Byrne NC *et al.* Design of a consensus-based geriatric assessment tailored for older chronic kidney disease patients: results of a pragmatic approach. *Eur Geriatr Med* 2021; 12: 931–42.
  53. Farrington K, Covic A, Nistor I *et al.* Clinical practice guideline on management of older patients with chronic kidney disease stage 3b or higher (eGFR < 45 mL/min/1.73 m<sup>2</sup>): a summary document from the European Renal Best Practice Group. *Nephrol Dial Transplant* 2017; 32: 9–16.
  54. Raj R, Thiruvengadam S, Ahuja KDK, Frandsen M, Jose M. Discussions during shared decision-making in older adults with advanced renal disease: a scoping review. *BMJ Open* 2019; 9: e031427. <https://doi.org/10.1136/bmjopen-2019-031427>.
  55. Brown EA, Farrington K. Geriatric assessment in advanced kidney disease. *Clin J Am Soc Nephrol* 2019; 14: 1091–3.
  56. de Labra C, Guimaraes-Pinheiro C, Maseda A, Lorenzo T, Millán-Calenti JC. Effects of physical exercise interventions in frail older adults: a systematic review of randomized controlled

- trials. *BMC Geriatr* 2015; 15: 154. <https://doi.org/10.1186/s12877-015-0155-4>.
57. Sherrington C, Fairhall NJ, Wallbank GK *et al.* Exercise for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2019; 2019: CD012424. <https://doi.org/10.1002/14651858.CD012424.pub2>.
  58. Bernier-Jean A, Beruni NA, Bondonno NP *et al.* Exercise training for adults undergoing maintenance dialysis. *Cochrane Database Syst Rev* 2022; 2022. <https://doi.org/10.1002/14651858.CD014653>.
  59. Alshammari B, Noble H, McAnaney H, Alshammari F, O'Halloran P. Factors associated with burden in caregivers of patients with end-stage kidney disease (a systematic review). *Healthcare* 2021; 9: 1212. <https://doi.org/10.3390/healthcare9091212>.
  60. Godlee F. Outcomes that matter to patients. *BMJ* 2012; 344: e318. <https://doi.org/10.1136/bmj.e318>.
  61. ICHOM. International Consortium for Health Outcomes Measurements. ICHOMorg, 2021.
  62. Spiers G, Boulton E, Corner L *et al.* What matters to people with multiple long-term conditions and their carers? *Postgraduate Medical Journal* 2021. <https://doi.org/10.1136/postgradmedj-2021-140825>.
- Received 14 April 2022; editorial decision 20 September 2022**
-



## ID NOW™ PLATFORM

KNOW FASTER SO YOU  
CAN ACT QUICKER

# NOW



Now, you can provide rapid molecular respiratory testing for COVID-19, influenza, RSV and strep A in any acute care setting, where and when it's needed most.



IDNOW.ABBOTT

## NOW

**IMPROVED WORKFLOW**  
with single patient swab for  
COVID-19 and influenza A & B