

This is a repository copy of *A synthetic biology approach to transgene expression in insects*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/216765/>

Version: Published Version

Article:

Leftwich, Philip, Purcell, Jessica, Anderson, Michelle orcid.org/0000-0003-1510-2942 et al. (4 more authors) (2024) A synthetic biology approach to transgene expression in insects. ACS Synthetic Biology. ISSN 2161-5063

<https://doi.org/10.1021/acssynbio.4c00250>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

A Synthetic Biology Approach to Transgene Expression in Insects

Philip T. Leftwich,^{*,†} Jessica C. Purcell,[†] Michelle A. E. Anderson, Rennos Fragkoudis, Sanjay Basu, Gareth Lycett, and Luke Alphey^{*}



Cite This: <https://doi.org/10.1021/acssynbio.4c00250>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: The ability to control gene expression is pivotal in genetic engineering and synthetic biology. However, in most nonmodel and pest insect species, empirical evidence for predictable modulation of gene expression levels is lacking. This knowledge gap is critical for genetic control systems, particularly in mosquitoes, where transgenic methods offer novel routes for pest control. Commonly, the choice of RNA polymerase II promoter (Pol II) is the primary method for controlling gene expression, but the options are limited. To address this, we developed a systematic approach to characterize modifications in translation initiation sequences (TIS) and 3' untranslated regions (UTR) of transgenes, enabling the creation of a toolbox for gene expression modulation in mosquitoes and potentially other insects. The approach demonstrated highly predictable gene expression changes across various cell lines and 5' regulatory sequences, representing a significant advancement in mosquito synthetic biology gene expression tools.

KEYWORDS: Promoter, mosquito, genetic engineering, insect, mRNA, translation



INTRODUCTION

The ability to control the strength of expression of transgenes in a species of interest has underpinned genetic engineering and synthetic biology from their conception. There is, however, a lack of robust, empirical evidence for predictable modulation of gene expression levels in most nonmodel and pest insect species.^{1,2} In mosquitoes, transgenic methods afford novel routes for pest control, however, genetic control systems depend on precise gene expression, so this lack of information is a critical gap in our technical capability. Commonly the selection of RNA polymerase II promoter (Pol II) can dictate the amount, timing, and spatial specificity of gene transcription. A limited number of promoters of viral origin are sometimes used, as these are active across a range of insect species, however there are applications for characterizing endogenous promoters of varying activity. The choice of promoter is often determined by a requirement for specific spatial and/or temporal regulation, with few options for controlling expression level by this route beyond bespoke analysis of endogenous promoters.

Other methods commonly employed for modulating gene expression involve modifications to the mRNA sequences of the 5' and 3' untranslated regions (UTR), flanking the coding sequence of a transgene. Although these regions do not contribute to the final protein, they play crucial roles in mRNA stability and translation efficiency. In particular, the translation initiation sequence (TIS), a short (~10nt) segment within the 5' UTR just upstream of the start codon, has been shown to significantly impact translation efficiency in both vertebrates and invertebrates.^{3–6} By using different TIS sequences,

predictable changes in transgene expression can be achieved.^{2,6,7}

The 3'UTR is more closely associated with mediating the termination of translation and ensuring efficient recycling of the translation complex, enabling multiple translations from the same mRNA molecule.^{8,9} In insect transgenesis, exogenous 3'UTR sequences are routinely used, including the viral-derived simian virus 40 (SV40) 3'UTR,¹¹ the P10 baculovirus 3' UTR from *Autographa californica nucleopolyhedrovirus* (AcNPV)¹² and the 3'UTR of the *K10* gene from *Drosophila melanogaster*.¹³

Despite the widespread use of these 3'UTR sequences, their relative efficacies are often based on anecdotal evidence. Strategically manipulating these untranslated regions provides a valuable approach to finely tune and optimize transgene expression in insect transgenesis.

We developed a systematic approach to characterizing TIS and 3'UTR modifications to transgenes to build a toolbox for modulating gene expression in mosquitoes, and potentially other insects, when combined with viral or endogenous culicine 5' regulatory DNA sequences—hereafter referred to as promoters (though strictly these DNA regions may also

Received: April 9, 2024

Revised: August 21, 2024

Accepted: August 21, 2024




Figure 1. Combinations of translation initiation sequences (TIS) and 3'UTRs produce highly replicable gene expression across a range of insect cell lines. Ratios of FF luciferase compared to a RL control were used to measure activity; UTRs are organized left to right by average relative activity and nested within TIS also organized left to right by average relative activity. Large symbols and error bars represent mean and associated approximate 95% confidence intervals estimated with a generalized linear mixed model with a Gamma error distribution, raw data is shown as small points.

contain enhancers, silencers, or binding sites for transcription factors required for proper regulation of gene expression). This toolbox provides an efficient way of expanding available Pol II promoters and affords routes to generating better regulation of activity of promoters across the species barrier. We initially tested the activities of the viral promoter HR5-IE1¹⁰ with a fully factorial combination of TIS and 3'UTR sequences in *Aedes aegypti*, *Aedes albopictus*, *Culex quinquefasciatus* and *Spodoptera frugiperda* cell lines. We then developed this further by taking the TIS/3'UTR combinations that produced the highest, lowest and median expression levels and demonstrated the highly replicable gene expression modulation effects in a panel of promoters.

We found that gene expression changes are highly predictable across a wide range of cell lines and promoter sequences. In conjunction with the characterization of several

endogenous culicine promoters, this represents a significant advance in the available gene expression tools for mosquito synthetic biology.

RESULTS AND DISCUSSION

We first measured the activity of five translation initiation sequences (TIS) (Table S1; BmLo, Syn21, Kozak, BmHi, and Lep)^{1–3,15} and three different 3'UTRs (K10, SV40, P10)^{11–13} in a fully factorial design all downstream of the HR5-IE1 promoter.¹⁴ In total, we produced 15 different constructs and tested these in five different insect cell lines, from three disease relevant Culicine mosquito species (*A. aegypti* (Aag2), *A. albopictus* (U4.4 and C6.36) and *C. quinquefasciatus* (Hsu)) and one Lepidopteran species (*S. frugiperda* (Sf9)).

We found a highly replicable pattern of gene expression modulation across all five tested cell lines (Figure 1). Averaged




Figure 2. Fine-scale modulation of transgene expression with TIS/3'UTR combinations is highly replicable across a range of synthetic and endogenous promoters in two Culicine mosquito cell lines: (A) Aag2 (*A. aegypti*) cells and (B) *A. albopictus*-derived U4.4 cells. Ratios of FF luciferase compared to a RL control were used to measure activity; TIS/3'UTR combinations are organized left to right by average relative activity, with the *x*-axis indicating the size of promoter fragments in base pairs (bp). Large solid symbols and error bars represent mean and associated approximate 95% confidence intervals estimated with a generalized linear mixed model with a Gamma error distribution; raw data is shown as open symbols. Lines connect promoters of the same origin.

across cell lines, the choice of TIS could produce up to a 2.55 relative-fold change in luciferase expression (95% CI 2.28–2.84; Table S2), while the choice of 3'UTR produced up to a 4.88 relative-fold change in luciferase expression (95% CI 4.52–5.26; Table S2).

The estimates from our analysis indicate that TIS sequences are mainly insensitive to cell type and behave remarkably consistently ($F_{16,518} = 11$, $P < 0.001$; Table S3). By contrast, the effect of the 3'UTR sequence on transgene expression was much more strongly affected by cell type ($F_{8,518} = 250$, $P < 0.001$; Table S3). Expression from constructs with P10 had notably higher expression than expected in Sf9 cells and much lower in both C6/36 and U4.4 cells. Significant interactions between TIS sequences and 3'UTR sequences were small ($F_{8,518} = 20$, $P < 0.001$; Table S3) therefore transcriptional activity appears to be primarily an additive effect when pairing

TIS and 3'UTR sequences. This makes the “plug and play” notion of pairing different synthetic components together highly attractive, as effects on transgene expression are highly predictable.

The TIS/3'UTR combinations with the lowest (BmLo-K10) and highest (Lep-P10) expressions (18.2 (95% CI 13–22) relative-fold expression difference); were the same across all cell lines (Figure 1). To generalize our results, we decided to expand the range of promoter sequences tested by taking the BmLo-K10, Lep-P10 and Kozak-SV40 combinations and testing them with additional promoter sequences. In total, we tested seven regulatory sequences from four endogenous promoters from Culicine mosquitoes: two variants of the *hsp83* promoter¹⁶ (1.4kb and 888bp) (AAEL011708), from *A. aegypti* with the large (c.4.2kb) intronic sequence of the 5'UTR truncated to retain minimal acceptor and donor

regions, three engineered variants of the *Polyubiquitin* promoter (AAEL003888) from *A. aegypti*,¹⁷ along with *Polyubiquitin* from *C. quinquefasciatus* (CPIJ010919) and we demonstrate the first use case for a new endogenous promoter *A. albopictus* derived *Polyubiquitin* (AALF002118). These, along with OpIE2 were tested in Aag2 and U4.4 cells.

As expected, promoters of viral origin behaved very consistently across both cell lines, while endogenous promoters responded in a more cell-specific manner (Figure 2, Table S4). The *hsp83* promoter sequences produced equivalent levels of gene expression to OpIE2 when in Aag2 cells, but this was lower in U4.4 cells. Shortening this sequence by removing c.500bp upstream produced no significant reduction in gene expression. *Polyubiquitin*-derived promoter sequences generally produced the highest levels of gene expression; fluctuations in the strength of expression across cell lines may reflect the evolutionary origins of each sequence (*C. quinquefasciatus* derived sequence displayed lower activity than *A. albopictus* for example).

The effects of TIS/3'UTR modification on gene expression were remarkably consistent for all promoter sequences. We found minimal differences in the responses between different regulatory sequences and TIS/3'UTR combinations (Promoter: $F_{8,1207} = 354.91$, $P < 0.001$; TIS/3'UTR: $F_{2,1207} = 509.96$, $P < 0.001$; Interaction effect: $F_{14,1193} = 7.26$, $P < 0.001$; Figure 2, Table S5), indicating that these act largely independently.

We have developed a straightforward method for modulating transgene expression in *Culicine* mosquitoes using a combinatorial approach that enables fine-scale manipulation. Our experiments demonstrated that TIS and 3'UTR sequences consistently produce highly predictable outcomes on transgene expression irrespective of promoter sequence or cell line. While this work was conducted in cultured cell lines, previous research strongly suggests that these findings will effectively translate to whole-organism transgene expression.^{1,17} We are confident that this will be a valuable resource for researchers in synthetic biology, genetic modification, and mosquito genetic control.

METHODS

Plasmids, Cells, Transfections, and Luciferase Assay.

Cells were seeded in 96-well plates 1 day before transfection with TransIT-PRO transfection kit (Mirus Bio, Madison, WI, US) according to manufacturer's recommendations. Master mixes were prepared for eight wells of a 96-well plate, as replicate wells per experimental construct. This was repeated in three replicate experiments. Per well, transfection amounts are listed for each cell line in Supporting Information. Complete plasmid sequences are currently available as genbank files on Github (see below) and will be available through NCBI upon publication.

Two days after transfection, cells were washed twice with ion-free PBS, lysed with 1× Passive Lysis Buffer then analyzed using the Dual-Luciferase Assay kit on a GloMax multi+ plate reader (Promega, Southampton, UK).

General cell maintenance and plasmid information is described in Supporting Information (Tables S6–S9).

ASSOCIATED CONTENT

Data Availability Statement

Scripts and raw data can be found at Github (<https://github.com/Philip-Leftwich/Pol2-promoters>). Complete information on analyses can be found in Supporting Information.

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acssynbio.4c00250>.

Table S1, model summaries for experiment one; Table S2, pairwise contrasts for experiment one; Table S3, ANOVA tables for experiment one; Table S4, model summaries for experiment two; Table S5, ANOVA tables for experiment two; Table S6, transfection amounts per cell line; Table S7, lysate volumes per experiment; Table S8, DNA sequences; Table S9, brief descriptors of plasmids; additional methods for cell culture, transfections, dual luciferase assays DNA sequences, and references (PDF)

AUTHOR INFORMATION

Corresponding Authors

Philip T. Leftwich – *Arthropod Genetics, The Pirbright Institute, Pirbright GU24 0HN, U.K.*; Present Address: P.T.L.: School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, U.K.; orcid.org/0000-0001-9500-6592; Email: p.leftwich@uea.ac.uk

Luke Alphey – *Arthropod Genetics, The Pirbright Institute, Pirbright GU24 0HN, U.K.*; Present Address: M.A.E.A. and L.A.: Department of Biology, University of York, Heslington, YO10 5DD, U.K.; orcid.org/0000-0002-2916-3802; Email: luke.alphey@york.ac.uk

Authors

Jessica C. Purcell – *Arthropod Genetics, The Pirbright Institute, Pirbright GU24 0HN, U.K.*

Michelle A. E. Anderson – *Arthropod Genetics, The Pirbright Institute, Pirbright GU24 0HN, U.K.*; Present Address: M.A.E.A. and L.A.: Department of Biology, University of York, Heslington, YO10 5DD, U.K.; orcid.org/0000-0003-1510-2942

Rennos Fragkoudis – *Arbovirus Pathogenesis Group, The Pirbright Institute, Pirbright GU24 0HN, U.K.*; *University of Edinburgh, Edinburgh Genome Foundry, Centre for Mammalian Synthetic Biology, Edinburgh EH9 3BF, U.K.*; orcid.org/0000-0002-8451-2665

Sanjay Basu – *Arthropod Genetics, The Pirbright Institute, Pirbright GU24 0HN, U.K.*

Gareth Lycett – *Liverpool School of Tropical Medicine, Liverpool L3 5QA, U.K.*; orcid.org/0000-0002-2422-053X

Complete contact information is available at: <https://pubs.acs.org/10.1021/acssynbio.4c00250>

Author Contributions

[†]Philip T. Leftwich and Jessica C. Purcell are joint first authors; Philip T. Leftwich, Jessica C. Purcell, and Luke Alphey conceived the study; Philip T. Leftwich, Jessica C. Purcell, and Michelle A. E. Anderson performed the experiments; Philip T. Leftwich analyzed the data; Philip T. Leftwich and Jessica C. Purcell wrote the manuscript; Rennos Fragkoudis, Sanjay Basu, and Gareth Lycett provided essential reagents for the

experiments; Philip T. Leftwich, Jessica C. Purcell, Michelle A.E. Anderson, and Luke Alphey revised the manuscript. All authors read and approved the final manuscript.

Notes

For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. The findings and conclusions contained within are those of the authors and do not necessarily reflect positions or policies of the Bill & Melinda Gates Foundation. The views, opinions and/or findings expressed are those of the authors and should not be interpreted as representing the official views or policies of the Department of Defense or the U.S. Government. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors declare the following competing financial interest(s): L.A. is an adviser to Synvect Inc. and Biocentis Ltd., with equity and/or financial interest in those companies. The other authors declare that they have no competing interests.

ACKNOWLEDGMENTS

This research was funded in part by research grants from the Wellcome Trust [110117/Z/15/Z and 226721/Z/22/Z] and the Bill & Melinda Gates Foundation [INV-008549] to L.A. and by the Defense Advanced Research Projects Agency (DARPA) Safe Genes Program [N66001-17-2-4054] to Kevin Esvelt at MIT. J.P. was supported by a studentship from The Pirbright Institute (BBS/E/1/00001985). L.A. was also supported through strategic funding from the UK Biotechnology and Biological Sciences Research Council (BBSRC) to The Pirbright Institute (BBS/E/1/00007033, BBS/E/1/00007038 and BBS/E/1/00007039).

REFERENCES

- (1) Pfeiffer, B. D.; Truman, J. W.; Rubin, G. M. Using translational enhancers to increase transgene expression in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 6626–6631.
- (2) Tatematsu, K.-I.; Uchino, K.; Sezutsu, H.; Tamura, T. Effect of ATG initiation codon context motifs on the efficiency of translation of mRNA derived from exogenous genes in the transgenic silkworm, *Bombyx mori*. *SpringerPlus* **2014**, *3*, 136.
- (3) Kozak, M. An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs. *Nucleic Acids Res.* **1987**, *15*, 8125–8148.
- (4) Kozak, M. Point mutations define a sequence flanking the AUG initiator codon that modulates translation by eukaryotic ribosomes. *Cell* **1986**, *44*, 283–292.
- (5) Cavener, D. R. Combinatorial control of structural genes in *Drosophila*: solutions that work for the animal. *Bioessays* **1987**, *7*, 103–107.
- (6) Cavener, D. R.; Ray, S. C. Eukaryotic start and stop translation sites. *Nucleic Acids Res.* **1991**, *19*, 3185–3192.
- (7) Horstick, E. J.; Jordan, D. C.; Bergeron, S. A.; Tabor, K. M.; Serpe, M.; Feldman, B.; Burgess, H. A. Increased functional protein expression using nucleotide sequence features enriched in highly expressed genes in zebrafish. *Nucleic Acids Res.* **2015**, *43*, No. e48.
- (8) Schuller, A. P.; Green, R. Roadblocks and resolutions in eukaryotic translation. *Nat. Rev. Mol. Cell Biol.* **2018**, *19*, 526–541.
- (9) Mayr, C. 3' UTRs Regulate Protein Functions by Providing a Nurturing Niche during Protein Synthesis. *Cold Spring Harb. Symp. Quant. Biol.* **2019**, *84*, 95–104.
- (10) Guarino, L. A.; Summers, M. D. Interspersed Homologous DNA of *Autographa californica* Nuclear Polyhedrosis Virus Enhances Delayed-Early Gene Expression. *J. Virol.* **1986**, *60*, 215–223.
- (11) Brand, A. H.; Perrimon, N. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **1993**, *118*, 401–415.
- (12) van Oers, M. M.; Vlak, J. M.; Voorma, H. O.; Thomas, A. A. M. Role of the 3' untranslated region of baculovirus p10 mRNA in high-level expression of foreign genes. *J. Gen. Virol.* **1999**, *80*, 2253–2262.
- (13) Prost, E.; Deryckere, F.; Roos, C.; Haenlin, M.; Pantesco, V.; Mohier, E. Role of the oocyte nucleus in determination of the dorsoventral polarity of *Drosophila* as revealed by molecular analysis of the K10 gene. *Genes Dev.* **1988**, *2*, 891–900.
- (14) Theilmann, D. A.; Stewart, S. Molecular analysis of the transactivating IE-2 gene of *Orgyia pseudotsugata* multicapsid nuclear polyhedrosis virus. *Virology* **1992**, *187*, 84–96.
- (15) Sano, K.-I.; Maeda, K.; Oki, M.; Maeda, Y. Enhancement of protein expression in insect cells by a lobster tropomyosin cDNA leader sequence. *FEBS Lett.* **2002**, *532*, 143–146.
- (16) Webster, S. H.; Scott, M. J. The *Aedes aegypti* (Diptera: Culicidae) hsp83 Gene Promoter Drives Strong Ubiquitous DsRed and ZsGreen Marker Expression in Transgenic Mosquitoes. *J. Med. Entomol.* **2021**, *58*, 2533–2537.
- (17) Anderson, M. A. E.; Leftwich, P. T.; Wilson, R.; Paladino, L. Z. C.; Basu, S.; Rooney, S.; Adelman, Z. N.; Alphey, L. *AePUB* promoter length modulates gene expression in *Aedes aegypti*. *Sci. Rep.* **2023**, *13*, 20352.