

Odd chromatic number of graph classes

Rémy Belmonte¹  | Ararat Harutyunyan² |

Noleen Köhler³  | Nikolaos Melissinos⁴ 

¹Université Gustave Eiffel, CNRS, LIGM, Marne-la-Vallée, France

²Université Paris-Dauphine, PSL University, CNRS UMR7243, LAMSADE, Paris, France

³School of Computing, University of Leeds, Leeds, UK

⁴Department of Theoretical Computer Science, Faculty of Information Technology, Czech Technical University in Prague, Prague, Czech Republic

Correspondence

Noleen Köhler, School of Computing, University of Leeds, Leeds, UK.
Email: n.koehler@leeds.ac.uk

Funding information

French National Research Agency under JCJC program (ASSK: ANR-18-CE40-0025-01); French National Research Agency under JCJC program (DAGDigDec: ANR-21-CE48-0012); CTU Global postdoc fellowship program

Abstract

A graph is called *odd* (respectively, *even*) if every vertex has odd (respectively, even) degree. Gallai proved that every graph can be partitioned into two even induced subgraphs, or into an odd and an even induced subgraph. We refer to a partition into odd subgraphs as an *odd colouring* of G . Scott proved that a connected graph admits an odd colouring if and only if it has an even number of vertices. We say that a graph G is k -odd colourable if it can be partitioned into at most k odd induced subgraphs. The *odd chromatic number* of G , denoted by $\chi_{\text{odd}}(G)$, is the minimum integer k for which G is k -odd colourable. We initiate the systematic study of odd colouring and odd chromatic number of graph classes. We first consider a question due to Scott, which states that every graph G of even order n has $\chi_{\text{odd}}(G) \leq c\sqrt{n}$, for some positive constant c , by proving that this is indeed the case if G is restricted to having girth at least seven. We also show that any graph G whose all components have even order satisfies $\chi_{\text{odd}}(G) \leq 2\Delta - 1$, where Δ is the maximum degree of G . Next, we show that certain interesting classes have bounded odd chromatic number. Our main results in this direction are that interval graphs, graphs of bounded modular-width all have bounded odd chromatic number. In particular, every even interval graph is

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Journal of Graph Theory* published by Wiley Periodicals LLC.

6-odd colourable, and every even graph is $3mw$ -odd colourable, where mw is the modular width of a graph.

KEYWORDS

colouring variant, graph classes, odd colouring, upper bounds, vertex partition problem

1 | INTRODUCTION

A graph is called *odd* (respectively *even*) if all its degrees are odd (respectively even). Gallai proved the following theorem (see [10], Problem 5.17 for a proof).

Theorem 1. *For every graph G , there exist:*

- a partition (V_1, V_2) of $V(G)$ such that $G[V_1]$ and $G[V_2]$ are both even;
- a partition (V'_1, V'_2) of $V(G)$ such that $G[V'_1]$ is odd and $G[V'_2]$ is even.

This theorem has two main consequences. The first one is that every graph contains an induced even subgraph with at least $|V(G)|/2$ vertices. The second is that every graph can be *even coloured* with at most two colours, that is, partitioned into two (possibly empty) sets of vertices, each of which induces an even subgraph of G . In both cases, it is natural to wonder whether similar results hold true when considering odd subgraphs.

The first question, known as the *odd subgraph conjecture* and mentioned already by Caro [4] as part of the graph theory folklore, asks whether there exists a constant $c > 0$ such that every graph G contains an odd subgraph with at least $|V(G)|/c$ vertices. In a recent breakthrough paper, Ferber and Krivelevich proved that the conjecture is true.

Theorem 2 [9]. *Every graph G with no isolated vertices has an odd induced subgraph of size at least $|V(G)|/10,000$.*

Note that the requirement that G does not have isolated vertices is necessary, as those cannot be part of any odd subgraph.

The second question is whether every graph can be partitioned into a bounded number of odd induced subgraphs. We refer to such a partition as an *odd colouring*, and the minimum number of parts required to odd colour a given graph G , denoted by $\chi_{\text{odd}}(G)$, as its *odd chromatic number*. This can be seen as a variant of proper (vertex) colouring, where one seeks to partition the vertices of a graph into odd subgraphs instead of independent sets. An immediate observation is that to be odd colourable, a graph must have all its connected components be of even order, as an immediate consequence of the handshake lemma. Scott [13] proved that this necessary condition is also sufficient. Therefore, graphs can generally be assumed to have all their connected components of even order, unless otherwise specified.

Motivated by this result, it is natural to ask how many colours are necessary to partition a graph into odd induced subgraphs. Unsurprisingly, on the computational side, it was shown by Belmonte and Sau [2] that the problem of deciding whether a graph is k -odd colourable is solvable in polynomial time when $k \leq 2$, and NP-complete otherwise, similarly

to the case of proper colouring. Scott showed [13], there exist graphs with arbitrarily large odd chromatic number. More precisely, he showed that there exist graphs of order n that require $\Theta(\sqrt{n})$ colours. In particular, the *subdivided clique*, i.e., the graph obtained from a complete graph on n vertices by subdividing every edge once requires exactly n colours. More generally, given any connected graph G , the graph H obtained from G by subdividing every edge once has $\chi_{\text{odd}}(H) = \chi(G)$, and H is odd colourable if and only if $|V(H)| = |V(G)| + |E(G)|$ is even.

As an upper bound, best known general bound is the following.

Theorem 3 Scott [13]. *Every connected graph G of even order n has $\chi_{\text{odd}}(G) \leq cn(\log \log n)^{-1/2}$, for some constant $c > 0$.*

Scott conjectured that this bound is far from being best possible, and that the aforementioned lower bound of \sqrt{n} is the right order of magnitude.

Conjecture 1 Scott [13]. *Every connected graph G of even order n has $\chi_{\text{odd}}(G) \leq (1 + o(1))c\sqrt{n}$, for some constant $c > 0$.*

A related conjecture about a general upper bound is due to Aashtab et al. [1], who conjectured that a Brooks-type result for graph colouring should extend to odd colouring.

Conjecture 2 Aashtab et al. [1]. *Every connected graph G of even order has $\chi_{\text{odd}}(G) \leq \Delta + 1$, where Δ is the maximum degree of G .*

In this work, we study the above two conjectures, which have not yet been given much attention. We first show that the conjecture of Aashtab et al. [1] holds if one is allowed $2\Delta - 1$ colours. Next, we use this to prove Scott's conjecture for sparse graphs. Then we study specific classes of graphs (such as interval graphs and graphs of bounded modular-width), showing that these have small odd chromatic numbers.

The rest of the work is structured as follows. In Section 3, we prove that every graph G of even order and maximum degree Δ has $\chi_{\text{odd}}(G) \leq 2\Delta - 1$, extending the result of Aashtab et al. on subcubic graphs to graphs of bounded degree. We actually prove a more general result, which provides additional corollaries for graphs of large girth. In particular, we obtain that planar graphs of girth 11 are 3-odd colourable. We also obtain that graphs of girth at least 7 are $\mathcal{O}(\sqrt{n})$ -odd colourable, hence obtaining Scott's conjecture for sparse graphs. While this bound is not constant, it is of particular interest since subdivided cliques have girth exactly 6. In Section 4, we prove that every graph with all connected components of even order satisfies $\chi_{\text{odd}}(G) \leq 3 \cdot mw(G)$, where $mw(G)$ denotes the modular width of G . This significantly generalizes a result on cographs from Belmonte and Sau [2] and provides an important step towards proving that graphs of bounded rank-width have bounded odd chromatic number, which in turn would imply that the ODD CHROMATIC NUMBER is FPT when parameterized by rank-width alone. Finally, we prove in Section 5 that every interval graph with all components of even order is 6-odd colourable. It was pointed out to us by Müller [11] that our proof, albeit with more tedious notation, yields the exact same bound for the larger class of AT-free graphs. Additionally, we show that every proper interval graph with all components of even order is 3-odd colourable, and this bound is tight. An overview of known results and open cases is provided in Figure 1 below.




FIGURE 1 Overview of known and open cases. [Color figure can be viewed at wileyonlinelibrary.com]

2 | PRELIMINARIES

For a positive integer i , we denote by $[i]$ the set containing every integer j such that $1 \leq j \leq i$. We consider a partition of a set X to be a tuple $\mathcal{P} = (P_1, \dots, P_k)$ of subsets of X such that $X = \bigcup_{i \in [k]} P_i$ and $P_i \cap P_j = \emptyset$, that is, we allow parts to be the empty set. Let $\mathcal{P} = (P_1, \dots, P_k)$ be a partition of X and $Y \subseteq X$. We let $\mathcal{P}|_Y$ be the partition of Y obtained from $(P_1 \cap Y, \dots, P_k \cap Y)$ by removing all empty parts. A partition (Q_1, \dots, Q_ℓ) of X is a coarsening of a partition (P_1, \dots, P_k) of X if for every P_i and every Q_j either $P_i \cap Q_j = \emptyset$ or $P_i \cap Q_j = P_i$, i.e., every Q_j is the union of P_i 's.

Every graph in this work is simple, undirected and finite. We use standard graph-theoretic notation, and we refer the reader to Diestel [7] for any undefined notation. The *degree* (resp. *open neighbourhood*) of a vertex $v \in V(G)$ is denoted by $d_G(v)$ (resp. $N_G(v)$). We denote the subgraph induced by S by $G[S]$. $G \setminus S = G[V(G) \setminus S]$. The *maximum* degree of any vertex of G is denoted by Δ . We denote paths and cycles by tuples of vertices. The *girth* of G is the length of a shortest cycle of G . Given two vertices u and v lying in the same connected component of G , we say an edge e *separates* u and v if they lie in different connected components of $G \setminus \{e\}$.

A graph is called *odd* (even, respectively) if every vertex has odd (respectively, even) degree. A partition (V_1, \dots, V_k) of $V(G)$ is a *k-odd colouring* of G if $G[V_i]$ induces an odd subgraphs of G for every $1 \leq i \leq k$. We say a graph is *k-odd colourable* if it admits a *k-odd colouring*. The *odd chromatic number* of G , denoted by $\chi_{\text{odd}}(G)$, is the smallest integer k such that G is *k-odd colourable*. The empty graph (i.e., the graph such that $V(G) = \emptyset$) is considered to be both even and odd. Since odd colouring exists only for graphs whose every component has even size and the colouring of each component does not affect the colouring of the others, we can consider each component separately. Therefore, it suffices to prove the statements for connected graphs of even order.

We would like to point out here that the definition of odd colouring used in this work is not to be confused with the one introduced by Petrusevski and Skrekovski [12], which is a specific type of proper colouring.

Modular width A set S of vertices is called a *module* if, for all $u, v \in S$, $N(u) \setminus S = N(v) \setminus S$. A partition $\mathcal{M} = (M_1, \dots, M_k)$ of $V(G)$ is a module partition of G if every M_i is a module in G . Without loss of generality, we further ask that any module partition \mathcal{M} of G , unless $G = K_1$, is nontrivial, that is, \mathcal{M} has at least two nonempty parts. Given two sets of vertices X and Y , we say that X and Y are *complete to each other* (*completely nonadjacent*, respectively) if $uv \in E(G)$ ($uv \notin E(G)$, respectively) for every $u \in X, v \in Y$. Note that for any two modules M and N in G , either M and N are nonadjacent or complete to each other. We let $G_{\mathcal{M}}$ be the module graph of \mathcal{M} , that is, the graph on vertex set \mathcal{M} with an edge between M_i and M_j if and only if M_i and M_j are complete to each other (nonadjacency between modules M_i, M_j in $G_{\mathcal{M}}$ corresponds to M_i and M_j being nonadjacent in G). We define the modular width of a graph G , denoted by $\text{mw}(G)$, recursively as follows. $\text{mw}(K_1) = 1$, the width of a module partition (M_1, \dots, M_k) of G is the maximum of k and $\text{mw}(G[M_i])$ for all $i \in [k]$ and $\text{mw}(G)$ is the minimum width of any module partitions of G .

3 | GRAPHS OF BOUNDED DEGREE AND GRAPHS OF LARGE GIRTH

In this section, we study Scott's conjecture (Conjecture 1) as well as the conjecture made by Aashtab et al. [1] which states that $\chi_{\text{odd}}(G) \leq \Delta + 1$ for any graph G . We settle Conjecture 1 for graphs of girth at least 7, and prove that $\chi_{\text{odd}}(G) \leq 2\Delta - 1$ for any graph G , thus obtaining a weaker version of the conjecture of Aashtab et al. To this end, we prove the following more general theorem, which implies both of the aforementioned results.

Theorem 4. *Let \mathcal{H} be a class of graphs such that:*

- $K_2 \in \mathcal{H}$
- \mathcal{H} is closed under vertex deletion and
- there is a $k \geq 2$ such that any connected graph $G \in \mathcal{H}$ satisfies at least one of the following properties:
 - (I) G has two pendant vertices u, v such that $N_G(u) = N_G(v)$ or
 - (II) G has two adjacent vertices u, v such that $d_G(u) + d_G(v) \leq k$.

Then any connected graph $G \in \mathcal{H}$ of even order has $\chi_{\text{odd}}(G) \leq k - 1$.

Proof. First notice that \mathcal{H} is well defined as K_2 has the desired properties. The proof is by induction on the number of vertices. Let $|V(G)| = 2n$.

For $n = 1$, since G is connected, we have that $G = K_2$ which is odd. Therefore, $\chi_{\text{odd}}(G) = 1 \leq k - 1$ (recall that $k \geq 2$). Let G be a graph of order $2n$. Notice that we only need to consider the case where G is connected as, otherwise, we can apply the inductive hypothesis to each of the components of G . Assume first that G has two pendant vertices u, v such that $N_G(u) = N_G(v) = \{w\}$. Then, since $G - \{u, v\}$ is connected and belongs to \mathcal{H} , by induction, there is an odd colouring of $G - \{u, v\}$ that uses at most $k - 1$ colours. Let (V_1, \dots, V_{k-1}) be a partition of $V(G) \setminus \{u, v\}$ such that $G[V_i]$ is odd for all $i \in [k - 1]$. We may assume that $w \in V_1$. We give a partition V'_1, \dots, V'_{k-1} of $V(G)$ by setting $V'_1 = V_1 \cup \{u, v\}$ and $V'_i = V_i$ for all $i \in [k] \setminus \{1\}$. Notice that for all $i \in [k - 1]$, $G[V'_i]$ is odd. Therefore, $\chi_{\text{odd}}(G) \leq k - 1$.

Thus, we assume that G has an edge $uv \in E(G)$ such that $d_G(u) + d_G(v) \leq k$. Note that we may assume that $k \geq 3$ for otherwise the theorem follows. We consider two cases; $G[V(G) \setminus \{u, v\}]$ is connected and $G[V(G) \setminus \{u, v\}]$ is disconnected.

Assume that $G[V(G) \setminus \{u, v\}]$ is connected. Since $G[V(G) \setminus \{u, v\}]$ has $|V(G) \setminus \{u, v\}| = 2n - 2$ and belongs to \mathcal{H} , by induction, there is an odd colouring of it that uses at most $k - 1$ colours. Let (V_1, \dots, V_{k-1}) be a partition of $V(G) \setminus \{u, v\}$, such that $G[V_i]$ is odd of all $i \in [k - 1]$. We give a partition of G into $k - 1$ odd graphs as follows. Since $|N_G(\{u, v\})| \leq k - 2$, there exists $\ell \in [k - 1]$ such that $V_\ell \cap N_G(\{u, v\}) = \emptyset$. We define a partition (U_1, \dots, U_{k-1}) of $V(G)$ as follows. For all $i \in [k - 1]$, if $i \neq \ell$, we define $U_i = V_i$, otherwise we set $U_i = V_i \cup \{u, v\}$. Notice that for all $i \neq \ell$, $G[U_i]$ is odd since $U_i = V_i$. Also, since $N_{G[U_\ell]}[u] = N_{G[U_\ell]}[v] = \{u, v\}$ and $G[V_\ell]$ is odd, we conclude that $G[U_\ell]$ is odd. Thus, $\chi_{\text{odd}}(G) \leq k - 1$.

Now, we consider the case where $G \setminus \{u, v\}$ is disconnected. First, we assume that there is at least one component in $G \setminus \{u, v\}$ of even order. Let U be the set of vertices of this component. By induction, $\chi_{\text{odd}}(G[U]) \leq k - 1$ and $\chi_{\text{odd}}(G \setminus U) \leq k - 1$. Furthermore, $|N_G(\{u, v\}) \cap U| \leq k - 3$ because $G \setminus \{u, v\}$ has at least two components. Let (U_1, \dots, U_{k-1}) be a partition of U such that $G[U_i]$ is odd for all $i \in [k - 1]$. Also, let (V_1, \dots, V_{k-1}) be a partition of $V(G) \setminus U$ such that $G[V_i]$ is odd for all $i \in [k - 1]$. We may assume that $V_i \cap \{u, v\} = \emptyset$ for all $i \in [k - 3]$. Since $|N_G(\{u, v\}) \cap U| \leq k - 3$, there are at least two indices $l, l' \in [k - 1]$ such that $U_l \cap N_G(\{u, v\}) = U_{l'} \cap N_G(\{u, v\}) = \emptyset$. We may assume that $l = k - 2$ and $l' = k - 1$. We define a partition (V'_1, \dots, V'_{k-1}) of $V(G)$ as follows. For all $i \in [k - 1]$ we define $V'_i = V_i \cup U_i$. We claim that $G[V'_i]$ is odd for all $i \in [k - 1]$. To show the claim, we consider two cases; either $V'_i \cap \{u, v\} = \emptyset$ or not. If $V'_i \cap \{u, v\} = \emptyset$, since the only vertices in $V(G) \setminus U$ that can have neighbours in U are v and u , we have that $G[V'_i]$ is odd. Indeed, this holds because $U_i \cap N_G(V_i) = \emptyset$ and both $G[U_i]$ and $G[V_i]$ are odd. If $V'_i \cap \{u, v\} \neq \emptyset$, then $i = k - 2$ or $i = k - 1$. In both cases, we know that $U_i \cap N_G(V_i) = \emptyset$ because the only vertices in $V(G) \setminus U$ that may have neighbours in U are v and u and we have assumed that u, v do not have neighbours in $U_{k-2} \cup U_{k-1}$. So, $G[V'_i]$ is odd because $U_i \cap N_G(V_i) = \emptyset$ and both $G[U_i]$ and $G[V_i]$ are odd.

Thus, we can assume that all components of $G \setminus \{u, v\}$ are of odd order. Let $\ell > 0$ be the number of components, denoted by V_1, \dots, V_ℓ , of $G \setminus \{u, v\}$ and note that ℓ must be even. We consider two cases, either for all $i \in [\ell]$, one of $G[V_i \cup \{u\}]$ or $G[V_i \cup \{v\}]$ is disconnected, or there is at least one $i \in [\ell]$ such that both $G[V_i \cup \{u\}]$ and $G[V_i \cup \{v\}]$ are connected.

In the first case, for each $V_i, i \in [\ell]$ we call w_i the vertex in $\{u, v\}$ such that $G[V_i \cup \{w_i\}]$ is connected. Note that w_i is uniquely determined, that is, only one of u and v can be w_i for each $i \in [\ell]$. Now, by induction, for all $i \in [\ell]$, $G[V_i \cup \{w_i\}]$ has $\chi_{\text{odd}}(G[V_i \cup \{w_i\}]) \leq k - 1$. Let, for each $i \in [\ell]$, $(V^i_1, \dots, V^i_{k-1})$ denote a partition of $V_i \cup \{w_i\}$ such that $G[V^i_j]$ be odd, for all $j \in [k - 1]$. Furthermore, we may assume that for each $i \in [\ell]$, if $v \in V_i \cup \{w_i\}$, then $v \in V^i_{k-2}$. Also, we can assume that for each $i \in [\ell]$, if $u \in V_i \cup \{w_i\}$, then $u \in V^i_{k-1}$. Finally, let $I = \{i \in [\ell] \mid w_i = v\}$ and $J = \{i \in [\ell] \mid w_i = u\}$.

We consider two cases. If $|I|$ is odd, then $|J|$ is odd since $\ell = |I| + |J|$ is even. Then, we claim that for the partition (U_1, \dots, U_{k-1}) of $V(G)$ where $U_i = \bigcup_{j \in [I]} V^j_i$ it holds that $G[U_i]$ is odd for all $i \in [k - 1]$. First notice that (U_1, \dots, U_{k-1}) is indeed a partition of $V(G)$. Indeed, the only vertices that may belong in more than one set are u and v . However, v belongs only to some sets V^i_{k-2} , and hence it is no set U_i except U_{k-2} . Similarly, u belongs

to no set U_i except U_{k-1} . Therefore, it remains to show that $G[U_i]$ is odd for all $i \in [k - 1]$. We will show that for any $i \in [k - 1]$ and for any $x \in U_i$, $|N_G(x) \cap U_i|$ is odd. Let $x \in U_i \setminus \{u, v\}$, for some $i \in [k - 1]$. Then we know that $N_G(x) \cap U_i = N_G(x) \cap V_i^j$ for some $j \in [\ell]$. Since $G[V_i^j]$ is odd for all $i \in [k - 1]$ and $j \in [\ell]$ we have that $|N_G(x) \cap U_i| = |N_G(x) \cap V_i^j|$ is odd. Therefore, we only need to consider u and v . Notice that $v \in U_{k-2} = \bigcup_{j \in [\ell]} V_{k-2}^j$ (respectively, $u \in U_{k-1} = \bigcup_{j \in [\ell]} V_{k-1}^j$). Also, v (respectively, u) is included in V_{k-2}^j (respectively, V_{k-1}^j) only if $j \in I$ (respectively, $j \in J$). Since $G[V_{k-2}^j]$ (respectively, $G[V_{k-1}^j]$) is odd for any $j \in [\ell]$ we have that $|N(v) \cap V_{k-2}^j|$ (respectively, $|N(u) \cap V_{k-1}^j|$) is odd for any $j \in I$ (resp. $j \in J$). Finally, since $|I|$ and $|J|$ are odd, we have that $|N_G(v) \cap U_{k-2}| = \sum_{j \in I} |N(v) \cap V_{k-2}^j|$ and $|N_G(u) \cap U_{k-1}| = \sum_{j \in J} |N(u) \cap V_{k-1}^j|$ are both odd. Therefore, for any $i \in [k - 1]$, $G[U_i]$ is odd and $\chi_{\text{odd}}(G) \leq k - 1$.

Now, suppose that both $|I|$ and $|J|$ are even. We consider the partition (U_1, \dots, U_{k-1}) of $V(G)$ where, for all $i \in [k - 3]$ $U_i = \bigcup_{j \in [\ell]} V_i^j$, $U_{k-2} = \bigcup_{j \in I} V_{k-2}^j \cup \bigcup_{j \in J} V_{k-1}^j$ and $U_{k-1} = \bigcup_{j \in I} V_{k-2}^j \cup \bigcup_{j \in J} V_{k-1}^j$. We claim that for this partition it holds that $G[U_i]$ is odd for all $i \in [k - 1]$. First notice that (U_1, \dots, U_{k-1}) is indeed a partition of $V(G)$. Indeed, this is clear for all vertices except for u and v . However, v only belongs to sets of type V_{k-2}^j for $i \in I$, and u only belongs to sets of type V_{k-1}^j for $i \in J$. Therefore, u or v belong to no set U_i except U_{k-1} . We will show that for any $i \in [k - 1]$ and $x \in U_i$, $|N_G(x) \cap U_i|$ is odd. Let $x \in U_i \setminus \{u, v\}$, for some $i \in [k - 1]$. Then we know that $N_G(x) \cap U_i = N_G(x) \cap V_i^j$ for some $j \in [\ell]$. Since $G[V_i^j]$ is odd for all $i \in [k - 1]$ and $j \in [\ell]$ we have that $|N_G(x) \cap U_i| = |N_G(x) \cap V_i^j|$ is odd. Therefore, we only need to consider u and v . Note that $u, v \in U_{k-1}$. Since both $|I|$ and $|J|$ are even and $U_{k-1} = \bigcup_{j \in I} V_{k-2}^j \cup \bigcup_{j \in J} V_{k-1}^j$, we have that $|N_G(v) \cap U_{k-1} \setminus \{u\}|$ and $|N_G(u) \cap U_{k-1} \setminus \{v\}|$ are both even. Finally, since $uv \in E(G)$ we have that $|N_G(u) \cap U_{k-1}|$ and $|N_G(v) \cap U_{k-1}|$ are both odd. Hence, $\chi_{\text{odd}}(G) \leq k - 1$.

Now we consider the case where there is at least one $i \in [\ell]$ where both $G[V_i \cup \{u\}]$ and $G[V_i \cup \{v\}]$ are connected. We define the following sets I and J . For each $i \in [\ell]$

- $i \in I$, if $G[V_i \cup \{u\}]$ is disconnected and
- $i \in J$, if $G[V_i \cup \{v\}]$ is disconnected.

Finally, for the rest of the indices, $i \in [\ell]$, which are not in $I \cup J$, it holds that both $G[V_i \cup \{u\}]$ and $G[V_i \cup \{v\}]$ are connected. Call this set of indices X and note that by assumption $|X| \geq 1$. Since $|I| + |J| + |X|$ is even, it is easy to see that there is a partition of X into two sets X_1 and X_2 such that both $I' := I \cup X_1$ and $J' := J \cup X_2$ have odd size. Let $V_I = \bigcup_{i \in I'} V_i$ and $V_J = \bigcup_{i \in J'} V_i$. Now, by induction, we have that $\chi_{\text{odd}}(G[V_I \cup \{v\}]) \leq k - 1$ and $\chi_{\text{odd}}(G[V_J \cup \{u\}]) \leq k - 1$. Assume that $(V_1^I, \dots, V_{k-1}^I)$ is a partition of V_I and $(V_1^J, \dots, V_{k-1}^J)$ is a partition of V_J such that for any $i \in [k - 1]$, $G[V_i^I]$ and $G[V_i^J]$ are odd. Without loss of generality, we may assume that $v \in V_1^I$ and $u \in V_{k-1}^J$. Since $|X| \geq 1$, note that both $d_G(u)$ and $d_G(v)$ are at least two, which implies that $d_G(u) \leq k - 2$ and $d_G(v) \leq k - 2$. Therefore, there exists $i_0 \in [k - 2]$ such that $N_G(v) \cap V_{i_0}^I = \emptyset$ and $j_0 \in [k - 1] \setminus \{1\}$ such that $N_G(v) \cap V_{j_0}^J = \emptyset$. We reorder the sets $V_i^I, i \in [k - 2]$, so that $i_0 = 1$ and we reorder the sets $V_i^J, i \in [k - 1] \setminus \{1\}$ so that $j_0 = k - 1$. Note that this reordering does not change the fact that $v \in V_1^I$ and $u \in V_{k-1}^J$. Consider the partition

(U_1, \dots, U_{k-1}) of $V(G)$, where $U_i = V_i^I \cup V_i^J$. We claim that for all $i \in [k-1]$, $G[U_i]$ is odd. Note that for any $x \in U_i$, we have $N_G(x) \cap U_i = N_G(x) \cap V_i^I$ or $N_G(x) \cap U_i = N_G(x) \cap V_i^J$. Since for any $i \in [k-1]$, $G[V_i^I]$ and $G[V_i^J]$ are odd we conclude that $G[U_i]$ is odd for any $i \in [k-1]$. \square

Notice that the class of graphs G of maximum degree Δ satisfies the requirements of Theorem 4. Indeed, this class is closed under vertex deletions and any connected graph in the class has least two adjacent vertices u, v such that $d_G(u) + d_G(v) \leq 2\Delta$. Therefore, the following corollary holds.

Corollary 1. *Any connected graph G of even order and maximum degree Δ has $\chi_{\text{odd}}(G) \leq 2\Delta - 1$.*

Next, we prove Conjecture 1 for graphs of girth at least seven.

Corollary 2. *Any connected graph G of even order and of girth at least 7 has $\chi_{\text{odd}}(G) \leq \frac{3\sqrt{n}}{2} + 1$ where $n = |V(G)|$.*

Proof. Let \mathcal{G}_7 be the class of graphs of girth at least 7. Note that \mathcal{G}_7 is closed under vertex deletion. Therefore, we need prove that any connected graph $G \in \mathcal{G}_7$ of even order that does not satisfy the property (I) of the Theorem 4 has at least two adjacent vertices u, v such that $d_G(u) + d_G(v) \leq 3\sqrt{n}/2 + 2$. Then, the corollary follows from the Theorem 4.

Claim 1. Let G be a graph in \mathcal{G}_7 of order n . If G does not have two pendant vertices u, v such that $N_G(u) = N_G(v)$, then it has two adjacent vertices u', v' such that $d_G(u') + d_G(v') \leq \frac{3\sqrt{n}}{2} + 2$.

Proof of Claim. Assume that for any two adjacent vertices u, v such that $d_G(u) + d_G(v) \geq 3\sqrt{n}/2 + 3$. Let G' be the graph we obtain after we remove all pendant vertices of G . Since each vertex of G had at most one pendant vertex we have that for any edge $uv \in E(G')$, $d_{G'}(u) + d_{G'}(v) \geq 3/2\sqrt{n} + 1$. Also notice that G' does not contain any pendant vertices as otherwise this vertex was attached to a pendant vertex of G and this gives us an edge $uv \in E$ such that $d_G(u) + d_G(v) = 3 \leq 3\sqrt{n}/2 + 2$.

Let w be a vertex such that $d_{G'}(w) \geq 3\sqrt{n}/4 + 1/2$. We consider all the vertices of distance at most 3 from w in G' . Let V_1 be the set of vertices of distance one from w , V_2 be the set of vertices of distance two from w and V_3 be the set of vertices of distance three from w . Notice that, since G has girth at least 7 we have that both V_1 and V_2 are independent sets, no two vertices in V_1 have a common neighbour in V_2 , and no two vertices in V_2 have a common neighbour in V_3 .

We will compute the minimum number of vertices in these sets. For any $j \in \{1, 2\}$, let $|V_j| = m_j$, $v_{j,i}$, $i \in [m_j]$, be the vertices of V_j and $d_{j,i} = d_{G'}(v_{j,i})$ for all $i \in [m_j]$. For each vertex $v_{1,i}$, $i \in [m_1]$, select $i' \in [m_2]$ such that $v_{2,i'} \in N_G(v_{1,i}) \setminus \{w\}$. We note that the selected vertices i' are necessarily distinct for each vertex $v_{1,i} \in V_1$. We have $|N_{G'}[\{v_{1,i}, v_{2,i'}\}] \setminus \{w\}| = d_{1,i} + d_{2,i'} - 1 \geq 3\sqrt{n}/2$. It follows that $|V(G)| \geq \sum_{i \in [m_1]} (3\sqrt{n}/2) = m_1(3\sqrt{n}/2) \geq 9n/8 + 3\sqrt{n}/4 > n$. This is a contradiction since G has n vertices. \square

One may wonder if graphs of sufficiently large girth may have bounded odd chromatic number. In fact, this is far from being true, which we show in the next proposition. Recall that the *chromatic number* $\chi(G)$ of a graph G is the smallest integer k such that $V(G)$ can be partitioned into k sets each of which is independent.

Proposition 1. *For every integer g and k , there exist graphs of even order and of girth at least g such that $\chi_{\text{odd}}(G) \geq k$.*

Proof. We use a classical result of Erdős [8], which states that for all sufficiently large n , there exists a n -vertex graph G of girth at least g and $\chi(G) \geq k$. Let G be such a graph, with n even. We may assume that G has no component of odd order (otherwise, we can add an edge between any pair of odd components without affecting the girth or decreasing the chromatic number). Let H be the graph obtained from G by subdividing each edge of G once. We claim that $\chi_{\text{odd}}(H) \geq k$. Suppose that $\chi_{\text{odd}}(H) \leq k - 1$ and let U_1, \dots, U_{k-1} be a partition of $V(H)$ such that $G[U_i]$ is odd for each $i \in [k - 1]$. Since $\chi(G) \geq k$, there must exist two adjacent vertices $u, v \in V(G)$ such that both $\{u, v\} \in U_i$ for some $i \in [k - 1]$. But we know that there is a vertex w_{uv} in H with $N_H(w_{uv}) = \{u, v\}$. Let U_j be the set containing w_{uv} . Then U_j is not odd, a contradiction. \square

Remark 1. In fact, by using a stronger result of Bollobás [3], it is possible to show that for every g , there is $\epsilon > 0$ such that for all even n sufficiently large, there exist connected graphs G of order n and girth at least g , with $\chi_{\text{odd}}(G) > n^\epsilon$.

Next, we obtain the following result for sparse planar graphs.

Corollary 3. *Any connected planar graph G of even order and of girth at least 11 has $\chi_{\text{odd}}(G) \leq 3$.*

Proof. Let \mathcal{G} be the class of planar graphs of girth at least 11. Notice that this class is closed under vertex deletion. We will show that for any graph $G \in \mathcal{G}$ at least one of the following properties holds:

- (I) G has two pendant vertices $u, v \in V(G)$ such that $N_G(u) = N_G(v)$ or
- (II) G has an edge $uv \in E(G)$ such that $d_G(u) + d_G(v) \leq 4$

Assume that G does not satisfy the property (I). We construct G' by deleting all pendant vertices of G . If the minimum degree of G' is 1, then the property (II) holds for G . Indeed, if G' has a pendant vertex u , then must have a pendant vertex v in G . Therefore, $d_G(u) + d_G(v) = 2 + 1 \leq 4$.

Assume that G' has minimum degree 2. Since G' is also planar and has girth at least 11 we can apply the Theorem 4.11 (Chang and Duh [5]), which states that there exists an edge $uv \in E(G')$ such that $d_{G'}(u) = d_{G'}(v) = 2$. We consider two cases: either one of u and v were attached to a pendant vertex v in G or none of them were attached to a pendant vertex of G . In the first case, we may assume that u is attached to a pendant vertex w of G . Then we have $d_G(u) + d_G(w) = 3 + 1 \leq 4$, therefore G satisfies the

property (II). In the latter case, both u and v have $d_G(u) = d_G(v) = 2$. Then G satisfies the property (II).

Now, by applying Theorem 4 to the class \mathcal{G} the corollary follows. \square

Remark 2. The upper bound presented in Corollary 3 is tight as C_{14} , the cycle of length 14, has $\chi_{\text{odd}}(C_{14}) = 3$.

4 | GRAPHS OF BOUNDED MODULAR-WIDTH

In this section, we consider graphs of bounded modular-width and show that we can upper bound the odd chromatic number by the modular-width of a graph.

Theorem 5. *For every graph G with all components of even order, $\chi_{\text{odd}}(G) \leq 3\text{mw}(G)$.*

The following is an easy consequence of Theorem 1 which will be useful to colour modules and gain control over the parity of parts in the case the module is of even size.

Remark 3. For every nonempty graph G of even order, there exists a partition (V_1, V_2, V_3) of $V(G)$ with $|V_2|, |V_3|$ being odd such that $G[V_1]$ is odd and $G[V_2], G[V_3]$ are even. This can be derived from Theorem 1 by taking an arbitrary vertex $v \in V(G)$, setting $V_3 := \{v\}$ and then using the existence of a partition (V_1, V_2) of $V(G) \setminus \{v\}$ such that $G[V_1]$ is odd and $G[V_2]$ is even.

To prove Theorem 5, we first show that every graph G is 3-colourable for which we have a module partition \mathcal{M} such that the module graph $G_{\mathcal{M}}$ exhibits a particular structure, that is, is either a star Lemma 1 or a special type of tree Lemma 2. (Figure 2).

Lemma 1. *For every connected graph G of even order with a module partition $\mathcal{M} = \{M_1, \dots, M_k\}$ such that $G_{\mathcal{M}}$ is a star, $\chi_{\text{odd}}(G) \leq 3$.*

Proof. Assume that in $G_{\mathcal{M}}$ the vertices M_2, \dots, M_k have degree 1. We refer to M_1 as the centre and to M_2, \dots, M_k as leaves of $G_{\mathcal{M}}$. We further assume that $|M_2|, \dots, |M_{\ell}|$ are odd and $|M_{\ell+1}|, \dots, |M_k|$ are even for some $\ell \in [k]$. We use the following two claims.

Claim 2. If $W \subseteq V(G)$ such that $G[W \cap M_i]$ is odd for every $i \in [k]$, then $G[W]$ is odd.

Proof of Claim. First observe that the degree of any vertex $v \in W \cap M_1$ in $G[W]$ is $d_{G[W \cap M_1]}(v) + \sum_{i=2}^k |W \cap M_i|$. Since $d_{G[W \cap M_1]}(v)$ is odd and $|W \cap M_i|$ is even for every $i \in \{2, \dots, k\}$ (which follows from $G[W \cap M_i]$ being odd by the handshake lemma) we get that $d_{G[W]}(v)$ is odd. For every $i \in \{2, \dots, k\}$ the degree of any vertex $v \in W \cap M_i$ in $G[W]$ is $d_{G[W \cap M_i]}(v) + |W \cap M_1|$ which is odd (again, because $|W \cap M_1|$ must be even). Hence $G[W]$ is odd. \square

Claim 3. If $W \subseteq V(G)$ such that $G[W \cap M_i]$ is even for every $i \in [k]$, $|W \cap M_1|$ is odd and $|\{i \in \{2, \dots, k\} : |W \cap M_i| \text{ is odd}\}|$ is odd, then $G[W]$ is odd.




FIGURE 2 Schematic illustration of the three cases in the proof of Lemma 1. Depicted is the module graph G_M along with a partition of the modules into sets V_1, V_2 and V_3 such that $G[V_i]$ is odd for $i \in [3]$. Specifically, the left figure depicts the case where the centre is of odd size, the middle figure depicts the case where the centre is of even size and there is no odd sized leaf and the right figure depicts the case where the centre is of even size and there is at least one odd sized leaf. [Color figure can be viewed at wileyonlinelibrary.com]

Proof of Claim. Since G_M is a star and M_1 its centre we get that the degree of any vertex $v \in W \cap M_i$ for any $i \in \{2, \dots, k\}$ is $d_{G[W \cap M_i]}(v) + |W \cap M_1|$. Since $|W \cap M_1|$ is odd and $d_{G[W \cap M_i]}(v)$ is even we get that every vertex $v \in W \cap M_i$ for every $i \in \{2, \dots, k\}$ has odd degree in $G[W]$. On the other hand, the degree of $v \in W \cap M_1$ is $d_{G[W \cap M_1]}(v) + \sum_{i=2}^k |W \cap M_i|$. Since $d_{G[W \cap M_1]}(v)$ is even and $|\{i \in \{2, \dots, k\} : |W \cap M_i| \text{ is odd}\}|$ is odd $d_{G[W]}(v)$ is odd. We conclude that $G[W]$ is odd. \square

First, consider the case that $|M_1|$ is odd. Since G is of even order this implies that there must be an odd number of leaves of G_M of odd size and hence ℓ is even. Using Theorem 1, we let (W_1^i, W_2^i) be a partition of M_i such that $G[W_1^i]$ is odd and $G[W_2^i]$ is even for every $i \in [k]$. Note that since $G[W_1^i]$ is odd $|W_1^i|$ has to be even and hence $|W_2^i|$ is odd if and only if $i \in [\ell]$. We define $V_1 := \cup_{i \in [k]} W_1^i$ and $V_2 := \cup_{i \in [k]} W_2^i$. Note that (V_1, V_2) is a partition of G . Furthermore, $G[V_1]$ is odd by Claim 2 and $G[V_2]$ is odd by Claim 3. For an illustration, we refer the reader to Figure 2.

Now consider the case that $|M_1|$ is even. We first consider the special case that $\ell = 1$, that is, there is no $i \in [k]$ such that $|M_i|$ is odd. In this case, we let (W_1^i, W_2^i, W_3^i) be a partition of M_i for $i \in \{1, 2\}$ such that $G[W_1^i]$ is odd, $G[W_2^i], G[W_3^i]$ are even and $|W_2^i|, |W_3^i|$ are odd which exists due to Remark 3. For $i \in \{3, \dots, k\}$ we let (W_1^i, W_2^i) be a partition of M_i such that $G[W_1^i]$ is odd and $G[W_2^i]$ is even which exists by Theorem 1. We define $V_1 := \cup_{i \in [k]} W_1^i, V_2 := \cup_{i \in [k]} W_2^i$ and $V_3 := W_3^1 \cup W_3^2$. As before we observe that

(V_1, V_2, V_3) is a partition of $V(G)$, $G[V_1]$ is odd by Claim 2 and $G[V_2], G[V_3]$ are even by Claim 3. For an illustration see Figure 2.

Lastly, consider the case that $|M_1|$ is even and $\ell > 1$. By Remark 3 there is a partition (W_1^1, W_2^1, W_3^1) of M_1 such that $G[W_1^1]$ is odd, $G[W_2^1], G[W_3^1]$ are even and $|W_2^1|, |W_3^1|$ are odd. For $i \in \{2, \dots, k\}$ we let (W_1^i, W_2^i) be a partition of M_i such that $G[W_1^i]$ is odd and $G[W_2^i]$ is even which exists by 1. We define $V_1 := \bigcup_{i \in [k]} W_1^i$, $V_2 := W_2^1 \cup \bigcup_{i=3}^k W_2^i$ and $V_3 := W_3^1 \cup W_2^2$. Note that (V_1, V_2, V_3) is a partition of $V(G)$. Furthermore, $G[V_1]$ is odd by Claim 2 and $G[V_3]$ is odd by 3. Additionally, since $|M_1|$ is even there is an even number of $i \in \{2, \dots, k\}$ such that $|M_i|$ is odd. Since for each $i \in \{2, \dots, k\}$ for which $|M_i|$ is odd, $|W_1^i|$ must be odd, we get that $|\{i \in \{2, \dots, k\} : |V_1 \cap M_i| \text{ is odd}\}|$ is odd (note that $V_1 \cap M_2 = \emptyset$ because $W_2^2 \subseteq V_3$). Hence we can use Claim 3 to conclude that $G[V_2]$ is odd. For an illustration see Figure 2. \square

Let G be a connected graph of even order with module partition $\mathcal{M} = (M_1, \dots, M_k)$ such that $G_{\mathcal{M}}$ is a tree. For an edge e of $G_{\mathcal{M}}$ we let X_e and Y_e be the two components of the graph obtained from $G_{\mathcal{M}}$ by removing e . We say that the tree $G_{\mathcal{M}}$ is colour propagating if the following properties hold.

- (i) $|\mathcal{M}| \geq 3$.
- (ii) Every non-leaf module has size one.
- (iii) $|\bigcup_{M \in V(X_e)} M|$ is odd for every edge $e \in E(G_{\mathcal{M}})$ which is not incident to any leaf of $G_{\mathcal{M}}$.

Lemma 2. *For every connected graph G of even order with a module partition $\mathcal{M} = (M_1, \dots, M_k)$ such that $G_{\mathcal{M}}$ is a colour propagating tree, $\chi_{\text{odd}}(G) \leq 2$.*

Proof. To find an odd colouring (V_1, V_2) of G , we first let (W_1^i, W_2^i) be a partition of M_i such that $G[W_1^i]$ is odd and $G[W_2^i]$ is even for every $i \in [k]$. The partitions (W_1^i, W_2^i) exist due to Theorem 1. Note that (ii) implies that for every module M_i which is not a leaf $|W_2^i| = 1$ and $W_1^i = \emptyset$. We define $V_1 := \bigcup_{i \in [k]} W_1^i$ and $V_2 := \bigcup_{i \in [k]} W_2^i$.

To argue that (V_1, V_2) is an odd colouring of G first consider any $v \in V(G)$ such that $v \in M_i$ for some leaf M_i of $G_{\mathcal{M}}$. Condition (i) implies that $G_{\mathcal{M}}$ must have at least three vertices and hence the neighbour M_j of M_i cannot be a leaf due to $G_{\mathcal{M}}$ being a tree. Hence $|M_j| = 1$ by (ii). Hence, if $v \in W_1^i$, then $d_{G[V_1]}(v) = d_{G[W_1^i]}(v)$ since $W_1^j = \emptyset$ and therefore $d_{G[V_1]}(v)$ is odd. Further, if $v \in W_2^i$, then $d_{G[V_2]}(v) = d_{G[W_2^i]}(v) + 1$ since $|W_2^j| = 1$ and hence $d_{G[V_2]}(v)$ is odd. Hence the degree of any vertex $v \in M_i$ is odd in $G[V_1], G[V_2]$ respectively.

Now consider any vertex $v \in V(G)$ such that $M_i = \{v\}$ for some non-leaf M_i of $G_{\mathcal{M}}$. Let $M_{i_1}, \dots, M_{i_\ell}$ be the neighbours of M_i in $G_{\mathcal{M}}$. Let e_j be the edge $M_i M_{i_j} \in E(G)$ for every $j \in [\ell]$. Without loss of generality, assume that $M_i \notin V(X_{e_j})$ for every $j \in [\ell]$. By (iii) we have that $|\bigcup_{M \in V(X_{e_j})} M|$ is odd whenever M_{i_j} is not a leaf in $G_{\mathcal{M}}$. Hence, by (ii), $|\bigcup_{M \in V(X_{e_j})} M| \equiv |M_{i_j}| \pmod{2}$ for every $j \in [\ell]$ for which M_{i_j} is not a leaf in $G_{\mathcal{M}}$. On the other hand, as a consequence of the handshake lemma we get that $|W_2^{i_j}|$ is odd if and only if $|M_{i_j}|$ is odd. Hence the following holds for the parity of the degree of v in $G[V_2]$.

$$d_{G[V_2]}(v) = |\{j \in [m] : d_{G_M}(M_{i_j}) \geq 2\}| + \sum_{\substack{j \in [m] \\ d_{G_M}(M_{i_j})=1}} |W_2^{i_j}| \equiv |V(G) \setminus M_i| \pmod{2}.$$

Since G has even order we conclude that $d_{G[V_2]}(v)$ is odd and hence (V_1, V_2) is an odd colouring of G . □

We now show that, given a graph G with module partition \mathcal{M} , we can decompose the graph in such a way that the module graph of any part of the decomposition is either a star or a colour propagating tree. Here we consider the module graph with respect to the module partition \mathcal{M} restricted to the part of the decomposition we are considering. To obtain the decomposition we use a spanning tree $G_{\mathcal{M}}$ and inductively find a non-separating star, i.e., a star whose removal does not disconnect the graph, or a colour propagating tree. To handle parity during this process we might separate a module into two parts of the decomposition.

Lemma 3. *For every graph G of even order and module partition $\mathcal{M} = (M_1, \dots, M_k)$ there is a partition $\widehat{\mathcal{M}}$ of $V(G)$ with at most $2k$ many parts such that there is a coarsening \mathcal{P} of $\widehat{\mathcal{M}}$ with the following properties. $|P|$ is even for every part P of \mathcal{P} . Furthermore, for every part P of \mathcal{P} we have that $\widehat{\mathcal{M}}|_P$ is a module partition of $G[P]$ and $G[P]_{\widehat{\mathcal{M}}|_P}$ is either a star (with at least two vertices) or a colour propagating tree.*

Proof. We use the following extensively throughout the proof.

Claim 4. If \mathcal{N} is a module partition of a graph H and $W \subseteq V(G)$ such that \mathcal{M}_W has at least two nonempty parts, then \mathcal{M}_W is a module partition of $G[W]$.

Proof of Claim. Assume that this is not the case and there is a part N of \mathcal{M}_W which is not a module in $G[W]$. By construction, there is a part N' of \mathcal{N} such that $N \subseteq N'$. Since N is not a module in $G[W]$ there are vertices $u, v \in N, w \in W \setminus N$ such that $uw \in E(G[W])$ and $vw \notin E(G[W])$. Since $N \subseteq N'$ this implies that N' cannot be a module in G , a contradiction. □

We use an induction on the number of modules in \mathcal{M} to find partitions $\widehat{\mathcal{M}}$ and \mathcal{P} . Observe that in case $G_{\mathcal{M}}$ is a star or a colour propagating tree we can set $\widehat{\mathcal{M}} := \mathcal{M}$ and $\mathcal{P} := (V(G))$ which satisfies the conditions of the statement. Hence assume that $G_{\mathcal{M}}$ is neither a star nor a colour propagating tree. We use the two following claims to conduct our inductive argument.

Claim 5. Let H be a graph with module partition $\mathcal{N} = (N_1, \dots, N_\ell)$ such that $H_{\mathcal{N}}$ is neither a star nor a colour propagating tree. If $H_{\mathcal{N}}$ is a tree, then there is a partition $\widehat{\mathcal{N}} = (\widehat{N}_1, \dots, \widehat{N}_\ell)$ of $V(G)$ with $\widehat{\ell} \leq \ell + 1$ and a coarsening $\mathcal{Q} = (Q_1, Q_2)$ of $\widehat{\mathcal{N}}$ with the following properties. $|Q_i|$ is even and $\widehat{\mathcal{N}}|_{Q_i}$ is a module partition of $H[Q_i]$ for $i \in [2]$. Furthermore, $H[Q_1]$ is connected and $H[Q_2]_{\widehat{\mathcal{N}}|_{Q_2}}$ is either a star or a colour propagating tree. Additionally, for any fixed index $i \in [\ell]$ we can enforce that $N_i \cap Q_1 \neq \emptyset$.

Proof of Claim. First observe that since $H_{\mathcal{N}}$ is a tree but neither a colour propagating tree nor a star we know that either

- there is a nonleaf vertex N in H_N with $|N| > 1$ which has at least one non-leaf neighbour or
- there is an edge $e \in E(H_N)$ not incident to any leaf of H_N and $|\bigcup_{N \in V(X_e)} N|$ is even.

Assume we have fixed $i \in [\ell]$ (N_i will be the part which is guaranteed to be partially contained in Q_1). For any part $N \neq N_i$ of \mathcal{N} we let e_N be an edge incident to N which separates N from N_i . Let X_N be the component of H_N after removing e_N which contains N . Let Z be the set of nonleaf parts $N \neq N_i$ in H_N such that either $|N| > 1$ or e_N is not incident to a leaf and X_N is of even order. Note that by our previous observation, we know that $Z \cup \{N_i\}$ cannot be empty. Finally, we in case $Z \neq \emptyset$ we let $N \in Z$ be a part with minimum $|V(X_N)|$ among all parts in Z . In case $Z = \emptyset$ we define $N := N_i$. In this case, we let e_N be an edge incident to N_i and some other non-leaf vertex and X_N the component containing N_i after removing e_N . Observe that in case $Z = \emptyset$ we get as an immediate consequence that $|N_i| > 1$ and $|X_N|$ is odd. Now observe that in any case, our choice of N guarantees that $|N'| = 1$ for every nonleaf part $N' \neq N$ of X_N and for every edge $e \in E(X_N)$ not incident to a leaf of X_N we have that $|\bigcup_{N' \in V(X_e)} N'|$ is odd. Furthermore, since N is not a leaf X_N has more than one vertex.

First, consider the case that $|N| = 1$. We set $\widehat{\mathcal{N}} := \mathcal{N}$, $Q_2 := \bigcup_{N' \in V(X_N)} N'$ and $Q_1 := V(H) \setminus Q_2$. Since $N \in Z$ and $|N| = 1$, we know that $|Q_1|$ and $|Q_2|$ are even. Furthermore, Claim 4 implies that \mathcal{M}_{Q_i} is a module partition of $H[Q_i]$ for $i \in [2]$. By construction $H[Q_1]$ is connected and $H[Q_2]_{\widehat{\mathcal{N}}|_{Q_2}}$ is a colour propagating tree. Lastly, observe that by choosing e_N to be an edge separating N from N_i (since $|N| = 1$ we get $N \neq N_i$) we get that $N_i \subseteq Q_1$ as required.

Now consider the case that $|N| > 1$. First, assume that all neighbours of N in X_N are leaves. In this case, let $N' \subseteq N$ such that $N' \neq \emptyset$ and $|N' \cup \bigcup_{N'' \in V(X_N)} N''|$ is even. Now we define $\widehat{\mathcal{N}}$ to be the partition obtained from \mathcal{N} by removing part N and adding N' and $N \setminus N'$. We further let $Q_2 := N' \cup \bigcup_{N'' \in V(X_N)} N''$ and $Q_1 := V(H) \setminus Q_2$. By construction $|Q_1|, |Q_2|$ are even. Furthermore, $\widehat{\mathcal{N}}|_{Q_1}$ must contain at least two parts since H_N is not a star. Hence $\widehat{\mathcal{N}}|_{Q_i}$ is a module partition of $H[Q_i]$ for $i \in [2]$ by Claim 4. Furthermore, $H(Q_1)$ is connected and $H[Q_2]_{\widehat{\mathcal{N}}|_{Q_2}}$ is a star. In case $N = N_i$ recall that X_N is of odd order and hence we can pick N' such that $N \setminus N' \neq \emptyset$ which implies $N_i \cap Q_1 \neq \emptyset$. Finally, in case $N \neq N_i$ we get $N_i \subseteq Q_1$ as in the previous case.

On the other hand, assume that N has at least one nonleaf neighbour N' in X_N . Choose an arbitrary vertex $n \in N$. We define $\widehat{\mathcal{N}}$ to be the partition obtained from \mathcal{N} by removing N and adding $\{n\}$ and $N \setminus \{n\}$. We additionally set $Q_2 := \{n\} \cup \bigcup_{N'' \in V(X_N)} N''$ and $Q_1 := V(H) \setminus Q_2$. Note that since NN' is an edge between non-leaf vertices we get that $|\bigcup_{N'' \in V(X_N)} N''|$ is odd and hence $|Q_1|$ and $|Q_2|$ must be even. Since $N \setminus \{n\} \neq \emptyset$ we get that \mathcal{M}_{Q_1} contains at least two parts and hence $\widehat{\mathcal{N}}|_{Q_1}$ is a module partition of $H[Q_i]$ for $i \in [2]$ by 4. Additionally, since $N \setminus \{n\} \neq \emptyset$ we have that $H(Q_1)$ must be connected. Finally, $H[Q_2]_{\widehat{\mathcal{N}}|_{Q_2}}$ is a colour propagating tree. The condition that $N_i \cap Q_1 \neq \emptyset$ is trivially satisfied in case $N = N_i$ and follows as before in case $N \neq N_i$. \square

Claim 6. Let T be any spanning tree of G_M . If there exists an edge $e \in E(G_M) \setminus E(T)$, then there is a partition $\widehat{\mathcal{N}} = (\widehat{\mathcal{N}}_1, \dots, \widehat{\mathcal{N}}_{\widehat{\ell}})$ of $V(G)$ with $\widehat{\ell} \leq k + 1$ and a coarsening

$\mathcal{Q} = (Q_1, Q_2)$ of $\widehat{\mathcal{N}}$ with the following properties. $|Q_i|$ is even and $\widehat{\mathcal{N}}|_{Q_i}$ is a module partition of $G[Q_i]$ for $i \in [2]$. Furthermore, $G[Q_1]$ is connected and $G[Q_2]|_{\widehat{\mathcal{N}}|_{Q_2}}$ is either a star or a colour propagating tree.

Proof of Claim. For any edge $e = MM' \in E(G_M) \setminus E(T)$ we let $e_1, e_2 \in E(T)$ such that e_1 is incident to M , e_2 is incident to M' , $(T \setminus \{e_1, e_2\}) \cup \{e\}$ has exactly two components and M is in the same component as M' in $(T \setminus \{e_1, e_2\}) \cup \{e\}$. For $e = MM' \in E(G_M) \setminus E(T)$ let C_e be the subgraph of G_M induced by the vertices of the component of $(T \setminus \{e_1, e_2\}) \cup \{e\}$ which contains M and M' . We now define e to be an edge minimizing $|V(C_e)|$. This means that C_e must be a tree. First consider the case that C_e is of even order and C_e is a colour propagating tree. First consider that $G_M \setminus C_e$ has at least two vertices. In this case, we can set $\widehat{\mathcal{N}} := \mathcal{M}$, $Q_2 := \bigcup_{M \in V(C_e)} M$ and $Q_1 := V(G) \setminus Q_2$ satisfying all requirements.

Hence assume that $G_M \setminus C_e$ consists of one vertex N . Hence in particular e_1, e_2 must be incident to N . Since C_e is of even order $|N|$ must be even. Partition N into two parts N', N'' of odd size and obtain $\widehat{\mathcal{N}}$ from \mathcal{M} by removing N and adding N' and N'' . Furthermore, since C_e is a colour propagating tree we get that $|\bigcup_{X \in V(X_e)} X|$ is odd where X_e is one of the two components of C_e after removing e . Now observe that since N' is odd and adjacent to precisely one module of X_e the graph $G_M[V(X_e) \cup N']$ is a colour propagating tree. Hence we can set $Q_2 := N' \cup \bigcup_{X \in V(X_e)} X$ and $Q_1 := V(G) \setminus Q_2$ which satisfies all requirements.

On the other hand, consider the case that C_e is not a colour propagating tree. First assume that C_e is of even order. Since C_e is not a colour propagating tree we can use Claim 5 on C_e with module partition $\mathcal{M}|_{V_e}$ where $V_e := \bigcup_{N \in V(C_e)} N$. We obtain a partition $\mathcal{N}' = (N'_1, \dots, N'_{\ell'})$ of V_e with $\ell' \leq |V(C_e)| + 1$ and a coarsening $\mathcal{Q}' = (Q'_1, Q'_2)$ of \mathcal{N}' such that $M_i \cap Q'_1 \neq \emptyset$ as in Claim 5 where M_i is one of the modules incident to e . We obtain $\widehat{\mathcal{N}}$ by removing all parts in $V(C_e)$ from \mathcal{M} and adding the parts from \mathcal{N}' . We further set $Q_2 := Q'_2$ and $Q_1 := V(G) \setminus Q_2$. Note that since $M_i \cap Q'_1 \neq \emptyset$, $G_M \setminus C_e$, $G[Q'_1]$ are connected and either e_1 or e_2 is incident to both M_i and some vertex in $G_M \setminus C_e$ we get that $G[Q_1]$ is connected. All other properties follow from Claim 5.

On the other hand, if C_e is of odd order, then either X_e or Y_e must be of even order where X_e, Y_e are the two connected components of C_e after removing e . Without loss of generality let X_e be of even size. Note that removing e and e_1 from G_M splits G_M into precisely two component of which one is X_e . In the case that X_e is a colour propagating tree or star we can set $\widehat{\mathcal{N}} := \mathcal{M}$, $Q_2 := \bigcup_{M \in V(X_e)} M$ and $Q_1 := V(G) \setminus Q_2$. On the other hand, if X_e is not a colour propagating tree, we can use the same argument as above only considering X_e in place of C_e . □

Note that since G_M is not a star or colour propagating tree the premise of either Claim 5 or Claim 6 must be satisfied. We obtain a partition $\widehat{\mathcal{N}} = (\widehat{\mathcal{N}}_1, \dots, \widehat{\mathcal{N}}_{\hat{\ell}})$ of $V(G)$ with $\hat{\ell} \leq k + 1$ and a coarsening $\mathcal{Q} = (Q_1, Q_2)$ of $\widehat{\mathcal{N}}$ as in the two claims. Since $\widehat{\mathcal{N}}|_{Q_2}$ must contain at least two modules we get that $\widehat{\mathcal{N}}|_{Q_1}$ has strictly less modules than \mathcal{M} . Let $k' < k$ be the number of modules of $\widehat{\mathcal{N}}|_{Q_1}$. Hence we can recursively obtain a partition \mathcal{M}' of $G[Q_1]$ with at most $2k'$ parts and a coarsening \mathcal{P}' of \mathcal{M}' with the following properties. $|P|$ is even, $\mathcal{M}'|_P$ is a module partition of $G[P]$ and $G[P]|_{\mathcal{M}'|_P}$ is either a star or a colour propagating tree for every part P of \mathcal{P}' . We obtain the partition $\widehat{\mathcal{M}}$ of $V(G)$ by adding all parts of $\widehat{\mathcal{N}}|_{Q_2}$ to \mathcal{M}' and the coarsening \mathcal{P} of $\widehat{\mathcal{M}}$ by adding P_2 to \mathcal{P}' . Note that the number of

parts of $\widehat{\mathcal{M}}$ is at most $2k' + (\widehat{l} - k') \leq 2k$. Hence $\widehat{\mathcal{M}}$ and \mathcal{P} satisfy the conditions of the statement. \square

Proof of Theorem 5. Without loss of generality assume that G is connected. Furthermore, let $k := \text{mw}(G)$ and $\mathcal{M} = (M_1, \dots, M_k)$ be a module partition of G . Let $\widehat{\mathcal{M}}$ be a partition of $V(G)$ with at most $2k$ parts and \mathcal{P} be a coarsening of $\widehat{\mathcal{M}}$ as in Lemma 3. First observe that $\widehat{\mathcal{M}}|_P$ must contain at least two parts for every part P of \mathcal{P} as $\widehat{\mathcal{M}}|_P$ is a module partition of $G[P]$. Since $\widehat{\mathcal{M}}$ has at most $2k$ parts and \mathcal{P} is a coarsening of $\widehat{\mathcal{P}}$ this implies that \mathcal{P} has at most k parts. Since $G[P]_{\widehat{\mathcal{M}}|_P}$ is either a star or a colour propagating tree we get that $\chi_{\text{odd}}(G[P]) \leq 3$ for every part P of \mathcal{P} by Lemma 1 and Lemma 2. Using a partition (W_1^P, W_2^P, W_3^P) of $G[P]$ such that $G[W_i^P]$ is odd for every $i \in [3]$ for every part P we obtain a global partition of G into at most $3k$ parts such that each part induces an odd subgraph. \square

Since deciding whether a graph is k -odd colourable can be solved in time $2^{\mathcal{O}(\text{krw}(G))}$ [[2], Theorem 6] and $\text{rw}(G) \leq \text{cw}(G) \leq \text{mw}(G)$, where $\text{cw}(G)$ denotes the clique-width of G and $\text{rw}(G)$ rank-width, we obtain the following as a corollary.

Corollary 4. *Given a graph G and a module partition of G of width m the problem of deciding whether G can be odd coloured with at most k colours can be solved in time $2^{\mathcal{O}(m^2)}$.*

5 | INTERVAL GRAPHS

In this section, we study the odd chromatic number of interval graphs and provide an upper bound in the general case as well as a tight upper bound in the case of proper interval graphs. We use the following lemma in both proofs.

Lemma 4. *Let G be a connected interval graph and $P = (p_1, \dots, p_k)$ a maximal induced path in G with the following property.*

(*) $\ell_{p_i} = \min\{\ell_v : v \in V(G)\}$ and for every $i \in [k - 1]$ we have that $r_{p_{i+1}} \geq r_v$ for every $v \in N_G(p_i)$.

Then every $v \in V(G)$ is adjacent to at least one vertex on P .

Proof. Towards a contradiction, assume that there is $v \in V(G)$ such that v is not adjacent to any vertex of P . Note that $v \notin \{p_1, \dots, p_k\}$. Furthermore, by the assumption that v is not adjacent to any vertex of P either $\ell_{p_i} \leq r_{p_i} < \ell_v$ or $\ell_v < \ell_{p_i} \leq r_{p_i}$ for every $i \in [k]$. Pick $i \in [k]$ to be the maximum index such that $r_{p_i} < \ell_v$. Observe that i is well defined as by property (*) $\ell_{p_i} = \min\{\ell_v : v \in V(G)\} \leq r_{p_i} < \ell_v$. First consider the case that $i < k$. But then $r_{p_i} < \ell_v \leq r_v < \ell_{p_{i+1}}$ which contradicts that p_i and p_{i+1} are adjacent. Hence $i = k$. Since G is connected there must be a path $Q = (q_1, \dots, q_\ell)$ from p_k to v . Let $j \in [\ell]$ be the last index such that $\ell_{q_j} \leq r_{p_k}$. Since $q_1 = p_k$ we know that q_j exists and is adjacent to

some vertex in P . Indeed $j < \ell$ as $r_{p_k} < \ell_v$ and $q_\ell = v$. Therefore q_{j+1} exists and further $q_j q_{j+1} \in E(G)$ and $\ell_{q_{j+1}} > r_{p_k}$ (by choice of j). We conclude that $r_{p_k} < r_{q_j}$. If q_j is adjacent to p_{k-1} , this contradicts the property (*). On the other hand, if q_j is not adjacent to p_{k-1} , then the set $\{v \in V(G) : p_{k-1}v \notin E(G), p_kv \in E(G)\}$ is not empty which contradicts the maximality of P . Hence v has to be adjacent to at least one vertex of P . \square

To prove that the odd chromatic number of proper interval graphs is bounded by three we essentially partition the graph into maximal even sized cliques greedily in a left to right fashion.

Proposition 2. *For every proper interval graph G with all components of even order, $\chi_{\text{odd}}(G) \leq 3$.*

Proof. We assume that G is connected. Fix an interval representation of G and denote the interval representing vertex $v \in V(G)$ by $I_v = [\ell_v, r_v]$ where $\ell_v, r_v \in \mathbb{R}$. Let $P = (p_1, \dots, p_k)$ be a maximal induced path in G as in Lemma 4. For every vertex $v \in V(G) \setminus \{p_1, \dots, p_k\}$ let $i_v \in [k]$ be the index such that p_{i_v} is the first neighbour of v on P . Note that this is well defined by Lemma 4. For $i \in [k]$ we let Y_i be the set with the following properties.

- (II1) $_i$ $\{v \in V(G) : i_v = i\} \subseteq Y_i \subseteq \{v \in V(G) : i_v = i\} \cup \{p_i, p_{i+1}\}$.
- (II2) $_i$ $p_i \in Y_i$ if and only if $|\{p_1, \dots, p_{i-1}\} \cup \bigcup_{j \in [i-1]} \{v \in V(G) : i_v = j\}|$ is even.
- (II3) $_i$ $p_{i+1} \in Y_i$ if and only if $|\{p_1, \dots, p_i\} \cup \bigcup_{j \in [i]} \{v \in V(G) : i_v = j\}|$ is odd.

First observe that (Y_1, \dots, Y_k) is a partition of $V(G)$ as (II2) $_i$ and (II3) $_i$ imply that every p_i is in exactly one set Y_i . Furthermore, $|Y_i|$ is even for every $i \in [k]$ since (II1) $_i$ and (II3) $_i$ imply that $|Y_i \cup \{p_1, \dots, p_i\} \cup \bigcup_{j \in [i-1]} \{v \in V(G) : i_v = j\}|$ is even and (II2) $_i$ implies that $|\{p_1, \dots, p_i\} \cup \bigcup_{j \in [i-1]} \{v \in V(G) : i_v = j\} \setminus Y_i|$ is even. Since $v \in V(G) \setminus \{p_1, \dots, p_k\}$ is not adjacent to p_{i_v-1} we get that $\ell_v \in I_{p_{i_v}}$. Since G is a proper interval graph this implies that $r_{p_{i_v}} \leq r_v$ and hence v is adjacent to p_{i_v+1} . Hence (II1) $_i$ implies that $G[Y_i]$ must be a clique since $Y_i \cap \{p_1, \dots, p_k\} \subseteq \{p_i, p_{i+1}\}$ for every $i \in [k]$. Furthermore, $N_G(Y_i)$ and Y_{i+3} are disjoint since $r_v \leq r_{p_{i+1}}$ for every $v \in Y_i$ by property (*) and $r_{p_{i+1}} < \ell_{p_{i+3}} \leq r_w$ for every $w \in Y_{i+3}$ since P is induced. Hence we can define an odd-colouring (V_1, V_2, V_3) of G in the following way. We let $V_j := \bigcup_{i \equiv j \pmod{3}} Y_i$ for $j \in [3]$. Note that since $N_G(Y_i) \cap Y_{i+3} = \emptyset$ we get that $d_{G[Y_i]}(v) = d_{G[V_j]}(v)$ for $i \equiv j \pmod{3}$ which is odd (as Y_i is a clique of even size). Hence $G[V_j]$ is odd for every $j \in [3]$. \square

Remark 4. The upper bound presented in Proposition 2 is tight. Consider the graph G consisting of a K_4 with two added pendant vertices u, w adjacent to different vertices of K_4 . Clearly, G is a proper interval graph and further $\chi_{\text{odd}}(G) = 3$.

We use a similar setup (i.e., a path P covering all vertices of the graph G) as in the proof of 2 to show our general upper bound for interval graphs. The major difference is that we are not guaranteed that sets of the form $\{p_i\} \cup \{v \in V(G) : i_v = i\}$ are cliques. To nevertheless find an odd colouring with few colours of such sets we use an odd/even colouring as in 1 of $\{v \in V(G) : i_v = i\}$ and the universality of p_i . Hence this introduces a factor of two on the number of colours. Furthermore, this approach prohibits us from moving the p_i around as in

the proof of Proposition 2. As a consequence, we get that the intervals of vertices contained in a set Y_i span a larger area of the real line than they do in the proof of Proposition 2. This makes the analysis more technical (Figure 3).

Theorem 6. For every interval graph G with all components of even order, $\chi_{\text{odd}}(G) \leq 6$.

Proof. We assume that G is connected. First, we fix an interval representation of G . We denote the interval representing vertex $v \in V(G)$ by $I_v = [\ell_v, r_v]$ where $\ell_v, r_v \in \mathbb{R}$. Let $P = (p_1, \dots, p_k)$ be a maximal induced path in G as in Lemma 4. Let $Y = V(G) \setminus \{p_1, \dots, p_k\}$. For every $v \in Y$ we define $i_v \in [k]$ to be the minimum index such that v is adjacent to p_{i_v} . Note that this is well defined by Lemma 4.

We now recursively define a partition (Y_1, \dots, Y_k) of Y such that for every $i \in [k]$ the following properties hold.

- (P1) _{i} Every vertex in Y_i is adjacent to p_i .
- (P2) _{i} If $|\{p_1, \dots, p_i\} \cup \{v \in Y : i_v \leq i\}|$ is even, then $\bigcup_{j=1}^i Y_j = \{v \in Y : i_v \leq i\}$.
- (P3) _{i} If $|\{p_1, \dots, p_i\} \cup \{v \in Y : i_v \leq i\}|$ is odd, then either $|\{v \in Y : i_v \leq i\} \setminus \bigcup_{j=1}^i Y_j| = 1$ or $\bigcup_{j=1}^i Y_j = \{v \in Y : i_v \leq i\}$ and $N_G(p_{i+1}) \cap Y_i = \emptyset$.
- (P4) _{i} If $i_w \leq i$ for $w \notin \bigcup_{j=1}^i Y_j$, then $w \in N_G(p_{i+1})$ and $i_w = \max\{i_v \leq i : v \in Y \cap N_G(p_{i+1})\}$.

Fix $i \in [k]$ and assume that we have defined Y_1, \dots, Y_{i-1} satisfying (P1) _{j} , (P2) _{j} , (P3) _{j} , (P4) _{j} for every $j \in [i-1]$. In the following, we show how to construct Y_i . Define $Y'_i := \{v \in Y \setminus \bigcup_{j=1}^{i-1} Y_j : i_v \leq i\}$. Note that $Y'_i \cup \bigcup_{j=1}^{i-1} Y_j = \{v \in Y : i_v \leq i\}$. In the case that either $|\{p_1, \dots, p_i\} \cup \{v \in Y : i_v \leq i\}|$ is even or $Y'_i \cap N_G(p_{i+1}) = \emptyset$ we set $Y_i := Y'_i$. Otherwise, pick $w \in Y'_i \cap N_G(p_{i+1})$ such that $i_w = \max\{i_v \leq i : v \in Y'_i \cap N_G(p_{i+1})\}$ and define




FIGURE 3 Schematic representation of the partition $(\hat{V}_1^{i-1}, \dots, \hat{V}_6^{i-1})$ in the case that $i_w = i - 2$ in the proof of Claim 9. Note that in the figure Y_i is coloured in the two colours of the indices $\hat{j}_1 = 3$ and $\hat{j}_2 = 2$ we obtain in this case. [Color figure can be viewed at wileyonlinelibrary.com]

$Y'_i := Y'_i \setminus \{w\}$. Note that w is well defined since we are considering the case that $Y'_i \cap N_G(p_{i+1}) \neq \emptyset$. Observe that properties $(P2)_i$ and $(P3)_i$ are true by construction of Y_i . To argue that property $(P1)_i$ is true we observe that by $(P4)_{i-1}$ every vertex $w \in Y'_i$ with $i_w < i$ has to be adjacent to p_i . Since in addition every vertex v with $i_v = i$ is adjacent to p_i by choice of i_v , property $(P1)_i$ holds. To argue that property $(P4)_i$ holds we observe that every vertex $v \in Y$ with $i_v = i$ is contained in Y'_i . Hence, if $\max\{i_v \leq i : v \in Y \cap N_G(p_{i+1})\} = i$, then we would choose w with $i_w = i$. In the case that $\max\{i_v \leq i : v \in Y \cap N_G(p_{i+1})\} < i$ then $(P4)_i$ follows directly from $(P4)_{i-1}$. This concludes the construction of the sets Y_1, \dots, Y_k . The following two claims allows us to reuse the colours used to colour Y_i for sets Y_{i+c}, Y_{i+2c}, \dots for some small constant c .

Claim 7. For every vertex $v \in Y$ it holds that $I_v \cap I_{p_i} = \emptyset$ for every $i \notin \{i_v, i_v + 1, i_v + 2\}$. In particular, $N_G(v) \cap \{p_1, \dots, p_k\}$ is contained in $\{p_{i_v}, p_{i_v+1}, p_{i_v+2}\}$ for every vertex $v \in Y$.

Proof of Claim. First observe that $I_v \cap I_{p_i} = \emptyset$ for every $i < i_v$ by definition of i_v . Since P is an induced path $r_{p_{i_v+1}} < \ell_{p_{i_v+3}}$. On the other hand, $r_v \leq r_{p_{i_v+1}}$ by property (*). Hence $r_v < \ell_{p_{i_v+3}} \leq \ell_{p_i}$ for every $i \geq i_v + 3$. Hence $I_v \cap I_{p_i} = \emptyset$ for every $i \geq i_v + 3$ concluding the proof of the statement. □

As a consequence of Claim 7 we get the following claim.

Claim 8. If $Y_i \subseteq \{v \in Y : i_v \geq i'\}$, then

- $N_G(\{p_i\})$ is disjoint from $\{p_j\} \cup Y_j$ for any $j \leq i - 3$ and
- $N_G(Y_i)$ is disjoint from $\{p_j\} \cup Y_j$ for any $j \leq i' - 2$.

Proof of Claim. From Claim 7 we get that no $v \in Y$ with $i_v \leq i - 3$ can be adjacent to $\{p_i\}$. Furthermore, P is an induced path so p_j is nonadjacent to p_i for every $j \leq i - 2$. Therefore, $N_G(\{p_i\})$ is disjoint from $\{p_j\} \cup Y_j$ for every $j \leq i - 3$.

To prove the second property, observe that the property (*) and P being an induced path imply that for every $v \in \{p_1, \dots, p_{i'-2}\} \cup \bigcup_{j=1}^{i'-2} Y_j$ we have that $r_v \leq r_{p_{i'-1}}$. Since every $w \in Y_i$ satisfies that $i_w \geq i'$ we get that every $v \in \{p_1, \dots, p_{i'-2}\} \cup \bigcup_{j=1}^{i'-2} Y_j$ cannot be adjacent to any vertex in Y_i (note that this is not true for p_i in case $i = i'$ as $\ell_{p_i} \leq r_{p_{i-1}}$). Since by construction of Y_1, \dots, Y_k for every $j \leq i' - 2$ we have $Y_j \subseteq \{v \in Y : i_v \leq i' - 2\}$ we get that $N_G(Y_i)$ is disjoint from Y_j for any $j \leq i' - 2$. □

Using the sets Y_1, \dots, Y_k , Claim 7 and Claim 8 we can now find an odd colouring of G . To colour G we use a recursive argument. In the i -th step we find a partition of the set $\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j$ into six (possibly empty) parts V_1^i, \dots, V_6^i with the following properties.

- (C1) _{i} If $|\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j|$ is even, then (V_1^i, \dots, V_6^i) is an odd colouring of $G[\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j]$.

- (C2)_i If $|\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j|$ is odd, then there is $j_i \in [6]$ such that $G[V_{j_i}^i]$ is odd for every $j \neq j_i, p_i \in V_{j_i}^i$ and in $G[V_{j_i}^i]$ every vertex apart from p_i has odd degree.
- (C3)_i $\{p_i\} \cup Y_i$ is contained in the union of at most two parts of the partition (V_1^i, \dots, V_6^i) .
- (C4)_i For every $j \in [6]$, any pair of vertices $v, w \in V_j^i$ can be separated in $G[V_j^i]$ by removing an edge of the path P if there are two indices $i' \neq i''$ such that $v \in \{p_{i'}\} \cup Y_{i'}$ and $w \in \{p_{i''}\} \cup Y_{i''}$.

Let us fix $i \in [k]$ and assume we have partitioned $\{p_1, \dots, p_{i-1}\} \cup \bigcup_{j=1}^{i-1} Y_j$ into six parts $V_1^{i-1}, \dots, V_6^{i-1}$ with properties (C1)_j, (C2)_j, (C3)_j and (C4)_j for every $j \leq i-1$. Our goal is to find a partition (W_1, W_2) of $\{p_i\} \cup Y_i$ and two indices $j_1 \neq j_2 \in [6]$ such that the partition obtained from $(V_1^{i-1}, \dots, V_6^{i-1})$ by adding W_1 to $V_{j_1}^{i-1}$ and W_2 to $V_{j_2}^{i-1}$ is a partition of $\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j$ with properties (C1)_i, (C2)_i, (C3)_i and (C4)_i.

To define the partition (W_1, W_2) we use a partition (W'_1, W'_2) of $G[Y_i]$ such that $G[W'_1]$ is odd and $G[W'_2]$ is even which exists due to Theorem 1. Note that in the case that Y_i is empty we simply obtain the partition (\emptyset, \emptyset) which is sufficient for our purpose. We define $W_1 := W'_1$ and $W_2 := W'_2 \cup \{p_i\}$. Observe that $G[W'_1]$ being odd implies that $|W'_1|$ is even by the handshake lemma. Hence $|W'_2|$ is odd if and only if $|Y_i|$ is odd. Since every vertex in W'_2 is adjacent to p_i by (P1)_i, we obtain that every vertex in W_2 has odd degree in $G[W_2]$ and p_i has odd degree in $G[W_2]$ if and only if $|Y_i|$ is odd. Note that we can get a colouring with 12 colours at this point without much further analysis. Obtaining a colouring with six colours requires careful analysis.

The following claim will provide us with possible choices for indices j_1 and j_2 . Note that the indices from the claim will not in every case be a suitable choice.

Claim 9. There is a partition $(\widehat{V}_1^{i-1}, \dots, \widehat{V}_6^{i-1})$ of $\{p_1, \dots, p_{i-1}\} \cup \bigcup_{j=1}^{i-1} Y_j$ with properties (C1)_j, (C2)_j, (C3)_j, (C4)_j for every $j \leq i-1$ and indices $\widehat{j}_1 \neq \widehat{j}_2 \in [6]$ such that $N_G(W_1)$ is disjoint from $\widehat{V}_{\widehat{j}_1}^{i-1}$ and $N_G(W_2)$ is disjoint from $\widehat{V}_{\widehat{j}_2}^{i-1}$.

Proof of Claim. First consider the case that $Y_i \subseteq \{v \in Y : i_v \in \{i, i-1\}\}$. In this case, we will set $\widehat{V}_j^{i-1} := V_j^{i-1}$ for every $j \in [6]$. Since $Y_i \subseteq \{v \in Y : i_v \in \{i, i-1\}\}$ we obtain using Claim 8 that both $N_G(W_1)$ and $N_G(W_2)$ are disjoint from any set $\{p_j\} \cup Y_j$ with $j \leq i-3$. Since (C3)_j, $j < i$ implies that $\{p_{i-2}, p_{i-1}\} \cup Y_{i-2} \cup Y_{i-1}$ is contained in at most four parts of the partition $(\widehat{V}_1^{i-1}, \dots, \widehat{V}_6^{i-1})$ we can find $\widehat{j}_1, \widehat{j}_2 \in [6], \widehat{j}_1 \neq \widehat{j}_2$ with the following properties. $\widehat{V}_{\widehat{j}_1}^{i-1}$ and $\widehat{V}_{\widehat{j}_2}^{i-1}$ do not contain any element from $\{p_{i-2}, p_{i-1}\} \cup Y_{i-2} \cup Y_{i-1}$. This choice guarantees that $N_G(W_1)$ is disjoint from $\widehat{V}_{\widehat{j}_1}^{i-1}$ and $N_G(W_2)$ is disjoint from $\widehat{V}_{\widehat{j}_2}^{i-1}$.

Now consider the case that Y_i contains a vertex w with $i_w < i-1$. Observe that in this case we get that $i_w = i-2$ as a consequence of Claim 7. Figure 3 illustrates the layout of intervals and the available colours we obtain in this case.

We know that at least one set out of W_1, W_2 is fully contained in $\{v \in Y : i_v = i\} \cup \{p_i\}$ (i.e., the one not containing w) by (P3)_i. Without loss of generality assume that this is true for W_1 (i.e., we do not use in the following argument that W_2 contains p_i). By Claim 8 we infer that $N_G(W_2) \subseteq N_G(\{p_i\} \cup Y_i)$ is disjoint from any set $\{p_j\} \cup Y_j$ with $j \leq i-4$. Furthermore, (C3)_j, $j < i$ implies that $\{p_{i-3}, p_{i-2}\} \cup Y_{i-3} \cup Y_{i-2}$ is contained in at most

four parts of the partition $(V_1^{i-1}, \dots, V_6^{i-1})$. Hence, $\{p_{i-3}, p_{i-2}, p_{i-1}\} \cup Y_{i-3} \cup Y_{i-2}$ is contained in at most five parts of the partition $(V_1^{i-1}, \dots, V_6^{i-1})$. Pick \hat{j}_2 in such a way that $V_{\hat{j}_2}^{i-1}$ is disjoint from $\{p_{i-3}, p_{i-2}, p_{i-1}\} \cup Y_{i-3} \cup Y_{i-2}$. In the following, we will argue that we can, after potentially modifying the partition $(V_1^{i-1}, \dots, V_6^{i-1})$, assume that $\{p_{i-1}\} \cup Y_{i-1}$ is also disjoint from $V_{\hat{j}_2}^{i-1}$.

First observe that $w \in Y_i$ with $i_w = i - 2$ implies that $Y_{i-1} \subseteq \{v \in Y : i_v = i - 1\}$ by $(P2)_{i-2}$ and $(P3)_{i-2}$. By Claim 8 we get that $N_G(Y_{i-1})$ is disjoint from $\{p_j\} \cup Y_j$ for any $j \leq i - 3$. Since $\{p_{i-2}\} \cup Y_{i-2}$ is contained in at most two parts of the partition $(V_1^{i-1}, \dots, V_6^{i-1})$ by $(C3)_j, j < i$ we can pick $j', j'' \in [6], j' \neq j''$ distinct from \hat{j}_2 such that $V_{j'}^{i-1}, V_{j''}^{i-1}$ are disjoint from $\{p_{i-2}\} \cup Y_{i-2}$. By $(C3)_{i-1}$ we know that $\{p_{i-1}\} \cup Y_{i-1}$ is contained in at most two parts of the partition $(V_1^{i-1}, \dots, V_6^{i-1})$. Assume $V_{m'}^{i-1}, V_{m''}^{i-1}$ are those two parts and $p_{i-1} \in V_{m'}^{i-1}$. In case $\{p_{i-1}\} \cup Y_{i-1}$ is contained in one part we let $m' = m''$. Define $M' := V_{m'}^{i-1} \cap \cup Y_{i-1}$. Observe that $m'' \neq \hat{j}_2$ as $V_{\hat{j}_2}^{i-1}$ does not contain p_{i-1} by choice of \hat{j}_2 . If $m' \neq \hat{j}_2$ then $V_{\hat{j}_2}^{i-1}$ is disjoint from $\{p_{i-3}, p_{i-2}, p_{i-1}\} \cup \cup_{j=i-3}^{i-1} Y_j$ and hence with setting $\widehat{V}_j^{i-1} := V_j^{i-1}$ for every $j \in [6]$ we get that $N_G(W_2)$ is disjoint from $\widehat{V}_{\hat{j}_2}^{i-1}$. Hence assume that $m' = \hat{j}_2$. Since $j' \neq j''$ we get that m'' is not equal to either j' or j'' . Assume $j' \neq m''$. We define $\widehat{V}_{j'}^{i-1} := V_{j'}^{i-1} \cup M', \widehat{V}_{\hat{j}_2}^{i-1} := V_{\hat{j}_2}^{i-1} \setminus M'$ and $\widehat{V}_j^{i-1} := V_j^{i-1}$ for $j \notin \{j', \hat{j}_2\}$. To see that the partition $(\widehat{V}_1^{i-1}, \dots, \widehat{V}_6^{i-1})$ of $\{p_1, \dots, p_{i-1}\} \cup \cup_{j=1}^{i-1} Y_j$ satisfies $(C1)_{i-1}, (C2)_{i-1}, (C3)_{i-1}, (C4)_{i-1}$ we make the following two observations. Since $V_{\hat{j}_2}^{i-1}$ is disjoint from $\{p_{i-3}, p_{i-2}, p_{i-1}\} \cup Y_{i-3} \cup Y_{i-2}$ and $N_G(M') \subseteq N_G(Y_{i-1})$ is disjoint from $\{p_j\} \cup Y_j$ for any $j \leq i - 3$ we get that M' is a component of $G[V_{\hat{j}_2}^{i-1}]$. Hence $d_{G[V_{\hat{j}_2}^{i-1}]}(v) = d_{G[\widehat{V}_{\hat{j}_2}^{i-1}]}(v)$ for any vertex $v \in V_{\hat{j}_2}^{i-1} \setminus M'$. Additionally, since $V_{j'}^{i-1}$ is disjoint from $\{p_{i-2}\} \cup Y_{i-2}$ and $N_G(M') \subseteq N_G(Y_{i-1})$ is disjoint from $\{p_j\} \cup Y_j$ for any $j \leq i - 3$ we get that M' is a connected component of $G[\widehat{V}_{j'}^{i-1}]$. Hence $d_{G[V_{j'}^{i-1}]}(v) = d_{G[\widehat{V}_{j'}^{i-1}]}(v)$ for any vertex $v \in V_{j'}^{i-1}$ and $d_{G[V_{\hat{j}_2}^{i-1}]}(v) = d_{G[\widehat{V}_{j'}^{i-1}]}(v)$ for any vertex $v \in M'$. This argument shows that $(C1)_{i-1}$ and $(C2)_{i-1}$ are satisfied. $(C4)_{i-1}$ follows from the observation that M' is a connected component of $G[\widehat{V}_{j'}^{i-1}]$ and $(V_1^{i-1}, \dots, V_6^{i-1})$ satisfying $(C4)_{i-1}$. Furthermore, $(C3)_{i-1}$ is trivially satisfied.

To choose \hat{j}_1 we first observe that $W_1 \subseteq \{p_j\} \cup \{v \in Y : i_v = i\}$ implies that $N_G(W_1)$ is disjoint from $\{p_j\} \cup Y_j$ for every $j \leq i - 3$ using Claim 8. Since $\{p_{i-2}, p_{i-1}\} \cup Y_{i-2} \cup Y_{i-1}$ is contained in at most four parts of the partition $(\widehat{V}_1^{i-1}, \dots, \widehat{V}_6^{i-1})$ by $(C3)_j, j < i$ we can choose $\hat{j}_1 \in [6]$ such that $\hat{j}_1 \neq \hat{j}_2$ and $N_G(W_1)$ is disjoint from $\widehat{V}_{\hat{j}_1}^{i-1}$ as required. \square

For the remainder of the argument we pick $\hat{j}_1, \hat{j}_2 \in [6]$ and $(\widehat{V}_1^{i-1}, \dots, \widehat{V}_6^{i-1})$ as in the statement of Claim 9. We define partition (V_1^i, \dots, V_6^i) in the following considering several different cases.

First consider the case that $|\{p_1, \dots, p_{i-1}\} \cup \cup_{j=1}^{i-1} Y_j|$ is even. We set $j_1 := \hat{j}_1$ and $j_2 := \hat{j}_2$ and define $V_{j_1}^i := \widehat{V}_{j_1}^{i-1} \cup W_1, V_{j_2}^i := \widehat{V}_{j_2}^{i-1} \cup W_2$ and $V_j^i := \widehat{V}_j^{i-1}$ for every $j \notin \{j_1, j_2\}$. Since $|\{p_1, \dots, p_{i-1}\} \cup \cup_{j=1}^{i-1} Y_j|$ is even we get that $G[\widehat{V}_j^{i-1}]$ is odd for every $j \in [6]$ by $(C1)_{i-1}$.

Since additionally $G[W_1]$ is odd and $N_G(W_1)$ is disjoint from \widehat{V}_j^{i-1} we get that $G[V_j^i]$ is odd. Furthermore, recall that every vertex $v \in W_2 \setminus \{p_i\}$ has odd degree in $G[W_2]$ and p_i has odd degree in $G[W_2]$ if and only if $|Y_i|$ is even. Since $|\{p_1, \dots, p_{i-1}\} \cup \bigcup_{j=1}^{i-1} Y_j|$ is even, $|\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j|$ is even if and only if $|Y_i|$ is even (and p_i has odd degree). Hence (V_1^i, \dots, V_6^i) satisfies $(C1)_i$ and $(C2)_i$. Additionally, $(C3)_i$ is true because $\{p_i\} \cup Y_i$ are contained in $V_{j_1}^i \cup V_{j_2}^i$. Lastly, $(C4)_i$ follows from $(C4)_{i-1}$ and the fact that $N_G(W_1)$ is disjoint from $\widehat{V}_{j_1}^{i-1}$ and $N_G(W_2)$ is disjoint from $\widehat{V}_{j_2}^{i-1}$.

Now consider the case that $|\{p_1, \dots, p_{i-1}\} \cup \bigcup_{j=1}^{i-1} Y_j|$ is odd. Assume $j_2 \in [6]$ is the index such that $p_{i-1} \in \widehat{V}_{j_2}^{i-1}$ and further set $j_1 := \widehat{j_1}$. We define $V_{j_1}^i := \widehat{V}_{j_1}^{i-1} \cup W_1$, $V_{j_2}^i := \widehat{V}_{j_2}^{i-1} \cup W_2$ and $V_j^i := \widehat{V}_j^{i-1}$ for every $j \notin \{j_1, j_2\}$. By $(C1)_{i-1}$ and $(C2)_{i-1}$ we directly conclude that $G[V_j^i]$ is odd for $j \notin \{j_1, j_2\}$.

Note that by Claim 9, we have that $N_G(W_1)$ is disjoint from $\widehat{V}_{j_1}^{i-1}$. As W_1 is therefore a connected component in $G[V_{j_1}^i]$ and both $G[W_1]$ and $G[\widehat{V}_{j_1}^{i-1}]$ are odd we get that $G[V_{j_1}^i]$ is odd. Furthermore, property $(C2)_{i-1}$ implies that $d_{G[\widehat{V}_{j_2}^{i-1}]}(p_{i-1})$ is even and $d_{G[\widehat{V}_{j_2}^{i-1}]}(v)$ is odd for every $v \in \widehat{V}_{j_2}^{i-1} \setminus \{p_{i-1}\}$. Additionally, $d_{G[W_2]}(v)$ is odd for every $v \in Y_i \setminus \{p_i\}$. To determine the degree of p_{i-1} in $G[V_{j_2}^i]$ observe that $(P3)_{i-1}$ ensures that $Y_i \subseteq \{v \in Y : i_v = i\}$ and hence p_{i-1} is nonadjacent to any $v \in Y_i$. Since p_{i-1} is adjacent to p_i we get $d_{G[V_{j_2}^i]}(p_{i-1}) = d_{G[\widehat{V}_{j_2}^{i-1}]}(p_{i-1}) + 1$ and hence the degree of p_{i-1} is odd in $G[V_{j_2}^i]$. To determine the degree of $v \in V_{j_2}^i \setminus \{p_{i-1}\}$, we first argue that $(C4)_i$ holds for the partition (V_1^i, \dots, V_6^i) . Let G' be the graph obtained from $G[\widehat{V}_{j_2}^{i-1}]$ by removing all edges of the path P . Then $(C4)_{i-1}$ implies that every $v \in \widehat{V}_{j_2}^{i-1} \cap (\{p_1, \dots, p_{i-2}\} \cup \bigcup_{j=1}^{i-2} Y_j)$ has to be in a different component of G' then any $w \in \widehat{V}_{j_2}^{i-1} \cap (\{p_{i-1}\} \cup Y_{i-1})$. Hence $r_v < \ell_w$ for any pair of vertices $v \in \widehat{V}_{j_2}^{i-1} \cap (\{p_1, \dots, p_{i-2}\} \cup \bigcup_{j=1}^{i-2} Y_j)$, $w \in \widehat{V}_{j_2}^{i-1} \cap (\{p_{i-1}\} \cup Y_{i-1})$ apart from the pair p_{i-2}, p_{i-1} . Since $Y_{i-1} \cap N_G(p_i) = \emptyset$ and $Y_i \subseteq \{v \in Y : i_v = i\}$ we further know that $r_v < \ell_w$ for every $v \in Y_{i-1}$, $w \in \{p_i\} \cup Y_i$. Combined we get that $r_v < \ell_w$ (and therefore v is nonadjacent to w) for any $v \in \widehat{V}_{j_2}^{i-1} \cap (\{p_1, \dots, p_{i-1}\} \cup \bigcup_{j=1}^{i-1} Y_j)$, $w \in \{p_i\} \cup Y_i$ apart from the pair p_{i-1}, p_i . Hence we argued that $(C4)_i$ holds for the partition (V_1^i, \dots, V_6^i) . Since both p_{i-1} and p_i are contained in $V_{j_2}^i$ the property $(C4)_i$ implies that $d_{G[V_{j_2}^i]}(w) = d_{G[W_2]}(w)$ for every $w \in Y_i$. Further, $(C4)_i$ implies that $d_{G[V_{j_2}^i]}(w) = d_{G[\widehat{V}_{j_2}^{i-1}]}(w)$ for every $w \in \widehat{V}_{j_2}^{i-1} \setminus \{p_{i-1}\}$. Lastly, $d_{G[V_{j_2}^i]}(p_i) = d_{G[W_2]}(p_i) + 1$ since p_i is adjacent to p_{i-1} . Furthermore, $|\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j|$ is even if and only if $|Y_i|$ is odd since $|\{p_1, \dots, p_{i-1}\} \cup \bigcup_{j=1}^{i-1} Y_j|$ is odd. Since $d_{G[W_2]}(p_i)$ is even if and only if $|Y_i|$ is odd we get that $d_{G[V_{j_2}^i]}(p_i)$ is odd if and only if $|\{p_1, \dots, p_i\} \cup \bigcup_{j=1}^i Y_j|$ is even as required. Therefore, $(C1)_i$ and $(C2)_i$ hold for (V_1^i, \dots, V_6^i) . Additionally, $(C3)_i$ is true by construction.

Finally, since $|\{p_1, \dots, p_k\} \cup \bigcup_{j=1}^k Y_j| = |V(G)|$ is even (V_1^k, \dots, V_6^k) is an odd colouring of G by $(C1)_k$. \square

Observe that the proof of Theorem 6 only relies on the fact that interval graphs have a so-called *dominating pair* (a pair of vertices u, v such that every path from u to v dominates all

vertices). Since this property also holds for all connected AT-free graphs [6], the following result can be shown analogously to Theorem 6.

Theorem 7 [11]. *For every AT-free graph G with all components of even order, $\chi_{\text{odd}}(G) \leq 6$.*

ACKNOWLEDGEMENTS

This study was funded by French National Research Agency under JCJC programs (DAGDigDec: ANR-21-CE48-0012 and ASSK: ANR-18-CE40-0025-01) and by the CTU Global postdoc fellowship program.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

ORCID

Rémy Belmonte  <https://orcid.org/0000-0001-8043-5343>

Noleen Köhler  <https://orcid.org/0000-0002-1023-6530>

Nikolaos Melissinos  <http://orcid.org/0000-0002-0864-9803>

REFERENCES

1. A. Aashtab, S. Akbari, M. Ghanbari, and A. Shidani, *Vertex partitioning of graphs into odd induced subgraphs*, *Discuss. Math. Graph Theory* **43** (2023), no. 2, 385–399. <https://doi.org/10.7151/dmgt.2371>
2. R. Belmonte and I. Sau, *On the complexity of finding large odd induced subgraphs and odd colorings*, *Algorithmica* **83** (2021), no. 8, 2351–2373. <https://doi.org/10.1007/s00453-021-00830-x>
3. B. Bollobás, *Chromatic number, girth and maximal degree*, *Discr. Math.* **24** (1978), no. 3, 311–314.
4. Y. Caro, *On induced subgraphs with odd degrees*, *Discret. Math.* **132** (1994), no. 1–3, 23–28. [https://doi.org/10.1016/0012-365X\(92\)00563-7](https://doi.org/10.1016/0012-365X(92)00563-7)
5. G. J. Chang and G. Duh, *On the precise value of the strong chromatic index of a planar graph with a large girth*, *Discret. Appl. Math.* **247** (2018), 389–397. <https://doi.org/10.1016/j.dam.2018.03.075>
6. D. G. Corneil, S. Olariu, and L. Stewart, *Asteroidal triple-free graphs*, *SIAM J. Discret. Math.* **10** (1997), no. 3, 399–430. <https://doi.org/10.1137/S0895480193250125>
7. R. Diestel, *Graph theory*, 4th ed., Springer, 2012.
8. P. Erdős, *Graph theory and probability*, *Can. J. Math.* **11** (1959), 34–38. <https://doi.org/10.4153/CJM-1959-003-9>
9. A. Ferber and M. Krivelevich, *Every graph contains a linearly sized induced subgraph with all degrees odd*, *Adv. Math.* **406** (2022), 108534.
10. L. Lovász, *Combinatorial problems and exercises*, North-Holland, 1993.
11. H. Müller, private communication.
12. M. Petruševski and R. Škrekovski, *Colorings with neighborhood parity condition*, *Discr. Appl. Math.* **321** (2022), 385–391.
13. A. D. Scott, *On induced subgraphs with all degrees odd*, *Graphs Comb.* **17** (2001), no. 3, 539–553. <https://doi.org/10.1007/s003730170028>

How to cite this article: R. Belmonte, A. Harutyunyan, N. Köhler, and N. Melissinos, *Odd chromatic number of graph classes*, *J. Graph Theory* (2024), 1–23. <https://doi.org/10.1002/jgt.23200>