

# Local stability of differential rotation in magnetized radiation zones and the solar tachocline

R. W. Dymott <sup>\*</sup>, A. J. Barker <sup>\*</sup>, C. A. Jones and S. M. Tobias

*Department of Applied Mathematics, School of Mathematics, University of Leeds, Leeds LS2 9JT, UK*

Accepted 2024 October 9. Received 2024 October 9; in original form 2024 September 10

## ABSTRACT

We study local magnetohydrodynamical instabilities of differential rotation in magnetized, stably stratified regions of stars and planets using a Cartesian Boussinesq model. We consider arbitrary latitudes and general shears (with gravity direction misaligned from this by an angle  $\phi$ ), to model radial ( $\phi = 0$ ), latitudinal ( $\phi = \pm 90^\circ$ ), and mixed differential rotations, and study both non-diffusive [including magnetorotational instability (MRI) and Solberg–Høiland instability] and diffusive instabilities [including Goldreich–Schubert–Fricke (GSF) and MRI with diffusion]. These instabilities could drive turbulent transport and mixing in radiative regions, including the solar tachocline and the cores of red giant stars, but their dynamics are incompletely understood. We revisit linear axisymmetric instabilities with and without diffusion and analyse their properties in the presence of magnetic fields, including deriving stability criteria and computing growth rates, wave vectors, and energetics, both analytically and numerically. We present a more comprehensive analysis of axisymmetric local instabilities than prior work, exploring arbitrary differential rotations and diffusive processes. The presence of a magnetic field leads to stability criteria depending upon angular velocity rather than angular momentum gradients. We find MRI operates for much weaker differential rotations than the hydrodynamic GSF instability, and that it typically prefers much larger length-scales, while the GSF instability is impeded by realistic strength magnetic fields. We anticipate MRI to be more important for turbulent transport in the solar tachocline than the GSF instability when  $\phi > 0$  in the Northern (and vice versa in the Southern) hemisphere, though the latter could operate just below the convection zone when MRI is absent for  $\phi < 0$ .

**Key words:** instabilities – MHD – waves – Sun: rotation – stars: interiors – stars: rotation.

## 1 INTRODUCTION

The evolution of angular momentum (AM) and its profile in a star is one of the most fundamental – yet poorly understood – aspects of stellar evolution (e.g. Maeder & Meynet 2000; Maeder 2009). Many unsolved problems in solar physics, including the development and maintenance of the differentially rotating solar convection zone, tachocline, and solar dynamo, as well as those of other stars, all likely require a better understanding of the mechanisms that shape the AM transport and hence profile of star throughout its life cycle. Current stellar evolution codes are too simplistic and do not correctly model the evolution of AM, as is evident from red giant and subgiant stars for example, whose core-envelope differential rotations inferred from asteroseismology are not well explained by existing stellar evolution models (e.g. Aerts, Mathis & Rogers 2019).

Some of the most important proposed mechanisms for AM transport and turbulent mixing in stars are magnetohydrodynamical (MHD) instabilities, which have not yet been fully explored (e.g. Zahn 1974; Maeder & Meynet 2000) and their non-linear evolution is particularly poorly understood. Here we analyse local MHD instabilities in stellar radiative zones, focussing on the stability properties of stably stratified, differentially rotating, and magnetized

shear flows – relevant, for example, to modelling the solar tachocline or the cores of red giant stars. Radiative zones are often thought of as quiescent regions, but they have already been shown to exhibit a range of local and global MHD instabilities that transport AM and lead to turbulent mixing. Examples of relevant instabilities in such regions include the hydrodynamic Goldreich–Schubert–Fricke (GSF) instability, a double-diffusive centrifugal instability of differential rotation enabled by thermal diffusion (Goldreich & Schubert 1967; Fricke 1968; Knobloch & Spruit 1982; Rashid, Jones & Tobias 2008; Barker, Jones & Tobias 2019, 2020; Park, Prat & Mathis 2020; Park et al. 2021; Dymott et al. 2023; Tripathi et al. 2024), and the more violent instabilities excited when the Solberg–Høiland (SH) criteria for non-diffusive instabilities are violated (Solberg 1936; Høiland 1941). Not only do these linear instabilities lead to turbulence with enhanced transport properties in their non-linear evolution, they also exhibit the emergence of long-term anisotropic quasi-stable structures such as ‘zonal jets’ (or layering in the AM; Barker et al. 2020; Dymott et al. 2023). The complex anisotropic nature of the long-term evolution of many of these instabilities suggests that modelling them in one-dimensional stellar evolution models simply as a one-dimensional diffusive process is probably insufficient. Indeed, in certain cases turbulent transport is known to be antidiffusive.

The presence of even a weak magnetic field is known to drastically modify the stability of differentially rotating flows (e.g. Chandrasekhar 1961; Acheson & Gibbons 1978; Balbus & Hawley

<sup>\*</sup> E-mail: [mmrwd@leeds.ac.uk](mailto:mmrwd@leeds.ac.uk) (RWD); [A.J.Barker@leeds.ac.uk](mailto:A.J.Barker@leeds.ac.uk) (AJB)

1991). Stability criteria with magnetic fields tend to involve angular velocity gradients – which typically require much weaker differential rotations to predict instability – rather than the AM gradients without fields; this is because the field can act as a tether between fluid particles and allow them to exchange AM. The magnetorotational instability (MRI) is one such manifestation when a weak magnetic field is introduced into a differentially rotating flow (e.g. Chandrasekhar 1961; Acheson & Gibbons 1978; Balbus & Hawley 1991, 1994; Balbus 1995, 2009; Spruit 1999; Ogilvie 2007; Oishi et al. 2020; Vasil et al. 2024). This can operate and drive turbulence even in many hydrodynamically stable flows. Its operation in stably stratified stellar interiors (i.e. radiation zones) in the presence of diffusive processes has been studied in some prior works (Menou, Balbus & Spruit 2004; Menou & Le Mer 2006; Parfrey & Menou 2007; Guilet & Müller 2015; Caleo & Balbus 2016; Caleo, Balbus & Tognelli 2016), but much remains to be explored of its linear properties, and especially its non-linear evolution in stars. Guilet & Müller (2015) performed linear analysis and numerical simulations of the MRI in a local stably stratified model of a proto-neutron star (with extra neutrino cooling). Our approach is broadly similar to theirs but we will study arbitrary local differential rotations. Global simulations in spherical geometry of the MRI (or Tayler instability, which is a current-driven instability that is also present in these) in stellar radiative zones have also been performed (Gaurat et al. 2015; Jouve, Gastine & Lignières 2015; Meduri, Lignières & Jouve 2019; Jouve, Lignières & Gaurat 2020), though these kinds of studies may not adequately capture all of the possible local instabilities. We choose to adopt a local model here, partly for simplicity and because such models are appropriate for studying small-scale MHD instabilities, and also because they can explore more realistic parameter regimes with numerical simulations (particularly with regards to smaller diffusivities) than global models would allow in non-linear regimes.

Here we introduce magnetic fields to build directly upon Barker et al. (2019, hereafter Paper I), Barker et al. (2020, hereafter Paper II), and Dymott et al. (2023, hereafter Paper III) that studied hydrodynamical instabilities in a local Cartesian representation of a small patch of a stably stratified, differentially rotating stellar or planetary radiation zone. A global ‘shellular’ (radial) differential rotation varying only with spherical radius was considered at the equator in Paper I (and an axisymmetric turbulence closure model was developed and verified for this case by Tripathi et al. 2024), and at a general latitude in Paper II. In Paper III, we generalized the model to consider an arbitrary differential rotation profile, which varies with both radius and latitude. Here we incorporate a poloidal magnetic field into this more general model. Following a similar approach, we perform an axisymmetric linear stability analysis here, which we will follow with complementary three-dimensional non-linear numerical simulations (which can consider the effects of more general field orientations) in future work. Our primary goals are to understand the properties of the GSF instability in the magnetic system, as well as the operation of the MRI, and to determine their potential roles in AM transport, chemical mixing, and dynamo generation. Our linear study is related to the one undertaken by Latter & Papaloizou (2018) for the vertical shear instability in astrophysical discs (e.g. Urpin & Brandenburg 1998; Nelson, Gressel & Umurhan 2013; Barker & Latter 2015), which is the name used for the GSF instability in that context.

The goal of this paper is to gain insights into how the presence of a locally uniform magnetic field affects the linear properties of local instabilities of differential rotation in stellar and planetary radiative zones. We do this by investigating the axisymmetric linear stability of the system (which we define in Section 2), both analytically and

numerically in Sections 3–6. We determine how the properties of the unstable modes depend on magnetic field strength  $B_0$  and magnetic Prandtl number  $\text{Pm} = \nu/\eta$  (the ratio of kinematic viscosity  $\nu$  to ohmic diffusivity  $\eta$ ). We will analyse the energetics of the various instabilities in our model and derive several new results before applying them to the solar tachocline and red giant stars in Section 7.


## 2 LOCAL CARTESIAN MODEL

### 2.1 The model and governing equations

We follow Paper I, Paper II, and Paper III and employ a local Cartesian model to study small-scale instabilities of differential rotation in a stably stratified region of a star or planet. We use coordinates  $(x, y, z)$ , where  $y$  is the local azimuthal coordinate, and  $x$  and  $z$  are two coordinates in the meridional plane. We adopt the Bousinesq approximation (Spiegel & Veronis 1960), which is expected to be valid for the local instabilities we study (see e.g. Paper I, for justification regarding studying GSF modes). The differential rotation is represented by a linear shear flow  $\mathbf{U}_0 = -Sxe_y$ , which in general varies with both spherical radius  $r$  and latitude  $\beta$ , and we have defined  $x$  to be aligned with the axis of variation of  $\mathbf{U}_0$ .  $S$  is the constant value locally of  $-\varpi|\nabla\Omega(r, \beta)|$  (where  $\Omega(r, \beta)$  is the angular velocity), and  $\varpi$  is the distance from the axis of rotation (cylindrical radius). The local effective gravity vector  $\mathbf{e}_g = (\cos\phi, 0, \sin\phi)$  defines the angle  $\phi$ , and the rotation axis lies along the vector  $\hat{\boldsymbol{\Omega}} = (\sin\Lambda, 0, \cos\Lambda)$ , thereby defining the angle  $\Lambda$ . We note that both of these are defined locally with respect to  $x$ . The latitude angle is then given by  $\beta = \Lambda + \phi$ , which measures the angle between the equator  $\hat{\boldsymbol{\Omega}}^\perp = (\cos\Lambda, 0, -\sin\Lambda)$  and the spherical radial direction  $\mathbf{e}_g$ . Our model is illustrated in Fig. 1, which shows the various angles involved.

We build upon the hydrodynamical studies of Paper I, Paper II, and Paper III by introducing a uniform static background poloidal magnetic field  $\mathbf{B}_0 = B_0\hat{\mathbf{B}}$  that is in equilibrium, satisfying the local analogue of Ferraro’s law of isorotation (Ferraro 1937). For this flow to be in equilibrium in the meridional/poloidal  $(x, z)$  plane, it must lie along  $z$  with  $\hat{\mathbf{B}} = (0, 0, 1)$ ,<sup>1</sup> being always perpendicular to variation of the shear flow  $\mathbf{U}_0$  locally. This permits a well-defined equilibrium state even if it may complicate interpretation of our model because the field is not purely radial or horizontal, depending on the value of  $\phi$ . The field is radial if  $\phi = \pm 90^\circ$  and it is latitudinal if  $\phi = 0^\circ, \pm 180^\circ$ . We do not consider toroidal/azimuthal fields in our local analysis, which are typically thought to be dominant in the solar tachocline, because they play no role for linear incompressible axisymmetric perturbations. A toroidal field would affect non-axisymmetric perturbations (e.g. Ogilvie & Pringle 1996) but analysing those (and their non-modal growth) is less straightforward, and it is likely that axisymmetric instabilities are the fastest growing ones in any case (e.g. Latter & Papaloizou 2018). Since we have adopted a local Cartesian model, as appropriate to explore small-scale instabilities in stellar radiative zones, we do not capture the effects of azimuthal magnetic fields on axisymmetric modes via hoop stresses, leading to azimuthal MRI (Hollerbach & Rüdiger 2005; Kirillov & Stefani 2010; Guseva et al. 2017; Mamatsashvili et al. 2019; Meduri, Jouve & Lignières 2024). This could be important on larger length-scales than those that we consider but it would require us to adopt a global model. An initially purely toroidal field can

<sup>1</sup>The MHD equations are invariant under the transformation  $\mathbf{B} \rightarrow -\mathbf{B}$ , so there is no loss of generality in considering  $B_0 \geq 0$ .



**Figure 1.** Illustration of the various vectors and corresponding angles in the  $(x, z)$  plane. The cylindrical radial direction (along the equator) is  $\hat{\Omega}^\perp$ , and the rotation axis is  $\hat{\Omega}$ . The local radial direction is (approximately) along the effective gravity direction  $\mathbf{e}_g$ , which is misaligned with respect to the  $x$ -direction when  $\phi$  is non-zero. The magnetic field in linear theory is always along  $z$  (therefore perpendicular to the shear in  $x$ ), which is the only direction in the meridional plane in which an equilibrium exists.

play a role non-linearly in local models even if the linear instability is axisymmetric however, so future non-linear simulations should explore these fields, also because of their possible role in driving non-axisymmetric instabilities.

The incompressible MHD equations governing perturbations to the shear flow  $\mathbf{U}_0$  and background stable stratification in the Boussinesq approximation, in the frame rotating at the rate  $\mathbf{\Omega}$ , are

$$D\mathbf{u} + 2\mathbf{\Omega} \times \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{U}_0 = -\nabla p + \theta \mathbf{e}_g + \mathbf{B} \cdot \nabla \mathbf{B} + \nu \nabla^2 \mathbf{u}, \quad (1)$$

$$D\theta + \mathcal{N}^2 \mathbf{u} \cdot \mathbf{e}_\theta = \kappa \nabla^2 \theta, \quad (2)$$

$$D\mathbf{B} = \mathbf{B} \cdot \nabla \mathbf{u} + \mathbf{B} \cdot \nabla \mathbf{U}_0 + \eta \nabla^2 \mathbf{B}, \quad (3)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (4)$$

$$\nabla \cdot \mathbf{u} = 0, \quad (5)$$

$$D \equiv \partial_t + \mathbf{u} \cdot \nabla + \mathbf{U}_0 \cdot \nabla. \quad (6)$$

Here  $\mathbf{u}$  is a velocity perturbation and  $\mathbf{B}$  is the total magnetic field. We define a temperature perturbation  $\theta$  having units of acceleration and related to the standard temperature perturbation  $\tilde{T}$  via  $\theta = \alpha g \tilde{T}$ , where  $\alpha$  is the thermal expansion coefficient and  $g$  is the local gravitational acceleration. We use Alfvén speed units for the magnetic field, such that the dimensional magnetic field is  $\mathbf{B}/\sqrt{\mu_0 \rho}$ , where  $\rho$  is the constant reference density that we henceforth set to unity

**Table 1.** Differential rotation profiles and magnetic field orientations (in the meridional plane) as  $\Lambda$  and  $\phi$  are varied. Here  $\beta$  is colatitude,  $z$  is distance along rotation axis,  $r$  is spherical radius, and  $\varpi$  is cylindrical radius.

$\Lambda$	$\phi$	Differential rotation	Magnetic field
0	–	$\Omega(\varpi)$ (cylindrical)	Arbitrary
$\pm 90^\circ$	–	$\Omega(z)$ (axial variation)	Arbitrary
–	0	$\Omega(r)$ (spherical/shellular)	Horizontal/latitudinal
–	$\pm 90^\circ$	$\Omega(\beta)$ (horizontal/latitudinal)	Radial
–	–	$\Omega(r, \beta)$ (arbitrary)	Arbitrary

and  $\mu_0$  is the vacuum permeability. Magnetic pressure is contained within the total pressure  $p$ . We consider constant kinematic viscosity  $\nu$ , thermal diffusivity  $\kappa$ , and ohmic diffusivity  $\eta$ . For reference, the basic state satisfies

$$2\mathbf{\Omega} \times \mathbf{U}_0 = -\nabla p_0 + \alpha g T \mathbf{e}_g, \quad (7)$$

$$0 = \kappa \nabla^2 T, \quad (8)$$

since  $\partial_t \mathbf{U}_0 = \mathbf{U}_0 \cdot \nabla \mathbf{U}_0 = \nabla^2 \mathbf{U}_0 = \mathbf{B}_0 \cdot \nabla \mathbf{B}_0 = \mathbf{0}$  and the equivalents of equations (3)–(5) are trivially satisfied.

A background temperature (entropy) profile  $T(x, z)$  has also been adopted, with uniform gradient (hence satisfying equation 8)  $\alpha g \nabla T = \mathcal{N}^2 \mathbf{e}_\theta$ , where  $\mathbf{e}_\theta = (\cos \Gamma, 0, \sin \Gamma)$ , where the buoyancy frequency  $\mathcal{N}^2 > 0$  in radiative zones. The effective gravity vector  $\mathbf{e}_g$  lies approximately in the spherical radial direction, and is inclined to  $x$  by an angle  $\phi$ . For clarity, we consider sufficiently slowly rotating stars that  $\mathbf{e}_g$  lies approximately along the spherical radial direction, and hence ‘radial’ will be assumed to be along  $\mathbf{e}_g$ , though the model itself does not require this restriction (and it would not be the correct interpretation in very rapidly rotating stars).

We expect the star to adjust rapidly to satisfy thermal wind balance, and enforcing this requirement eliminates the angle  $\Gamma$  as a free parameter. This means that  $\mathbf{U}_0$  and its thermal state satisfy the thermal wind equation (TWE),


$$2\mathbf{\Omega} S \sin \Lambda = \mathcal{N}^2 \sin(\Gamma - \phi), \quad (9)$$

which is derived from the azimuthal component of the vorticity equation for the basic state, i.e. the curl of equation (7). This is unaffected by our magnetic field  $\mathbf{B}_0$ . The representation of various global differential rotation profiles and magnetic field orientations in our local model is summarized in Table 1. We also illustrate the various angles in our problem in Fig. 2. See Paper III for further details of the non-magnetic model.

We use units defined by the rotational time-scale,  $\Omega^{-1}$ , and the length-scale

$$d = \left( \frac{\nu \kappa}{\mathcal{N}^2} \right)^{1/4}. \quad (10)$$

The fastest growing hydrodynamic (GSF) modes typically have wavelengths  $O(d)$ . With this choice of length, the buoyancy time-scale  $\mathcal{N}^{-1}$  is equal to the geometric mean of the viscous ( $d^2/\nu$ ) and thermal ( $d^2/\kappa$ ) diffusion time-scales (see e.g. Radko 2013, for other double-diffusive problems). Note that with the addition of a magnetic field it is not at all clear that unstable modes will necessarily have length-scales  $O(d)$ , and in fact we will show that MRI modes may have much larger scales. However, for comparison with Paper I, Paper II, and Paper III and for comparing GSF and MRI modes, we continue to adopt this choice of units here.



**Figure 2.** Illustration of the key vectors and corresponding angles in the  $(x, z)$  plane. The cylindrical radial direction (along the equator) is along  $\hat{\Omega}^\perp$ , and the rotation axis is along  $\hat{\Omega}$ . The local radial direction is (approximately) along the effective gravity direction  $\mathbf{e}_g$ , which is misaligned with respect to the  $x$ -direction when  $\phi$  is non-zero. The magnetic field is along  $z$ .

### 3 LINEAR THEORY

We consider linear perturbations to our flow  $\mathbf{U}_0$ , thermal state, and magnetic field  $\mathbf{B}_0$ . From equations (1)–(6), such velocity ( $\mathbf{u}$ ), magnetic ( $\mathbf{B}$ ), pressure ( $p$ ), and temperature ( $\theta$ ) perturbations are described by (where we have avoided introducing hats on perturbations)

$$D\mathbf{u} + 2\hat{\Omega} \times \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{U}_0 = -\nabla p + \theta \mathbf{e}_g + \mathbf{B}_0 \cdot \nabla \mathbf{B} + \nu \nabla^2 \mathbf{u}, \quad (11)$$

$$D\theta + \mathcal{N}^2 \mathbf{u} \cdot \mathbf{e}_\theta = \kappa \nabla^2 \theta, \quad (12)$$

$$D\mathbf{B} = \mathbf{B}_0 \cdot \nabla \mathbf{u} + \mathbf{B} \cdot \nabla \mathbf{U}_0 + \eta \nabla^2 \mathbf{B}, \quad (13)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (14)$$

$$\nabla \cdot \mathbf{u} = 0, \quad (15)$$

$$D \equiv \partial_t + \mathbf{U}_0 \cdot \nabla. \quad (16)$$

Note that we have defined our field and flow to satisfy  $\mathbf{B}_0 \cdot \nabla \mathbf{U}_0 = 0$  so that the basic state is an equilibrium configuration. Note that this restriction was *not* made in many prior works, including Balbus & Hawley (1994), Menou et al. (2004), and Menou & Le Mer (2006), but it is necessary to have a well-defined steady basic state. It is unclear whether results obtained for any other poloidal field configuration (with a time-dependent basic state) are valid. We might expect results in such cases to only be approximately valid if growth times are sufficiently small compared with the time-scale for the evolution of the basic state but not when the instability grows weakly.

#### 3.1 Dispersion relation for axisymmetric modes

We consider axisymmetric modes with meridional wave vectors  $\mathbf{k} = (k_x, 0, k_z) = k(\cos \theta_k, 0, -\sin \theta_k)$  with magnitudes  $k = \sqrt{k_x^2 + k_z^2}$  and angles  $\theta_k$ , since axisymmetric modes are likely to be the fastest growing (e.g. Latter & Papaloizou 2018), and we define

$\hat{\mathbf{k}} = \mathbf{k}/k$ . These permit complex exponential solutions proportional to  $\exp(ik_x x + ik_z z + st)$ . We define the complex growth rate  $s = \sigma + i\omega$ , where the growth (decay) rate  $\sigma \in \mathbb{R}$  and the oscillation frequency  $\omega \in \mathbb{R}$ . We manipulate equations (11)–(16) for such perturbations and define the modified growth rates  $s_\nu = s + \nu k^2$ ,  $s_\kappa = s + \kappa k^2$ , and  $s_\eta = s + \eta k^2$ , to obtain the quintic dispersion relation

$$s_\eta^2 s_\nu^2 s_\kappa + 2s_\eta s_\nu s_\kappa \omega_\Lambda^2 + s_\kappa \omega_\Lambda^4 + a s_\eta^2 s_\kappa + s_\kappa \xi + b(s_\eta^2 s_\nu + s_\eta \omega_\Lambda^2) = 0, \quad (17)$$

where

$$a = \frac{2}{\varpi} (\hat{\mathbf{k}} \cdot \hat{\Omega})(\hat{\mathbf{k}} \cdot (\nabla \ell)^\perp) \quad (18)$$

$$= \frac{2\Omega}{k^2} (s_\Lambda k_x + c_\Lambda k_z)(2\Omega k_x s_\Lambda + (2\Omega c_\Lambda - \mathcal{S})k_z) \quad (19)$$

$$= \frac{2\Omega |\nabla \ell|}{\varpi} s_{\Lambda-\theta_k} s_{\gamma-\theta_k}, \quad (20)$$

$$b = \mathcal{N}^2 (\hat{\mathbf{k}} \cdot \mathbf{e}_\theta^\perp)(\hat{\mathbf{k}} \cdot \mathbf{e}_g^\perp) \quad (21)$$

$$= \frac{\mathcal{N}^2}{k^2} (k_z c_\Gamma - k_x s_\Gamma)(k_z c_\phi - k_x s_\phi) \quad (22)$$

$$= \mathcal{N}^2 s_{\theta_k+\phi} s_{\theta_k+\Gamma}, \quad (23)$$

$$\xi = -2(\hat{\mathbf{k}} \cdot \hat{\Omega}) \mathcal{S} \omega_\Lambda^2 \hat{k}_z = 2\mathcal{S} \Omega s_{\Lambda-\theta_k} s_{\theta_k} \omega_\Lambda^2, \quad (24)$$

and

$$\omega_\Lambda^2 = (\mathbf{B}_0 \cdot \mathbf{k})^2 = k^2 B_0^2 s_{\theta_k}^2 \quad (25)$$

is the squared Alfvén frequency. In the above  $c_\Lambda$  and  $s_\Lambda$  refer to  $\cos \Lambda$  and  $\sin \Lambda$  for brevity, and similarly for trigonometric functions with other arguments (though  $s_\nu$ ,  $s_\kappa$ , and  $s_\eta$  always represent modified growth rates instead). We have also defined the local AM gradient

$$\nabla \ell = \varpi(2\Omega c_\Lambda - \mathcal{S}, 0, -2\Omega s_\Lambda), = |\nabla \ell|(c_\gamma, 0, -s_\gamma), \quad (26)$$

which has magnitude

$$|\nabla \ell|^2 = \varpi^2 (\mathcal{S}^2 + 4\Omega(\Omega - \mathcal{S}c_\Lambda)). \quad (27)$$

The normal to this is

$$(\nabla \ell)^\perp = \varpi(2\Omega s_\Lambda, 0, 2\Omega c_\Lambda - \mathcal{S}) = |\nabla \ell|(s_\gamma, 0, c_\gamma). \quad (28)$$

We also define the vector perpendicular to the effective gravity,

$$\mathbf{e}_g^\perp = (-s_\phi, 0, c_\phi), \quad (29)$$

and the normal to stratification surfaces,

$$\mathbf{e}_\theta^\perp = (-s_\Gamma, 0, c_\Gamma). \quad (30)$$

The baroclinic shear (along the rotation axis) is

$$\hat{\Omega} \cdot (\nabla \ell) = -\mathcal{S} \varpi s_\Lambda = |\nabla \ell| s_{\gamma-\Lambda}. \quad (31)$$

The dispersion relation (17) can be expanded out as a quintic equation

$$s^5 + c_1 s^4 + c_2 s^3 + c_3 s^2 + c_4 s + c_5 = 0, \quad (32)$$

where the coefficients  $c_1$  to  $c_5$  are given by

$$c_1 = k^2(2\eta + 2\nu + \kappa), \quad (33)$$

$$c_2 = k^4(\eta^2 + 2\eta\kappa + 4\eta\nu + 2\nu\kappa + \nu^2) + 2\omega_\Lambda^2 + a + b, \quad (34)$$

$$c_3 = k^6(\eta^2\kappa + 2\eta\nu^2 + 2\eta^2\nu + \kappa\nu^2 + 4\eta\kappa\nu) + 2\omega_\Lambda^2 k^2(\eta + \nu + \kappa) + a k^2(2\eta + \kappa) + b k^2(2\eta + \nu), \quad (35)$$

$$c_4 = k^8(2\eta v^2\kappa + 2\eta^2\nu\kappa + \eta^2v^2) + 2\omega_A^2k^4(\eta\nu + \eta\kappa + \nu\kappa) + \omega_A^4 + ak^4(2\eta\kappa + \eta^2) + \xi + bk^4(2\eta\nu + \eta^2) + b\omega_A^2, \quad (36)$$

$$c_5 = k^{10}\eta^2v^2\kappa + 2\omega_A^2k^6\eta\nu\kappa + \omega_A^4k^2\kappa + a\eta^2\kappa k^6 + \xi k^2\kappa + bk^6\eta^2\nu + b\omega_A^2k^2\eta. \quad (37)$$

### 3.2 Non-diffusive (in)stability

Non-diffusive modes, i.e. those with  $\nu = \kappa = \eta = 0$ , are described by the reduced dispersion relation

$$s^4 + (2\omega_A^2 + a + b)s^2 + (\omega_A^4 + b\omega_A^2 + \xi) = 0, \quad (38)$$

ignoring neutral modes with  $s = 0$ . Note that the only appearance of the magnetic field is through the combination  $\mathbf{B}_0 \cdot \mathbf{k} = B_0 k_z$  in  $\omega_A$ , therefore the adiabatic growth rate is independent of  $B_0$  if arbitrary  $k_z$  are permitted. This can be solved to give

$$s^2 = \frac{-(2\omega_A^2 + a + b) \pm \sqrt{4a\omega_A^2 - 4\xi + (a + b)^2}}{2}. \quad (39)$$

If we take  $B_0 = 0$ , then this reduces to the adiabatic dispersion relation in Paper III (equation 27),  $s^2 = -(a + b)$ . We would also obtain  $s^2 = -(a + b)$  in the diffusive case if we took  $k \rightarrow 0$  in equation (32), since  $\omega_A^2 \rightarrow 0$  in this limit, thereby eliminating the influence of magnetic fields on such modes.

Since  $s^2$  is always negative if  $a + b$  is positive, with  $s$  being purely imaginary, then the system is SH stable. The discriminant  $\Delta = 4a\omega_A^2 - 4\xi + (a + b)^2$  in equation (38) is always positive, so the roots for  $s^2$  are always real. Hence, non-diffusive oscillatory instabilities cannot occur. To see this, note that from equations (18) and (28) we have

$$\hat{\mathbf{k}} \cdot (\nabla\ell)^\perp = \varpi(2(\hat{\mathbf{k}} \cdot \boldsymbol{\Omega}) - \hat{k}_z \mathcal{S}), \quad (40)$$

and using the definition of  $\xi$  (equation 24),

$$\Delta = 4a\omega_A^2 - 4\xi + (a + b)^2 = (a + b)^2 + 16\omega_A^2(\hat{\mathbf{k}} \cdot \boldsymbol{\Omega})^2, \quad (41)$$

which being the sum of squares must be non-negative.

The criterion for onset of direct instability (real roots) occurs when (for neutral stability  $s = 0$ )

$$\omega_A^4 + b\omega_A^2 + \xi = 0, \quad (42)$$

and instabilities occur when this term is negative. So the only way to destabilize a hydrodynamically SH stable configuration without diffusion is for the left-hand side of equation (42) to be negative, which corresponds with a direct instability, the MRI. MRI works best with a weak field (or on large length-scales), meaning the stabilizing term  $\omega_A^4$  is small compared with the others, and when the fluid is neutrally rather than stably stratified ( $b = 0$ ). Then, MRI just requires a mode with a  $\mathbf{k}$  which makes  $\xi$  negative. In the weak field or small wavenumber case,  $\omega_A^4 \rightarrow 0$  faster than the remaining terms in equation (42), so for non-zero  $B_0$ , instability occurs if

$$b - 2(\hat{\mathbf{k}} \cdot \boldsymbol{\Omega})\mathcal{S}\hat{k}_z < 0. \quad (43)$$

Hence in the special case of cylindrical differential rotation,  $\Omega(\varpi)$ , we have  $\mathcal{N}^2 - 2\Omega\mathcal{S} < 0$ . If the stabilizing effects of buoyancy are absent, the stability criterion in the weak field case is  $-2\Omega\mathcal{S} < 0$ , which involves angular velocity rather than AM gradients (e.g. Balbus & Hawley 1998). So  $\mathcal{S} > 0$  is required for instability (to MRI), which is generally much easier to satisfy than Rayleigh's criterion for centrifugal instability, which requires  $\mathcal{S} > 2$  in the hydrodynamic case, implying outwardly decreasing AM.

The three quantities  $a$ ,  $b$ , and  $\xi$  that appear in equation (38) correspond to three different instability mechanisms. If there is no stratification or magnetic field, angular-momentum-driven instability occurs if  $\mathbf{k}$  can be chosen so that  $a < 0$ . In the case of cylindrical rotation this is just Rayleigh's criterion that instability occurs when the AM decreases outwards. More generally, using equation (20),  $a < 0$  when the wave vector  $\mathbf{k}$  lies in the wedge between  $\boldsymbol{\Omega}^\perp$  and  $\nabla\ell$ . The quantity  $b$  is associated with the stratification, see equation (23), and baroclinic instability  $b < 0$  occurs when the wave vector  $\mathbf{k}$  lies in the wedge between  $\mathbf{e}_g$  and  $\mathbf{e}_\theta$ . When the stratification is very strong, the TWE makes this wedge angle small, so baroclinic instability is weak. Then in a convectively stable region  $b$  will be large and positive unless the wave vector is nearly parallel to gravity, i.e. horizontal flow along the isobars. The quantity  $\xi$ , equation (24), is associated with MRI instability. From equation (42) if  $b > 0$  we need  $\xi < 0$  for instability. This happens if the wave vector  $\mathbf{k}$  lies outside the wedge between  $\hat{\boldsymbol{\Omega}}^\perp$  and the  $x$ -axis.

#### 3.2.1 Marginal stability to stratified non-diffusive MRI

In order to find the non-diffusive unstable modes for weak fields, we substitute  $\mathbf{k} = k(\cos\theta_k, 0, -\sin\theta_k)$  into equation (43) and solve for the marginal stability lines, giving

$$\mathcal{N}^2 \sin(\theta_k + \Gamma) \sin(\theta_k + \phi) + 2\Omega\mathcal{S} \sin(\Lambda - \theta_k) \sin\theta_k = 0. \quad (44)$$

In the strongly stratified limit,  $\mathcal{N}^2 \gg |\Omega\mathcal{S}|$  and  $\Gamma \sim \phi$  (from equation 9),

$$\mathcal{N}^2 \sin^2(\theta_k + \phi) - 2\Omega\mathcal{S} \sin(\theta_k - \Lambda) \sin\theta_k = 0. \quad (45)$$

If  $\mathcal{N}^2 \gg 2|\Omega\mathcal{S}|$ , this can only be satisfied when  $\sin^2(\theta_k + \phi) \sim 0$ , hence  $\theta_k + \phi \approx n\pi$  where  $n \in \mathbb{N}$ . Note that  $\theta_k$  is defined below the  $x$ -axis so  $-\theta_k$  is the angle above it. This means that  $-\theta_k = \phi = \Gamma$  when  $n = 0$ , indicating that  $\mathbf{k}$  lies along  $\mathbf{e}_\theta$  or  $\mathbf{e}_g$ , so that fluid motions are along stratification (or constant pressure) surfaces, i.e. parallel to  $\mathbf{e}_\theta^\perp \sim \mathbf{e}_g^\perp$ . Hence instability is possible for a wedge of wave vector angles around  $\mathbf{e}_\theta$  (e.g. Balbus 1995). If  $\phi > 0$  and the box is in the Northern hemisphere,  $\mathbf{e}_g$  lies outside the MRI-stable wedge between  $\hat{\boldsymbol{\Omega}}^\perp$  and the  $x$ -axis, see Fig. 2, and we expect MRI instability. However, if  $\phi < 0$  in the Northern hemisphere,  $\mathbf{e}_g$  lies inside the MRI-stable wedge (recall  $\Lambda + \phi > 0$  in the Northern hemisphere so  $\Lambda > |-\phi|$ ), and so no MRI will occur, though there may be unstable GSF modes.

#### 3.2.2 Fastest growing non-diffusive modes

We now find the wave vector magnitude  $k$  and orientation  $\theta_k$  corresponding to the maximum growth rate, and in turn identify the dominant mode. To find the fastest growing mode we first maximize over  $k^2$ , and then maximize over the angle  $\theta_k$ . Note  $a$  and  $b$  only depend on  $\theta_k$  and not on the magnitude  $k$ , so  $\partial a/\partial k^2$  and  $\partial b/\partial k^2$  are both zero. We also have

$$\frac{\partial \omega_A^2}{\partial k^2} = \frac{\omega_A^2}{k^2} \quad \text{and} \quad \frac{\partial \xi}{\partial k^2} = \frac{\xi}{k^2}. \quad (46)$$

To obtain the fastest growing mode properties, we differentiate equation (38) with respect to both  $k^2$  and  $\theta_k$  and require  $\partial_{k^2} s = \partial_{\theta_k} s = 0$ . Setting the  $k^2$  derivative of equation (38) to zero gives

$$s^2 = -\frac{\mathcal{N}^2}{2} s_{\phi+\theta_k} s_{\Gamma+\theta_k} - \Omega\mathcal{S} s_{\theta_k} s_{\Lambda-\theta_k} - k^2 B_0^2 s_{\theta_k}^2. \quad (47)$$

This is clearly maximized for weak fields or for modes with  $k \rightarrow 0$  where the last term vanishes, since that provides a stabilizing effect,

though we must have non-zero  $B_0$  to obtain this result. Now we can maximize over  $\theta_k$  to obtain

$$0 = -\frac{\mathcal{N}^2}{2} s_{\Gamma+\phi+2\theta_k} - \Omega S s_{\Lambda-2\theta_k} - k^2 B_0^2 s_{2\theta_k}. \quad (48)$$

In the strongly stratified limit the dominant term is usually the first one involving  $\mathcal{N}^2$ , which is stabilizing, unless we choose a specific range of  $\theta_k$ . This means that in order to maximize the growth rate we need to minimize this term. Indeed the magnitude of this term is smallest when  $\theta_k \approx -(\Gamma + \phi)/2$ , i.e. when the wave vector is approximately halfway between  $\mathbf{e}_g$  and  $\mathbf{e}_\theta$ . Note that in the strongly stratified limit the TWE implies that  $\phi \approx \Gamma$  and hence this term approximately vanishes for  $\theta_k \approx -\phi \approx -\Gamma$ . For such wave vectors that minimize the stabilizing effects of buoyancy,

$$s^2 = \Omega S s_\phi s_{\Lambda+\phi} - B_0^2 \omega_A^2 s_\phi^2. \quad (49)$$

Note that the magnetic term that only occurs in the non-weak field limit is always negative, meaning that in the adiabatic regime a stronger magnetic field should decrease the maximum growth rate of the instability. In the weak field case, where we can ignore the second term, we are left with

$$s^2 = 2\Omega S s_\phi s_{\Lambda+\phi}. \quad (50)$$

Hence, we require both  $\phi$  and  $\Lambda + \phi$  to have the same sign, either both in the Northern or Southern hemisphere for onset of instability.

### 3.3 Diffusive instabilities

#### 3.3.1 Small Pr/Pm limit: very efficient thermal diffusion

In the limit of very efficient thermal diffusion relative to viscous and ohmic diffusion, we would expect equation (38) to approximately apply for sufficiently large wavelength instabilities (with smallish  $k$ , for which viscous and ohmic diffusion are relatively unimportant) but with  $b = 0$ . To show that this is indeed the case, if we consider equation (17), set  $v = \eta = 0$  and then consider the limit  $\kappa \rightarrow \infty$ . This means that in equation (17)  $\eta k^2 \ll s$  and  $\nu k^2 \ll s$  but  $\kappa k^2 \gg s$ , which is like considering the joint limits  $\text{Pr}/\text{Pm} \rightarrow 0$  and  $\text{Pr} \rightarrow 0$  with all other quantities  $O(1)$ . We obtain the dispersion relation

$$s^4 + (2\omega_A^2 + a)s^2 + (\omega_A^4 + \xi) = 0. \quad (51)$$

This is the same as equation (38) with  $b = 0$  and describes MRI modes satisfying the unstratified ( $b = 0$ ) non-diffusive dispersion relation with non-zero field.<sup>2</sup> The fastest growing modes (maximizing over  $k^2$ , i.e. setting  $\partial_{k^2} s = 0$ ) in the limit of weak fields or small  $k$  (for which  $\omega_A^4$  can be ignored relative to the other terms) satisfy

$$s^2 = -\frac{\xi}{2\omega_A^2} = (\hat{\mathbf{k}} \cdot \boldsymbol{\Omega}) S \frac{k_z}{k} = S \Omega s_{\theta_k - \Lambda} s_{\theta_k}. \quad (52)$$

In this limit instability occurs for any  $S > 0$  (though strictly the approximations for which this limit applies are then no longer valid). The growth rate is maximized over  $\theta_k$  when  $\partial_{\theta_k} s^2 = 0$ , giving

$$s_{\Lambda-2\theta_k} = 0 \quad \Rightarrow \quad \theta_k = \frac{\Lambda}{2} - n \frac{\pi}{2} \quad (53)$$

<sup>2</sup>This is analogous to what was found for the hydrodynamic case in the limit  $\text{Pr} \rightarrow 0$  and  $\text{RiPr} \rightarrow 0$ , where the fastest growing mode growth rates were described by  $s^2 = -a$ , the adiabatic unstratified dispersion relation (Paper II; Paper III). However, the dispersion relation here requires the presence of non-vanishing magnetic field.

for  $n \in \mathbb{N}$ , i.e. for modes with orientations halfway between the rotation axis (along  $\hat{\boldsymbol{\Omega}}$ ) and the angular velocity gradient (along  $x$ ) when  $n = 1$ . For cylindrical differential rotation ( $\Lambda = 0$ ), this implies  $\theta_k = \pm(\pi/2)$ , and hence wave vectors are along  $z$ , as expected (e.g. Balbus & Hawley 1991). On the other hand, when  $\Lambda = -30^\circ$ ,  $\theta_k = -105^\circ$  (indicating  $105^\circ$  above the  $x$ -axis), and when  $\Lambda = 60^\circ$ ,  $\theta_k = -60^\circ$  (indicating  $60^\circ$  above the  $x$ -axis). This is consistent with our later Figs 3–5 for the largest Pm considered, and is most evident for the strongest magnetic fields plotted there.

For even stronger fields or larger wavenumbers, there is a stabilizing effect of magnetic tension through the  $\omega_A^4$  term in the dispersion relation. Fields are sufficiently strong when  $\omega_A^2 = B_0^2 k^2 s_{\theta_k}^2 \sim 2\Omega S s_{\theta_k - \Lambda} s_{\theta_k}$ , and hence typically for  $k^2 \sim 2\Omega S / B_0^2$ . In addition, larger  $k$  modes would be increasingly affected by ohmic diffusion and viscosity.

#### 3.3.2 Diffusive modes in the small shear (small $S/\Omega$ ) limit

In Paper III, we computed the curves showing the lowest value of the shear  $S$  for which instability is possible as a function of the angle  $\phi$ , the angle between the shear and gravity directions (see fig. 6 in Paper III). Apart from the exceptional case on the equator, there is a finite minimum  $S$  below which no instability occurs. This is no longer the case when a magnetic field is added. There is then a whole range of  $\phi$  for which the system is unstable for arbitrarily small  $S$ . This is quite surprising, as both GSF and MRI instability are driven by the shear, so one might imagine that reducing the shear towards zero would eliminate them. What happens is that the growth rate does tend to zero as  $S$  is reduced, but it can always remain positive, so the critical value of  $S$  for instability can be zero.

To establish this, we consider the case where  $S/\Omega \rightarrow 0$ , and seek modes with small but positive growth rate  $s \sim O(S)$ . We now consider the ordering of the terms in the quintic dispersion relation equation (17). We retain the diffusive terms, choosing  $O(\eta k^2) \sim O(\nu k^2) \sim O(\kappa k^2) \sim O(S)$ .  $a$  is  $O(\Omega^2)$  and we choose the magnetic field strength so that  $\omega_A^2 \sim O(S\Omega)$ , which makes  $\xi \sim O(S^2\Omega^2)$ . If we take  $\mathcal{N}^2 \sim O(\Omega^2)$  or larger, it appears we have an inconsistency, because then the dominant term in equation (17) is  $b s_\eta \omega_A^2 \sim O(S^2\Omega^3)$ , whereas the remaining terms are  $O(S^3\Omega^2)$  or smaller. However, the TWE (equation 9) implies that  $\Gamma$  and  $\phi$  are almost aligned if  $S/\Omega \ll 1$ . If we choose our wavenumber  $\mathbf{k} = k(c_{\theta_k}, 0, -s_{\theta_k})$  so that  $\theta_k = \pi - (\Gamma + \phi)/2$ , aligned in the limit  $S/\Omega \rightarrow 0$  with both  $\mathbf{e}_\theta$  and  $\mathbf{e}_g$ , then  $b = \mathcal{N}^2 s_{\theta_k + \Gamma} s_{\theta_k + \phi} = -\mathcal{N}^2 s_{\Gamma/2 - \phi/2}^2$ , and now the TWE (equation 9) implies that  $b \sim O(S^2)$ , much smaller than  $O(\Omega^2)$ . Numerical solutions of equation (17) show that the critical wavenumber  $\mathbf{k}$  is indeed aligned with  $\mathbf{e}_\theta$  and  $\mathbf{e}_g$ . Now the inconsistency in the  $b$  term in equation (17) is removed, because this term is now negligible compared with the  $O(S^3\Omega^2)$  terms. We obtain


$$a s_\eta^2 s_\kappa + (\omega_A^4 + \xi) s_\kappa = 0, \quad (54)$$

where


$$\begin{aligned} \omega_A^2 &= B_0^2 k^2 \sin^2 \phi, & a &= 4\Omega^2 \sin^2 \beta, \\ \xi &= -2k^2 S \Omega B_0^2 \sin \beta \sin^3 \phi, \end{aligned} \quad (55)$$

and  $\beta = \Lambda + \phi$  is the latitude. Dividing by  $s_\kappa$ , which must be positive for growing modes,


$$a s_\eta^2 + \omega_A^4 + \xi = 0, \quad \text{implying } s = -\eta k^2 + \sqrt{\frac{-\omega_A^4 - \xi}{a}}. \quad (56)$$



**Figure 3.** Logarithm of the growth rate  $\log_{10}(\sigma/\Omega)$  of axisymmetric perturbations plotted on the  $(k_x, k_z)$  plane according to equation (17), for various  $B_0$  and  $Pm$ , with  $\phi = 30^\circ$ ,  $\Lambda = -30^\circ$ , i.e. a mixed radial/latitudinal shear at the equator with latitude  $\Lambda + \phi = 0^\circ$ . Parameters are  $\mathcal{N}^2/\Omega^2 = 10$ ,  $\text{Pr} = 10^{-2}$ , and  $S/\Omega = 2$ . We vary the strength of the magnetic field from  $B_0 = 0$  to  $B_0 = 5$  down each column, and vary  $Pm$  from  $Pm = 0.01$  to  $Pm = 1$  along each row. GSF modes are primarily confined within the wedge bounded by  $\hat{\Omega}^\perp$  and  $\nabla\ell$  (red lines). However, as the field strength increases (downwards) the wave vector orientation is shifted to correspond more with MRI. Increasing  $B_0$  at fixed  $Pm$  decreases both the maximum growth rate and the size of the unstable region on the  $(k_x, k_z)$  plane. Reducing magnetic diffusivity by increasing  $Pm$  on the other hand seems to have the opposite effect and enhances the destabilizing effects of the field, leading to a larger region of instability as well as a stronger destabilization of the dominant mode.



**Figure 4.** Logarithm of the growth rate  $\log_{10}(\sigma/\Omega)$  of axisymmetric perturbations plotted on the  $(k_r, k_z)$  plane according to equation (17), for various  $B_0$  and  $Pm$ , with  $\phi = -30^\circ$  and  $\Lambda = 60^\circ$ , i.e. a mixed radial/latitudinal shear at latitude  $\Lambda + \phi = 30^\circ$ . Parameters are  $\mathcal{N}^2/\Omega^2 = 10$ ,  $Pr = 10^{-2}$ , and  $\mathcal{S}/\Omega = 2$ . We vary the strength of the magnetic field from  $B_0 = 0$  to  $B_0 = 5$  down each column, and vary  $Pm$  from  $Pm = 0.01$  to  $Pm = 1$  along each row. GSF unstable modes are primarily confined to within the wedge bounded by  $\hat{\Omega}^\perp$  and  $\nabla\ell$  (red lines) for weak fields, but this direction is modified when the MRI takes over. We observe a secondary set of unstable oscillatory modes, consisting of weakly destabilized magneto-inertial-gravity waves. These grow more weakly than the primary lobes but they are less affected by the stabilizing effects of the magnetic field and even operate (as in the  $B_0 = 5$ ,  $Pm = 0.01$  case) when the primary lobes have been stabilized by it.



**Figure 5.** Logarithm of the growth rate  $\log_{10}(\sigma/\Omega)$  of axisymmetric perturbations plotted on the  $(k_x, k_z)$  plane according to equation (17), for various  $B_0$  and  $Pm$ , with  $\phi = 60^\circ$  and  $\Lambda = -30^\circ$ , i.e. a mixed radial/latitudinal shear at latitude  $\Lambda + \phi = 30^\circ$ . Parameters are  $\mathcal{N}^2/\Omega^2 = 10$ ,  $Pr = 10^{-2}$ , and  $S/\Omega = 2$ . We vary the strength of the magnetic field from  $B_0 = 0$  to  $B_0 = 5$  down each column, and vary  $Pm$  from  $Pm = 0.01$  to  $Pm = 1$  along each row. When  $B_0 = 0$  the system is adiabatically unstable since it violates the SH criterion. This is visually characterized by a tendency for the fastest growing modes to occur even as  $k \rightarrow 0$ , suggesting that the presence of diffusion leads to the preference of the largest possible wavelengths in this regime. This is in comparison to the GSF and MRI modes cases where the fastest growing modes here have a unique non-zero wavenumber and hence a preferred wavelength in real space.

From equation (55)  $a > 0$ , so for instability we must have  $\xi < 0$ , and so for  $\mathcal{S} > 0$  the angle  $\phi$  and the latitude  $\beta$  must have the same sign for instability, so in the Northern hemisphere,  $\beta > 0$ , we must have  $\phi > 0$  for our small  $\mathcal{S}$  modes to be unstable. In the solution equation (56), the largest term as  $k \rightarrow 0$  is the  $\xi$  term, so that for negative  $\xi$  there is always a small enough  $k$  that makes  $s > 0$ , so that that however small  $\mathcal{S}$  is, it is always possible to find a growing mode. This behaviour is illustrated below in Fig. 12 for latitude  $30^\circ$  north showing instability for very small  $\mathcal{S}$  when  $\phi > 0$ . For very large  $\mathcal{N}^2$ , as we might expect in stars, and for small  $\mathcal{S}$  (weak differential rotations), the inertial term  $a$  is negligible. Since  $b$  would then in general dominate all other terms in equation (17), for instability we must choose modes for which fluid motion is along stratification surfaces, meaning  $\mathbf{k}$  must be parallel or antiparallel to gravity. For small  $\mathcal{S}$ ,  $\mathbf{e}_g$  is very close to  $\mathbf{e}_\theta$ . Why cannot you wipe out the effect of the stratification by thermal diffusion in this limit as you can with GSF (for example)? For thermal diffusion to work,  $k$  must be reasonably large, and GSF modes are stabilized by the magnetic field at large (or moderate)  $k$ . At small  $\mathcal{S}$ , MRI is stabilized by the  $\omega_A^4$  term too because  $\xi$  is proportional to  $\mathcal{S}$  unlike  $\omega_A^4$ . Only small  $k$  (or small  $B_0$ ) reduces  $\omega_A^4$  relative to  $\xi$ , and having small  $k$  does not allow efficient enough thermal diffusion for instability. Hence, the fundamental MRI mechanism that operates for small  $\mathcal{S}$  is typically non-diffusive.

As  $\mathcal{N}^2$  is reduced,  $\mathcal{S}$  is made larger, or the field is reduced, then GSF modes with a larger  $k$  can operate, and there are more possibilities for instability. We will return to this point later when analysing instability in the solar tachocline.

#### 4 NUMERICAL LINEAR RESULTS

In this section, we solve the dispersion relation and graphically analyse the properties of the possible instabilities in our system. In Figs 3–5, we probe effects of varying the magnetic field strength  $B_0$  and magnetic Prandtl number  $\text{Pm}$  for three different configurations with different latitudes  $\beta = \Lambda + \phi$  and orientations of the shear with respect to gravity  $\phi$ . We fix  $\text{Pr} = 0.01$  small, but motivated by parameters accessible with non-linear numerical simulations, and  $S = 2$  (following Paper II; Paper III), since the latter choice would be marginally stable according to Rayleigh’s criterion for cylindrical differential rotation. We present pseudo-colour plots of the base 10 logarithm of the growth rate of an axisymmetric perturbation in Fourier space  $(k_x, k_z)$  in these figures. Overplotted in red are the lines  $\hat{\Omega}^\perp$  and  $\nabla\ell$ , within which the direct GSF instability occurs, and in light blue are the directions of buoyancy (more specifically, the normal to stratification surfaces) and gravity,  $\mathbf{e}_\theta$  and  $\mathbf{e}_g$ , respectively. For comparison the hydrodynamic cases with  $B_0 = 0$  are shown in the top row of each figure. The magnetic field strength  $B_0$  is increased within the set  $[0, 1, 2.5, 5]$  with each successive row, and  $\text{Pm}$  is increased within  $[0.01, 0.1, 1]$  with each successive column.

We identify two sets of ‘lobes’ of instability operating in the system. The dominant sets are bounded by  $\nabla\ell$  and  $\hat{\Omega}^\perp$  in the hydrodynamic case, and they correspond to the dominant direct instability. This is either the double-diffusive GSF or the adiabatic SH instability (see section 3.2 of Paper III for the explicit conditions required for the latter to operate). The fastest growing modes typically have growth rates  $O(1)$ , which we note is comparable to  $\Omega^{-1}$ , given our unit of time, and are initially (in the hydrodynamic case) observed to lie along the line that is approximately halfway between  $\nabla\ell$  and  $\hat{\Omega}^\perp$ . This wedge is perpendicular to the physical wedge within which the unstable mode displacements and velocity perturbations arise due

to incompressibility. The second set of smaller lobes, when present, contains oscillatory modes, which are weakly growing internal magneto-inertia-gravity waves that propagate and are destabilized within the wedges bounded by  $\mathbf{e}_g$  and  $\mathbf{e}_\theta$ .

The introduction of non-zero  $B_0$  has observable effects on the orientation, strength, and structure of the unstable region in parameter space. Increasing the strength of the field for a fixed  $\text{Pm}$  has a tendency to force the modes into alignment with the preferred direction for MRI modes, and to shift them to larger scales (smaller  $k$  magnitudes), both as predicted in Section 3.3. Note that we do not observe the MRI modes to be well aligned with  $\mathbf{e}_\theta$  here, as explained in Section 3.2. This is likely due to the effects of thermal diffusion in eliminating the stabilizing effects of buoyancy forces on MRI modes when  $\text{Pr}/\text{Pm} = \eta/\kappa$  and  $\text{Pr}$  are both small, which is the case here when  $\text{Pm} \geq 0.01$  as we fix  $\text{Pr} = 0.01$ . The addition of a field seems to impose a stabilizing effect on the hydrodynamically unstable GSF modes, ultimately resulting from the stabilizing effects of magnetic tension (see also Latter & Papaloizou 2018), and so the majority of cases exhibit a smaller growth rate for the dominant instability as  $B_0$  is increased.

Magnetic diffusion counteracts the effects of magnetic fields, and this is clearest for small  $\text{Pm}$  in the left panels of Figs 3–5. Cases with  $\text{Pm} = 1$  have the weakest ohmic diffusion, and cases with  $\text{Pm} = 0.01$  have much more efficient ohmic than viscous diffusion. Small  $\text{Pm}$  allows magnetic cases to return to the hydrodynamic limit and larger  $\text{Pm}$  (closer to unity, in our case) therefore exhibit the strongest magnetic effects for a given  $B_0$ . When  $\text{Pm} = 1$ , instability is possible outside the hydrodynamic region contained within the lines  $\hat{\Omega}^\perp$  and  $\nabla\ell$ . This can be seen most clearly in the rightmost bottom panel of these figures, where magnetic effects are strongest (and magnetic diffusion is weakest). The direction of the preferred modes in that case is better described by the unstratified (due to rapid thermal diffusion) MRI in Section 3.3.


The oscillatory modes seem to be only very marginally modified by the magnetic field, as is seen most clearly in Fig. 4. This suggests that the internal inertia-gravity waves observed to be destabilized in Paper III within the wedge between  $\mathbf{e}_g$  and  $\mathbf{e}_\theta$  continue to be weakly destabilized magneto-inertia-gravity waves.

#### 5 PARAMETER DEPENDENCE OF FASTEST GROWING MODE

After displaying the properties of all unstable axisymmetric modes on the  $(k_x, k_z)$  plane in our system as  $B_0$  and  $\text{Pm}$  are varied, we now turn to explore the variation in the fastest growing mode optimized over  $k_x$  and  $k_z$  as the parameters are varied. We primarily consider  $S = 2$  and  $N^2 = 10$  at four latitudes ( $\beta = \phi + \Lambda = 0^\circ, 30^\circ, 60^\circ$ , and  $90^\circ$ ).

##### 5.1 Non-diffusive instabilities

We first explore non-diffusive (stratified) instabilities by solving the quartic dispersion relation in equation (38) numerically (using `fminsearch` on  $-\Re[s]$  in MATLAB), to determine the properties of the fastest growing mode, which we present in Fig. 6 as a function of the direction of the differential rotation  $\phi$ , for three different latitudes  $\beta = 30^\circ, 60^\circ$ , and  $90^\circ$ . The left panel shows the growth rate  $\sigma$  of the fastest growing mode, the middle panel shows the corresponding wavenumber magnitude  $k$ , and the right panel shows the wave vector orientation  $\theta_k = \tan^{-1}(-k_z/k_x)$ . Note that the magnetic field only appears in equation (38) through powers of  $\omega_A$ , therefore the results



**Figure 6.** A selection of figures comparing the properties of the fastest growing non-diffusive ( $\nu = \kappa = \eta = 0$ ) modes with an imposed magnetic field for  $S = 2$  and  $N^2 = 10$  for latitudes  $30^\circ$ ,  $60^\circ$ , and  $90^\circ$ . Left panel: Maximum growth rate  $\sigma$  as a function of  $\phi$ . The blue dotted curve is where the fastest growing mode is hydrodynamic and the magnetic field plays no role, which prefers modes with  $k \rightarrow 0$ . The solid curves are where the corresponding  $k$  is non-zero and magnetic field affects the growth rate. Middle panel:  $k$  when it is finite and there is instability. Right panel: Corresponding wave vector orientation  $\theta_k$ , which is well-defined for all growing modes.

in the left and right panels of Fig. 6 are independent of the magnetic field, whereas the middle panel shows results for  $B_0 = 1$  but  $k$  can be straightforwardly scaled to consider any  $B_0$  since the  $y$ -axis can be interpreted as  $B_0 k$  (and  $k_z$  can be obtained using the corresponding  $\theta_k$ ). This means that for strong fields, instability prefers small  $k$ .

The blue lines in Fig. 6 are results at a latitude of  $30^\circ$ . Blue dashed lines are where the non-diffusive hydrodynamic SH instability operates, and blue lines where the magnetic field modifies the instability over the hydrodynamic case (shown in fig. 4b in Paper III). Values of  $\phi$  for which there are no blue or red line are non-diffusively stable. When the non-diffusive hydrodynamic SH modes are unstable in red, in the range approximately  $\phi \in [60^\circ, 150^\circ]$ , there is no preferred  $k$ , only a preferred wave vector orientation. In this limit, the blue lines suggest the wave vector  $k \rightarrow 0$  for this range of  $\phi$ . The magnetic field widens the unstable region to below  $\phi = 0$  from  $\phi \approx 30^\circ$ .

For latitudes  $60^\circ$  (green lines) and  $90^\circ$  (red lines), there are no purely hydrodynamically unstable non-diffusive modes, but it can be seen by comparison with fig. 4 in Paper III that the field widens the unstable range of  $\phi$ . The growth rate has a similar maximum value to the hydrodynamic case, with  $\sigma \sim 1$  for the maximal  $\phi$ .  $\phi \approx -30^\circ$  to  $180^\circ$  are typically the most unstable configurations, and they also have the largest wavelength (smallest  $k$ ) instabilities, whereas  $\phi$  approximately between  $-150^\circ$  and  $-30^\circ$  are typically non-diffusively stable. Note that we have obtained a new region of instability near to  $\phi \approx -180^\circ$  that we omitted from fig. 4 in Paper III but is also present in the hydrodynamic case.

## 5.2 Diffusive instabilities


In stellar radiation zones, rapid thermal diffusion means that  $\text{Pr} \ll 1$ ,  $\text{Pm} \ll 1$ , and  $\text{Pr}/\text{Pm} = \eta/\kappa \ll 1$ . Hence, the stratified non-diffusive instabilities we have just analysed are likely to be modified by thermal diffusion. To explore this, we solve the full triply diffusive dispersion relation equation (17) with  $\text{Pr} = 10^{-2}$  and  $\text{Pm} = 0.1$ , for which  $\text{Pr}/\text{Pm} = 0.1$  and is therefore small. Though these are small values they are not in the regime of stellar interiors – however these values are accessible for non-linear calculations. We show the growth rate (left panels), wave vector magnitudes (middle panels), and orientations  $\theta_k$  (right panels) as a function of  $\phi$  for various field strengths  $B_0 \in [0, 1, 2.5, 5, 10]$  in Figs 7, 8, 9, 10 for latitudes  $0^\circ, 30^\circ, 60^\circ$ , and  $90^\circ$ , respectively. These demonstrate the effects of

a magnetic field on linear growth rates over the complete range of differential rotation configurations (value of  $\phi$ ) with  $S = 2$ .


The equatorial case in Fig. 7 is symmetric about  $\phi = 0$  and is adiabatically (non-diffusively) stable for  $S = 2$  for any  $\phi$  and  $B_0$ . The hydrodynamic  $B_0 = 0$  case is stable when  $\phi = 0$ , corresponding to cylindrical rotation at the equator, but it becomes destabilized by even a weak magnetic field. This destabilization is seen at all latitudes and is a result of a change in the stability criteria governing instability here. In the hydrodynamic case, we require a violation of Rayleigh’s criterion, which requires AM to decrease outwards on isobars for instability, whereas in the magnetic case this criteria can – for certain field strengths – correspond to an MRI mode that requires angular velocity to decrease along isobars instead, which is much easier to satisfy. Within close proximity to the cylindrically rotating profile ( $-15^\circ \lesssim \phi \lesssim 15^\circ$ ), the magnetic instability operating is significantly more unstable than the hydrodynamic case. Increases in field strength of up to  $B_0 \approx 2.5$  increase linear growth rates of the dominant modes, paired with a decrease in their wavelengths (increase in  $k$ ), and a significant deviation in orientation from the hydrodynamic case there (towards  $\theta_k \sim 90^\circ$ , implying  $k$  is along  $z$ ). Outside of this region (particularly for  $\phi$  outside of  $-60^\circ \lesssim \phi \lesssim 60^\circ$ ), the field inhibits growth of the hydrodynamic GSF modes and increases their wavelength (reduces its  $k$ ). For these parameters it is clear that the magnetic field typically has a stabilizing effect except for close to cylindrical rotation profiles.

Fig. 8 shows the same results for latitude  $\Lambda + \phi = 30^\circ$ . Here the symmetry about  $\phi = 0$  seen at the equator is broken and varying  $\phi$  has more complicated effects. Cylindrical differential rotation corresponds here with  $\phi = 30^\circ$  (since then  $\Lambda = 0^\circ$ ), and we observe that it is stable when  $B_0 = 0$  but is destabilized by the addition of a magnetic field, with more magnetized cases becoming more unstable until the growth rate becomes independent of  $B_0$  for  $B_0 \gtrsim 2.5$ . After  $\phi \approx 60^\circ$  there is very good agreement between all cases. This is when the non-diffusive hydrodynamic SH instability operates (as seen in Fig. 6), which prefers  $k \rightarrow 0$ , and magnetic fields have little effect on it.

There is a notable change in the  $\phi$  range of non-zero  $|k|$  values as  $B_0$  is increased, which goes from  $169^\circ \lesssim \phi \lesssim -157^\circ \cup -135^\circ \lesssim \phi \lesssim 28^\circ$  in the hydrodynamic case (note that the boundary between  $180^\circ$  and  $-180^\circ$  is continuous due to symmetry) to  $168^\circ \lesssim \phi \lesssim -160^\circ \cup -100^\circ \lesssim \phi \lesssim 60^\circ$  in the strongest  $B_0 = 10$  case. Note that



**Figure 7.** Properties of the fastest growing modes for various values of the magnetic field  $B_0$  with  $S = 2$ ,  $\text{Pr} = 10^{-2}$ ,  $N^2 = 10$ , and  $\text{Pm} = 0.1$ , for different rotation profiles (values of  $\phi$ ) at the equator. The hydrodynamically stable case of  $\phi = 0^\circ$ , corresponding to cylindrical rotation, is destabilized by the magnetic field. Within close proximity of cylindrical rotation ( $-15^\circ \lesssim \phi \lesssim 15^\circ$ ) increases in field strength of up to roughly  $B_0 = 2.5$  increase the growth rate. This is paired with a decrease in the wavelength of this mode and deviation in orientation from the hydrodynamic case, where  $\theta_k$  tends to align itself more so with the orientation of the field. For other  $\phi$ , the field tends to stabilize the instability over the hydrodynamic case, reducing its maximum growth rate and wavenumber  $k$ .



**Figure 8.** Properties of the fastest growing modes for various values of the magnetic field  $B_0$  with  $S = 2$ ,  $\text{Pr} = 10^{-2}$ ,  $N^2 = 10$ , and  $\text{Pm} = 0.1$ , for different rotation profiles (values of  $\phi$ ) at a latitude  $\beta = \Lambda + \phi = 30^\circ$ . The addition of a magnetic field significantly alters the linear growth rate of the diffusive modes, and typically acts to reduce both the growth rate  $\sigma$  and wavenumber  $k$ , but it does not affect the adiabatically unstable region for  $\phi \in [60^\circ, 170^\circ]$ . The effect of the magnetic field depends on both field strength ( $B_0$ ) and differential rotation profile ( $\phi$ ). Nearly cylindrical differential rotations  $\phi \approx 30^\circ$  ( $\Lambda = 0$ ) that are hydrodynamically stable are heavily destabilized by the addition of a magnetic field, which corresponds to onset of the MRI.


in any region where more than one  $B_0$  has well-defined  $|k|$  values, the smaller  $B_0$  always has the shorter wavelength. In regions where the magnetic instability operates the field again acts to force  $\theta_k$  into alignment with the preferred direction for unstratified MRI modes discussed in Section 3.3 for these parameters. Regions where the dominant mode switches from one form of instability to another can also be seen by a discontinuity in  $\theta_k$ , as seen in panel (c).

At latitude  $\Lambda + \phi = 60^\circ$ , shown in Fig. 9, the effects of the field are in many ways similar to Fig. 8. The range of unstable  $\phi$  values does however decrease with increasing  $B_0$ , with the smallest range of instability being  $-145^\circ \lesssim \phi \lesssim -100^\circ$  in the hydrodynamic case up to  $-155^\circ \lesssim \phi \lesssim -59^\circ$  at  $B_0 = 10$ . However, the field is destabilizing, leading to larger growth rates in the range  $15^\circ \lesssim \phi \lesssim 160^\circ$ , where the largest  $B_0$  is the most unstable.


Similar results are found at latitude  $\Lambda + \phi = 90^\circ$  in Fig. 10. Modes with  $20^\circ \lesssim \phi \lesssim 150^\circ$  are destabilized by the field, with nearly cylindrical rotation profiles ( $\Lambda \sim 0^\circ$ ) being most strongly destabilized by the field. Cylindrical rotation is marginally stable in the hydrodynamic case but is the most unstable configuration for any  $B_0 > 1$  plotted here, and grows faster than any hydrodynamic case in this figure.

### 5.2.1 Variation of $B_0$ , $\text{Pm}$ , and $\text{Pr}$


In Fig. 11, we explore the dependence of our results on  $\text{Pm}$ ,  $\text{Pr}$ , and  $B_0$  at a latitude of  $30^\circ$ , using the same parameters otherwise as in Fig. 8. Our prior results indicate that increasing magnetic diffusion can mitigate the effects of the field and bring predictions closer to the hydrodynamic case when GSF unstable. We observe in Fig. 11 that increasing  $B_0$  inhibits instability for  $\phi < 0$  in the Northern hemisphere, but that smaller  $\text{Pm}/\text{Pr}$  (more efficient magnetic diffusion) can mitigate this, as seen by comparing the solid and dashed blue lines with the green dashed line. Fig. 11 indicates that there does not appear to be a single parameter neatly describing the competing influences of magnetic field and diffusion on instability. For example, increasing  $B_0$  tends to enhance instability around  $\phi \sim 0$  when comparing the solid red and dashed red lines (with the same  $\text{Pr}$  and  $\text{Pm}$ ) but it has the opposite effect for the solid blue and dashed blue lines. Decreasing  $\text{Pr}$ , thereby reducing viscosity relative to thermal diffusion, enhances instability for  $\phi < 0$  (compare the pink dashed and solid blue lines). Varying  $\text{Pm}$  can have different effects depending on other parameters, but we see reducing  $\text{Pm}$  tends to enhance instability for  $\phi < 0$  when comparing the green and blue



**Figure 9.** Properties of the fastest growing modes for various values of the magnetic field  $B_0$  with  $S = 2$ ,  $\text{Pr} = 10^{-2}$ ,  $N^2 = 10$ , and  $\text{Pm} = 0.1$ , for different rotation profiles (values of  $\phi$ ) at a latitude  $\beta = \Lambda + \phi = 60^\circ$ .



**Figure 10.** Properties of the fastest growing modes for various values of the magnetic field  $B_0$  with  $S = 2$ ,  $\text{Pr} = 10^{-2}$ ,  $N^2 = 10$ , and  $\text{Pm} = 0.1$ , for different rotation profiles (values of  $\phi$ ) at a latitude  $\beta = \Lambda + \phi = 90^\circ$ .




**Figure 11.** Properties of the fastest growing modes for various values of the magnetic field  $B_0$ ,  $\text{Pm}$ , and  $\text{Pr}$  with  $S = 2$ ,  $\text{Pr} = 10^{-2}$ , and  $N^2 = 10$ , for different rotation profiles (values of  $\phi$ ) at a latitude  $\beta = \Lambda + \phi = 30^\circ$ .

dashed lines (with  $B_0 = 10$ ), but inhibit it when comparing the solid blue and red lines. (Fixing  $\text{Pr}/\text{Pm} = \eta/\kappa$  is also clearly not the sole parameter of importance.)

### 5.2.2 Critical value of $S$

Finally, we determine numerically the critical value of  $S$  for instability ( $S_{\text{crit}}$ ), once again by optimizing over  $k_x$  and  $k_z$ . We show results in Fig. 12 for  $S_{\text{crit}}$  as a function of  $\phi$  for  $B_0 = 0.1$  (solid lines),  $B_0 = 1$  (dashed lines), and  $B_0 = 10$  (dotted lines) at a latitude

$30^\circ$ , along with the corresponding wave vector magnitude  $k$  and orientation  $\theta_k$ . Our numerical results confirm the arguments presented in Section 3.3.2. In complete contrast to the hydrodynamic case, we find MRI occurs for any  $S > 0$  for  $\phi > 0$ , such that  $S_{\text{crit}} = 0$  for such differential rotations. Note that if we had chosen a negative latitude value,  $S_{\text{crit}} = 0$  would have occurred for  $\phi < 0$ . The corresponding wavenumber  $k$  also becomes arbitrarily small, implying arbitrarily large wavelength instabilities according to our local model. When  $\phi < 0$ , the dominant instability is primarily the hydrodynamic GSF instability, weakly modified by magnetic fields. This has a preferred



**Figure 12.** Critical value of  $S$  for instability (top), and the corresponding wave vector magnitude (middle) and orientation (bottom), for  $B_0 = 0.1, 1$ , and  $10$ , with  $S = 2$ ,  $\text{Pr} = 10^{-2}$ ,  $N^2 = 10$ , and  $\text{Pm} = 0.1$ , for different rotation profiles (values of  $\phi$ ) at latitude  $\Lambda + \phi = 30^\circ$ . For  $\phi > 0$ , there is instability for any  $S > 0$  for  $B_0 \neq 0$ , consistent with results obtained in Section 3, due to the operation of the MRI (shown in red). This instability prefers arbitrarily small wavenumbers. For  $\phi < 0$ , the instability is similar to the hydrodynamic GSF instability (shown in blue; cf. fig. 6 in Paper III), and exhibits a preferred  $k \sim d$ . The magnetic field weakens operation of the GSF instability for  $\phi < 0$ .

$k = O(1)$  (in units of  $d^{-1}$ ) when it operates, and it is weakly inhibited by the presence of the magnetic field. We expect to find similar results – in terms of the modification of the hydrodynamic results shown in fig. 6 of Paper III – for different latitudes and field strengths. Stronger fields would widen operation of the MRI and inhibit the GSF modes further. However, GSF modes may still be the dominant instability for differential rotations with  $\phi < 0$ .

Overall, the addition of a magnetic field tends to inhibit diffusive rotational instabilities by reducing  $\sigma$  for  $\phi \lesssim 0$ , and to promote (increase  $\sigma$ ) instability for  $\phi \gtrsim 0$ , particularly for nearly cylindrical differential rotations ( $\Lambda \sim 0$ , where  $\phi$  equals the latitude). The wavelength of the dominant instability is typically affected by the strength of the field, with stronger fields generally exciting larger wavelengths (smaller  $k$ s). The orientation of the mode also differs from the hydrodynamic prediction for strong enough fields. The effects of magnetic fields on diffusive rotational instabilities are therefore complex, but in nearly all cases the field strongly modifies the growth rate or wavenumber of the dominant mode. We may thus expect magnetic fields to substantially modify turbulent transport in stellar radiative regions.

## 6 ENERGETICS OF THE INSTABILITIES

### 6.1 Derivation of the energy equations and evaluation for linear modes

In this section, we analyse the energetics of the instabilities in our model. This helps identify the physical mechanisms and energy sources that drive the various instabilities and quantify the role of magnetic fields. In order to derive the total energy equation we must first calculate equations that govern the different types of energy in our system, namely, kinetic, thermal/potential, and magnetic. We start with the equations governing the evolution of perturbations given by equations (11)–(16). To obtain volume-averaged energy equations we take the product of the relevant equation and quantity there (scalar product of equation 11 with  $\mathbf{u}$  and equation 13 with  $\mathbf{B}$  for kinetic and magnetic energies, and multiplication of equation 12 by  $\theta$  for thermal energy) and volume average. We denote volume averages by  $\langle \cdot \rangle$  where  $\langle \cdot \rangle = \frac{1}{V} \iiint \cdot dV$ , where  $V$  is the volume of our box, which for linear modes is taken to be a single wavelength of the dominant mode. We define the kinetic, magnetic, and thermal energies of our perturbations according to (assuming  $\mathcal{N}^2 > 0$ )

$$\mathcal{K} = \frac{1}{2} \langle |\mathbf{u}|^2 \rangle, \quad \mathcal{M} = \frac{1}{2} \langle |\mathbf{B}|^2 \rangle, \quad \mathcal{P} = \frac{1}{2} \left\langle \frac{|\theta|^2}{\mathcal{N}^2} \right\rangle. \quad (57)$$

For the kinetic energy equation, we obtain

$$\partial_t \mathcal{K} = -\langle \mathbf{u} \cdot (\mathbf{u} \cdot \nabla) \mathbf{U}_0 \rangle + \langle \theta \mathbf{u} \cdot \mathbf{e}_g \rangle + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{u} \rangle + \langle \mathbf{u} \cdot (\mathbf{B}_0 \cdot \nabla) \mathbf{B} \rangle. \quad (58)$$

Note that  $\langle \mathbf{u} \cdot (\mathbf{u} \cdot \nabla) \mathbf{u} \rangle = \langle \mathbf{u} \cdot (\mathbf{U}_0 \cdot \nabla) \mathbf{u} \rangle = \langle \mathbf{u} \cdot \nabla p \rangle = 0$  using the chain rule, incompressibility and the divergence theorem (applying periodic boundary conditions), and noting that the Coriolis force does no work ( $\mathbf{u} \cdot (2\boldsymbol{\Omega} \times \mathbf{u}) = 0$ ). We then substitute  $\mathbf{U}_0 = -Sx\mathbf{e}_y$  to yield

$$\partial_t \mathcal{K} = S \langle u_x u_y \rangle + \langle \theta \mathbf{u} \cdot \mathbf{e}_g \rangle + \langle \mathbf{u} \cdot (\mathbf{B}_0 \cdot \nabla) \mathbf{B} \rangle + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{u} \rangle. \quad (59)$$

This indicates that the kinetic energy of perturbations can grow by extracting kinetic energy from the shear flow/differential rotation (first term), from conversions of thermal to kinetic energy (second term), from conversions of magnetic to kinetic energy (third term),

and that it is dissipated by viscosity (fourth term, which can be shown to be negative definite through an integration by parts).

In a similar manner we can obtain the magnetic energy equation, noting that  $\langle \mathbf{B} \cdot (\mathbf{u} \cdot \nabla) \mathbf{B} \rangle = \langle \mathbf{B} \cdot (\mathbf{U}_0 \cdot \nabla) \mathbf{B} \rangle = 0$ , giving

$$\partial_t \mathcal{M} = \langle \mathbf{B} \cdot (\mathbf{B}_0 \cdot \nabla) \mathbf{u} \rangle - S \langle B_x B_y \rangle + \eta \langle \mathbf{B} \cdot \nabla^2 \mathbf{B} \rangle. \quad (60)$$

This indicates that magnetic energy of perturbations can grow from conversion of kinetic to magnetic energy (first term), from extracting kinetic energy from the background shear flow/differential rotation (second term), and that it is dissipated ohmically (third term; negative definite). Note that the term  $\langle \mathbf{B} \cdot (\mathbf{B}_0 \cdot \nabla) \mathbf{u} \rangle$  in equation (60) can be shown to be equivalent with  $-\langle \mathbf{u} \cdot (\mathbf{B}_0 \cdot \nabla) \mathbf{B} \rangle$  in equation (59) using integration by parts, which indicates that these just convert kinetic to magnetic energy and vice versa, and do not inject total energy into the system.

The final energy equation that we need to consider is the one governing thermal energy. We obtain

$$\partial_t \mathcal{P} = -\langle \theta \mathbf{u} \cdot \mathbf{e}_\theta \rangle + \kappa \left\langle \frac{\theta \nabla^2 \theta}{\mathcal{N}^2} \right\rangle, \quad (61)$$

using the results  $\langle \theta (\mathbf{u} \cdot \nabla) \theta \rangle = \langle \theta (\mathbf{U}_0 \cdot \nabla) \theta \rangle = 0$ . This shows that thermal energy grows through conversion from kinetic to thermal energy (first term) and that it is dissipated by thermal diffusion (second term; negative definite). Hence, the equation for the total energy,  $\mathcal{E} = \mathcal{K} + \mathcal{M} + \mathcal{P}$ , is

$$\begin{aligned} \partial_t \mathcal{E} = & S \langle (u_x u_y) - \langle B_x B_y \rangle \rangle + \langle \theta \mathbf{u} \cdot (\mathbf{e}_g - \mathbf{e}_\theta) \rangle \\ & + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{u} \rangle + \eta \langle \mathbf{B} \cdot \nabla^2 \mathbf{B} \rangle + \kappa \left\langle \frac{\theta \nabla^2 \theta}{\mathcal{N}^2} \right\rangle. \end{aligned} \quad (62)$$

This indicates that the total energy of perturbations can grow only via extraction of kinetic energy from the shear flow/differential rotation into perturbation kinetic or magnetic energies (first two terms), or via the baroclinic term that extracts potential energy from the basic state into kinetic and thermal energies (last term on first the line), if and only if these contributions exceed the sum of the viscous, ohmic, and thermal dissipations (terms on the bottom line).

We can use these results to analyse the energy sources contributing to the instabilities described by equation (17). To do this, we calculate each of the terms in these energy equations for a single axisymmetric Fourier mode with a wave vector  $\mathbf{k} = (k_x, 0, k_z)$ . This can be used to understand better both the driving forces of the instability and the momentum transporting properties of the instability. We first express  $\langle u_x u_y \rangle$  for a single mode with  $u_x = \Re [\hat{u}_x \exp(i\mathbf{k} \cdot \mathbf{x} + st)]$  and  $u_y = \Re [\hat{u}_y \exp(i\mathbf{k} \cdot \mathbf{x} + st)]$ . Using the properties of complex numbers,  $\Re(A)\Re(B) = \frac{1}{2}\Re(AB + AB^*)$ , where  $*$  denotes the complex conjugate, this can be written as

$$\begin{aligned} \langle u_x u_y \rangle = & \frac{1}{2} \Re (\hat{u}_x \hat{u}_y \exp(2st + 2i\mathbf{k} \cdot \mathbf{x}) + \exp(2\Re[s]t) \hat{u}_x \hat{u}_y^*) \\ = & \frac{1}{2} \exp(2\Re[s]t) \Re(\hat{u}_x \hat{u}_y^*), \end{aligned} \quad (63)$$

upon applying the periodic boundary conditions (thereby eliminating the first term on the top line).

For a single linear mode we can use equations (11)–(16) to relate  $\hat{u}_y$  to  $\hat{u}_x$  (and similarly for all other variables) to obtain

$$\hat{u}_y = \frac{s_\eta}{s_\nu s_\eta + \omega_A^2} \left( S \left( 1 + \frac{\omega_A^2}{s_\eta^2} \right) - 2\Omega \left( c_\Lambda + \frac{k_x}{k_z} s_\Lambda \right) \right) \hat{u}_x, \quad (64)$$

and we also note that  $\hat{u}_z = -(k_x/k_z)\hat{u}_x$ . Substituting this into equation (63), and using  $\Re[\hat{u}_x \hat{u}_y^*] = \Re[\hat{u}_x^* \hat{u}_y]$ , gives the concise form for

the  $xy$  component of the Reynolds stress:

$$\langle u_x u_y \rangle = \frac{|\hat{u}_x|^2}{2} \Re \left[ \frac{s_\eta \left( S \left( 1 + \frac{\omega_A^2}{s_\eta^2} \right) - 2\Omega \left( c_\Lambda + \frac{k_x}{k_z} s_\Lambda \right) \right)}{s_\nu s_\eta + \omega_A^2} \right]. \quad (65)$$

We also have

$$\begin{aligned} \hat{B}_y = & \frac{i\omega_A}{s_\eta} \left[ \hat{u}_y - \frac{S}{s_\eta} \hat{u}_x \right] \\ = & \frac{i\omega_A \hat{u}_x}{s_\eta} \left[ -\frac{S}{s_\eta} + \frac{S \left( 1 + \frac{\omega_A^2}{s_\eta^2} \right) - 2\Omega \left( c_\Lambda + s_\Lambda \frac{k_x}{k_z} \right)}{s_\nu + \frac{\omega_A^2}{s_\eta}} \right], \end{aligned} \quad (66)$$

along with  $\hat{B}_z = -(k_x/k_z)\hat{B}_x$ , which allows the  $xy$  component of the Maxwell stress for a single mode to be written as

$$\begin{aligned} \langle B_x B_y \rangle = & \frac{1}{2} \Re [\hat{B}_x \hat{B}_y^*] \exp(2\Re[s]t) \\ = & \Re \left[ \frac{\omega_A^2 |\hat{u}_x|^2}{2s_\eta^2} \left( -\frac{S}{s_\eta} + \frac{S \left( 1 + \frac{\omega_A^2}{s_\eta^2} \right) - 2\Omega \left( c_\Lambda + s_\Lambda \frac{k_x}{k_z} \right)}{s_\nu + \frac{\omega_A^2}{s_\eta}} \right) \right]. \end{aligned} \quad (67)$$

The third and final term that can inject energy into the system is

$$\langle \theta \mathbf{u} \cdot (\mathbf{e}_g - \mathbf{e}_\theta) \rangle = \langle \theta [u_x (c_\phi - c_\Gamma) + u_z (s_\phi - s_\Gamma)] \rangle, \quad (69)$$

which is essentially a measure of the extent that baroclinicity (i.e. non-coincidence of constant density and pressure surfaces) drives the instability. We use

$$\hat{\theta} = -\frac{\mathcal{N}^2}{s_\kappa} \left( c_\Gamma - \frac{k_x}{k_z} s_\Gamma \right) \hat{u}_x, \quad (70)$$

along with incompressibility to write

$$\begin{aligned} \langle \theta \mathbf{u} \cdot (\mathbf{e}_g - \mathbf{e}_\theta) \rangle = & \frac{1}{2} \Re \left[ \frac{\mathcal{N}^2}{s_\kappa} \left( \frac{k_x}{k_z} s_\Gamma - c_\Gamma \right) \left( (c_\phi - c_\Gamma) - \frac{k_x}{k_z} (s_\phi - s_\Gamma) \right) \right] |\hat{u}_x|^2. \end{aligned} \quad (71)$$

Note that this term vanishes when  $\mathbf{e}_g = \mathbf{e}_\theta$ , and is thus unimportant when  $s_\Lambda = 0$  (no differential rotation along the rotation axis), and it is small compared to the other terms in the strongly stratified limit for which  $\mathcal{N}^2 \gg 2\Omega s_\Lambda$ .

The kinetic energy is

$$\mathcal{K} = \frac{1}{2} \langle |\mathbf{u}|^2 \rangle = \frac{1}{4} [|\hat{u}_x|^2 + |\hat{u}_y|^2 + |\hat{u}_z|^2] \exp(2\Re[s]t), \quad (72)$$

which can be expressed in terms of  $|\hat{u}_x|^2$  using the above results. The magnetic energy is

$$\mathcal{M} = \frac{1}{2} \langle |\mathbf{B}|^2 \rangle = \frac{1}{4} [|\hat{B}_x|^2 + |\hat{B}_y|^2 + |\hat{B}_z|^2] \exp(2\Re[s]t), \quad (73)$$

which can also be expressed in terms of  $|\hat{u}_x|^2$  using the following expressions:

$$|\hat{B}_x|^2 = \frac{\omega_A^2}{|s_\eta|^2} |\hat{u}_x|^2, \quad (74)$$

$$|\hat{B}_y|^2 = -\frac{\omega_A^2}{|s_\eta|^2} \left| -\frac{S}{s_\eta} + \frac{S \left( 1 + \frac{\omega_A^2}{s_\eta^2} \right) - 2\Omega \left( c_\Lambda + s_\Lambda \frac{k_x}{k_z} \right)}{s_\nu + \frac{\omega_A^2}{s_\eta}} \right|^2 |\hat{u}_x|^2, \quad (75)$$

$$|\hat{B}_z|^2 = \frac{\omega_A^2}{|s_\eta|^2} \left( \frac{k_x}{k_z} \right)^2 |\hat{u}_x|^2. \quad (76)$$

The thermal energy for a single mode can be expressed as

$$\left\langle \frac{|\theta|^2}{2N^2} \right\rangle = \frac{N^2}{4|s_x|^2} \left( c_\Gamma - \frac{k_x}{k_z} s_\Gamma \right)^2 |\hat{u}_x|^2. \quad (77)$$

Using these expressions we can determine the energetic contributions to the growth rate for a single Fourier mode by noting that

$$2\Re[s] = \partial_t \ln \mathcal{E} = \frac{1}{\mathcal{E}} \partial_t \mathcal{E}, \quad (78)$$

where the right-hand side contains all six terms in equation (62) and is independent of the mode amplitude since  $|\hat{u}_x|^2$  cancels in both the numerator and denominator. This can also be used as a check of our codes by ensuring that the growth rate  $\Re[s]$  is predicted to close to machine precision by using the linear relations between the components that we have just derived. Once this has been confirmed we can compute the contribution of each of the first three possible driving terms on the right-hand side of equation (62) to the growth rate to determine whether a given instability is driven by Reynolds stresses, Maxwell stresses, or the baroclinic term.

## 6.2 Numerical analysis of linear mode energetics

### 6.2.1 Unstable modes energetics: variation with $B_0$

We present results from computing the contributions to the growth rate from the three source terms on the right-hand side of equation (62). In particular, we determine the contributions to the growth rate from the Reynolds stress, Maxwell stress, and baroclinic driving terms in equations (65), (68), and (71) as a visual tool to understand better the mechanisms driving the various instabilities, as well as the role of the magnetic field. Each of these are divided by  $2\mathcal{E}$  in order to compute their contribution to  $\sigma$  for the reason explained in equation (78). All of the figures in this section use our standard choice of parameters,  $\text{Pr} = 10^{-2}$ ,  $N^2 = 10$ , and  $S = 2$  unless stated otherwise.

Figs 13 and 14 show pseudo-colour plots for various  $\Lambda$  and  $\phi$  of the growth rate (first row) along with the contributions to it from Reynolds stresses (second row), Maxwell stresses (third row), and baroclinic source terms (fourth row) on the  $(k_x, k_z)$  plane, for various magnetic field strengths  $B_0 = 1, 2.5, 5$ . Rows two to four represent the first three terms in equation (62), the sum of these, together with the three diffusive terms (not plotted) in equation (62) has been verified to match the growth rate  $\sigma$  to machine precision. In contrast to Figs 3–5, they use a linear colour scale since the various contributions plotted can take either sign, as we observe in these figures. Overall, these figures allow us to explore how variations in field strength for ( $B_0 = 1, 2.5, 5$ ) and rotation profile (through  $\Lambda$  and  $\phi$ ) alter the instabilities whilst simultaneously probing which energy source terms are responsible.

In Fig. 13, we first analyse the configuration at the equator with mixed shear ( $\Lambda = -30^\circ$ ,  $\phi = 30^\circ$ ) explored earlier in Fig. 3. This configuration is GSF unstable in the hydrodynamic case and remains unstable for weak fields. Strong fields tend to inhibit instability for  $k \sim 1$  and to shrink the unstable lobes, in addition to changing their orientation. For  $B_0 \leq 2.5$ , Reynolds stresses are the primary drivers of instability for most  $(k_x, k_z)$ , indicating that unstable modes are primarily driven by extracting kinetic energy from the differential rotation. As  $B_0$  is increased further, Maxwell stresses play an increasingly important role, until they dominate for  $B_0 = 5$ , indicating

that shear flow kinetic energy is extracted and input into perturbation magnetic energy. The different locations of the peaks in Reynolds and Maxwell stresses – and the increasingly stabilizing effects (negative values shown) of Maxwell stresses where the Reynolds stresses are maximal – are consistent with the changes in orientations of the unstable lobes as  $B_0$  is increased, from initially being between  $\hat{\Omega}^\perp$  and  $\nabla\ell$  to become closer to  $\nabla\ell$  for the strongest fields. For this latitude and flow, baroclinic driving terms are typically subdominant, but they still contribute non-negligibly to driving instabilities for weaker fields. The effect of the field in reducing the maximum growth rate observed in Fig. 7 is also confirmed here.

We next look at a case with latitude  $30^\circ$  with mixed shear ( $\Lambda = 60^\circ$ ,  $\phi = -30^\circ$ ) in Fig. 14 as first studied in Fig. 4. This configuration is GSF unstable hydrodynamically and remains unstable for weak fields. We saw from Fig. 8 that the field acts to monotonically stabilize the system with increasing  $B_0$ , which is consistent with Fig. 14. We again observe that the primary lobes of instability are driven primarily by Reynolds stresses for  $B_0 \leq 2.5$ , but become increasingly driven by Maxwell stresses for stronger fields. We also observe the positive Reynolds stress contributions are mainly confined to within the hydrodynamically unstable wedge delineated by the lines  $\hat{\Omega}^\perp$  and  $\nabla\ell$ , and are maximal approximately halfway between these. The increasing importance of Maxwell stresses and the shift in orientation of the lobes indicates the transition in the dominant instability from GSF to MRI. Notice that the Maxwell stress generally has a preferred wave vector magnitude, evident by the darkest red (most unstable) modes being located in the centre of the lobes. We also observe the unstable region shrinking as the MRI enables instability for smaller and smaller  $k$  for appropriately oriented modes. The baroclinic term is unimportant for the primary lobes, as is indicated by the bottom panels.


The secondary lobes evident in Fig. 14 are hydrodynamically unstable oscillatory modes within the wedge defined by  $e_g$  and  $e_\theta$ . The bottom panels of this figure confirm that these modes are baroclinically driven since  $\sigma$  approximately equals its baroclinic contribution, with Reynolds and Maxwell stresses playing negligible roles in driving them. The growth rates and unstable mode wave vectors are mostly unaffected by the magnetic field, except that these become weakly destabilized magneto-inertial-gravity waves rather than inertia-gravity waves when the field is sufficiently strong.

We have found similar trends as  $B_0$  is varied are found for  $\Lambda$  and  $\phi$  that are adiabatically SH unstable in the hydrodynamic case, and for cases at the poles that are hydrodynamically adiabatically stable.


### 6.2.2 Fastest growing mode energetics: variation with $B_0$

We next turn to analyse how the energetic contributions vary with  $B_0$  for the fastest growing modes, obtained by optimizing over  $k_x$  and  $k_z$  for each case. Results are shown in Fig. 15 for various latitudes and differential rotations. We study both  $\text{Pm} = 1$  and  $\text{Pm} = 0.1$  in order to investigate the role magnetic diffusivity plays in these results.


Panel (a) of Fig. 15 analyses a case with  $\phi = 60^\circ$  and  $\Lambda = -30^\circ$  that is adiabatically SH unstable in the hydrodynamic case. We find the growth rate in this case is essentially independent of  $B_0$ , as predicted from Figs 6 and 8. The primary result of changing  $B_0$  is to decrease the range of unstable  $k_x$  and  $k_z$  as we have confirmed in Fig. 5. This case is driven by the Reynolds stress for all  $B_0$  considered, since the red symbols provide a larger contribution to the total growth rate, for both  $\text{Pm}$  plotted. Magnetic diffusion does not play an important role here, confirmed by the negligible role of  $\text{Pm}$ . The baroclinic driving term is the secondary contributor to instability for



**Figure 13.** Energetic contributions to instability on the  $(k_x, k_z)$  plane for  $\Lambda = -30^\circ$  and  $\phi = 30^\circ$ ,  $B_0 = 1, 2.5$ , and  $5$  (increasing in columns as we go from left to right) all with  $S = 2$ ,  $N^2 = 10$ ,  $\text{Pr} = 0.01$ , and  $\text{Pm} = 0.1$ . Top row: Growth rate. Second row: Reynolds stress contribution. Third row: Maxwell stress contribution. Fourth row: Baroclinic contribution. Stable modes with  $\Re[s] \leq 0$  are indicated in white for clarity.



**Figure 14.** Energetic contributions to instability on the  $(k_x, k_z)$  plane for  $\Lambda = 60^\circ$  and  $\phi = -30^\circ$ ,  $B_0 = 1, 2.5,$  and  $5$  (increasing in columns as we go from left to right) all with  $S = 2, N^2 = 10, \text{Pr} = 0.01,$  and  $\text{Pm} = 0.1$ . Top row: Growth rate. Second row: Reynolds stress contribution. Third row: Maxwell stress contribution. Fourth row: Baroclinic contribution. Stable modes with  $\Re[s] \leq 0$  are indicated in white for clarity.



**Figure 15.** Energetic contributions to instability for the fastest growing mode (optimized over  $k_x$  and  $k_z$ ) as a function of the magnetic field strength  $B_0$  for various  $\Lambda$  and  $\phi$  cases. These show the growth rate, and the contributions to it from Reynolds stresses ( $\langle u_x u_y \rangle$ ), Maxwell stresses ( $\langle b_x b_y \rangle$ ), and baroclinic source terms ( $\langle \theta u \cdot (e_y - e_\theta) \rangle$ ) against field strength  $B_0$ . All panels show  $\text{Pm} = 0.1$  and  $\text{Pm} = 1$ , and the other parameters are  $\text{Pr} = 10^{-2}$ ,  $\mathcal{N}^2 = 10$ , and  $\mathcal{S} = 2$ .

the smaller  $B_0$ , but it appears that it may be superseded for  $B_0 \gtrsim 25$  by Maxwell stresses. There is a jump from one unstable mode to another around  $B_0 = 12$  where baroclinic and Maxwell stress terms approximately balance, where the  $k$  and  $\theta_k$  values instantly switch.

Panel (b) of Fig. 15 analyses a case with cylindrical differential rotation ( $\Lambda = 0$ , which is neutrally hydrodynamically GSF stable for all latitudes with  $S = 2$ ). This case is significantly destabilized by even weak magnetic fields as we have seen in Figs 8, 9, and 10, due to the MRI. This instability is driven by an approximately equal combination of Reynolds and Maxwell stresses, which perfectly coincide for large  $B_0$ . This can be termed ‘Alfvénization’ of the instability for sufficiently strong fields. The complete lack of any baroclinic driving is evident in this figure, and is also observed in panel (f) which also has  $\Lambda = 0$ , as is expected for any cylindrical rotation profile.

Panel (c) probes instabilities at the poles by considering  $\phi = 60^\circ$  and  $\Lambda = 30^\circ$ . Figs 5 and 10 indicated that this latitude is widely unstable to adiabatic magnetic instabilities, in stark comparison to the hydrodynamic results (Paper III), which found no adiabatic instability there. This instability is again the MRI, and it is driven by an approximately equal balance of Reynolds and Maxwell stresses indicating Alfvénization once again. The growth rate increases by around 35 per cent between  $B_0 = 0$  and  $B_0 = 5$ , after which increases in  $B_0$  lead to only marginal increases in  $\sigma$ .

Panel (d) explores a shellular rotation profile with  $\phi = 0$  and  $\Lambda = 60^\circ$ . These cases were explored hydrodynamically in Paper II. This figure indicates that the instability is initially driven almost entirely by Reynolds stresses when  $B_0 \sim 0$ , but Maxwell stresses dominate for  $B_0 \gtrsim 5$ . The introduction of magnetic fields weakens the instability and reduces  $\sigma$  (after a small rise for  $B_0 \sim 1$ ) over the  $B_0 = 0$  case. A plateau is reached for  $\sigma$  by  $B_0 \gtrsim 15$ , where the instability is primarily driven by Maxwell stresses. Once again, the baroclinic driving term is very weak in this case for any  $B_0$ .

Panel (e) shows the behaviour of the fastest growing mode from the parameters of Fig. 14 with  $\phi = -30^\circ$  and  $\Lambda = 60^\circ$ . As  $B_0$  is increased  $\sigma$  is drastically reduced. Up to  $B_0 \approx 5$  for  $\text{Pm} = 0.1$ , and  $B_0 = 3$  for  $\text{Pm} = 1$ , Reynolds stresses are the dominant contributor to the growth rate, but as the growth rate decreases with increasing field strength Maxwell stresses become the dominant contributor with these lines converging towards each other. We may achieve Alfvénization again for sufficiently large  $B_0$ , but this is not observed by  $B_0 = 25$ . As was observed in Fig. 4 the  $\text{Pm} = 1$  case is consistently more unstable than the  $\text{Pm} = 0.1$  case, however, as the growth rate tends to zero this difference becomes marginal.

In this section, we have analysed the unstable mode energetics as  $B_0$ ,  $\text{Pm}$ , and the properties of the differential rotation were varied. We have found that the fastest growing modes are always driven predominantly by a combination of Reynolds and Maxwell stresses for non-zero  $B_0$  and that baroclinic driving is negligible except for the subdominant secondary lobes. For strong enough magnetic fields, in many cases in which the MRI operates, the contributions of Reynolds and Maxwell stresses equalize. Overall, these results confirm that even a weak magnetic field can drastically alter the stability of differentially rotating flows in stellar radiation zones.

## 7 APPLICATIONS TO THE SUN AND RED GIANT STARS

We now turn to estimate parameter values for the solar tachocline as a potential application of this work. Recall that we defined our

length-scale  $d$  as

$$d = \left( \frac{\nu\kappa}{\mathcal{N}^2} \right)^{1/4}, \quad (79)$$

since this describes the scales of the dominant hydrodynamic GSF modes. In the solar tachocline (e.g. Gough 2007; Caleo et al. 2016), we find  $\nu = 2.7 \times 10^1 \text{ cm}^2 \text{ s}^{-1}$ ,  $\kappa = 1.4 \times 10^7 \text{ cm}^2 \text{ s}^{-1}$ , hence  $\text{Pr} = 2 \times 10^{-6}$  and  $\mathcal{N} = 8 \times 10^{-4} \text{ s}^{-1}$ . This produces a length-scale  $d \approx 49.3 \text{ m}$ .<sup>3</sup> The linear GSF modes thus have very short length-scales approx  $10^{-5}$  times smaller than the tachocline thickness. The dimensional wavenumber  $k_{\text{dim}} = k/d$ , using our dimensionless wavenumber  $k$ . Note that  $\eta = 4.1 \times 10^2 \text{ cm}^2 \text{ s}^{-1}$  in the tachocline, so  $\text{Pm} = 0.065$  and  $\text{Pr}/\text{Pm} = 3 \times 10^{-5}$  there. Hence, we are in the regime of rapid thermal diffusion relative to viscous and ohmic diffusion in the tachocline, as we discussed in Sections 3.3 and 3.3.2.

The magnetic field strength and structure in the tachocline is highly uncertain. Nevertheless, any poloidal magnetic field is probably in the range 0.5 G to 5 kG (e.g. Mestel & Weiss 1987, and we are not aware of substantially stronger subsequent constraints). The field there is likely to be mostly toroidal, but only poloidal fields enter our stability analysis for axisymmetric modes. The arguments of Gough & McIntyre (1998) for the maintenance of the tachocline also suggest a minimum poloidal field of 1 G is required there.

Our dimensionless magnetic field  $\mathbf{B}$  is written in Alfvén speed units; therefore, it has units  $d\Omega$  where  $\Omega = 2\pi/P_{\text{rot}}$ , and  $P_{\text{rot}} = 27 \text{ d}$  is the Sun’s mean rotation period. The corresponding physical magnetic field magnitude  $B_{\text{dim}}$  from the dimensional Alfvén speed  $V_A = B_{\text{dim}}/\sqrt{\mu_0\rho}$ ,


$$B_{\text{dim}} = B_0 d\Omega\sqrt{\mu_0\rho} \approx 2.1 \times 10^{-6} B_0 \text{ T} \approx 0.021 B_0 \text{ G}, \quad (80)$$

using  $\Omega = 2.7 \times 10^{-6} \text{ s}^{-1}$  (implying  $\mathcal{N}^2/\Omega^2 \approx 8.7 \times 10^4$ ),  $\rho = 210 \text{ kg m}^{-3}$ , and  $\mu_0 = 4\pi \times 10^{-7}$  in SI units. This means that a field of 1 G corresponds to a dimensionless  $B_0 \approx 46$  in our units if  $d$  is the relevant length-scale. Note that  $d$  was defined based on the diffusive hydrodynamic GSF modes, and we have found the MRI to potentially have much larger wavelengths. However, the small length-scales which GSF modes prefer do make them very vulnerable to even a rather weak magnetic field.

On the other hand if we want to consider a field of 1 kG, this requires  $B_0 = 4.6 \times 10^4$  in dimensionless units, which is much larger than we have so far considered here. The fields we have primarily explored in this work are at the weaker end, with  $B_0 \lesssim 25$ , corresponding to fields weaker than approximately 0.5 G in the tachocline. This choice was partly made to permit us to explore the modification of hydrodynamic diffusive rotational instabilities by a weak field, and was partly made because we found that for larger  $B_0$  the GSF mode is primarily stabilized and the dominant instability by far is the MRI.

Motivated by the values in the solar tachocline, we compute the linear growth rates, wavenumbers, and orientations numerically and display them in Fig. 16 at a latitude of  $30^\circ$  for a moderately strong field with  $B_0 = 10^3$  for the solar-like values  $S = 0.2$ ,  $\text{Pm} = 0.05$ , and  $\text{Pr} = 10^{-6}$  (so that  $\text{Pr}/\text{Pm} = 2 \times 10^{-5}$ ). We consider three different values of  $N^2 \in [10, 10^3, 10^5]$  to account for the variation of values in the solar radiation zone as we approach the radiative–convective boundary. Deep down, the higher value is appropriate (and corresponds to the value in Gough 2007), but  $N^2$  passes through

<sup>3</sup>Please note the unfortunate typo in Paper I, where this was written as km instead! No other values in Paper I need modifying and slightly different numbers were used from stellar models for the various parameters there than the ones we quote here.



**Figure 16.** Properties of the fastest growing modes for tachocline parameter values at a latitude  $\beta = \Lambda + \phi = 30^\circ$ . We assume  $B_0 = 1000$ ,  $S = 0.2$ ,  $\text{Pm} = 0.05$ ,  $\text{Pr} = 10^{-6}$ , and explore  $N^2 \in [10, 10^3, 10^5]$ . Note the growth rate is on a log-scale, because for  $\phi > 0$  we get MRI that grows much faster than the GSF modes that may exist for  $\phi < 0$ .

zero as the convection zone is approached, motivating the smaller values we consider here (see e.g. fig. 1 in Barker & Ogilvie 2010). We observe that MRI operates for  $\phi > 0$  with modest growth rates  $\sigma \sim 0.1$  and  $k \sim 0.0005$  (corresponding to wavelengths of order 600–60 000 km, much smaller than the tachocline thickness), and that it is not affected by variations in  $N^2$ . This is because the unstable modes orient themselves to avoid doing work against gravity such that  $b \sim 0$ . On the other hand, the instabilities for  $\phi < 0$  are substantially modified by varying  $N^2$ . These instabilities are GSF modes inhibited by magnetic tension. For  $N^2 = 10^5$ , relevant for deeper parts of the solar tachocline and radiative interior, GSF is eliminated by the magnetic field. This agrees with some of the conclusions of Caleo & Balbus (2016) and Caleo et al. (2016). However, closer to the radiative/convective interface in the tachocline region itself, our smaller values of  $N^2$  are appropriate. For  $N^2 = 10$  and  $10^3$ , the GSF instability operates but with a much weaker growth rate than the MRI modes in operation when  $\phi > 0$ . Fig. 16 suggests that MRI may be more important than GSF for turbulent transport in the solar radiative interior whenever  $\phi > 0$ , but that more weakly growing GSF modes could operate for local rotation profiles with  $\phi < 0$ .

In the core of red giant stars, whose core-envelope differential rotations remain poorly understood, as considered in Paper I and using the numbers there,  $d \sim 100$  m and  $\Omega \sim 10^{-7} \text{ s}^{-1}$ . This produces  $B_{\text{dim}} \sim 1.12 B_0 \sqrt{\rho/10^5} \text{ g cm}^{-3} \text{ G}$ . Hence in that problem  $B_{\text{dim}} \sim B_0 \text{ G}$  in the cores of red giant stars. Since there have been constraints on fields in these from asteroseismology of order 40–610 kG (Deheuvels et al. 2023), this suggest we should consider  $B_0 \gtrsim 10^3$  in red giant stars also. Hence, MRI is expected to be more important than GSF, depending on the rotation profile (particularly for  $\phi > 0$ ), but perhaps not for  $\phi < 0$ .

## 8 CONCLUSIONS

We have presented a comprehensive theoretical analysis of local diffusive instabilities of differential rotation in magnetized radiation zones of stars and planets, building upon the hydrodynamical studies of Paper I, Paper II, and Paper III. Understanding the properties of these instabilities, and ultimately their non-linear behaviour, is essential because they have been proposed to play important roles in AM transport and chemical mixing in stars (e.g. Caleo et al. 2016; Aerts et al. 2019), and they may even play a role in the solar dynamo (Parfrey & Menou 2007; Vasil et al. 2024), but many aspects of them are currently very poorly understood. Our focus has been

on the effects of a poloidal magnetic field on the properties of linear axisymmetric instabilities of differential rotation, which are governed by a quintic dispersion relation first derived by Menou et al. (2004). We have performed a detailed analysis of the dispersion relation, first for non-diffusive instabilities, reproducing prior work on the stratified MRI (e.g. Balbus 1995), before comprehensively analysing diffusive instabilities in various limits analytically and numerically (see also Caleo & Balbus 2016; Caleo et al. 2016).

In strongly stably stratified regions of stars, the fastest growing mode displacements are along stratification (i.e. approximately spherical) surfaces and correspond with operation of the MRI. However, rapid thermal diffusion can eliminate the stabilizing effects of buoyancy if  $\text{Pr}/\text{Pm}$  and  $\text{Pr}$  are sufficiently small. In this limit MRI operates and can change the properties of the unstable modes depending on the differential rotation. We have obtained new analytical and numerical results on the various instabilities in this triply diffusive system as a function of the differential rotation profile and magnetic field strength.

Our analytical and numerical results have highlighted that even a weak magnetic field can considerably modify the local instabilities of differentially rotating flows (e.g. Balbus & Hawley 1998, and many prior works). We have found that for differential rotations with (angle from the local angular velocity gradient to the effective gravity direction)  $\phi > 0$  in the Northern hemisphere (and vice versa in the Southern hemisphere because the relevant parameter is the sign of  $\beta\phi$ ), MRI may dominate over the magnetic modification of hydrodynamic GSF instabilities. However, for  $\phi < 0$  there, hydrodynamic GSF modes could still be important though they are weakened by magnetic tension for moderately strong fields. We found that even weak fields destabilize hydrodynamically stable regions in parameter space, particularly for nearly cylindrical differential rotation profiles.

We have analysed in detail the properties of axisymmetric modes, including how the growth rates and wave vectors depend on the strength of the magnetic field, magnetic Prandtl number  $\text{Pm}$ , and local differential rotation profile. We have analysed in detail the energetics of the various instabilities in our system, first by deriving the energy equation and then by evaluating the various source terms for linear axisymmetric modes. These consist of Reynolds stresses, Maxwell stresses, and baroclinic driving terms. We find that the MRI is typically driven by Reynolds and Maxwell stresses in approximately equal proportions (so-called Alfvénization) in a wide range of cases.

We believe that it is important to set up a meaningful time-independent magnetic equilibrium to properly analyse MHD instabilities. We take a different viewpoint to many prior works that

attempted to model arbitrary field configurations without ensuring Ferraro's law of isorotation was satisfied (e.g. Balbus & Hawley 1994; Menou et al. 2004; Menou & Le Mer 2006; Parfrey & Menou 2007; Caleo et al. 2016). In our model, we ensured our basic state was an equilibrium state and verified the local analogue of Ferraro's law of isorotation. This is analogous to the original works of Goldreich & Schubert (1967) and Fricke (1968) having an additional degree of freedom because they ignored the constraint of thermal wind balance (e.g. Acheson & Gibbons 1978; Busse 1981). Similar issues have also plagued studies of the effects of magnetic fields on the vertical shear instability in astrophysical discs (e.g. Urpin & Brandenburg 1998; Latter & Papaloizou 2018, in which the latter authors take the same viewpoint as us).

Future work should explore the non-linear evolution of the instabilities we have analysed here (building upon Paper I; Paper II; Paper III; Tripathi et al. 2024), as well as the role of compositional gradients on both the linear (Knobloch & Spruit 1983) and non-linear properties of this problem. Further global simulations tailored to study these instabilities would also be worthwhile.

## ACKNOWLEDGEMENTS

RWD was supported by an STFC studentship (2443617). AJB was supported by STFC grants ST/S000275/1 and ST/W000873/1. SMT would like to acknowledge support of funding from the European Union Horizon 2020 Framework Programme (grant agreement no. D5SDLV-786780). Additionally, we would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programmes DYT2 and 'Anti-diffusive dynamics: from sub-cellular to astrophysical scales', supported by EPSRC grant no. EP/R014604/1. We would like to thank the referee for a constructive and timely report.

## DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

## REFERENCES

- Acheson D. J., Gibbons M. P., 1978, *Phil. Trans. R. Soc. A*, 289, 459  
 Aerts C., Mathis S., Rogers T. M., 2019, *ARA&A*, 57, 35  
 Balbus S. A., 1995, *ApJ*, 453, 380  
 Balbus S. A., 2009, *MNRAS*, 395, 2056  
 Balbus S. A., Hawley J. F., 1991, *ApJ*, 376, 214  
 Balbus S. A., Hawley J. F., 1994, *MNRAS*, 266, 769  
 Balbus S. A., Hawley J. F., 1998, *Rev. Mod. Phys.*, 70, 1  
 Barker A. J., Latter H. N., 2015, *MNRAS*, 450, 21  
 Barker A. J., Ogilvie G. I., 2010, *MNRAS*, 404, 1849  
 Barker A. J., Jones C. A., Tobias S. M., 2019, *MNRAS*, 487, 1777 (Paper I)  
 Barker A. J., Jones C. A., Tobias S. M., 2020, *MNRAS*, 495, 1468 (Paper II)  
 Busse F. H., 1981, *Geophys. Astrophys. Fluid Dyn.*, 17, 215  
 Caleo A., Balbus S. A., 2016, *MNRAS*, 457, 1711  
 Caleo A., Balbus S. A., Tognelli E., 2016, *MNRAS*, 460, 338

- Chandrasekhar S., 1961, *Hydrodynamic and Hydromagnetic Stability*. Dover Publications, New York  
 Deheuvels S., Li G., Ballot J., Lignières F., 2023, *A&A*, 670, L16  
 Dymott R. W., Barker A. J., Jones C. A., Tobias S. M., 2023, *MNRAS*, 524, 2857 (Paper III)  
 Ferraro V. C. A., 1937, *MNRAS*, 97, 458  
 Fricke K., 1968, *Z. Astrophys.*, 68, 317  
 Gaurat M., Jouve L., Lignières F., Gastine T., 2015, *A&A*, 580, A103  
 Goldreich P., Schubert G., 1967, *ApJ*, 150, 571  
 Gough D. O., 2007, in Hughes D. W., Rosner R., Weiss N. O., eds, *The Solar Tachocline*. Cambridge Univ. Press, Cambridge, p. 3  
 Gough D. O., McIntyre M. E., 1998, *Nature*, 394, 755  
 Guilet J., Müller E., 2015, *MNRAS*, 450, 2153  
 Guseva A., Willis A. P., Hollerbach R., Avila M., 2017, *ApJ*, 849, 92  
 Høiland E., 1941, *Avhandlinger Norske Videnskaps-Akademi i Oslo, I, math.-naturv. Klasse*, 11, 1  
 Hollerbach R., Rüdiger G., 2005, *Phys. Rev. Lett.*, 95, 124501  
 Jouve L., Gastine T., Lignières F., 2015, *A&A*, 575, A106  
 Jouve L., Lignières F., Gaurat M., 2020, *A&A*, 641, A13  
 Kirillov O. N., Stefani F., 2010, *ApJ*, 712, 52  
 Knobloch E., Spruit H. C., 1982, *A&A*, 113, 261  
 Knobloch E., Spruit H. C., 1983, *A&A*, 125, 59  
 Latter H. N., Papaloizou J., 2018, *MNRAS*, 474, 3110  
 Maeder A., 2009, *Physics, Formation and Evolution of Rotating Stars*. Springer-Verlag, Berlin  
 Maeder A., Meynet G., 2000, *ARA&A*, 38, 143  
 Mamatsashvili G., Stefani F., Hollerbach R., Rüdiger G., 2019, *Phys. Rev. Fluids*, 4, 103905  
 Meduri D. G., Lignières F., Jouve L., 2019, *Phys. Rev. E*, 100, 013110  
 Meduri D. G., Jouve L., Lignières F., 2024, *A&A*, 683, A12  
 Menou K., Le Mer J., 2006, *ApJ*, 650, 1208  
 Menou K., Balbus S. A., Spruit H. C., 2004, *ApJ*, 607, 564  
 Mestel L., Weiss N. O., 1987, *MNRAS*, 226, 123  
 Nelson R. P., Gressel O., Umurhan O. M., 2013, *MNRAS*, 435, 2610  
 Ogilvie G. I., 2007, in Hughes D. W., Rosner R., Weiss N. O., eds, *The Solar Tachocline*. Cambridge Univ. Press, Cambridge, p. 299  
 Ogilvie G. I., Pringle J. E., 1996, *MNRAS*, 279, 152  
 Oishi J. S., Vasil G. M., Baxter M., Swan A., Burns K. J., Lecoanet D., Brown B. P., 2020, *Proc. R. Soc. A*, 476, 20190622  
 Parfrey K. P., Menou K., 2007, *ApJ*, 667, L207  
 Park J., Prat V., Mathis S., 2020, *A&A*, 635, A133  
 Park J., Prat V., Mathis S., Bugnet L., 2021, *A&A*, 646, A64  
 Radko T., 2013, *Double-Diffusive Convection*. Cambridge Univ. Press, Cambridge  
 Rashid F. Q., Jones C. A., Tobias S. M., 2008, *A&A*, 488, 819  
 Solberg H., 1936, *Astrophysica Norvegica*, 1, 237  
 Spiegel E. A., Veronis G., 1960, *ApJ*, 131, 442  
 Spruit H. C., 1999, *A&A*, 349, 189  
 Tripathi B., Barker A. J., Fraser A. E., Terry P. W., Zweibel E. G., 2024, *ApJ*, 966, 195  
 Urpin V., Brandenburg A., 1998, *MNRAS*, 294, 399  
 Vasil G. M., Lecoanet D., Augustson K., Burns K. J., Oishi J. S., Brown B. P., Brummell N., Julien K., 2024, *Nature*, 629, 769  
 Zahn J.-P., 1974, in Ledoux P., Noels A., Rodgers A. W., eds, *Proc. IAU Symp. 59, Stellar Instability and Evolution*. Reidel, Dordrecht, p. 185

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.