



This is a repository copy of *Studying contact replays: investigating mechanisms, management and game exposures (SCRIMMAGE) for brain health in the Australasian National Rugby League: a protocol for a database design.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/219053/>

Version: Published Version

Article:

Gardner, A.J., Iverson, G.L., Bloomfield, P. et al. (21 more authors) (2024) Studying contact replays: investigating mechanisms, management and game exposures (SCRIMMAGE) for brain health in the Australasian National Rugby League: a protocol for a database design. *BMJ Open Sport & Exercise Medicine*, 10. e002216. ISSN 2055-7647

<https://doi.org/10.1136/bmjsem-2024-002216>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Studying Contact Replays: Investigating Mechanisms, Management and Game Exposures (SCRIMMAGE) for brain health in the Australasian National Rugby League: a protocol for a database design

Andrew J Gardner,^{1,2} Grant L Iverson,^{3,4,5,6} Paul Bloomfield,⁷ Sharron Flahive,⁸ James Brown,^{2,9,10} Suzi Edwards ,¹ Gordon W Fuller,¹¹ Mazdak Ghajari,¹² Prashant Jhala,¹ Ben Jones ,^{2,13,14,15,16} Christopher R Levi,¹⁷ Warren McDonald,¹⁸ Shreya McLeod,¹⁹ Cameron Owen,² Georgia Page,²⁰ Kenneth L Quarrie,^{21,22} Oliver Smith,¹ Peter Stanwell,¹ Daniel Tadmor,² Timana Tahu,¹ Douglas P Terry,^{23,24,25} Campbell Thomson,¹ Ross Tucker ,²⁶ Lauren V Fortington ²⁷

To cite: Gardner AJ, Iverson GL, Bloomfield P, *et al.* Studying Contact Replays: Investigating Mechanisms, Management and Game Exposures (SCRIMMAGE) for brain health in the Australasian National Rugby League: a protocol for a database design. *BMJ Open Sport & Exercise Medicine* 2024;**10**:e002216. doi:10.1136/bmjsem-2024-002216

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjsem-2024-002216>).

Accepted 13 September 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Andrew J Gardner;
andrew.gardner@sydney.edu.au

ABSTRACT

Concussions in contact sports are challenging for athletes, health professionals and sporting bodies to prevent, detect and manage. Design of interventions for primary prevention, early recognition of concussion and continuing to improve postconcussion management are essential for protecting athletes and promoting brain health. Over the last decade, there have been advancements in video technology for analysing head impact events and improvements in the clinical management of concussions. This study protocol describes how researchers, clinicians and staff from the Australasian National Rugby League (NRL) have brought these advancements together and developed a database of videos with head impact events and clinical outcomes. The intended outputs from this work will enhance the understanding of head impact events in NRL, from biomechanical and gameplay factors to concussion and return to play outcomes. Publishing this protocol increases the transparency of this large-scale effort to better identify head impacts and their relationship to concussions and player movement behaviour to contextualise these variables to generate new knowledge and support the reproducibility of these emerging findings. Between 2017 and 2023, over 5250 head contact cases were recorded in the database, from which >1700 head injury assessments were performed, and >600 concussions were diagnosed. Future studies using these data are planned to inform both primary and secondary injury prevention initiatives, such as risk analysis and prediction of game scenarios that result in concussion, as well as investigation of features and factors that help to inform the duration of recovery and return to play.

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Initiating the medical evaluation of a potential concussion during sports events occurs when there is suspicion of a possible injury.
- ⇒ Clinical suspicion may arise from observing a substantial head impact, observable concussive signs and/or the disclosure of symptoms by athletes.
- ⇒ This is difficult in a fast-paced contact sport (ie, professional rugby league match) because neurological signs may not be witnessed, and/or the athlete may minimise symptoms.
- ⇒ Video-supported identification of head impact events is now widely used on the sideline of professional rugby league games to facilitate the immediate recognition of a potential concussion.
- ⇒ The quick removal of an athlete suspected of having sustained a concussion from play and the initiation of a medical evaluation is designed to promote clinical recovery and encourage a safe and gradual return to play.

WHAT THIS STUDY ADDS

- ⇒ This protocol describes the development of a research database comprising a 7-year series of head impact event videos with linked findings of subsequent head injury assessments and clinical outcomes for athletes in the National Rugby League.
- ⇒ A future research programme to advance the understanding of sport-related concussion is outlined.

INTRODUCTION

A sport-related concussion is a mild traumatic brain injury that results in clinical signs and symptoms that can vary from person

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Data-driven insights from this research programme are intended to support new knowledge and robust decision-making related to the primary prevention, timely identification and medical management of concussions.

to person.¹ In the absence of objective tools and tests, both the identification and diagnosis of concussion are challenging. Standardised tests that include subjective reporting of symptoms and objective measures of neurological functioning, such as those in the Sport Concussion Assessment Tool fifth Edition (SCAT5),² assist health-care professionals in the diagnostic process and support clinical decisions. The initiation of a medical evaluation for concussion, such as a head injury assessment (HIA), largely relies on the observation of signs in an athlete that are consistent with a potential traumatic brain injury (eg, following an impact, the player is unresponsive, slow to get up or ataxic) and/or athlete disclosure of symptoms (eg, describing symptoms to the medical team or sports trainer). The reliance on player disclosure or observation by officials and/or health professionals to initiate the medical evaluation process can lead to missed cases during games, particularly because of the fast pace of matches and deliberate minimisation of symptoms by athletes who hope to continue playing.^{3,4} Video analysis has emerged as a useful adjunct to support the identification of potentially concussive head impacts because it enables the replay of incidents from different angles and at various playback speeds (eg, slow motion).^{5,6}

Video analysis has been used to better understand head impact events for over two decades. For example, in 2000, video playback was used to identify the contact site and calculate the kinematic data of the impact of medically confirmed concussion cases in professional rugby league, rugby union and Australian football.⁷ This research was an early example of using match videos to analyse head injuries. Still, the study was limited by its input data (ie, video sampled at 25 Hz) and perspective error of two-dimensional positional video data. Since that study was published, the potential to use video analysis in research to inform policy and practice has advanced substantially.

Previous rugby league research has linked signs and symptoms of concussion to video-identified on-field head impact events, demonstrating that valuable biomechanical, contextual and medical insights can emerge from studies involving video review.^{8–10} Earlier rugby league research was focused on specific, investigator-initiated questions or was limited in sample size: our research programme's initial work published in 2015 only sampled three clubs from a single NRL season (2013),⁸ while work published in 2021 used data from two NRL seasons (2017–2018).¹¹

To date, the video-supported identification of head impact events in the NRL and the subsequent clinical decisions and outcomes has been made on a case-by-case

basis for players at the time of their injury to inform their clinical management. This study protocol describes creating a central database for head impact events, HIAs and clinical outcomes. Publishing this protocol increases transparency related to this large-scale effort to better identify head impacts, concussions and their contextual variables surrounding these gameplay situations to generate new knowledge and support the reproducibility of these emerging findings. Study registration and prespecification of a protocol are also recommended by consensus research recommendations (eg, Strengthening the Reporting of OBservational Studies in Epidemiology),¹² and research and funding agencies (eg, National Institutes of Health). This protocol also allows other research teams to align data collection and coding of information to facilitate future data comparison across settings.

METHODS

The described database was created from seven NRL Premiership seasons, 2017 through 2023 inclusive. Data collection concluded at the end of the 2023 season. Coding of the variables from video footage is in progress at the time of publication.

Patient and public involvement

This protocol was developed with NRL medical and performance staff, representatives from the NRL Research Committee and NRL players. Together with an international multidisciplinary research team, these stakeholders have driven the study goals and data collection to ensure the research leads to insights that can improve player safety in the sport.

In accordance with the NRL and Rugby League Players Association Collective Bargaining Agreement, all athletes consented a priori to Rugby League Research Committee endorsed research using deidentified injury data. For this study, professional player contracts vary, and the players' Collective Bargaining Agreement has changed over time, but these factors have not impacted the design or conduct of this research.¹³

Setting

In Australia, the National Rugby League (NRL) Premiership is the highest rugby league club competition for men. The number of teams and matches has varied since 2017 (first inclusion year for this study; online supplemental table 1). In 2023, 17 teams (16 from Australia and 1 from New Zealand) competed across 27 rounds, with all teams having three 'byes' (non-playing weeks).

Each match has 2 teams, each with 13 players on the field and 4 regular substitute players on the interchange bench. An interchange can take place at any time during a match. From the 2016 season to the present (2024), each team can use up to eight interchanges per game. In the fifth year of the study (2021), an additional medical substitute ('18th player replacement') was introduced that can be activated when 2 players in the same team

are unable to continue in the same game following a concussion and/or game-limiting physical injury due to foul play from the opposing team (where an opponent is either permanently removed from the game or sent to the 'sin bin', a 10 min exclusion from the game where the player must leave the field of play and sideline, usually the dressing room).

Three rule changes have been introduced in the NRL during the observation seasons that are relevant to the occurrence of head impact events, HIAs and concussions. First, in the mid-season of 2020, a new rule for ruck infringements was introduced with 'six again' instead of a penalty. The six again rule restarts the tackle count for the attacking team and was intended to increase the speed of play without the need for game stoppage for a penalty. Second, since 2022, the off-site Injury Surveillance Bunker has included an Injury Surveillance Medical Officer (medical doctor independent of teams), supported by two video spotters who review play in real-time to identify potential head impact events. The Injury Surveillance Bunker Medical Officer will notify a specialist camera operator who has radio contact with the referee to stop the game if they believe the player has demonstrated signs of concussion. Finally, shoulder charge definitions and rulings have varied since first being banned in 2012. A shoulder charge, in which a defending player contacts/bumps an opposition player with the upper arm/shoulder while not attempting to tackle or steal the ball, is considered high-risk for injury and was outlawed from the game.

The in-game medical evaluation of a player suspected of having sustained a concussion referred to as the HIA in the Australian NRL, enables the removal of a player from the match for up to 15 min without the replacement player counting towards the team's limited interchanges. During the HIA, the club doctor conducts a formal medical assessment for concussion using standardised testing (ie, SCAT). In the NRL, an HIA may be initiated when: (i) a player shows signs or discloses symptoms of a concussion; (ii) it is deemed necessary by the club medical doctor and club sports trainer following a brief on-field assessment or (iii) it is recommended by an independent medical doctor with access to multiple camera angles to review any incident from an off-site medical bunker. The independent doctor will contact the team doctor to initiate an HIA if a concerning head impact event is identified by video review. The club sports trainer is responsible for signalling to the referee that a player is being removed from play for an HIA by patting the top of their head with one hand. Video playback of gameplay is available on the sideline at each NRL and NRL Women's Premiership (NRLW) game to support the team doctor by allowing immediate review of gameplay, mechanism of head impact and the postimpact event behaviour of the athlete.

Case definitions

Head impact event

A 'head impact event', defined as direct or indirect contact resulting in sudden contact or change of direction to the head, was the primary criterion for including cases. The operator of the sideline injury surveillance system identifies head impact events. All identified head impact events are included in the database. Not all head impact events require an HIA; medical assessment is only initiated when there is cause for further assessment and follow-up. The recording of a head impact event has evolved over time. Initially (ie, during the 2017–2020 seasons), the recording of head impact events came from one of two sources: (i) *sideline surveillance*—operators of team surveillance system or club medical staff can tag a player who had an impact to the head but was not removed from play for HIA or (ii) the *injury surveillance bunker*—the independent medical doctor in the bunker can tag a player who had an impact to the head but was not removed from play for HIA. From the 2021 season, all surveillance was centralised in the *injury surveillance bunker*. That is, the sideline was only equipped with a monitor, and all operations were managed via the injury surveillance bunker. In the injury surveillance bunker, the spotter tags events that are reviewed by the independent doctor and communication, if required, is made between the injury surveillance bunker and the team medical staff (figure 1).

Head injury assessment

Includes all athletes removed from play under the NRL HIA protocol by the team or independent medical doctor for an in-game medical evaluation.

Concussion

Includes all athletes who enter the NRL concussion management protocols. (ie, cases that were either diagnosed in-game or postgame).

Inclusion criteria and sample size

The video data collected as part of this research programme is a census sample of the identified head impact events, the HIAs and the medically diagnosed concussions from a game. The HIA results and the return-to-play information are available for most of the overall sample.

Data acquisition and storage

Over the course of the research programme, the video data provision process has changed. Initially (ie, 2017–2019), notification of any head impact events and the time at which they occurred was provided by email to the lead researcher by the Chief Medical Officer of the NRL. Access to video footage of the head impact event was obtained through the NRL's Digital Press Pass subscription.

Subsequently (2020–2023), in-game email alerts for each head impact event were provided to the lead researcher. Video clips of the events were available for




Figure 1 Visualisation of injury surveillance bunker (ISB) and its relationship with the sideline injury surveillance (SIS) and independent bunker doctor (diagram reproduced from an internal document, shared with permission from National Rugby League).

review through NRL stored systems. The lead researcher accessed and downloaded the relevant video clips, securely storing them on university-managed hardware and software, with additional encryption protection, and backing them up to secure online platforms accessible only to authorised research team members.

Data processing

Once the case videos were obtained, the coding of player, match, tackle and other event characteristics is completed. The variables of interest are presented in online supplemental tables 2–4, with a description and rationale for their inclusion provided below. Each case is coded by at least one of four possible reviewers. At least two reviewers code video signs of concussion to calculate inter-rater reliability. Each case requires approximately 5 min to review.

Player and game characteristics

The collection and analysis of data on player position and role allowed for a better understanding of patterns related to those who experience head impacts, sustain injuries and the phases of play involved. The player and game variables and options for selection are presented in online supplemental table 2. The incidence of concussion varies by playing position, with forwards tending to have a lower risk of concussion than backs.¹⁴ There is evidence of a difference in concussion risk for players in an offensive or defensive phase of play, although the direction (who is at greater risk) is inconsistent.^{9 15} Characteristics of play, such as time elapsed, may be studied in the future for insight into events and timing of higher risk. Research in rugby unions suggests that different injuries and risks are experienced at various periods in a match.^{16–18} To enable exploration of the timing of head

impact events, the time on the match clock for each case was reported and categorised.

Event and tackle features

Examining the on-field positioning and the first point of physical contact characteristics during head impact events may help to identify potential prevention strategies. To this end, a range of precontact, at-contact and postcontact variables are coded. The intention is to analyse the movement and positions of both the tackler and ball carrier to determine if there are common patterns or movement behaviours across different scenarios that led to a head impact event. The variables and potential responses included in the database are presented in online supplemental table 3. These were selected and refined from several studies in rugby league and rugby union that have investigated tackle technique, head injury, and video assessment.^{9–11 19–22}

Signs and symptoms of concussive injury

On-field signs of a potential concussion were recorded (online supplemental table 4), in alignment with the items outlined by the international consensus definitions of video signs of concussion in professional sports,²³ and a study of concussion signs specific to NRL players.²⁴ The symptoms the player experienced during the HIA were collected via self-report using SCAT5. The SCAT5 is relevant for this study, as testing was conducted from 2017 to 2023; SCAT6 was published in 2023.²⁵

Clinical reports (HIA and concussion outcomes)

All HIAs are performed by club medical staff. The results of the in-game HIA are retrospectively added to the database at the conclusion of each season. This ensures that coding of player and game characteristics, event and

tackle features and the video signs of potential concussive injury are independent of (ie, blind to) the eventual clinical outcomes. The administration and storage of the HIA data was provided via a smartphone application from CSx (2017–2020) and subsequently by HitIQ (2021–2023).

PLANNED APPLICATIONS

The early and accurate identification of concussions, along with data-informed primary and secondary prevention strategies, supports an evidence-based approach to managing the care of rugby league athletes and promoting brain health. Stakeholders within the NRL, including players, teams, administrators, medical staff and researchers, have supported the information collection detailed in this study protocol. We have provided a descriptive overview of the video footage and clinical outcomes collected and compiled over 7 years. The subsequent processing and preparation of the database will facilitate multiple studies relying on data from >5250 head impact events, including >1700 HIAs and >600 concussions.

Screening efficiency

The intention of the procedural changes introduced by the NRL, and subsequently captured in this database, was to improve concussion recognition and management of player welfare. With increased knowledge about player, game and tackle characteristics that result in clinical outcomes (eg, diagnosed concussion), there is potential to reduce ‘false alarms’ (ie, removing players from the game for HIA when it is not indicated). The occurrence of a head impact event can be likened to a medical screening test used to identify those players who need to undergo a diagnostic test (ie, an HIA). For certain health conditions and in many healthcare settings, diagnostic testing of everyone is considered inefficient and costly, so screening is used as a preliminary step to identify those with an elevated risk of a condition, which in turn justifies the conduct of a diagnostic test (often more invasive and costly). Therefore, one of the major aims of the Studying Contact Replays: Investigating Mechanisms, Management and Game Exposures in the Australasian National Rugby League (SCRIMAGE) study is to evaluate and improve screening efficiency. This will be realised through studies identifying if there are common characteristics or behaviours of the player(s), game and/or tackle, or a particular combination of factors consistently associated with specific outcomes (eg, cleared to return to play or confirmed diagnosis).

Observable signs

While video review can assist in identifying potential concussive events, previous studies have shown there are still challenges in discerning signs of head impact from observation. For example, in a recent study involving a video review of 25 medically diagnosed concussions from the Queensland Rugby League, none of the six international consensus video signs of possible concussion were

seen in 28% of these cases.²⁶ Similarly, 53% of concussions in the NHL did not have any video signs of concussion.²⁷ These prior findings were from databases with relatively small case numbers (<50 and <1000), and it is envisaged our larger number of cases will allow opportunities for diverse studies.

Concussion incidence and changes in gameplay

Another major aim of SCRIMAGE is to evaluate if there have been changes in the incidence or clinical features of concussion over the last 7 years. We will analyse the number of cases by different units of time (cases per number of players and cases per hours of play), and for different categories (eg, by player role as ball carrier or tackler). Alongside these studies, we will explore if there have been changes in gameplay characteristics (eg, types of tackles, number of players involved in tackles, speed of the game) that might have influenced outcomes (eg, increased severity of concussion or provided protection against reduced incidence).

Inter-rater and intra-rater reliability

Due to the volume of data, complete coding (100% of variables) for most cases has only been performed by one reviewer. At least two reviewers have coded all video signs of potential concussion for all cases. Methodologically focused research will be undertaken to continue refining the methods and processes of coding to maximise data reliability. For example, investigating the qualifications and training required for video review and coding is important. High inter-rater reliability was reported in an earlier study between expert reviewers who both had at least one season working at an NRL club with responsibility for concussion management (including identifying and assessing athletes with suspected concussions).²⁸ These expert reviewers were compared with those of ‘naïve’ reviewers with limited understanding of rugby league and no experience in concussion management. Low agreement between ‘naïve’ reviewers was reported.²⁸

Novel analyses

Alongside traditional methods of disease frequency and comparisons (ie, reported as incidence, prevalence and relative risks; reliability testing), novel approaches to these data will be explored. We intend to use machine learning (a subtype of artificial intelligence) to identify patterns and insights from the data. One example is to explore the three-dimensional kinematics of players’ movement before and during contact through pose estimate (eg, model-based image matching).^{29 30} In addition, computational modelling will be used to estimate brain strain and its association with clinical outcomes.³¹

Strengths and limitations

This paper describes the construction of a head impact event database that includes >80 variables from

6 years of professional rugby league competition. All cases have multiple video angles with high-definition footage provided from up to 10 or more cameras, and a subset of these have medical data gathered from players who undergo subsequent diagnostic screens. The study is focused only on professional NRL players, so findings of these cases will not be immediately generalisable to other settings, such as community rugby league. Separate work is underway for the NRLW Premiership.

Training injuries are not currently included. Recent data from men's teams in the European Super League suggest concussions can occur in training but are less frequent compared with match play.³² This may be an accurate assumption based on the scheduled training session (ie, the drills) and the level of intensity being less than match play. However, injury surveillance and medical support are often not as thorough during training, and some injuries may go unrecognised unless the athlete self-reports symptoms. Potentially, HIA or concussion outcomes from training could be included in the database. However, video data from training is unavailable and thus could not be included. Finally, we do not have records of players' data from their junior years, lower grades or any code-switches, as head impacts are limited to NRL career matches only.

CONCLUSION

Continuing to develop an understanding of the risk factors and mechanisms contributing to concussions in athletes is critical for reducing these injuries during practices and competitions. This multidisciplinary collaborative research programme identifies key variables and data sources to develop a large dataset linking player, game and tackle features with head impact events and associated clinical variables. Planned studies will inform both primary and secondary prevention efforts in the NRL.

Author affiliations

¹Sydney School of Health Sciences, Faculty of Medicine and Health, University of Sydney, Camperdown, New South Wales, Australia

²Carnegie Applied Rugby Research (CARR) Centre, Carnegie School of Sport, Leeds Beckett University, Leeds, UK

³Department of Physical Medicine and Rehabilitation, Harvard Medical School, Boston, Massachusetts, USA

⁴Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Charlestown, Massachusetts, USA

⁵Department of Physical Medicine and Rehabilitation, Schoen Adams Research Institute at Spaulding Rehabilitation, Charlestown, Massachusetts, USA

⁶Mass General for Children Sports Concussion Program, Boston, Massachusetts, USA

⁷Manly Sea Eagles Rugby League Club, Brookvale, New South Wales, Australia

⁸National Rugby League, Sydney, New South Wales, Australia

⁹The Division of Sport and Exercise Medicine (DiSEM), Department of Exercise, Sport and Lifestyle Medicine, Faculty of Medicine and Health Sciences, Stellenbosch University, Stellenbosch, South Africa

¹⁰Division of Physiological Sciences, Department of Human Biology, Faculty of Health Sciences, the University of Cape Town and the Sports Science Institute of South Africa, Cape Town, South Africa

¹¹School of Health and Related Research, The University of Sheffield, Sheffield, UK

¹²HEAD Lab, Dyson School of Design Engineering, Imperial College London, London, UK

¹³England Performance Unit, Rugby Football League, Red Hall, Manchester, UK

¹⁴Division of Physiological Sciences and Health through Physical Activity, Lifestyle and Sport Research Centre, Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

¹⁵Premiership Rugby, London, UK

¹⁶School of Behavioural and Health Sciences, Faculty of Health Sciences, Australian Catholic University, Brisbane, Queensland, Australia

¹⁷Hunter New England Local Health District, New Lambton Heights, New South Wales, Australia

¹⁸Rugby Australia, Moore Park, New South Wales, Australia

¹⁹Discipline of Physiotherapy, School of Allied Health, Faculty of Health Sciences, Australian Catholic University, Brisbane, Queensland, Australia

²⁰School of Medicine and Public Health, The University of Newcastle, Callaghan, New South Wales, Australia

²¹Sports Performance Research Institute New Zealand (SPRINZ), Auckland University of Technology, Auckland, New Zealand

²²Auckland Bioengineering Institute (ABI), The University of Auckland, Auckland, New Zealand

²³Vanderbilt University Medical Center, Nashville, Tennessee, USA

²⁴Vanderbilt Sports Concussion Center, Nashville, Tennessee, USA

²⁵Center for Cognitive Neurosurgical Studies, Nashville, Tennessee, USA

²⁶Institute of Sport and Exercise Medicine, Department of Sport Science, University of Stellenbosch, Stellenbosch, South Africa

²⁷Edith Cowan University, Joondalup, Western Australia, Australia

X Suzi Edwards @Suzi_Edwards, Ben Jones @23benjones and Ross Tucker @Scienceofsport

Contributors AG and LVF drafted the manuscript together with input from all authors throughout study design and drafting. All authors have read and approved the final manuscript. AG is the study guarantor.

Funding This study was funded by National Health and Medical Research Council (Early Career Researcher Fellowship (AJG), Investigator Grant (AJG)).

Competing interests AG has a clinical practice in neuropsychology involving individuals who have sustained sport-related concussion (including current and former athletes). He has been a contracted concussion consultant to Rugby Australia. He is a member of the World Rugby Concussion Working Group, and a member of the Australian Football League Concussion Scientific Advisory Committee. He has received travel funding or been reimbursed by professional sporting bodies, and commercial organisations for discussing or presenting sport-related concussion research at meetings, scientific conferences, workshops and symposiums. Previous grant funding includes the NSW Sporting Injuries Committee, the Brain Foundation (Australia), an Australian-American Fulbright Commission Postdoctoral Award, a Hunter New England Local Health District, Research, Innovation and Partnerships Health Research & Translation Centre and Clinical Research Fellowship Scheme and the Hunter Medical Research Institute (HMRI), supported by Jennie Thomas, and the HMRI, supported by Anne Greaves. He is supported by a National Health and Medical Research Council (NHMRC) Investigator Grant. He acknowledges unrestricted philanthropic support from the Tooth Foundation, and the National Rugby League (NRL) for research in retired professional rugby league players. GI has been reimbursed by the government, professional scientific bodies and commercial organisations for discussing or presenting research related to mild traumatic brain injury and sport-related concussion at meetings, scientific conferences and symposiums. He has a clinical and consulting practice in forensic neuropsychology, including expert testimony, involving individuals who have sustained mild TBIs (including athletes). He has received research funding from several test publishing companies, including ImPACT Applications, CNS Vital Signs and Psychological Assessment Resources (PAR). He has received research funding from the National Football League. He has also received research funding from the Harvard Integrated Programme to Protect and Improve the Health of National Football League Players Association Members. He acknowledges unrestricted philanthropic support from ImPACT Applications, the Mooney-Reed Charitable Foundation, NRL, Boston Bolts, Heinz Family Foundation and Schoen Adams Research Institute at Spaulding Rehabilitation. PB was the past NRL chief medical officer (2015–2021) and initiated and developed the NRL injury surveillance bunker and NRL concussion rules. He is a past club medical officer (2000–2013) and current club medical officer with the Manly Sea Eagles NRL

club. He also provides medical consultancy support to the Rugby League Players Association. SF is the NRL chief medical officer. JB has received research funding from World Rugby, the international governing body of rugby union. In addition, JB has received salary contributions and the coverage of travel costs from World Rugby. SE is an accredited sport scientist (AsPS2) and leads a tackle re-education programme that is partially funded by a NHMRC Ideas 2021 grant (202718). GF has previously received funding from World Rugby to attend research meetings and was a previous member of the Concussion in Sport expert panel. MG has received a Sports and Wellbeing Analytics/Cellbond Impact Solutions/Royal Academy of Engineering Senior Research Fellowship. PJ has no disclosures/conflicts of interest. BJ has secured research funding from World Rugby, Rugby Football Union, Scottish Rugby, Premiership Rugby, Rugby Football League, Catapult Sports, Prevent Biometrics, HitIQ, Leeds Rhinos Rugby League, Yorkshire Carnegie Rugby Union, Bath Rugby, Wasps Rugby. BJ is employed in a consultancy capacity by the Rugby Football League, and Premiership Rugby as their research lead. CRL serves as an honorary consultant neurologist to the NRL, providing a pro bono second opinion to current players regarding concussion assessment and management for the player's club doctor. WMcD is Rugby Australia's chief medical officer. SMcL has no disclosures/conflicts of interest. CO has a research fellowship that is part-funded by the Rugby Football League. GP has no disclosures/conflicts of interest. KLQ has been employed by New Zealand Rugby since 2000 and currently occupies the role of Chief Scientist, New Zealand Rugby. He was a member of World Rugby's Scientific Committee from 2013 to 2024 and has contributed to various World Rugby working groups focussed on player welfare issues from 2011 to the time of publication. He has received funding for travel and accommodation to attend World Rugby's medical meetings. OS and PS have no disclosures/conflicts of interest. DT has a PhD research programme that is part-funded by Leeds Rhinos rugby club. He also works as a medical doctor for multiple sports teams in the UK. TT is employed as the Senior Manager for Indigenous elite pathways at the NRL. He is also employed as a researcher by the University of Sydney through a 2021 NHMRC Ideas grant. DPT serves as a scientific advisor for HitIQ. He previously consulted for REACT Neuro. He has a consulting practice in forensic neuropsychology, including expert testimony, involving individuals who have sustained mild TBIs (including former athletes). He receives funding from Amgen and Football Research. CT has no disclosures/conflicts of interest. RT is a consultant to World Rugby. World Rugby is the governing body for the sport of Rugby Union and responsible for law changes in the sport. LVF is supported through research grant funding at Edith Cowan University. She has previously received project funding from several sports, health and government agencies including: Australian Football League; Australian Institute of Health and Welfare; Australian Institute of Sport; Combat Sport Commission Western Australia; Cricket Australia; Defence Science WA; Exercise & Sports Science Australia; Injury Matters; IOC; KidSafe WA; Rugby Australia; State Government of Victoria; VicSport.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the 'Methods' section for further details.

Patient consent for publication Not applicable.

Ethics approval The research design was approved by the University of Newcastle Human Ethics Committee (H-2012-0344). This work was conducted in accordance with the Declaration of Helsinki and the National Statement on the ethical conduct of human research and in line with national privacy laws. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data sharing not applicable as no datasets generated and/or analysed for this study. This article provides a description of the setup of a research database. No research data are currently available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Suzi Edwards <http://orcid.org/0000-0002-5790-0232>
 Ben Jones <http://orcid.org/0000-0002-4274-6236>
 Ross Tucker <http://orcid.org/0000-0002-4294-4822>
 Lauren V Fortington <http://orcid.org/0000-0003-2760-9249>

REFERENCES

- Silverberg ND, Iverson GL, ACRM Brain Injury Special Interest Group Mild TBI Task Force members, *et al*. The American Congress of Rehabilitation Medicine Diagnostic Criteria for Mild Traumatic Brain Injury. *Arch Phys Med Rehabil* 2023;104:1343–55.
- Echemendia RJ, Meeuwisse W, McCrory P, *et al*. The Sport Concussion Assessment Tool 5th Edition (SCAT5): Background and rationale. *Br J Sports Med* 2017;51:848–50.
- Longworth T, McDonald A, Cunningham C, *et al*. Do rugby league players under-report concussion symptoms? A cross-sectional study of elite teams based in Australia. *BMJ Open Sport Exerc Med* 2021;7:e000860.
- Tadmor D, Till K, Phillips G, *et al*. I won't let you down; why 20% of Men's and Women's Super League players underreported suspected concussions. *J Sci Med Sport* 2023;26:688–93.
- Della Villa F, Buckthorpe M, Grassi A, *et al*. Systematic video analysis of ACL injuries in professional male football (soccer): injury mechanisms, situational patterns and biomechanics study on 134 consecutive cases. *Br J Sports Med* 2020;54:1423–32.
- Waldén M, Krosshaug T, Bjørneboe J, *et al*. Three distinct mechanisms predominate in non-contact anterior cruciate ligament injuries in male professional football players: a systematic video analysis of 39 cases. *Br J Sports Med* 2015;49:1452–60.
- McIntosh AS, McCrory P, Comerford J. The dynamics of concussive head impacts in rugby and Australian rules football. *Med Sci Sports Exerc* 2000;32:1980–4.
- Gardner AJ, Iverson GL, Quinn TN, *et al*. A preliminary video analysis of concussion in the National Rugby League. *Brain Inj* 2015;29:1182–5.
- King D, Hume PA, Clark T. Nature of tackles that result in injury in professional rugby league. *Res Sports Med* 2012;20:86–104.
- McLeod S, Tucker R, Edwards S, *et al*. A case-control study of tackle based head impact event (HIE) risk factors from the first three seasons of the National Rugby League Women's competition. *Front Sports Act Living* 2023;5:1080356.
- Gardner AJ, Iverson GL, Edwards S, *et al*. A Case-Control Study of Tackle-Based Head Injury Assessment (HIA) Risk Factors in the National Rugby League. *Sports Med Open* 2021;7:84.
- von Elm E, Altman DG, Egger M, *et al*. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008;61:344–9.
- National Health and Medical Research Council, Australian Research Council, Australian Research Council, Universities Australia. National Statement on Ethical Conduct in Human Research. Canberra: National Health and Medical Research Council, 2023. Available: <https://www.nhmrc.gov.au/file/19531/download?token=rpY3-bU5>
- Gardner AJ, Iverson GL, Williams WH, *et al*. A systematic review and meta-analysis of concussion in rugby union. *Sports Med* 2014;44:1717–31.
- Gardner AJ, Iverson GL, Stanwell P, *et al*. A Video Analysis of Use of the New 'Concussion Interchange Rule' in the National Rugby League. *Int J Sports Med* 2016;37:267–73.
- Bathgate A, Best JP, Craig G, *et al*. A prospective study of injuries to elite Australian rugby union players. *Br J Sports Med* 2002;36:265–9.
- Falkenmire A, Manvell J, Callister R, *et al*. Injury incidence, characteristics and timing in amateur male rugby union: a prospective cohort study. *J Hum Sport Exerc* 2019;15.
- Roberts SP, Stokes KA, Williams S, *et al*. Injury in Starting and Replacement Players from Five Professional Men's Rugby Unions. *Sports Med* 2024;54:2157–67.
- Edwards S, Lee R, Fuller G, *et al*. 3D Biomechanics of Rugby Tackle Techniques to Inform Future Rugby Research Practice: a Systematic Review. *Sports Med Open* 2021;7:39.
- Cross MJ, Tucker R, Raftery M, *et al*. Tackling concussion in professional rugby union: a case-control study of tackle-based risk factors and recommendations for primary prevention. *Br J Sports Med* 2019;53:1021–5.



- 21 Hopkinson M, Bissas A, Nicholson G, *et al.* A video analysis framework for the rugby league tackle. *Sci Med Foot* 2022;6:15–28.
- 22 McIntosh AS, Savage TN, McCrory P, *et al.* Tackle characteristics and injury in a cross section of rugby union football. *Med Sci Sports Exerc* 2010;42:977–84.
- 23 Davis GA, Makdissi M, Bloomfield P, *et al.* International consensus definitions of video signs of concussion in professional sports. *Br J Sports Med* 2019;53:1264–7.
- 24 Gardner AJ, Howell DR, Iverson GL. A video review of multiple concussion signs in National Rugby League match play. *Sports Med Open* 2018;4:5.
- 25 Echemendia RJ, Brett BL, Broglio S, *et al.* Introducing the Sport Concussion Assessment Tool 6 (SCAT6). *Br J Sports Med* 2023;57:619–21.
- 26 Gardner AJ, Lang MA, Edwards S, *et al.* Head injury assessment in youth men's rugby league players: An evaluation of game play characteristics and video review of potential concussion signs. *J S A M S Plus* 2024;4:100072.
- 27 Echemendia RJ, Bruce JM, Meeuwisse W, *et al.* Can visible signs predict concussion diagnosis in the National Hockey League? *Br J Sports Med* 2018;52:1149–54.
- 28 Gardner AJ, Levi CR, Iverson GL. Observational Review and Analysis of Concussion: a Method for Conducting a Standardized Video Analysis of Concussion in Rugby League. *Sports Med Open* 2017;3:26.
- 29 Tierney GJ, Joodaki H, Krosshaug T, *et al.* Assessment of model-based image-matching for future reconstruction of unhelmeted sport head impact kinematics. *Sports Biomech* 2018;17:33–47.
- 30 Koga H, Nakamae A, Shima Y, *et al.* Hip and Ankle Kinematics in Noncontact Anterior Cruciate Ligament Injury Situations: Video Analysis Using Model-Based Image Matching. *Am J Sports Med* 2018;46:333–40.
- 31 Ghajari M, Hellyer PJ, Sharp DJ. Computational modelling of traumatic brain injury predicts the location of chronic traumatic encephalopathy pathology. *Brain (Bacau)* 2017;140:333–43.
- 32 Whitehead S, Owen C, Brown J, *et al.* Training injuries in elite men's senior and academy (Super League) rugby league; an analysis of 224,000 exposure-hours. *J Sci Med Sport* 2024;27:624–30.