



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/219685/>

Version: Accepted Version

Article:

Vicedo, Benoit, Caudrelier, Vincent and Singh, Anup Anand (2024) Lagrangian multiform for cyclotomic Gaudin models. SIGMA. ISSN: 1815-0659

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Lagrangian Multiform for Cyclotomic Gaudin Models

Vincent CAUDRELIER ^a, Anup Anand SINGH ^a and Benoît VICEDO ^b

^{a)} *School of Mathematics, University of Leeds, Leeds LS2 9JT, UK*

E-mail: v.caudrelier@leeds.ac.uk, anupandsingh@gmail.com

^{b)} *Department of Mathematics, University of York, York YO10 5DD, UK*

E-mail: b.vicedo@york.ac.uk

Received May 28, 2024, in final form November 07, 2024; Published online November ??, 2024

<https://doi.org/10.3842/SIGMA.2024.10?>

Abstract. We construct a Lagrangian multiform for the class of cyclotomic (rational) Gaudin models by formulating its hierarchy within the Lie dialgebra framework of Semenov-Tian-Shansky and by using the framework of Lagrangian multiforms on coadjoint orbits. This provides the first example of a Lagrangian multiform for an integrable hierarchy whose classical r -matrix is non-skew-symmetric and spectral parameter-dependent. As an important by-product of the construction, we obtain a Lagrangian multiform for the *periodic* Toda chain by choosing an appropriate realisation of the cyclotomic Gaudin Lax matrix. This fills a gap in the landscape of Toda models as only the open and infinite chains had been previously cast into the Lagrangian multiform framework. A slightly different choice of realisation produces the so-called discrete self-trapping (DST) model. We demonstrate the versatility of the framework by *coupling* the periodic Toda chain with the DST model and by obtaining a Lagrangian multiform for the corresponding integrable hierarchy.

Key words: Lagrangian multiforms; integrable systems; classical r -matrix; Gaudin models

2020 Mathematics Subject Classification: 17B80; 37J35; 70H06

1 Introduction

A characteristic property of an integrable system is that its equations of motion can be seen as members of a hierarchy of compatible equations. Traditional Lagrangians fail to capture this notion of commuting (Hamiltonian) flows, or its discrete analog known as multidimensional consistency [9, 25]. This obstacle to describing integrable hierarchies variationally was overcome for the first time in [23] by introducing a new object – a Lagrangian multiform – together with a generalised variational principle applied to an appropriate generalisation of a classical action, and noticing a fundamental property of the multiform: the so-called closure relation. Since its introduction, this idea has been developed in several directions. Its connections with more traditional features of integrability (Lax pairs and Hamiltonian structures, for instance) have been established in the various realms of integrable systems: discrete and continuous finite-dimensional systems [11, 27, 36, 43], continuous infinite-dimensional systems – field theories in $1 + 1$ dimensions [13, 14, 15, 28, 32, 33, 37] and in $2 + 1$ dimensions [26, 34] – and semi-discrete systems [35]. The relations between discrete and continuous multiforms were explored in [40]. The concept has even been extended to non-commuting flows in [12].

In general, hierarchies of models in d spacetime dimensions are described by a Lagrangian multiform which is a d -form integrated over a hypersurface of dimension d in a so-called multi-time space of dimension greater than d to yield an action functional depending not only on the field configurations but also on the hypersurface itself. This last point is the crucial new ingredient of the generalised variational principle used in Lagrangian multiform theory. One postulates a principle of least action which must be valid for *any* hypersurface embedded in the

multi-time space. This captures the commutativity of the flows variationally and was adopted as a definition of pluri-Lagrangian systems, see [27, 28] and references therein. In Lagrangian multiform theory, there is an additional postulate which leads to the *closure relation*. The latter has been shown [11, 36] to be equivalent to the Poisson involutivity of Hamiltonians, the Liouville criterion for integrability.

The generalised variational principle produces equations that come in two flavours: 1) Euler–Lagrange equations associated with each of the coefficients of the Lagrangian multiform which form a collection of Lagrangian densities; 2) Corner or structure equations on the Lagrangian coefficients themselves which select possible models and ensure the compatibility of the various equations of motion imposed on a common set of fields. This paper deals with a finite-dimensional system, so let us illustrate the ingredients for this case. The basic objects are a Lagrangian 1-form

$$\mathcal{L}[q] = \sum_{k=1}^N \mathcal{L}_k[q] dt_k$$

and the related generalised action

$$S[q, \Gamma] = \int_{\Gamma} \mathcal{L}[q],$$

where Γ is a curve in the multi-time \mathbb{R}^N with (time) coordinates t_1, \dots, t_N , and q denotes generic configuration coordinates. For instance, q could be a position vector in \mathbb{R}^d for some d , or as will be the case for us, an element of a (matrix) Lie group. The notations $\mathcal{L}[q]$ and $\mathcal{L}_k[q]$ mean that these quantities depend on q and a finite number of derivatives of q with respect to the times t_1, \dots, t_N . In this paper, we restrict ourselves only to the case of first derivatives and simply write \mathcal{L}_k for the Lagrangian coefficients. The application of the generalised variational principle leads to the following multi-time Euler–Lagrange equations [36]:

$$\frac{\partial \mathcal{L}_k}{\partial q} - \partial_{t_k} \frac{\partial \mathcal{L}_k}{\partial q_{t_k}} = 0, \tag{1.1}$$

$$\frac{\partial \mathcal{L}_k}{\partial q_{t_\ell}} = 0, \quad \ell \neq k, \tag{1.2}$$

$$\frac{\partial \mathcal{L}_k}{\partial q_{t_k}} = \frac{\partial \mathcal{L}_\ell}{\partial q_{t_\ell}}, \quad k, \ell = 1, \dots, N. \tag{1.3}$$

Note that (1.1) is simply the standard Euler–Lagrange equation for each \mathcal{L}_k . Equations (1.2) and (1.3) are the corner equations. It turns out that these corner equations will, in fact, be identically satisfied by our Lagrangian multiforms, but in general they can represent non-trivial equations for q . The closure relation then stipulates that

$$d\mathcal{L}[q] = 0 \quad \Leftrightarrow \quad \partial_{t_k} \mathcal{L}_j - \partial_{t_j} \mathcal{L}_k = 0$$

on solutions of (1.1)–(1.3).

Gaudin models are a general class of integrable systems associated with Lie algebras with a nondegenerate invariant bilinear form. They were first introduced for the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ by M. Gaudin in [19] as quantum integrable spin chains with long-range interactions, and then generalised for arbitrary semi-simple Lie algebras in [20]. At both the classical and quantum levels, the integrable structure of this model, associated with a Lie algebra \mathfrak{g} , is underpinned by the rational skew-symmetric solution

$$r_{\underline{12}}^0(\lambda, \mu) = \frac{C_{\underline{12}}}{\mu - \lambda} \tag{1.4}$$

of the classical Yang–Baxter equation¹ (CYBE)

$$[r_{\underline{12}}(\lambda, \mu), r_{\underline{13}}(\lambda, \nu)] + [r_{\underline{12}}(\lambda, \mu), r_{\underline{23}}(\mu, \nu)] + [r_{\underline{32}}(\nu, \mu), r_{\underline{13}}(\lambda, \nu)] = 0.$$

The rational classical r -matrix (1.4) takes values in $\mathfrak{g} \otimes \mathfrak{g}$ and depends on spectral parameters $\lambda, \mu \in \mathbb{C}$, and the tensor Casimir $C_{\underline{12}} \in \mathfrak{g} \otimes \mathfrak{g}$ appearing in (1.4) is defined as

$$C_{\underline{12}} = I_a \otimes I^a,$$

where $\{I_a\}$ and $\{I^a\}$ are dual bases of \mathfrak{g} with respect to a fixed nondegenerate invariant bilinear form. The main property of $C_{\underline{12}}$ which gives it its name is $[C_{\underline{12}}, X_{\underline{1}} + X_{\underline{2}}] = 0$ for all $X \in \mathfrak{g}$ and where we use the standard tensorial notation $X_{\underline{1}} \equiv X \otimes \mathbb{1}$ and $X_{\underline{2}} \equiv \mathbb{1} \otimes X$.

The cyclotomic Gaudin model of interest in this paper arises as a specialisation of a general procedure which can be traced back to reduction group ideas [24], first applied in the form of an averaging procedure [29] to the rational classical r -matrix (1.4) in order to produce the trigonometric and elliptic r -matrices. This was generalised in various ways, for instance, in [4, 5, 6] in the context of Sklyanin’s (linear) Poisson algebra

$$\{U_{\underline{1}}(\lambda), U_{\underline{2}}(\mu)\} = [r_{\underline{12}}^0(\lambda, \mu), U_{\underline{1}}(\lambda) + U_{\underline{2}}(\mu)], \quad (1.5)$$

where, as above, the index denotes which factor in the tensor product $\mathfrak{g} \otimes \mathfrak{g}$ the \mathfrak{g} -valued function U sits in, that is, $U_{\underline{1}}(\lambda) \equiv U(\lambda) \otimes \mathbb{1}$ and $U_{\underline{2}}(\mu) \equiv \mathbb{1} \otimes U(\mu)$.

The idea is to use a Lie algebra automorphism appropriately extended to a loop algebra automorphism. For our purpose, we use an automorphism σ of order T on \mathfrak{g} and define

$$\phi: U(\lambda) \mapsto \omega \sigma^{-1}(U(\omega\lambda)),$$

where ω is a T -th root of unity. Thus, ϕ is an automorphism of the Poisson algebra (1.5). This leads us to consider the fixed point subalgebra generated by

$$L(\lambda) = \frac{1}{T} \sum_{k=0}^{T-1} \omega^{-k} \sigma^k U(\omega^{-k}\lambda), \quad \sigma(L(\lambda)) = \omega L(\omega\lambda). \quad (1.6)$$

The Poisson algebra for $L(\lambda)$ closes into

$$\{L_{\underline{1}}(\lambda), L_{\underline{2}}(\mu)\} = [r_{\underline{12}}(\lambda, \mu), L_{\underline{1}}(\lambda)] - [r_{\underline{21}}(\mu, \lambda), L_{\underline{2}}(\mu)], \quad (1.7)$$

where

$$r_{\underline{12}}(\lambda, \mu) = \frac{1}{T} \sum_{k=0}^{T-1} \frac{\sigma_{\underline{1}}^k C_{\underline{12}}}{\mu - \omega^{-k}\lambda} \quad (1.8)$$

is a non-skew-symmetric solution of CYBE. These facts are special cases of [10, Propositions 2.2 and 4.1]. It is well known [8] that the Poisson algebra (1.7) ensures that the quantities $\text{Tr} L(\lambda)^p$ Poisson commute and the equations of motion generated by them with respect to (1.7) take the Lax form. More precisely, the generating Hamiltonian

$$H(\mu) = \frac{1}{p+1} \text{Tr} L(\mu)^{p+1}, \quad p \in \mathbb{Z}_{\geq 0},$$

satisfies

$$\{L(\lambda), H(\mu)\} = [M(\lambda, \mu), L(\lambda)],$$

¹Note that this is the general version of the CYBE applicable to non-skew-symmetric matrices.

where by definition $\{L(\lambda), H(\mu)\} = \{L^a(\lambda), H(\mu)\}I_a$, and

$$M_{\underline{1}}(\lambda, \mu) = \text{Tr}_{\underline{2}} r_{\underline{1}\underline{2}}(\lambda, \mu) L_{\underline{2}}(\mu)^p.$$

This yields an infinite number of Hamiltonians by specifying p and extracting coefficients at the poles of $L(\mu)$. Of course, only a finite number of these Hamiltonians are independent when acting on a finite-dimensional phase space. They generate mutually compatible flows forming an integrable hierarchy.

If one applies this construction to a Lax matrix of rational Gaudin model type, with poles in the finite set $D = \{0, \zeta_1, \dots, \zeta_N, \infty\} \subset \mathbb{C}P^1$,

$$U(\lambda) = \sum_{n=0}^{N_0-1} \frac{U_0^{(n)}}{\lambda^{n+1}} + \sum_{r=1}^N \sum_{n=0}^{N_r-1} \frac{U_r^{(n)}}{(\lambda - \zeta_r)^{n+1}} + \sum_{n=0}^{N_\infty} U_\infty^{(n)} \lambda^n, \quad U_s^{(n)} \in \mathfrak{g}, \quad (1.9)$$

then one obtains the so-called cyclotomic Gaudin model [31, 41, 42] and its associated integrable hierarchy. Note that in (1.9), we have anticipated that the pole at 0 and at infinity behave differently in (1.6) compared to the poles at $\lambda = \zeta_r \neq 0, \infty$. Although we will present the general algebraic setup, for simplicity, in our examples we will only consider the case where $U(\lambda)$ has simple poles at all ζ_r , $r \in \{1, \dots, N\}$, and double poles at the origin and at infinity. This will be sufficient for our application to the periodic Toda chain and the discrete self-trapping (DST) model. The corresponding Lax matrix of the cyclotomic Gaudin model then takes the form

$$L(\lambda) = \frac{X_0^{(0)}}{\lambda} + \frac{X_0^{(1)}}{\lambda^2} + \frac{1}{T} \sum_{r=1}^N \sum_{k=0}^{T-1} \frac{\sigma^k X_r}{\lambda - \omega^k \zeta_r} + X_\infty \quad (1.10)$$

with $\sigma X_0^{(j)} = \omega^{-j} X_0^{(j)}$ and $\sigma X_\infty = \omega X_\infty$.

Our goal is to give a Lagrangian multiform description of the cyclotomic Gaudin hierarchy. In a nutshell, what is required is to interpret $L(\lambda)$ as an element of an appropriate coadjoint orbit so that we can write our Lagrangian multiform following the construction introduced in [11]. The latter is based on using the Lie dialgebra formalism [30] which is reviewed in Section 2. In Section 3, we show how to construct the required algebraic ingredients for the cyclotomic Gaudin model. This is applied in Section 3.4, where we obtain a Lagrangian multiform for the cyclotomic Gaudin hierarchy and discuss its properties.

Section 4 deals with two special realisations of the cyclotomic Gaudin model: the periodic Toda chain (Section 4.1) and the integrable case of the DST model (Section 4.2). We derive Lagrangian multiforms for the hierarchies corresponding to these two theories using our results from Section 3. The construction of the periodic Toda multiform, in particular, complements the results of [11] where a Lagrangian multiform for the open Toda chain was derived, while in [35] the infinite Toda chain was considered within a semi-discrete Lagrangian multiform. Finally, in Section 4.3, we illustrate how to couple the periodic Toda and DST hierarchies together in a straightforward manner and derive a Lagrangian multiform for this coupled hierarchy. We end with concluding remarks in Section 5.

2 Lagrangian multiforms on coadjoint orbits

In this section, we briefly review the results of [11] where a Lagrangian multiform was obtained which produces a hierarchy of equations of motion in Lax form as its multi-time Euler–Lagrange equations. We will, in fact, be interested in applying this framework in the infinite-dimensional setting for which we refer more precisely to [11, Section 7]. In order to avoid confusion with the finite-dimensional Lie algebra \mathfrak{g} used throughout the rest of this paper, in this section we

denote by \mathfrak{g} the infinite-dimensional Lie algebra of interest. The construction of the specific Lie algebra \mathfrak{g} relevant for the cyclotomic Gaudin model will be the subject of the next section.

The framework of Lie dialgebras for constructing hierarchies of compatible Lax equations involves a number of ingredients. For the purpose of this paper, we will actually only require a particular instance of this general framework corresponding to the so-called Adler–Kostant–Symes scheme [2, 21, 38], so we will mainly focus on this case to ease the comparison with later sections. The ingredients are:

1. A Lie algebra \mathfrak{g} , with Lie group \mathbb{G} , which we assume is equipped with a nondegenerate invariant bilinear pairing

$$\langle \cdot, \cdot \rangle: V \times \mathfrak{g} \longrightarrow \mathbb{C} \quad (2.1)$$

with some representation V of \mathbb{G} , allowing one to use V as a model for the dual space \mathfrak{g}^* of \mathfrak{g} . We will denote the representation of \mathbb{G} by $\text{Ad}^*: \mathbb{G} \times V \rightarrow V$, $(\varphi, \xi) \mapsto \text{Ad}_{\varphi}^* \xi$ and the corresponding Lie algebra representation by $\text{ad}^*: \mathfrak{g} \times V \rightarrow V$, $(X, \xi) \mapsto \text{ad}_X^* \xi$, and refer to these as the coadjoint representations.

Note that if \mathfrak{g} were finite-dimensional then a good choice of representation space V would be the algebraic dual \mathfrak{g}' of \mathfrak{g} , in which case (2.1) is given by the canonical pairing. In the infinite-dimensional setting we are considering, the algebraic dual is too big. When \mathfrak{g} is equipped with a nondegenerate invariant symmetric bilinear form $(\cdot, \cdot): \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ then one possible replacement for the algebraic dual is afforded by the smooth dual $(\mathfrak{g}, \cdot) \subset \mathfrak{g}'$ which is canonically isomorphic to \mathfrak{g} itself. However, in our present setting it turns out that the smooth dual will not be the appropriate notion of dual space, which is why we need the above more general working definition for the dual space \mathfrak{g}^* .

2. A linear map $R: \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying the *modified* classical Yang–Baxter equation (mCYBE)

$$[R(X), R(Y)] - R([R(X), Y] + [X, R(Y)]) = -[X, Y], \quad \forall X, Y \in \mathfrak{g}. \quad (2.2)$$

This is the crucial ingredient providing us with the Lie dialgebra $(\mathfrak{g}, \mathfrak{g}_R)$ where \mathfrak{g}_R is the vector space \mathfrak{g} equipped with a second Lie bracket $[X, Y]_R = \frac{1}{2}([R(X), Y] + [X, R(Y)])$. In the Adler–Kostant–Symes scheme, this is determined by a direct sum decomposition²

$$\mathfrak{g} = \mathfrak{g}_+ \dot{+} \mathfrak{g}_- \quad (2.3)$$

into complementary subalgebras $\mathfrak{g}_{\pm} \subset \mathfrak{g}$. A solution to (2.2) is then given by $R = P_+ - P_-$, where $P_{\pm}: \mathfrak{g} \rightarrow \mathfrak{g}_{\pm}$ denote the projections onto these respective subalgebras. In this case, the Lie algebra \mathfrak{g}_R is the direct sum of Lie algebras $\mathfrak{g}_R = \mathfrak{g}_+ \oplus \mathfrak{g}_-$. Now (2.3) induces a corresponding direct sum decomposition of the model V for the dual space \mathfrak{g}^* , namely

$$V = V_+ \dot{+} V_-, \quad (2.4)$$

where V_{\pm} is the orthogonal complement of \mathfrak{g}_{\pm} with respect to the pairing (2.1). We will denote by $\mathcal{P}_{\pm}: V \rightarrow V_{\pm}$ the projections onto these respective subspaces.

Let \mathbb{G}_{\pm} denote the Lie subgroups of \mathbb{G} associated with \mathfrak{g}_{\pm} , and \mathbb{G}_R denote the Lie group associated with \mathfrak{g}_R , so that (at least locally) $\mathbb{G}_R \simeq \mathbb{G}_+ \times \mathbb{G}_-$. It will be useful to note for later that the coadjoint representation of \mathbb{G}_R on $V \equiv \mathfrak{g}^*$ can be expressed in terms of that of the subgroups $\mathbb{G}_{\pm} \subset \mathbb{G}$ as

$$\text{Ad}_{\varphi}^{R*} \xi = \mathcal{P}_-(\text{Ad}_{\varphi_+}^* \xi) + \mathcal{P}_+(\text{Ad}_{\varphi_-}^* \xi) \quad (2.5)$$

for any $\varphi = (\varphi_+, \varphi_-) \in \mathbb{G}_R$ and $\xi \in \mathfrak{g}^*$.

²Throughout the text, we use $\dot{+}$ to denote a direct sum of vector spaces, and \oplus to denote a direct sum of Lie algebras. This is a convenient convention borrowed from [30].

3. A collection of independent Ad^* -invariant functions on \mathfrak{g}^* , say H_k , for $k = 1, \dots, N$. It follows from their Ad^* -invariance that these functions are in involution with respect to the Lie–Poisson bracket on \mathfrak{g}^* given by

$$\{f, g\}_R(\xi) = \langle \xi, [\nabla f(\xi), \nabla g(\xi)]_R \rangle, \quad f, g \in C^\infty(\mathfrak{g}^*),$$

and generate a hierarchy of compatible flows on the Poisson manifold $(\mathfrak{g}^*, \{, \}_R)$ which take the form of Lax equations

$$\partial_{t_k} L := \{L, H_k\}_R = \text{ad}_{M_k(L)}^* L, \quad M_k(L) = \frac{1}{2} R \nabla H_k(L). \quad (2.6)$$

When \mathfrak{g} and V are subspaces of a common ambient Lie algebra and the coadjoint action $\text{ad}^*: \mathfrak{g} \times V \rightarrow V$ is given by a Lie bracket in this ambient Lie algebra, as will be the case for us in Section 3 below, the Lax equations (2.6) take the more familiar form

$$\partial_{t_k} L = [M_k(L), L].$$

This is the content of Semenov-Tian-Shansky’s theorem, see, for instance, [30].

The symplectic leaves of the Poisson manifold $(\mathfrak{g}^*, \{, \}_R)$ are given by the coadjoint orbits

$$\mathcal{O}_\Lambda = \{ \text{Ad}_\varphi^{R*} \Lambda \mid \varphi \in \mathbb{G}_R \} \quad \text{for } \Lambda \in \mathfrak{g}^*, \quad (2.7)$$

which therefore provide natural phase spaces for integrable systems. The key idea of [11] was to define a Lagrangian multiform on such a coadjoint orbit which produces the hierarchy (2.6) as its Euler–Lagrange equations. Specifically, let $L = \text{Ad}_\varphi^{R*} \Lambda$ and define

$$\mathcal{L}[\varphi] = \sum_{k=1}^N \mathcal{L}_k dt_k, \quad \mathcal{L}_k = \langle L, \partial_{t_k} \varphi \cdot_R \varphi^{-1} \rangle - H_k(L), \quad (2.8)$$

where \cdot_R denotes the multiplication in \mathbb{G}_R . Then, we have the following.

Theorem 2.1 ([11]). *The Lagrangian 1-form (2.8) satisfies the corner equations (1.2)–(1.3) of the multi-time Euler–Lagrange equations. The standard Euler–Lagrange equations (1.1) associated with the Lagrangian coefficients \mathcal{L}_k give the hierarchy of compatible Lax equations (2.6)*

$$\partial_{t_k} L = \text{ad}_{M_k(L)}^* L, \quad k = 1, \dots, N. \quad (2.9)$$

The closure relation³ holds: on solutions of (2.9) we have

$$\partial_{t_k} \mathcal{L}_j - \partial_{t_j} \mathcal{L}_k = 0, \quad j, k = 1, \dots, N.$$

In short, the machinery of [11] provides a systematic way of constructing a Lagrangian multiform for a hierarchy of Lax equations (2.6) on a coadjoint orbit (2.7). To apply this result to the cyclotomic Gaudin model we must identify its Lax matrix (1.10) as an element of an appropriate coadjoint orbit $\mathcal{O}_\Lambda \subset \mathfrak{g}^*$, as in (2.7), for a suitable Lie algebra \mathfrak{g} and R -matrix. In particular, we will need to build a suitable model V for the dual space \mathfrak{g}^* via a corresponding pairing $\langle \cdot, \cdot \rangle: V \times \mathfrak{g} \rightarrow \mathbb{C}$. This will be the central goal of the next section.

³The closure relation for the Lagrangian one-form \mathcal{L} is, in fact, equivalent to the involutivity of the Hamiltonians H_k with respect to the Lie–Poisson R -bracket $\{, \}_R$. We do not discuss this result here and refer the interested reader to [11, Corollary 3.4].

3 Cyclotomic Gaudin model

The purpose of this section is to formulate a Lagrangian multiform for the cyclotomic Gaudin model using the machinery developed in [11]. As outlined at the end of Section 2, the first immediate task is to introduce the relevant infinite-dimensional Lie algebra \mathfrak{g} and a suitable model V for its dual space \mathfrak{g}^* by constructing a nondegenerate invariant bilinear pairing as in (2.1). This will be achieved in Section 3.1, which culminates in a description of the relevant coadjoint orbits $\mathcal{O}_\Lambda \subset \mathfrak{g}^*$ in Lemma 3.5. In Section 3.2, we specialise to the case of simple poles at $\omega^k \zeta_r$, $k \in \{0, \dots, T-1\}$, $r \in \{1, \dots, N\}$, and double poles at the origin and at infinity, and then describe the Lax matrix (1.10) of the cyclotomic Gaudin model as an element of this coadjoint orbit. In Section 3.3, we then describe the Lax equations as flows on this coadjoint orbit, as in (2.6). Finally, in Section 3.4, we put everything together to obtain the Lagrangian multiform for the cyclotomic Gaudin model.

3.1 Lie dialgebra framework

Let us fix $T \in \mathbb{Z}_{\geq 1}$ and pick a primitive T -th root of unity $\omega \in \mathbb{C}^\times$. We can then define

$$\Gamma := \{1, \omega, \omega^2, \dots, \omega^{T-1}\},$$

a copy of the cyclic group \mathbb{Z}_T of order T that acts on \mathbb{C} by multiplication. Further, we introduce the finite set of points $D = \{0, \zeta_1, \dots, \zeta_N, \infty\} \subset \mathbb{C}P^1$ including the point at infinity such that the Γ -orbits of the points ζ_1, \dots, ζ_N are pairwise disjoint, that is,

$$\Gamma \zeta_r \cap \Gamma \zeta_s = \emptyset \quad \text{for all } 1 \leq r \neq s \leq N.$$

Note that unlike the non-zero finite points ζ_1, \dots, ζ_N , the origin and the point at infinity are fixed under the action of Γ .

Let \mathfrak{g} be a finite-dimensional Lie algebra over \mathbb{C} with an automorphism σ of order T . For simplicity, we will only consider matrix Lie algebras in this work, with the Lie bracket being the commutator, and a nondegenerate invariant bilinear pairing given by the trace. We also assume that the automorphism σ preserves the trace, that is, $\text{Tr}(\sigma(x)\sigma(y)) = \text{Tr}(xy)$ for any $x, y \in \mathfrak{g}$. The eigenspaces of σ ,

$$\mathfrak{g}^{(k)} = \{X \in \mathfrak{g} \mid \sigma(X) = \omega^k X\}, \quad k \in \{0, \dots, T-1\}, \quad (3.1)$$

form a \mathbb{Z}_T -gradation of \mathfrak{g} , namely

$$\mathfrak{g} = \mathfrak{g}^{(0)} \dot{+} \dots \dot{+} \mathfrak{g}^{(T-1)} \quad \text{with} \quad [\mathfrak{g}^{(k_1)}, \mathfrak{g}^{(k_2)}] = \mathfrak{g}^{(k_1+k_2 \bmod T)}. \quad (3.2)$$

From now on, we will simply write $\mathfrak{g}^{(n)}$ to mean $\mathfrak{g}^{(n \bmod T)}$. Further, let us define the local parameters

$$\lambda_0 = \lambda, \quad \lambda_r = \lambda - \zeta_r \quad \text{for } r \in \{1, \dots, N\}, \quad \lambda_\infty = \frac{1}{\lambda}. \quad (3.3)$$

It will be convenient to also introduce an additional set of local parameters $\lambda_{r,k} = \lambda - \omega^k \zeta_r$ for all $k \in \{0, \dots, T-1\}$ and $r \in \{1, \dots, N\}$. We also introduce an index set $S = \{0, 1, \dots, N, \infty\}$ where 0 and ∞ denote labels for indices rather than points in $\mathbb{C}P^1$.

Let us denote by $\mathcal{F}_{D'}$ the algebra of rational functions in the formal variable λ with values in \mathfrak{g} that are regular outside $D' = \{\omega^k \zeta_r \mid k \in \{0, \dots, T-1\}, r \in \{1, \dots, N\}\} \cup \{0, \infty\}$. Two subspaces of this algebra will be of relevance here: the subspace $\mathcal{F}_{D'}^\Gamma$ of *equivariant functions* and the subspace $\Omega_{D'}^\Gamma$ of *equivariant one-forms* that we define as

$$\mathcal{F}_{D'}^\Gamma := \{f \in \mathcal{F}_{D'} \mid \sigma(f(\lambda)) = f(\omega\lambda)\}, \quad \Omega_{D'}^\Gamma := \{g \in \mathcal{F}_{D'} \mid \sigma(g(\lambda)) = \omega g(\omega\lambda)\}.$$

In general, equivariant functions and equivariant one-forms take the forms

$$f = \sum_{n=0}^{M_0-1} \frac{X_0^{(n)}}{\lambda^{n+1}} + \sum_{r=1}^N \sum_{k=0}^{T_r-1} \sum_{n=0}^{M_r-1} \frac{\sigma^k X_r^{(n)}}{(\omega^{-k}\lambda - \zeta_r)^{n+1}} + \sum_{n=0}^{M_\infty} \frac{X_\infty^{(n)}}{\lambda_\infty^n},$$

$$g = \sum_{n=0}^{N_0-1} \frac{Y_0^{(n)}}{\lambda^{n+1}} + \sum_{r=1}^N \sum_{k=0}^{T_r-1} \sum_{n=0}^{N_r-1} \frac{\omega^{-k} \sigma^k Y_r^{(n)}}{(\omega^{-k}\lambda - \zeta_r)^{n+1}} + \sum_{n=0}^{N_\infty} \frac{Y_\infty^{(n)}}{\lambda_\infty^n},$$

respectively, with

$$\begin{aligned} X_0^{(n)} &\in \mathfrak{g}^{(-n-1)}, & X_r^{(n)} &\in \mathfrak{g} \quad \text{for } r \in \{1, \dots, N\}, & X_\infty^{(n)} &\in \mathfrak{g}^{(n)}, \\ Y_0^{(n)} &\in \mathfrak{g}^{(-n)}, & Y_r^{(n)} &\in \mathfrak{g} \quad \text{for } r \in \{1, \dots, N\}, & Y_\infty^{(n)} &\in \mathfrak{g}^{(n+1)}. \end{aligned} \quad (3.4)$$

One can check that the subspace $\mathcal{F}_{D'}^\Gamma$ is, in fact, a Lie subalgebra of $\mathcal{F}_{D'}$. The subspace $\Omega_{D'}^\Gamma$, on the other hand, contains the Lax matrix (1.10) of the model. To construct the cyclotomic Gaudin multiform à la [11], we will realise this Lax matrix as an element of a certain coadjoint orbit. To do so, let us define the loop algebra of formal Laurent series in the local parameter λ_r with coefficients in \mathfrak{g} , for each $r \in S$,

$$\tilde{\mathfrak{g}}_r = \mathfrak{g} \otimes \mathbb{C}((\lambda_r)), \quad (3.5)$$

with Lie bracket

$$[X\lambda_r^i, Y\lambda_r^j] = [X, Y]\lambda_r^{i+j}, \quad X, Y \in \mathfrak{g}, \quad (3.6)$$

and then consider the direct sum

$$\tilde{\mathfrak{g}}_D = \bigoplus_{r \in S} \tilde{\mathfrak{g}}_r.$$

The Lie bracket of two elements $X, Y \in \tilde{\mathfrak{g}}_D$ is defined component-wise

$$[X, Y]_r(\lambda_r) = [X_r(\lambda_r), Y_r(\lambda_r)].$$

Of interest to us here are two particular subspaces of $\tilde{\mathfrak{g}}_D$,

$$\tilde{\mathfrak{g}}_D^{(0)} = \tilde{\mathfrak{g}}_0^{\Gamma,0} \oplus \bigoplus_{r=1}^N \tilde{\mathfrak{g}}_r \oplus \tilde{\mathfrak{g}}_\infty^{\Gamma,0} \quad \text{and} \quad \tilde{\mathfrak{g}}_D^{(1)} = \tilde{\mathfrak{g}}_0^{\Gamma,1} \oplus \bigoplus_{r=1}^N \tilde{\mathfrak{g}}_r \oplus \tilde{\mathfrak{g}}_\infty^{\Gamma,1},$$

where the twisted spaces attached to the origin and infinity are defined as

$$\begin{aligned} \tilde{\mathfrak{g}}_0^{\Gamma,k} &= \{X_0(\lambda_0) \in \tilde{\mathfrak{g}}_0 \mid \sigma(X_0(\lambda_0)) = \omega^k X_0(\omega\lambda_0)\}, \\ \tilde{\mathfrak{g}}_\infty^{\Gamma,k} &= \{X_\infty(\lambda_\infty) \in \tilde{\mathfrak{g}}_\infty \mid \sigma(X_\infty(\lambda_\infty)) = \omega^k X_\infty(\omega^{-1}\lambda_\infty)\} \end{aligned} \quad (3.7)$$

for $k = 0, 1$. Elements of $\tilde{\mathfrak{g}}_D^{(0)}$ and $\tilde{\mathfrak{g}}_D^{(1)}$ are tuples of the form

$$X = \left(\sum_{n=-\infty}^{M_0-1} \frac{X_0^{(n)}}{\lambda^{n+1}}, \sum_{n=-\infty}^{M_1-1} \frac{X_1^{(n)}}{(\lambda - \zeta_1)^{n+1}}, \dots, \sum_{n=-\infty}^{M_N-1} \frac{X_N^{(n)}}{(\lambda - \zeta_N)^{n+1}}, \sum_{n=-\infty}^{M_\infty} \frac{X_\infty^{(n)}}{\lambda_\infty^n} \right), \quad (3.8)$$

and

$$Y = \left(\sum_{n=-\infty}^{N_0-1} \frac{Y_0^{(n)}}{\lambda^{n+1}}, \sum_{n=-\infty}^{N_1-1} \frac{Y_1^{(n)}}{(\lambda - \zeta_1)^{n+1}}, \dots, \sum_{n=-\infty}^{N_N-1} \frac{Y_N^{(n)}}{(\lambda - \zeta_N)^{n+1}}, \sum_{n=-\infty}^{N_\infty} \frac{Y_\infty^{(n)}}{\lambda_\infty^n} \right), \quad (3.9)$$

respectively, where the coefficients $X_r^{(n)}$ and $Y_r^{(n)}$, $r \in S$, satisfy the conditions in (3.4). Note that each coefficient in (3.8) and (3.9) is a \mathfrak{g} -valued Laurent series, i.e., an element of (3.5) for $r \in S$. This is clear from changing variables in each sum from n to $-n$. The reason for the unusual choice of range for the indices n in each of the above sums is to make the expressions for the maps in (3.12) defined below slightly more transparent.

The subspace $\tilde{\mathfrak{g}}_D^{(0)}$ is a Lie subalgebra of $\tilde{\mathfrak{g}}_D$ which defines the desired infinite-dimensional Lie algebra $\mathfrak{g} = \tilde{\mathfrak{g}}_D^{(0)}$ in the notation of Section 2. Continuing the identification of the different ingredients from the Lie dialgebra framework listed in Section 2, notice that $V = \tilde{\mathfrak{g}}_D^{(1)}$ is clearly a representation of the Lie algebra $\mathfrak{g} = \tilde{\mathfrak{g}}_D^{(0)}$ since we have $[\tilde{\mathfrak{g}}_r^{\Gamma,0}, \tilde{\mathfrak{g}}_r^{\Gamma,1}] \subset \tilde{\mathfrak{g}}_r^{\Gamma,1}$ for $r \in \{0, \infty\}$. In the notation of Section 2, the representation $\text{ad}^*: \mathfrak{g} \times V \rightarrow V$ is given explicitly by $(X, Y) \mapsto [X, Y]$. The next proposition identifies $V = \tilde{\mathfrak{g}}_D^{(1)}$ as a suitable model for the dual space \mathfrak{g}^* .

Proposition 3.1. *The bilinear pairing $\langle \cdot, \cdot \rangle: \tilde{\mathfrak{g}}_D^{(1)} \times \tilde{\mathfrak{g}}_D^{(0)} \rightarrow \mathbb{C}$ defined by*

$$\langle Y, X \rangle = T \sum_{r=1}^N \text{Res}_{\lambda_r=0} \text{Tr}(Y_r(\lambda_r)X_r(\lambda_r))d\lambda + \sum_{r \in \{0, \infty\}} \text{Res}_{\lambda_r=0} \text{Tr}(Y_r(\lambda_r)X_r(\lambda_r))d\lambda \quad (3.10)$$

for any $Y \in \tilde{\mathfrak{g}}_D^{(1)}$ and $X \in \tilde{\mathfrak{g}}_D^{(0)}$, is nondegenerate and invariant under the adjoint action of $\tilde{\mathfrak{g}}_D^{(0)}$.

Proof. Recall that the trace $\text{Tr}: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$, $(x, y) \mapsto \text{Tr}(xy)$ is a nondegenerate invariant bilinear pairing on \mathfrak{g} and invariant under the action of σ . Given any $x \in \mathfrak{g}^{(m)}$ and $y \in \mathfrak{g}^{(n)}$, we have $\text{Tr}(xy) = \text{Tr}(\sigma(x)\sigma(y)) = \omega^{m+n} \text{Tr}(xy)$. Therefore, $\text{Tr}(xy) = 0$ for all $x \in \mathfrak{g}^{(m)}$ and $y \in \mathfrak{g}^{(n)}$ if $m+n \not\equiv 0 \pmod{T}$. Now, since $\text{Tr}: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ is nondegenerate on \mathfrak{g} , given $y \in \mathfrak{g}^{(n)}$, there is $x \in \mathfrak{g}$ such that $\text{Tr}(xy) \neq 0$. From (3.2), we have $x = x^{(0)} + \dots + x^{(T-1)}$, and from the previous result $\text{Tr}(xy) = \text{Tr}(x^{(m)}y)$ with $m+n \equiv 0 \pmod{T}$. Hence, $\text{Tr}(x^{(m)}y) \neq 0$ and we get a nondegenerate pairing between the subspaces $\mathfrak{g}^{(m)}$ and $\mathfrak{g}^{(n)}$ with $m+n \equiv 0 \pmod{T}$.

Now given any non-zero element $Y \in \tilde{\mathfrak{g}}_D^{(1)}$, it has a non-zero component $Y_r^{(m)} \lambda_r^{-m-1}$ for some $m \in \mathbb{Z}$ and $r \in S$. If $r = 0$ then $Y_0^{(m)} \in \mathfrak{g}^{(-m)}$ by (3.4), and we can find an $X_0^{(-m-1)} \in \mathfrak{g}^{(m)}$ which pairs non-trivially with it under the trace. So, $X_0^{(-m-1)} \lambda_0^m$ pairs non-trivially with Y . Likewise, if $r = \infty$ then $Y_\infty^{(m)} \in \mathfrak{g}^{(m+1)}$ by (3.4) and we can find an $X_\infty^{(-m-1)} \in \mathfrak{g}^{(-m-1)}$ which pairs non-trivially with it under the trace so that $X_\infty^{(-m-1)} \lambda_\infty^{m+1}$ pairs non-trivially with Y . And if $r \in \{1, \dots, N\}$ then pick any $X_r^{(-m-1)} \in \mathfrak{g}$ which pairs non-trivially with $Y_r^{(m)} \in \mathfrak{g}$ under the trace so that $X_r^{(-m-1)} \lambda_r^m$ pairs non-trivially with Y . Therefore, the bilinear pairing $\langle \cdot, \cdot \rangle$ is nondegenerate in the first argument. A similar argument establishes the nondegeneracy in the second argument.

Finally, the invariance of the bilinear pairing $\langle \cdot, \cdot \rangle$ under the adjoint action of $\tilde{\mathfrak{g}}_D^{(0)}$ follows from the definition (3.6) of the Lie bracket in $\tilde{\mathfrak{g}}_D$ and the invariance of the trace under the adjoint action of \mathfrak{g} . ■

Next, we turn to the identification of the subalgebras $\mathfrak{g}_\pm \subset \mathfrak{g}$ and the corresponding subspaces $V_\pm \subset V$ in the notation of Section 2. Crucially for us, the spaces $\mathcal{F}_{D'}^\Gamma$ and $\Omega_{D'}^\Gamma$ embed into $\tilde{\mathfrak{g}}_D^{(0)}$ and $\tilde{\mathfrak{g}}_D^{(1)}$, respectively,

$$\begin{aligned} \iota_\lambda: \mathcal{F}_{D'}^\Gamma &\hookrightarrow \tilde{\mathfrak{g}}_D^{(0)}, & f &\mapsto (\iota_{\lambda_0} f, \iota_{\lambda_1} f, \dots, \iota_{\lambda_N} f, \iota_{\lambda_\infty} f), \\ \iota_\lambda: \Omega_{D'}^\Gamma &\hookrightarrow \tilde{\mathfrak{g}}_D^{(1)}, & g &\mapsto (\iota_{\lambda_0} g, \iota_{\lambda_1} g, \dots, \iota_{\lambda_N} g, \iota_{\lambda_\infty} g), \end{aligned} \quad (3.11)$$

where, for each $r \in S$, $\iota_{\lambda_r} f$ and $\iota_{\lambda_r} g$ respectively denote the formal Laurent expansion of f and g about the point ζ_r . It will also be useful to define left inverses for these embeddings:

$$\begin{aligned} \pi_\lambda^{(0)}: \tilde{\mathfrak{g}}_D^{(0)} &\longrightarrow \mathcal{F}_{D'}^\Gamma, \\ \pi_\lambda^{(1)}: \tilde{\mathfrak{g}}_D^{(1)} &\longrightarrow \Omega_{D'}^\Gamma, \end{aligned} \quad (3.12)$$

defined explicitly as

$$\begin{aligned} \pi_\lambda^{(0)} & \left(\sum_{n=-\infty}^{M_0-1} \frac{X_0^{(n)}}{\lambda^{n+1}}, \sum_{n=-\infty}^{M_1-1} \frac{X_1^{(n)}}{(\lambda - \zeta_1)^{n+1}}, \dots, \sum_{n=-\infty}^{M_N-1} \frac{X_N^{(n)}}{(\lambda - \zeta_N)^{n+1}}, \sum_{n=-\infty}^{M_\infty} \frac{X_\infty^{(n)}}{\lambda^n} \right) \\ & = \sum_{n=0}^{M_0-1} \frac{X_0^{(n)}}{\lambda^{n+1}} + \sum_{r=1}^N \sum_{k=0}^{T-1} \sum_{n=0}^{M_r-1} \frac{\sigma^k X_r^{(n)}}{(\omega^{-k} \lambda - \zeta_r)^{n+1}} + \sum_{n=0}^{M_\infty} \frac{X_\infty^{(n)}}{\lambda^n}, \end{aligned}$$

and

$$\begin{aligned} \pi_\lambda^{(1)} & \left(\sum_{n=-\infty}^{N_0-1} \frac{Y_0^{(n)}}{\lambda^{n+1}}, \sum_{n=-\infty}^{N_1-1} \frac{Y_1^{(n)}}{(\lambda - \zeta_1)^{n+1}}, \dots, \sum_{n=-\infty}^{N_N-1} \frac{Y_N^{(n)}}{(\lambda - \zeta_N)^{n+1}}, \sum_{n=-\infty}^{N_\infty} \frac{Y_\infty^{(n)}}{\lambda^n} \right) \\ & = \sum_{n=0}^{N_0-1} \frac{Y_0^{(n)}}{\lambda^{n+1}} + \sum_{r=1}^N \sum_{k=0}^{T-1} \sum_{n=0}^{N_r-1} \frac{\omega^{-k} \sigma^k Y_r^{(n)}}{(\omega^{-k} \lambda - \zeta_r)^{n+1}} + \sum_{n=0}^{N_\infty} \frac{Y_\infty^{(n)}}{\lambda^n}. \end{aligned}$$

We will also need to define the following subspaces of $\tilde{\mathfrak{g}}_D^{(0)}$ and $\tilde{\mathfrak{g}}_D^{(1)}$, respectively:

$$\tilde{\mathfrak{g}}_{D+}^{(0)} = \tilde{\mathfrak{g}}_{0+}^{\Gamma,0} \oplus \bigoplus_{r=1}^N \tilde{\mathfrak{g}}_{r+} \oplus \tilde{\mathfrak{g}}_{\infty+}^{\Gamma,1} \quad \text{and} \quad \tilde{\mathfrak{g}}_{D+}^{(1)} = \tilde{\mathfrak{g}}_{0+}^{\Gamma,1} \oplus \bigoplus_{r=1}^N \tilde{\mathfrak{g}}_{r+} \oplus \tilde{\mathfrak{g}}_{\infty+}^{\Gamma,1},$$

where for $k = 0, 1$, we introduced (cf. (3.5) and (3.7))

$$\begin{aligned} \tilde{\mathfrak{g}}_{0+}^{\Gamma,k} & = \tilde{\mathfrak{g}}_0^{\Gamma,k} \cap \mathfrak{g} \otimes \mathbb{C}[[\lambda_0]], \\ \tilde{\mathfrak{g}}_{r+} & = \mathfrak{g} \otimes \mathbb{C}[[\lambda_r]], \quad r \in \{1, \dots, N\}, \\ \tilde{\mathfrak{g}}_{\infty+}^{\Gamma,k} & = \tilde{\mathfrak{g}}_\infty^{\Gamma,k} \cap \mathfrak{g} \otimes \lambda_\infty \mathbb{C}[[\lambda_\infty]]. \end{aligned}$$

Here we denoted by $\mathfrak{g} \otimes \mathbb{C}[[\lambda_r]]$ the algebra of formal Taylor series in λ_r with coefficients in \mathfrak{g} , for $r \neq \infty$, and by $\mathfrak{g} \otimes \lambda_\infty \mathbb{C}[[\lambda_\infty]]$ the algebra of formal Taylor series in λ_∞ with coefficients in \mathfrak{g} without constant term.

Coming back to the identification of the Lie dialgebra ingredients from Section 2, the next proposition identifies the desired decomposition (2.3) of $\mathfrak{g} = \tilde{\mathfrak{g}}_D^{(0)}$ into complementary subalgebras $\mathfrak{g}_\pm \subset \mathfrak{g}$. Explicitly, we have the identifications $\mathfrak{g}_+ = \tilde{\mathfrak{g}}_{D+}^{(0)}$ and $\mathfrak{g}_- = \iota_\lambda \mathcal{F}_{D'}^\Gamma$. We also identify a decomposition (2.4) of our model $V = \tilde{\mathfrak{g}}_D^{(1)}$ for the dual space \mathfrak{g}^* into complementary subspaces $V_\pm \subset V$, where explicitly $V_+ = \tilde{\mathfrak{g}}_{D+}^{(1)}$ and $V_- = \iota_\lambda \Omega_{D'}^\Gamma$, but will show only later in Proposition 3.4 that this decomposition of V is the desired one induced by that of \mathfrak{g} .

Proposition 3.2. *The spaces $\tilde{\mathfrak{g}}_D^{(0)}$ and $\tilde{\mathfrak{g}}_D^{(1)}$ admit the vector space decompositions*

$$\tilde{\mathfrak{g}}_D^{(0)} = \tilde{\mathfrak{g}}_{D+}^{(0)} \dot{+} \iota_\lambda \mathcal{F}_{D'}^\Gamma, \quad (3.13)$$

$$\tilde{\mathfrak{g}}_D^{(1)} = \tilde{\mathfrak{g}}_{D+}^{(1)} \dot{+} \iota_\lambda \Omega_{D'}^\Gamma, \quad (3.14)$$

respectively. Moreover, the subspaces $\tilde{\mathfrak{g}}_{D+}^{(0)}$ and $\iota_\lambda \mathcal{F}_{D'}^\Gamma$ are Lie subalgebras of $\tilde{\mathfrak{g}}_D^{(0)}$.

Proof. To any $X = (X_0, X_1, \dots, X_N, X_\infty) \in \tilde{\mathfrak{g}}_D^{(0)}$, we associate an equivariant function $f = \pi_\lambda^{(0)} X$. Notice that for all $r \in S$, $X_r - \iota_{\lambda_r} f$ is a formal Taylor series in λ_r . It follows that X splits uniquely as the direct sum of the tuples $(X_0 - \iota_{\lambda_0} f, X_1 - \iota_{\lambda_1} f, \dots, X_N - \iota_{\lambda_N} f, X_\infty - \iota_{\lambda_\infty} f) \in \tilde{\mathfrak{g}}_{D+}^{(0)}$ and $(\iota_{\lambda_0} f, \iota_{\lambda_1} f, \dots, \iota_{\lambda_N} f, \iota_{\lambda_\infty} f) \in \iota_\lambda \mathcal{F}_{D'}^\Gamma$, as required.

One can repeat the above steps (with the map $\pi_\lambda^{(1)}$ acting on some $Y \in \tilde{\mathfrak{g}}_D^{(1)}$) to prove that (3.14) defines a vector space decomposition as well. \blacksquare

Let P_+ and P_- denote the projectors onto the subspaces $\widetilde{\mathfrak{g}}_{D^+}^{(0)}$ and $\iota_\lambda \mathcal{F}_{D'}^\Gamma$, respectively, relative to the decomposition in (3.13), and \mathcal{P}_+ and \mathcal{P}_- the projectors onto $\widetilde{\mathfrak{g}}_{D^+}^{(1)}$ and $\iota_\lambda \Omega_{D'}^\Gamma$, respectively, relative to the decomposition in (3.14), in line with the notation from Section 2. As recalled in Section 2, the linear map

$$R = P_+ - P_- \quad (3.15)$$

is a solution of the mCYBE on $\widetilde{\mathfrak{g}}_D^{(0)}$. It is also useful to define the maps

$$R_\pm = \frac{1}{2}(R \pm \text{id}),$$

which can be related to the projectors onto $\widetilde{\mathfrak{g}}_{D^+}^{(0)}$ and $\iota_\lambda \mathcal{F}_{D'}^\Gamma$, relative to direct sum decomposition (3.13) as

$$R_\pm = \pm P_\pm.$$

The linear map R is related to the non-skew-symmetric r -matrix (1.8) underlying the cyclotomic Gaudin model, as one would naturally expect from the construction. More precisely, the expression (1.8) provides the kernel of the linear map (3.15) with respect to the bilinear pairing (3.10). To show this, we will use standard tensor product space notation as follows. Given any rational function $U_{\underline{12}}(\lambda, \mu)$ in the parameters λ and μ such that $\iota_\mu \iota_\lambda U_{\underline{12}}, \iota_\lambda \iota_\mu U_{\underline{12}} \in \widetilde{\mathfrak{g}}_D^{(0)} \otimes \widetilde{\mathfrak{g}}_D^{(1)}$, where the first tensor factor in the formally completed tensor product is the loop algebra $\widetilde{\mathfrak{g}}_D^{(0)}$ with the loop parameter λ , and the second tensor factor is the space $\widetilde{\mathfrak{g}}_D^{(1)}$ with the loop parameter μ , we introduce the following shorthand notations:

$$\begin{aligned} \langle \iota_\mu \iota_\lambda U_{\underline{12}}, X_{\underline{2}} \rangle_{\underline{2}} &= T \sum_{r=1}^N \text{Res}_{\mu_r=0} \text{Tr}_{\underline{2}}(\iota_{\mu_r} \iota_\lambda U_{\underline{12}}(\lambda, \mu) X_{r_{\underline{2}}}(\mu_r)) d\mu \\ &\quad + \sum_{r \in \{0, \infty\}} \text{Res}_{\mu_r=0} \text{Tr}_{\underline{2}}(\iota_{\mu_r} \iota_\lambda U_{\underline{12}}(\lambda, \mu) X_{r_{\underline{2}}}(\mu_r)) d\mu, \end{aligned}$$

and

$$\begin{aligned} \langle \iota_\lambda \iota_\mu U_{\underline{12}}, X_{\underline{2}} \rangle_{\underline{2}} &= T \sum_{r=1}^N \text{Res}_{\mu_r=0} \text{Tr}_{\underline{2}}(\iota_\lambda \iota_{\mu_r} U_{\underline{12}}(\lambda, \mu) X_{r_{\underline{2}}}(\mu_r)) d\mu \\ &\quad + \sum_{r \in \{0, \infty\}} \text{Res}_{\mu_r=0} \text{Tr}_{\underline{2}}(\iota_\lambda \iota_{\mu_r} U_{\underline{12}}(\lambda, \mu) X_{r_{\underline{2}}}(\mu_r)) d\mu \end{aligned}$$

for any $X \in \widetilde{\mathfrak{g}}_D^{(0)}$, where the parameters μ_r are defined analogously to the parameters λ_r in (3.3), and the linear map ι_μ is defined as in (3.11), returning the tuple of formal Laurent expansions in μ_r , $r \in S$.

Proposition 3.3. *For all $X \in \widetilde{\mathfrak{g}}_D^{(0)}$, the linear maps R_+ and R_- satisfy*

$$R_+(X) = \langle \iota_\mu \iota_\lambda r_{\underline{12}}, X_{\underline{2}} \rangle_{\underline{2}} \quad \text{and} \quad R_-(X) = \langle \iota_\lambda \iota_\mu r_{\underline{12}}, X_{\underline{2}} \rangle_{\underline{2}}, \quad (3.16)$$

respectively, where

$$r_{\underline{12}}(\lambda, \mu) = \frac{1}{T} \sum_{k=0}^{T-1} \frac{\sigma_1^k C_{\underline{12}}}{\mu - \omega^{-k} \lambda}.$$

Proof. In what follows, to treat the origin on the same footing as the points ζ_r , $r \in \{1, \dots, N\}$, it will be convenient to introduce the notation $\zeta_0 = 0$. We have

$$\begin{aligned} & \iota_{\lambda_s} \left(\frac{1}{T} \sum_{k=0}^{T-1} \frac{\sigma_{\mathbf{1}}^k C_{\mathbf{12}}}{\mu - \omega^{-k} \lambda} \right) \\ &= \begin{cases} \frac{1}{T} \sum_{k=0}^{T-1} \sum_{m=0}^{\infty} \frac{\omega^{-km} (\lambda - \zeta_s)^m \sigma_{\mathbf{2}}^{-k} C_{\mathbf{12}}}{(\mu - \omega^{-k} \zeta_s)^{m+1}} & \text{for } s \in \{0, 1, \dots, N\}, \\ -\frac{1}{T} \sum_{k=0}^{T-1} \sum_{m=0}^{\infty} \frac{\omega^{k(m+1)} \mu^m \sigma_{\mathbf{2}}^{-k} C_{\mathbf{12}}}{\lambda^{m+1}} & \text{for } s = \infty, \end{cases} \end{aligned} \quad (3.17)$$

and

$$\begin{aligned} & \iota_{\mu_r} \left(\frac{1}{T} \sum_{k=0}^{T-1} \frac{\sigma_{\mathbf{1}}^k C_{\mathbf{12}}}{\mu - \omega^{-k} \lambda} \right) \\ &= \begin{cases} -\frac{1}{T} \sum_{k=0}^{T-1} \sum_{m=0}^{\infty} \frac{\omega^{k(m+1)} (\mu - \zeta_r)^m \sigma_{\mathbf{1}}^k C_{\mathbf{12}}}{(\lambda - \omega^k \zeta_r)^{m+1}} & \text{for } r \in \{0, 1, \dots, N\}, \\ \frac{1}{T} \sum_{k=0}^{T-1} \sum_{m=0}^{\infty} \frac{\omega^{-km} \lambda^m \sigma_{\mathbf{1}}^k C_{\mathbf{12}}}{\mu^{m+1}} & \text{for } r = \infty. \end{cases} \end{aligned} \quad (3.18)$$

It follows that both $\iota_{\mu} \iota_{\lambda} r_{\mathbf{12}}$ and $\iota_{\lambda} \iota_{\mu} r_{\mathbf{12}}$ are elements of $\tilde{\mathfrak{g}}_D^{(0)} \otimes \tilde{\mathfrak{g}}_D^{(1)}$ where the loop parameter λ is in the first tensor factor and μ in the second. Now, pick an arbitrary $X \in \tilde{\mathfrak{g}}_D^{(0)}$ and let $X = W + Z$ be its decomposition relative to (3.13) where the two components can be written explicitly as

$$W = \left(\sum_{n=0}^{\infty} W_0^{(n)} \lambda^n, \sum_{n=0}^{\infty} W_1^{(n)} (\lambda - \zeta_1)^n, \dots, \sum_{n=0}^{\infty} W_N^{(n)} (\lambda - \zeta_N)^n, \sum_{n=1}^{\infty} W_{\infty}^{(n)} \lambda^n \right) \in \tilde{\mathfrak{g}}_{D+}^{(0)},$$

and

$$Z = \iota_{\lambda} \left(\sum_{n=0}^{N_0-1} \frac{Z_0^{(n)}}{\lambda^{n+1}} + \sum_{r=1}^N \sum_{k=0}^{T-1} \sum_{n=0}^{N_r-1} \frac{\sigma^k Z_r^{(n)}}{(\omega^{-k} \lambda - \zeta_r)^{n+1}} + \sum_{n=0}^{N_{\infty}} \frac{Z_{\infty}^{(n)}}{\lambda^n} \right) \in \iota_{\lambda} \mathcal{F}_{D'}^{\Gamma}.$$

Using the expansion (3.17), we get

$$\begin{aligned} & T \sum_{r=1}^N \text{Res}_{\mu_r=0} \text{Tr}_{\mathbf{2}} (\iota_{\mu_r} \iota_{\lambda_s} r_{\mathbf{12}}(\lambda, \mu) W_{r\mathbf{2}}(\mu_r)) d\mu \\ & \quad + \sum_{r \in \{0, \infty\}} \text{Res}_{\mu_r=0} \text{Tr}_{\mathbf{2}} (\iota_{\mu_r} \iota_{\lambda_s} r_{\mathbf{12}}(\lambda, \mu) W_{r\mathbf{2}}(\mu_r)) d\mu \\ &= \sum_{m=0}^{\infty} W_s^{(m)} (\lambda - \zeta_s)^m, \quad \text{when } s \in \{0, 1, \dots, N\}, \end{aligned}$$

and

$$\begin{aligned} & T \sum_{r=1}^N \text{Res}_{\mu_r=0} \text{Tr}_{\mathbf{2}} (\iota_{\mu_r} \iota_{\lambda_s} r_{\mathbf{12}}(\lambda, \mu) W_{r\mathbf{2}}(\mu_r)) d\mu \\ & \quad + \sum_{r \in \{0, \infty\}} \text{Res}_{\mu_r=0} \text{Tr}_{\mathbf{2}} (\iota_{\mu_r} \iota_{\lambda_s} r_{\mathbf{12}}(\lambda, \mu) W_{r\mathbf{2}}(\mu_r)) d\mu \\ &= \sum_{m=1}^{\infty} W_{\infty}^{(m)} \lambda^m, \quad \text{when } s = \infty. \end{aligned}$$

So, we find that

$$\langle \iota_\mu \iota_\lambda r_{\underline{12}}, W_{\underline{2}} \rangle_{\underline{2}} = W. \quad (3.19)$$

To evaluate $\langle \iota_\mu \iota_\lambda r_{\underline{12}}, Z_{\underline{2}} \rangle_{\underline{2}}$, we note that for an arbitrary equivariant rational function $f \in \mathcal{F}_D^\Gamma$ and equivariant rational one-form $g \in \Omega_{D'}^\Gamma$, we have the following relation:

$$\begin{aligned} \operatorname{Res}_{\lambda_r=0} \operatorname{Tr}(g(\lambda)f(\lambda))d\lambda &= \operatorname{Res}_{\lambda_r=0} \operatorname{Tr}(\sigma^k(g(\lambda))\sigma^k(f(\lambda)))d\lambda \\ &= \operatorname{Res}_{\lambda_r=0} \operatorname{Tr}(\omega^k g(\omega^k \lambda) f(\omega^k \lambda))d\lambda \\ &= \operatorname{Res}_{\lambda_r, k=0} \operatorname{Tr}(g(\lambda)f(\lambda))d\lambda \end{aligned} \quad (3.20)$$

for all $k \in \{0, \dots, T-1\}$. This allows us to rewrite each entry of the tuple $\langle \iota_\mu \iota_\lambda r_{\underline{12}}, Z_{\underline{2}} \rangle_{\underline{2}}$ as

$$\begin{aligned} & T \sum_{r=1}^N \operatorname{Res}_{\mu_r=0} \operatorname{Tr}_{\underline{2}}(\iota_{\mu_r} \iota_{\lambda_s} r_{\underline{12}}(\lambda, \mu) Z_{r\underline{2}}(\mu_r)) d\mu \\ & \quad + \sum_{r \in \{0, \infty\}} \operatorname{Res}_{\mu_r=0} \operatorname{Tr}_{\underline{2}}(\iota_{\mu_r} \iota_{\lambda_s} r_{\underline{12}}(\lambda, \mu) Z_{r\underline{2}}(\mu_r)) d\mu \\ & = \sum_{r=1}^N \sum_{k=0}^{T-1} \operatorname{Res}_{\mu_r, k=0} \operatorname{Tr}_{\underline{2}}(\iota_{\mu_r} \iota_{\lambda_s} r_{\underline{12}}(\lambda, \mu) Z_{r\underline{2}}(\mu_r)) d\mu \\ & \quad + \sum_{r \in \{0, \infty\}} \operatorname{Res}_{\mu_r=0} \operatorname{Tr}_{\underline{2}}(\iota_{\mu_r} \iota_{\lambda_s} r_{\underline{12}}(\lambda, \mu) Z_{r\underline{2}}(\mu_r)) d\mu, \quad s \in S, \end{aligned}$$

which is a sum over all the residues of a meromorphic one-form on $\mathbb{C}P^1$. Hence, we deduce

$$\langle \iota_\mu \iota_\lambda r_{\underline{12}}, Z_{\underline{2}} \rangle_{\underline{2}} = 0. \quad (3.21)$$

From (3.19) and (3.21), we then have

$$\langle \iota_\mu \iota_\lambda r_{\underline{12}}, X_{\underline{2}} \rangle_{\underline{2}} = W = P_+(X) = R_+(X).$$

Since $X \in \tilde{\mathfrak{g}}_D^{(0)}$ was arbitrary, this establishes the first relation in (3.16). Similarly, using the expansion (3.18), we find $\langle \iota_\lambda \iota_\mu r_{\underline{12}}, W_{\underline{2}} \rangle_{\underline{2}} = 0$ and $\langle \iota_\lambda \iota_\mu r_{\underline{12}}, Z_{\underline{2}} \rangle_{\underline{2}} = -Z$ from which we conclude that

$$\langle \iota_\lambda \iota_\mu r_{\underline{12}}, X_{\underline{2}} \rangle_{\underline{2}} = -Z = -P_-(X) = R_-(X).$$

Again, since $X \in \tilde{\mathfrak{g}}_D^{(0)}$ was arbitrary, this establishes the second relation in (3.16). \blacksquare

It follows from Proposition 3.3 that the kernel of the linear map $R = P_+ - P_-$ and that of the identity map $\operatorname{Id} = P_+ + P_-$ (with respect to the bilinear pairing of Proposition 3.1) are

$$(\iota_\mu \iota_\lambda + \iota_\lambda \iota_\mu) r_{\underline{12}}(\lambda, \mu) \quad \text{and} \quad (\iota_\mu \iota_\lambda - \iota_\lambda \iota_\mu), r_{\underline{12}}(\lambda, \mu),$$

respectively.

Recall that in Proposition 3.2 we identified the Lax matrix (1.9) of the cyclotomic Gaudin model, or rather its image under the embedding ι_λ in (3.11), as living in the subspace $V_- = \iota_\lambda \Omega_{D'}^\Gamma$ of our model $V = \tilde{\mathfrak{g}}_D^{(1)}$ for the dual space \mathfrak{g}^* . The next proposition establishes that this subspace $V_- \subset V$ is the orthogonal complement of $\iota_\lambda \mathcal{F}_{D'}^\Gamma \subset \tilde{\mathfrak{g}}_D^{(0)}$, i.e., $\mathfrak{g}_- \subset \mathfrak{g}$, with respect to the nondegenerate bilinear pairing of Proposition 3.1. This completes the proof of the claim that the decomposition $V = V_+ \dot{+} V_-$ obtained in Proposition 3.2 is the one induced from the corresponding decomposition of the Lie algebra $\mathfrak{g} = \mathfrak{g}_+ \dot{+} \mathfrak{g}_-$.

Proposition 3.4. *An element $Y \in \tilde{\mathfrak{g}}_D^{(1)}$ lies in $\iota_\lambda \Omega_{D'}^\Gamma$ if and only if $\langle Y, X \rangle = 0$ for all $X \in \iota_\lambda \mathcal{F}_{D'}^\Gamma$. Moreover, $Y \in \tilde{\mathfrak{g}}_D^{(1)}$ lies in $\tilde{\mathfrak{g}}_{D+}^{(1)}$ if and only if $\langle Y, X \rangle = 0$ for all $X \in \tilde{\mathfrak{g}}_{D+}^{(0)}$.*

Proof. This is a particular case of the Γ -equivariant strong residue theorem [41, Appendix A]. We recall the proof here in the present setting for completeness. We start by proving that if $Y \in \iota_\lambda \Omega_{D'}^\Gamma$, then $\langle Y, X \rangle = 0$ for all $X \in \iota_\lambda \mathcal{F}_{D'}^\Gamma$. Let us pick elements $f \in \mathcal{F}_{D'}^\Gamma$ and $g \in \Omega_{D'}^\Gamma$, and let $X = \iota_\lambda f$ and $Y = \iota_\lambda g$. Since $g(\lambda)f(\lambda)d\lambda$ is a meromorphic one-form on $\mathbb{C}P^1$, by the residue theorem we have

$$\sum_{r=1}^N \sum_{k=0}^{T-1} \text{Res}_{\lambda_{r,k}=0} \text{Tr}(\iota_{\lambda_{r,k}} g(\lambda) \iota_{\lambda_{r,k}} f(\lambda)) d\lambda + \sum_{r \in \{0, \infty\}} \text{Res}_{\lambda_r=0} \text{Tr}(\iota_{\lambda_r} g(\lambda) \iota_{\lambda_r} f(\lambda)) d\lambda = 0.$$

Using the relation (3.20) valid for any $f \in \mathcal{F}_{D'}^\Gamma$ and $g \in \Omega_{D'}^\Gamma$, we may rewrite the first term on the left-hand side as T times a sum over the residues at the points ζ_r for $r \in S$, so that

$$T \sum_{r=1}^N \text{Res}_{\lambda_r=0} \text{Tr}(\iota_{\lambda_r} g(\lambda) \iota_{\lambda_r} f(\lambda)) d\lambda + \sum_{r \in \{0, \infty\}} \text{Res}_{\lambda_r=0} \text{Tr}(\iota_{\lambda_r} g(\lambda) \iota_{\lambda_r} f(\lambda)) d\lambda = 0. \quad (3.22)$$

By definition of the bilinear pairing from Proposition 3.1, we thus have $\langle Y, X \rangle = 0$, as desired.

Let us now prove the converse. Namely, let $Y \in \tilde{\mathfrak{g}}_D^{(1)}$ be arbitrary and suppose that $\langle Y, X \rangle = 0$ for all $X \in \iota_\lambda \mathcal{F}_{D'}^\Gamma$. We must show that, in fact, $Y(\lambda) \in \iota_\lambda \Omega_{D'}^\Gamma$. It follows from Proposition 3.2 that $Y = (Y_0, Y_1, \dots, Y_N, Y_\infty) \in \tilde{\mathfrak{g}}_D^{(1)}$ has a unique decomposition as a direct sum of the tuples $\mathcal{P}_+(Y) = (\mathcal{P}_{0+}(Y_0), \mathcal{P}_{1+}(Y_1), \dots, \mathcal{P}_{N+}(Y_N), \mathcal{P}_{\infty+}(Y_\infty)) \in \tilde{\mathfrak{g}}_{D+}^{(1)}$ and $\mathcal{P}_-(Y) = \iota_\lambda g$ for some $g \in \Omega_{D'}^\Gamma$. Now, let $X = \iota_\lambda f$ for some $f \in \mathcal{F}_{D'}^\Gamma$. Since $\langle Y, X \rangle = 0$, writing this out explicitly means

$$\begin{aligned} T \sum_{r=1}^N \text{Res}_{\lambda_r=0} \text{Tr}(\mathcal{P}_{r+}(Y_r(\lambda_r)) \iota_{\lambda_r} f(\lambda)) d\lambda + \sum_{r \in \{0, \infty\}} \text{Res}_{\lambda_r=0} \text{Tr}(\mathcal{P}_{r+}(Y_r(\lambda_r)) \iota_{\lambda_r} f(\lambda)) d\lambda \\ + T \sum_{r=1}^N \text{Res}_{\lambda_r=0} \text{Tr}(\iota_{\lambda_r} g(\lambda) \iota_{\lambda_r} f(\lambda)) d\lambda + \sum_{r \in \{0, \infty\}} \text{Res}_{\lambda_r=0} \text{Tr}(\iota_{\lambda_r} g(\lambda) \iota_{\lambda_r} f(\lambda)) d\lambda = 0. \end{aligned}$$

From (3.22), we have that the last two terms on the left-hand side vanish on their own. Therefore, we get

$$T \sum_{r=1}^N \text{Res}_{\lambda_r=0} \text{Tr}(\mathcal{P}_{r+}(Y_r(\lambda_r)) \iota_{\lambda_r} f(\lambda)) d\lambda + \sum_{r \in \{0, \infty\}} \text{Res}_{\lambda_r=0} \text{Tr}(\mathcal{P}_{r+}(Y_r(\lambda_r)) \iota_{\lambda_r} f(\lambda)) d\lambda = 0.$$

One can then show, along the same lines as the argument in the proof of Proposition 3.1, that if $\mathcal{P}_+(Y) \neq 0$ then by picking a suitable $f \in \mathcal{F}_{D'}^\Gamma$ adapted to $\mathcal{P}_+(Y) \in \tilde{\mathfrak{g}}_{D+}^{(1)}$, one can ensure that the expression on the left-hand side is non-zero, which is a contradiction. Therefore, we conclude that $\mathcal{P}_+(Y) = 0$, and hence $Y \in \iota_\lambda \Omega_{D'}^\Gamma$, as required. The proof of the ‘‘moreover’’ part is completely analogous. ■

The upshot of Proposition 3.4 is that we are now exactly in the setting recalled in points 1 and 2 of Section 2. In particular, the coadjoint representation $\text{Ad}^{R*}: \mathbb{G}_R \times V \rightarrow V$ of the group \mathbb{G}_R on our model $V = \tilde{\mathfrak{g}}_D^{(1)}$ for the dual space \mathfrak{g}^* is given by formula (2.5). This will be used below in Lemma 3.5 to give a concise description of the desired coadjoint orbit for the Lax matrix of the cyclotomic Gaudin model.

Before stating the lemma, we first give an explicit description of the infinite-dimensional Lie group $\mathbb{G}_+ = \tilde{G}_{D+}^{(0)}$ associated with the Lie algebra $\mathfrak{g}_+ = \tilde{\mathfrak{g}}_{D+}^{(0)}$. Its elements are of the form

$$\varphi_+ = (\varphi_{0+}, \varphi_{1+}, \dots, \varphi_{N+}, \varphi_{\infty+}), \quad (3.23)$$

where $\varphi_{r+}(\lambda_r)$ is a Taylor series in the local parameter λ_r with values in G , the matrix Lie group associated with the Lie algebra \mathfrak{g} ,

$$\begin{aligned} \varphi_{r+}(\lambda_r) &= \sum_{n=0}^{\infty} \phi_r^{(n)} \lambda_r^n, \quad r \neq \infty, \\ \varphi_{\infty+}(\lambda_\infty) &= \mathbf{1} + \sum_{n=1}^{\infty} \phi_\infty^{(n)} \lambda_\infty^n. \end{aligned} \quad (3.24)$$

The Lie group $\mathbb{G}_+ = \tilde{G}_{D+}^{(0)}$ has a natural action on $V = \tilde{\mathfrak{g}}_D^{(1)}$ given by conjugation. This defines the coadjoint representation $\text{Ad}^*: \mathbb{G}_+ \times V \rightarrow V$ from Section 2 explicitly as

$$(\varphi_+, Y) \mapsto \text{Ad}_{\varphi_+}^* Y = (\varphi_{0+} Y_0 \varphi_{0+}^{-1}, \varphi_{1+} Y_1 \varphi_{1+}^{-1}, \dots, \varphi_{N+} Y_N \varphi_{N+}^{-1}, \varphi_{\infty+} Y_\infty \varphi_{\infty+}^{-1}).$$

We are now in a position to give an explicit description of the \mathbb{G}_R -coadjoint orbit where the Lax matrix of the cyclotomic Gaudin model lives and which will act as our phase space.

Lemma 3.5. *The orbit of the coadjoint action of $\mathbb{G}_R = (\tilde{G}_D^{(0)})_R$ on an element $\iota_\lambda \Lambda \in V_- = \iota_\lambda \Omega_D^\Gamma$ has the explicit form*

$$\mathcal{O}_\Lambda = \{ \mathcal{P}_-(\text{Ad}_{\varphi_+}^* \iota_\lambda \Lambda) \mid \varphi_+ \in \tilde{G}_{D+}^{(0)} \}.$$

Proof. By definition, the coadjoint orbit of $(\tilde{G}_D^{(0)})_R$ on any $\iota_\lambda \Lambda \in \iota_\lambda \Omega_D^\Gamma$ is given by

$$\mathcal{O}_\Lambda = \{ \text{Ad}_{\varphi}^{R*} \iota_\lambda \Lambda \mid \varphi \in (\tilde{G}_D^{(0)})_R \}.$$

Using the explicit form (2.5) for the coadjoint action of \mathbb{G}_R , we have

$$\text{Ad}_{\varphi}^{R*} \iota_\lambda \Lambda = \mathcal{P}_-(\text{Ad}_{\varphi_+}^* \iota_\lambda \Lambda) + \mathcal{P}_+(\text{Ad}_{\varphi_-}^* \iota_\lambda \Lambda) = \mathcal{P}_-(\text{Ad}_{\varphi_+}^* \iota_\lambda \Lambda),$$

where the last equality follows from the fact that $\iota_\lambda \Lambda \in V_- = \iota_\lambda \Omega_D^\Gamma$, so that $\text{Ad}_{\varphi_-}^* \iota_\lambda \Lambda \in V_-$ also and hence $\mathcal{P}_+(\text{Ad}_{\varphi_-}^* \iota_\lambda \Lambda) = 0$. \blacksquare

It will be useful in practice to express the action of the projector \mathcal{P}_- on $Y \in \tilde{\mathfrak{g}}_D^{(1)}$ as

$$\mathcal{P}_-(Y) = \iota_\lambda \circ \pi_\lambda^{(1)}(Y). \quad (3.25)$$

In the remaining sections, we will put this setup to use to describe the Lax matrix of the cyclotomic Gaudin model (1.9) as a point in a coadjoint orbit \mathcal{O}_Λ for some suitable $\Lambda \in \Omega_D^\Gamma$, and then introduce the ingredients from point 3 in Section 2 to derive the associated Lax equations.

3.2 Lax matrix

Our algebraic setup covers the case of the cyclotomic Gaudin model with arbitrary multiplicities. However, as mentioned at the start of this section, from now on we will restrict to the case with simple poles at all $\omega^k \zeta_r$, $k \in \{0, \dots, T-1\}$, $r \in \{1, \dots, N\}$, and double poles at the origin and at infinity, since this is the setting required for our examples in Section 4. The discussion in the remaining sections is easily generalised to the case of arbitrary multiplicities.

To describe a coadjoint orbit $\mathcal{O}_\Lambda \in \iota_\lambda \Omega_{D'}^\Gamma$, from Lemma 3.5 corresponding to the Lax matrix (1.9) of the cyclotomic Gaudin model, we fix a non-dynamical element $\Lambda \in \Omega_{D'}^\Gamma$ with the same pole structure as (1.9), namely we introduce

$$\Lambda(\lambda) = \frac{\Lambda_0^{(0)}}{\lambda} + \frac{\Lambda_0^{(1)}}{\lambda^2} + \frac{1}{T} \sum_{r=1}^N \sum_{k=0}^{T-1} \frac{\sigma^k \Lambda_r}{\lambda - \omega^k \zeta_r} + \Lambda_\infty \in \Omega_{D'}^\Gamma.$$

According to Lemma 3.5 and formula (3.25), the corresponding coadjoint orbit \mathcal{O}_Λ then consists of elements of the form

$$\begin{aligned} \mathcal{P}_- \left(\varphi_+ \iota_\lambda \left(\frac{\Lambda_0^{(0)}}{\lambda} + \frac{\Lambda_0^{(1)}}{\lambda^2} + \frac{1}{T} \sum_{r=1}^N \sum_{k=0}^{T-1} \frac{\sigma^k \Lambda_r}{\lambda - \omega^k \zeta_r} + \Lambda_\infty \right) \varphi_+^{-1} \right) \\ = \iota_\lambda \circ \pi_\lambda^{(1)} \left(\frac{A_0^{(0)}}{\lambda} + \frac{A_0^{(1)}}{\lambda^2}, \frac{1}{T} \frac{A_1}{\lambda - \zeta_1}, \dots, \frac{1}{T} \frac{A_N}{\lambda - \zeta_N}, A_\infty \right) \\ = \iota_\lambda \left(\frac{A_0^{(0)}}{\lambda} + \frac{A_0^{(1)}}{\lambda^2} + \frac{1}{T} \sum_{r=1}^{T-1} \sum_{k=0}^{T-1} \frac{\sigma^k A_r}{\lambda - \omega^k \zeta_r} + A_\infty \right) \\ \equiv \iota_\lambda L(\lambda), \end{aligned} \tag{3.26}$$

where, recalling the definitions (3.23) and (3.24), we have set

$$\begin{aligned} A_0^{(0)} &= \phi_0^{(0)} \Lambda_0^{(0)} \phi_0^{(0)-1} + [\phi_0^{(1)} \phi_0^{(0)-1}, \phi_0^{(0)} \Lambda_0^{(1)} \phi_0^{(0)-1}], \\ A_0^{(1)} &= \phi_0^{(0)} \Lambda_0^{(1)} \phi_0^{(0)-1}, \\ A_r &= \phi_r \Lambda_r \phi_r^{-1}, \quad r \in \{1, \dots, N\}, \\ A_\infty &= \Lambda_\infty, \end{aligned} \tag{3.27}$$

with $\phi_r^{(0)}$ denoted by ϕ_r , for $r \in \{1, \dots, N\}$, to simplify notations. Notice that we have the relations $\sigma \phi_0^{(0)} = \phi_0^{(0)}$ and $\sigma \phi_0^{(1)} = \omega \phi_0^{(1)}$ for the field elements which ensure that $A_0^{(0)} \in \mathfrak{g}^{(0)}$ and $A_0^{(1)} \in \mathfrak{g}^{(-1)}$. This gives us the desired parameterisation of the cyclotomic Gaudin Lax matrix

$$L(\lambda) = \frac{A_0^{(0)}}{\lambda} + \frac{A_0^{(1)}}{\lambda^2} + \frac{1}{T} \sum_{r=1}^N \sum_{k=0}^{T-1} \frac{\sigma^k A_r}{\lambda - \omega^k \zeta_r} + A_\infty \tag{3.28}$$

viewed as an element of the coadjoint orbit \mathcal{O}_Λ .

3.3 Lax equations

To derive Lax equations for the Lax matrix (3.28) of the cyclotomic Gaudin model, let us return to the hierarchy of Lax equations (2.6) induced by the family of Hamiltonians in involution with respect to $\{, \}_R$. In our current setup, $\{, \}_R$ is the Lie–Poisson bracket on $\tilde{\mathfrak{g}}_D^{(1)}$ associated with the linear map $R = P_+ - P_-$. Invariant functions on $\tilde{\mathfrak{g}}_D^{(1)}$ take the form

$$H_{p,r}: Y \in \tilde{\mathfrak{g}}_D^{(1)} \longmapsto \text{Res}_{\lambda_r=0} (\ell_{p,r}(\lambda_r) \text{Tr}(Y_r(\lambda_r)^{p+1})) d\lambda, \quad p \geq 1, \quad r \in S, \tag{3.29}$$

where $\ell_{p,r}(\lambda_r) \in \mathbb{C}((\lambda_r))$ is a collection of Laurent polynomials for $p \geq 1$ and $r \in S$. It follows from Proposition 3.1 that for these functions to be non-trivial, the Laurent polynomials $\ell_{p,r}(\lambda_r)$ for $r \in \{0, \infty\}$ should be chosen such that $\ell_{p,r}(\lambda_r) Y_r(\lambda_r)^p \in \tilde{\mathfrak{g}}_r^{\Gamma,0}$, while $\ell_{p,r}(\lambda_r)$ for $r \in \{1, \dots, N\}$ can be any Laurent polynomials. Let us then choose

$$\begin{aligned} \ell_{p,r}(\lambda_r) &= \iota_{\lambda_r} \frac{\lambda^p}{p+1} \quad \text{for } r \in \{0, \infty\}, \\ \ell_{p,r}(\lambda_r) &= \iota_{\lambda_r} \frac{T \lambda^p}{p+1} \quad \text{for } r \in \{1, \dots, N\}. \end{aligned} \tag{3.30}$$

The restriction of the functions $H_{p,r}$ to $\iota_\lambda L$ are Hamiltonians (in involution) of the model and generate the elementary (pairwise-commuting) flows $\partial_{t_r^p}$. However, by virtue of the choice (3.30) we made, it follows from (3.20) that $\sum_{r \in S} H_{p,r}(\iota_\lambda L) = 0$ by the residue theorem, for each $p \geq 1$.

In what follows, it will thus be sufficient to focus on the Hamiltonians $H_{p,r}$ for $r \neq \infty$ and look at the associated equations of motion they produce through (2.6). Using $R_\pm = \frac{1}{2}(R \pm \text{id})$, these equations can be rewritten as

$$\partial_{t_r^p} \iota_\lambda L = [R_\pm \nabla H_{p,r}(\iota_\lambda L), \iota_\lambda L]. \quad (3.31)$$

As it suffices to calculate only one of the two expressions $R_+ \nabla H_{p,r}(\iota_\lambda L)$ and $R_- \nabla H_{p,r}(\iota_\lambda L)$, let us compute the latter. The gradient of $H_{p,r}(\iota_\lambda L)$ is an element of $\tilde{\mathfrak{g}}_D^{(0)}$ and satisfies

$$\lim_{\epsilon \rightarrow 0} \frac{H_{p,r}(\iota_\lambda L + \epsilon \eta) - H_{p,r}(\iota_\lambda L)}{\epsilon} = \langle \eta, \nabla H_{p,r}(\iota_\lambda L) \rangle \quad (3.32)$$

for all $\eta \in \tilde{\mathfrak{g}}_D^{(1)}$. Using Proposition 3.2, we may decompose this gradient as follows:

$$\nabla H_{p,r}(\iota_\lambda L) = N_r^{(p)} + \iota_\lambda h_r^{(p)}, \quad N_r^{(p)} \in \tilde{\mathfrak{g}}_{D+}^{(0)}, \quad h_r^{(p)} \in \mathcal{F}_{D'}^\Gamma,$$

and rewrite (3.31) as

$$\partial_{t_r^p} \iota_\lambda L = [R_- \nabla H_{p,r}(\iota_\lambda L), \iota_\lambda L] = -[P_- (\nabla H_{p,r}(\iota_\lambda L)), \iota_\lambda L] = -[\iota_\lambda h_r^{(p)}, \iota_\lambda L].$$

Since ι_λ is an embedding which also clearly commutes with the Lie brackets $[\cdot, \cdot]: \mathcal{F}_{D'}^\Gamma \times \Omega_{D'}^\Gamma \rightarrow \Omega_{D'}^\Gamma$ and $[\cdot, \cdot]: \tilde{\mathfrak{g}}_D^{(0)} \times \tilde{\mathfrak{g}}_D^{(1)} \rightarrow \tilde{\mathfrak{g}}_D^{(1)}$, the above equation implies

$$\partial_{t_r^p} L = -[h_r^{(p)}, L]. \quad (3.33)$$

In order to calculate $h_r^{(p)} \in \mathcal{F}_{D'}^\Gamma$, by virtue of Proposition 3.4, it is sufficient to restrict η in (3.32) to live in $\tilde{\mathfrak{g}}_{D+}^{(1)}$. We then have

$$\langle \eta, N_r^{(p)} \rangle = 0 \quad \text{for all } \eta \in \tilde{\mathfrak{g}}_{D+}^{(1)}, \quad N_r^{(p)} \in \tilde{\mathfrak{g}}_{D+}^{(0)},$$

and the right-hand side of (3.32) becomes

$$T \sum_{s=1}^N \text{Res}_{\lambda_s=0} \text{Tr}(\eta_s(\lambda_s) \iota_{\lambda_s} h_r^{(p)}) + \sum_{s \in \{0, \infty\}} \text{Res}_{\lambda_s=0} \text{Tr}(\eta_s(\lambda_s) \iota_{\lambda_s} h_r^{(p)}),$$

while for the left-hand side we find

$$\lim_{\epsilon \rightarrow 0} \frac{H_{p,r}(\iota_\lambda L + \epsilon \eta) - H_{p,r}(\iota_\lambda L)}{\epsilon} = (p+1) \text{Res}_{\lambda_r=0} \text{Tr}(\ell_{p,r}(\lambda_r) \eta_r(\lambda_r) \iota_{\lambda_r} L^p), \quad r \in S,$$

for any $\eta_s(\lambda_s) \in \tilde{\mathfrak{g}}_{s+}$, $s \in \{1, \dots, N\}$, and $\eta_s(\lambda_s) \in \tilde{\mathfrak{g}}_{s+}^{\Gamma,1}$, $s \in \{0, \infty\}$. By definition of $\ell_{p,r}(\lambda_r)$ in (3.30), this implies

$$(\iota_{\lambda_s} h_r^{(p)})_- = \begin{cases} 0 & \text{for } s \neq r, \\ (\iota_{\lambda_r} \lambda^p \iota_{\lambda_r} L^p)_- & \text{for } s = r \end{cases}$$

for all $r \in S$, where X_- denotes the principal part of a Laurent series X . Plugging the equivariant functions $h_r^{(p)}$ obtained from the above conditions into (3.33) gives a hierarchy of Lax equations corresponding to our choice of invariant functions $H_{p,r}$ in (3.29).

Explicitly, for $p = 1$, we get the Lax equations

$$\partial_{t_1^r} L = -[h_r^{(1)}, L] \quad \text{with} \quad h_r^{(1)} = \begin{cases} \frac{A_0^{(1)}}{\lambda} & \text{for } r = 0, \\ \frac{1}{T} \sum_{k=0}^{T-1} \frac{\omega^k \zeta_r \sigma^k A_r}{\lambda - \omega^k \zeta_r} & \text{for } r \in \{1, \dots, N\}. \end{cases} \quad (3.34)$$

Finally, we turn to the primary goal of this paper which is to give a Lagrangian multiform for the cyclotomic Gaudin model that will provide a variational description of the hierarchy of Lax equations in (3.33). Having just described the cyclotomic Gaudin model within the Lie dialgebra framework, we now have all the necessary ingredients to achieve this goal: the non-dynamical element $\Lambda \in \Omega_D^\Gamma$ that fixes the phase space, the field element $\varphi_+ \in \widehat{G}_{D^+}^{(0)}$ containing the dynamical degrees of freedom, the linear map R that equips the phase space with the required Poisson structure, and the invariant functions $H_{p,r}$ which induce non-trivial equations of motion with respect to this Poisson structure.

3.4 Lagrangian multiform for the cyclotomic Gaudin hierarchy

We can now define a Lagrangian 1-form on the coadjoint orbit \mathcal{O}_Λ of the cyclotomic Gaudin model as

$$\mathcal{L} = \sum_{p=1}^M \sum_{r \in S \setminus \{\infty\}} \mathcal{L}_{p,r} dt_p^r,$$

using the expression (2.8) for the Lagrangian coefficients. It will be useful to recall the notations in (3.27) associated with the parameterisation of the cyclotomic Gaudin Lax matrix. The elementary times t_k that appear in (2.8) are now naturally labelled by a pair of indices $p \geq 1$ and $r \in S$, namely we now have elementary times t_p^r , associated with the corresponding Hamiltonians (3.29). Explicitly, we have the following.

Theorem 3.6. *The Lagrangian coefficients of the cyclotomic Gaudin multiform \mathcal{L} take the form*

$$\mathcal{L}_{p,r} = \sum_{s=1}^N \text{Tr}(A_s \partial_{t_p^r} \phi_s \phi_s^{-1}) + \text{Tr}(A_0^{(0)} \partial_{t_p^r} \phi_0^{(0)} \phi_0^{(0)-1}) - H_{p,r}(\iota_\lambda L), \quad (3.35)$$

with the potential part

$$H_{p,r}(\iota_\lambda L) = \text{Res}_{\lambda_r=0}(\ell_{p,r}(\lambda_r) \text{Tr}(\iota_{\lambda_r} L^{p+1})) d\lambda, \quad r \in \{0, 1, \dots, N\}, \quad (3.36)$$

where $\ell_{p,r}(\lambda_r)$ are the Laurent polynomials

$$\ell_{p,0} = \iota_{\lambda_0} \frac{\lambda^p}{p+1} \quad \text{and} \quad \ell_{p,r} = \iota_{\lambda_r} \frac{T \lambda^p}{p+1} \quad \text{for } r \in \{1, \dots, N\}.$$

The Lagrangian 1-form \mathcal{L} satisfies the corner equations (1.2)–(1.3) of the multi-time Euler–Lagrange equations, while the standard Euler–Lagrange equations for $\mathcal{L}_{p,r}$ give the hierarchy of Lax equations in (3.33). Further, on solutions of (3.33), we have the closure relation

$$\partial_{t_q^s} \mathcal{L}_{p,r} - \partial_{t_p^r} \mathcal{L}_{q,s} = 0$$

for all possible combinations of (p, r) and (q, s) in $\mathbb{Z}_{\geq 1} \times S$.

Proof. Let us start by reinterpreting the formula in (2.8) in the present context of the cyclotomic Gaudin model. First, note that on the coadjoint orbit \mathcal{O}_Λ , where the Lagrangian 1-form \mathcal{L} lives, the role of the Lax matrix is played by the image of L in $\widetilde{\mathfrak{g}}_D^{(1)}$ under the embedding ι_λ , given by (3.26). Next, the bilinear pairing used to define the kinetic part is the one constructed in Proposition 3.1. Furthermore, in the Adler–Kostant–Symes scheme where (locally) $\mathbb{G}_R \simeq \mathbb{G}_+ \times \mathbb{G}_-$, we explicitly have $\partial_{t_p^r} \varphi \cdot_R \varphi^{-1} = \partial_{t_p^r} \varphi_+ \varphi_+^{-1} + \partial_{t_p^r} \varphi_- \varphi_-^{-1}$, and since $\iota_\lambda L \in V_- = \iota_\lambda \Omega_{D'}^\Gamma$, it follows from Proposition 3.4 that only the $\partial_{t_p^r} \varphi_+ \varphi_+^{-1}$ piece contributes to the kinetic term. Finally, the potential part is simply the restriction of the invariant functions $H_{p,r}$ in (3.29) to $\iota_\lambda L$. Therefore, in the present setup, the Lagrangian coefficients in (2.8) can be expressed as

$$\begin{aligned} \mathcal{L}_{p,r} &= \langle \iota_\lambda L, \partial_{t_p^r} \varphi_+ \varphi_+^{-1} \rangle - H_{p,r}(\iota_\lambda L) \\ &= \langle \iota_\lambda \Lambda, \varphi_+^{-1} \partial_{t_p^r} \varphi_+ \rangle - H_{p,r}(\iota_\lambda L), \quad r \in \{0, 1, \dots, N\}, \end{aligned} \quad (3.37)$$

where in the second step we used the fact (3.26) that $\iota_\lambda L = \mathcal{P}_-(\varphi_+(\iota_\lambda \Lambda) \varphi_+^{-1})$. The kinetic term can be written out more explicitly in terms of the component fields (3.24) as

$$\begin{aligned} \langle \iota_\lambda \Lambda, \varphi_+^{-1} \partial_{t_p^r} \varphi_+ \rangle &= T \sum_{s=1}^N \operatorname{Res}_{\lambda_s=0} \operatorname{Tr}(\iota_{\lambda_s} \Lambda \varphi_{s+}(\lambda_s)^{-1} \partial_{t_p^r} \varphi_{s+}(\lambda_s)) d\lambda \\ &\quad + \sum_{s \in \{0, \infty\}} \operatorname{Res}_{\lambda_s=0} \operatorname{Tr}(\iota_{\lambda_s} \Lambda \varphi_{s+}(\lambda_s)^{-1} \partial_{t_p^r} \varphi_{s+}(\lambda_s)) d\lambda \\ &= \sum_{s=1}^N \operatorname{Tr}(A_s \partial_{t_p^r} \phi_s \phi_s^{-1}) + \operatorname{Tr}(A_0^{(0)} \partial_{t_p^r} \phi_0^{(0)} \phi_0^{(0)-1}) \\ &\quad + \partial_{t_p^r} \operatorname{Tr}(\phi_0^{(1)} \phi_0^{(0)-1} A_0^{(1)}) + \frac{1}{2} \partial_{t_p^r} \operatorname{Tr}(A_\infty (\phi_\infty^{(1)})^2). \end{aligned}$$

The last two terms associated to the poles at the origin and at infinity are total derivatives and do not contribute to the multi-time Euler–Lagrange equations since dropping them amounts to changing the Lagrangian 1-form \mathcal{L} by $d \operatorname{Tr}(\phi_0^{(1)} \phi_0^{(0)-1} A_0^{(1)} + \frac{1}{2} A_\infty (\phi_\infty^{(1)})^2)$ which is a total horizontal differential. Discarding these two terms, we are left with the required expression for the Lagrangian coefficients.

Since the Lagrangian coefficients $\mathcal{L}_{p,r}$ in (3.37) are of the form (2.8), it follows directly from Theorem 2.1 that the Lagrangian 1-form \mathcal{L} satisfies all the required conditions and the closure relation. \blacksquare

To close this section, let us present the explicit expressions for the first set of Lagrangian coefficients $\mathcal{L}_{1,r}$ with $r \in \{0, 1, \dots, N\}$. The kinetic terms are obtained by simply substituting $p = 1$ in the kinetic part in (3.35), while the potential terms defined by (3.36) read

$$\begin{aligned} H_{1,0} &= \frac{1}{2} \operatorname{Tr}(A_0^{(0)2}) - \sum_{r=1}^N \frac{\operatorname{Tr}(A_0^{(1)} A_r)}{\zeta_r} + \operatorname{Tr}(A_0^{(1)} A_\infty), \\ H_{1,r} &= \operatorname{Tr}(A_0^{(0)} A_r) + \frac{\operatorname{Tr}(A_0^{(1)} A_r)}{\zeta_r} + \frac{1}{2T} \sum_{k=0}^{T-1} \operatorname{Tr}(A_r \sigma^k A_r) \\ &\quad + \frac{1}{T} \sum_{s \neq r} \sum_{k=0}^{T-1} \frac{\operatorname{Tr}(A_r \sigma^k A_s) \zeta_r}{\zeta_r - \omega^k \zeta_s} + \operatorname{Tr}(A_r A_\infty) \zeta_r, \quad r \in \{1, \dots, N\}. \end{aligned}$$

Upon varying $\mathcal{L}_{1,r}$ with respect to ϕ_s , $s = 1, \dots, N$, $\phi_0^{(0)}$, and $\phi_0^{(1)}$, it can be checked that the associated Euler–Lagrange equations correspond to the set of Lax equations for $p = 1$ in (3.34), as it should be from Theorem 3.6.

4 Realisations of the cyclotomic Gaudin model

In this section, we study two different realisations of the cyclotomic Gaudin model – the periodic Toda chain and the discrete self-trapping (DST) model – with the objective of describing their corresponding hierarchies variationally. We will then go on to show how our framework allows for a straightforward coupling of integrable hierarchies, by using the approach devised in [15, Section 7] to couple together hierarchies of integrable field theories.

In what follows, we will work with the cyclotomic Gaudin model associated with the Lie algebra $\mathfrak{g} := \mathfrak{gl}_T(\mathbb{C})$ and the automorphism $\sigma \in \text{Aut } \mathfrak{g}$ defined by $\sigma(E_{ij}) = \omega^{j-i} E_{ij}$, for every $i, j = 1, \dots, T$. Here ω is a primitive T -th root of unity, and by E_{ij} , $i, j = 1, \dots, T$, we denote the standard basis of $\mathfrak{gl}_T(\mathbb{C})$ taking the indices i and j modulo T by convention. The eigenspaces of σ defined by (3.1) are then given by $\mathfrak{g}^{(n)} = \text{span}\{E_{i,i+n}\}_{i=1}^T$.

Let us also note the following useful identity that we shall frequently make use of

$$\frac{z_1^{T-1-[l]} z_2^{[l]}}{z_1^T - z_2^T} = \frac{1}{T} \sum_{k=0}^{T-1} \frac{\omega^{-kl}}{z_1 - \omega^k z_2} \quad (4.1)$$

for any $z_1, z_2 \in \mathbb{C}$ and $l \in \mathbb{Z}$, where $[l] \in \{0, \dots, T-1\}$ is such that $l = [l] \pmod{T}$.

4.1 Periodic Toda chain

The periodic Toda chain [39] describes a system of particles connected by “exponential springs” together with a periodic boundary condition, and has been extensively studied in the Hamiltonian formalism. See, for instance, [18] for the widely used Flaschka change of coordinates, and [3] for a proof of its integrability.

Lax matrix. We will work with the Lax matrix

$$L_{\text{Toda}}(\lambda) = \begin{pmatrix} p_1 \lambda^{-1} & 1 & 0 & \dots & a_T \lambda^{-2} \\ a_1 \lambda^{-2} & p_2 \lambda^{-1} & 1 & \dots & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & & a_{i-1} \lambda^{-2} & p_i \lambda^{-1} & 1 & 0 \\ \vdots & & & \ddots & & \vdots \\ 0 & \dots & & a_{T-2} \lambda^{-2} & p_{T-1} \lambda^{-1} & 1 \\ 1 & \dots & & 0 & a_{T-1} \lambda^{-2} & p_T \lambda^{-1} \end{pmatrix}, \quad (4.2)$$

where $a_i = e^{q_i - q_{i+1}}$, and q_i, p_i are the canonical coordinates satisfying the canonical Poisson bracket relations $\{p_i, q_j\} = \delta_{ij}$ for $i, j = 1, \dots, T$. We also have the periodic boundary conditions $(p_0, q_0) = (p_T, q_T)$ and $(p_{T+1}, q_{T+1}) = (p_1, q_1)$.

The standard Lax matrix for the periodic Toda chain (see [7, Section 6], for instance)

$$\tilde{L}_{\text{Toda}}(\lambda) = \begin{pmatrix} p_1 & a_1^{1/2} & 0 & \dots & a_T^{1/2} \lambda^{-1} \\ a_1^{1/2} & p_2 & a_2^{1/2} & \dots & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & & a_{i-1}^{1/2} & p_i & a_i^{1/2} & 0 \\ \vdots & & & \ddots & & \vdots \\ 0 & \dots & & a_{T-2}^{1/2} & p_{T-1} & a_{T-1}^{1/2} \\ a_T^{1/2} \lambda & \dots & & 0 & a_{T-1}^{1/2} & p_T \end{pmatrix}$$

is related to $L_{\text{Toda}}(\lambda)$ by conjugation by the diagonal matrix $Q = \text{diag}(e^{-q_1/2}\lambda^{-1}, \dots, e^{-q_T/2}\lambda^{-T})$ and multiplication by an overall factor of λ^{-1} , together with a change of λ -dependence, as follows:

$$L_{\text{Toda}}(\lambda) = \lambda^{-1} Q \tilde{L}_{\text{Toda}}(\lambda^T) Q^{-1}. \quad (4.3)$$

The Poisson bracket of the Lax matrix $\tilde{L}_{\text{Toda}}(\lambda)$ can be written as

$$\{\tilde{L}_{\text{Toda}_{\underline{1}}}(\lambda), \tilde{L}_{\text{Toda}_{\underline{2}}}(\mu)\} = [\tilde{r}_{\underline{1}\underline{2}}(\lambda, \mu), \tilde{L}_{\text{Toda}_{\underline{1}}}(\lambda) + \tilde{L}_{\text{Toda}_{\underline{2}}}(\mu)],$$

where $\tilde{r}_{\underline{1}\underline{2}}(\lambda, \mu)$ is the skew-symmetric r -matrix

$$\tilde{r}_{\underline{1}\underline{2}}(\lambda, \mu) = \frac{1}{2} \frac{\mu + \lambda}{\mu - \lambda} \sum_{i=1}^T E_{ii} \otimes E_{ii} + \frac{1}{\mu - \lambda} \left(\mu \sum_{i < j} + \lambda \sum_{i > j} \right) E_{ij} \otimes E_{ji}.$$

Under the gauge transformation (4.3), on using the identity (4.1), we find that the Lax matrix $L_{\text{Toda}}(\lambda)$ satisfies the Poisson bracket

$$\{L_{\text{Toda}_{\underline{1}}}(\lambda), L_{\text{Toda}_{\underline{2}}}(\mu)\} = [r_{\underline{1}\underline{2}}(\lambda, \mu), L_{\text{Toda}_{\underline{1}}}(\lambda)] - [r_{\underline{2}\underline{1}}(\mu, \lambda), L_{\text{Toda}_{\underline{2}}}(\mu)],$$

where $r_{\underline{1}\underline{2}}(\lambda, \mu)$ is the non-skew-symmetric cyclotomic r -matrix

$$r_{\underline{1}\underline{2}}(\lambda, \mu) = \frac{1}{T} \sum_{k=0}^{T-1} \frac{\omega^{k(j-i)}}{\mu - \omega^{-k}\lambda} E_{ij} \otimes E_{ji} \quad (4.4)$$

that we have been working with. This explains our choice of Lax matrix (4.2) as opposed to the more traditional one. As just proved, it satisfies the Poisson algebra of the cyclotomic $\mathfrak{gl}_T(\mathbb{C})$ -Gaudin model and therefore allows us to obtain the periodic Toda chain as a certain realisation of that model. Specifically, the Lax matrix $L_{\text{Toda}}(\lambda)$ can be seen as a realisation of the cyclotomic Gaudin Lax matrix with double poles at the origin and at infinity, that is,

$$L_{\text{Toda}}(\lambda) = \frac{J_0^{(0)}}{\lambda} + \frac{J_0^{(1)}}{\lambda^2} + J_\infty, \quad (4.5)$$

where

$$J_0^{(0)} = \sum_{i=1}^T p_i E_{ii}, \quad J_0^{(1)} = \sum_{i=1}^T e^{q_i - q_{i+1}} E_{i+1, i}, \quad J_\infty = \sum_{i=1}^T E_{i, i+1}.$$

Orbit realisation. Set $D = \{0, \infty\}$, and choose the non-dynamical element

$$\Lambda_{\text{Toda}}(\lambda) = \frac{\Lambda_0^{(1)}}{\lambda^2} + \Lambda_\infty \in \Omega_{D'}^\Gamma,$$

where

$$\Lambda_0^{(1)} = \sum_{i=1}^T E_{i+1, i} \in \mathfrak{g}^{(-1)}, \quad \Lambda_\infty = \sum_{i=1}^T E_{i, i+1} \in \mathfrak{g}^{(1)}.$$

The group elements $\varphi_+ = (\varphi_{0+}, \varphi_{\infty+})$, defined by (3.24), contain the dynamical degrees of freedom. From (3.27), we know that the components of $L_{\text{Toda}}(\lambda)$ can now be expressed as

$$J_0^{(0)} = [\phi_0^{(1)} \phi_0^{(0)-1}, J_0^{(1)}], \quad J_0^{(1)} = \phi_0^{(0)} \Lambda_0^{(1)} \phi_0^{(0)-1}, \quad J_\infty = \Lambda_\infty. \quad (4.6)$$

This gives us a parametrisation of the Lax matrix $L_{\text{Toda}}(\lambda)$ in (4.5) as an element of the coadjoint orbit $\mathcal{O}_\Lambda^{\text{Toda}}$. Since $\sigma(\varphi_{0+}(\lambda)) = \varphi_{0+}(\omega\lambda)$, we have $\sigma\phi_0^{(n)} = \omega^n\phi_0^{(n)}$. In particular, $\phi_0^{(0)}$ and $\phi_0^{(1)}$ have the form

$$\phi_0^{(0)} = \sum_{i=1}^T u_i E_{ii}, \quad \phi_0^{(1)} = \sum_{i=1}^T v_i E_{i,i+1}.$$

For convenience, define $(u_0, v_0) = (u_T, v_T)$ and $(u_{T+1}, v_{T+1}) = (u_1, v_1)$ to encode the periodic boundary conditions. Then, from (4.6), we get

$$J_0^{(0)} = \sum_{i=1}^T \left(\frac{v_i}{u_i} - \frac{v_{i-1}}{u_{i-1}} \right) E_{ii}, \quad J_0^{(1)} = \sum_{i=1}^T \frac{u_{i+1}}{u_i} E_{i+1,i}, \quad J_\infty = \sum_{i=1}^T E_{i,i+1}.$$

Defining

$$p_i = \frac{v_i}{u_i} - \frac{v_{i-1}}{u_{i-1}} \quad \text{and} \quad q_i = -\ln u_i \quad \text{for } i = 1, \dots, T,$$

we now have the desired realisation of the coefficients of the Lax matrix $L_{\text{Toda}}(\lambda)$. The coadjoint orbit $\mathcal{O}_\Lambda^{\text{Toda}}$ where $L_{\text{Toda}}(\lambda)$ lives is parameterised by $a_i, p_i, i = 1, \dots, T$, satisfying $\prod_i^T a_i = 1$ and $\sum_i^T p_i = 0$.

Lax equations. Let us now look at the Lax equations associated with the invariant functions on $\tilde{\mathfrak{g}}_D^{(1)}$ defined in (3.36) for the general case. The only functions we need to consider are

$$H_{p,0} = \frac{1}{p+1} \text{Res}_{\lambda=0} (\lambda^p \text{Tr}(\iota_{\lambda_0} L_{\text{Toda}}^{p+1})) d\lambda.$$

The set of functions $H_{p,\infty}$ are not independent: we have $H_{p,\infty} = -H_{p,0}$ for all $p \geq 1$. The Lax equations for the periodic Toda chain with respect to the elementary times t_p^0 are given by

$$\partial_{t_p^0} \iota_\lambda L_{\text{Toda}} = [R_\pm \nabla H_{p,0}(\iota_\lambda L_{\text{Toda}}), \iota_\lambda L_{\text{Toda}}].$$

For $p = 1$, these take the form of the set of Lax equations in (3.34):

$$\partial_{t_1^0} L_{\text{Toda}} = [M_{1,0}, L_{\text{Toda}}] \quad \text{with} \quad M_{1,0} = -\frac{J_0^{(1)}}{\lambda}. \quad (4.7)$$

Taking residues on both sides in (4.7) gives the equations of motion for p_i, q_i for $i = 1, \dots, T$. To get the equations of motion for q_i it is most convenient to “undo” the dressing and write the corresponding equation as

$$[\partial_{t_1^0} \phi_0^{(0)} \phi_0^{(0)-1} - J_0^{(0)}, J_0^{(1)}] = 0.$$

This tells us that the diagonal matrix

$$\phi_0^{(0)-1} (\partial_{t_1^0} \phi_0^{(0)} \phi_0^{(0)-1} - J_0^{(0)}) \phi_0^{(0)}$$

must commute with $\Lambda_0^{(1)}$, and is therefore equal to $\alpha \mathbf{1}$. Using the freedom $\phi_0^{(0)} \rightarrow \phi_0^{(0)} g$, where g is diagonal, to set $\det \phi_0^{(0)} = 1$, we see that $\alpha = 0$ by taking the trace of

$$\partial_{t_1^0} \phi_0^{(0)} \phi_0^{(0)-1} - J_0^{(0)} = \alpha \mathbf{1}.$$

Thus, we have the desired Toda equations

$$\begin{aligned} \partial_{t_1^0} q_i &= -p_i, \\ \partial_{t_1^0} p_i &= e^{q_i - q_{i+1}} - e^{q_{i-1} - q_i}. \end{aligned} \quad (4.8)$$

Lagrangian description. Using Theorem 3.6, we can now write a Lagrangian multiform for the periodic Toda hierarchy as

$$\mathcal{L}_{\text{Toda}} = \sum_{p=1}^M \mathcal{L}_{p,0} dt_p^0$$

with

$$\mathcal{L}_{p,0} = - \sum_{i=1}^T p_i \partial_{t_p^0} q_i - \frac{1}{p+1} \text{Res}_{\lambda=0} (\lambda^p \text{Tr}(L_{\text{Toda}}^{p+1})) d\lambda.$$

For $p=1$, this gives us the Lagrangian coefficients for the periodic Toda chain with respect to the time t_1^0 :

$$\mathcal{L}_{1,0} = - \sum_{i=1}^T p_i \partial_{t_1^0} q_i - \frac{1}{2} \sum_{i=1}^T p_i^2 - \sum_{i=1}^T e^{q_i - q_{i+1}}.$$

This is the expected *phase space* Lagrangian that our method produces and which gives Hamilton's equations (4.8) for periodic Toda. It corresponds to the (tangent bundle) Lagrangian

$$\mathcal{L}_{1,0} = \frac{1}{2} \sum_{i=1}^T (\partial_{t_1^0} q_i)^2 - \sum_{i=1}^T e^{q_i - q_{i+1}}.$$

4.2 DST model

The discrete self-trapping (DST) equation was introduced in [17] to describe the dynamics of small molecules, which then led to detailed studies of the DST dimer using different methods. The DST model we cast into our framework here is a generalisation of the dimer case to arbitrary (finite) degrees of freedom. This general case first appeared in [16] where its relationship with the periodic Toda chain was also hinted at. Our motivation here being different, we do not delve into this connection between the two theories. The interested reader is referred to [22] where this aspect was explored further.

Lax matrix. We work here with the following avatar of the Lax matrix of the DST model

$$L_{\text{DST}}(\lambda) = \frac{1}{\lambda} \sum_{i=1}^T c_i E_{ii} + \frac{1}{T} \sum_{i,j=1}^T \sum_{k=0}^{T-1} \frac{\omega^{k(j-i)} x_i X_j E_{ij}}{\lambda - \omega^k \zeta_1} + \sum_{i=1}^T E_{i,i+1},$$

where c_i , for $i=1, \dots, T$, are complex parameters, and x_i, X_i are the canonical coordinates satisfying the canonical Poisson bracket relations $\{X_i, x_j\} = \delta_{ij}$ for $i, j=1, \dots, T$, and the periodic conditions $(X_0, x_0) = (X_T, x_T)$ and $(X_{T+1}, x_{T+1}) = (X_1, x_1)$.

The DST Lax matrix in [22, equation (3.8)] is given as

$$\widehat{L}_{\text{DST}}(\mu) = \sum_{i,j=1}^T \frac{b^{T+i-j} x_i X_j E_{ij}}{\mu - b^T} + \mu E_{T1} + \sum_{i \geq j} b^{i-j} x_i X_j E_{ij} + \sum_{i=1}^T c_i E_{ii} + \sum_{i=1}^{T-1} E_{i,i+1},$$

where $b, c_i \in \mathbb{C}$ are parameters of the model, and μ is the spectral parameter of the Lax matrix. In the present setup where we realise the DST model as a cyclotomic Gaudin model, the role of the parameter b is played by the location of the pole ζ_1 on $\mathbb{C}P^1 \setminus \{0, \infty\}$. One obtains the Lax matrix $L_{\text{DST}}(\lambda)$ by conjugating $\widehat{L}_{\text{DST}}(\lambda^T)$ by the diagonal matrix $\mathcal{D} = \text{diag}(\lambda^{-1}, \dots, \lambda^{-T})$ followed by an overall multiplication by λ^{-1} , together with a change of λ -dependence, that is,

$$L_{\text{DST}}(\lambda) = \lambda^{-1} \mathcal{D} \widehat{L}_{\text{DST}}(\lambda^T) \mathcal{D}^{-1}, \quad (4.9)$$

and then using the identity (4.1).

The Poisson bracket of the Lax matrix $\widehat{L}_{\text{DST}}(\lambda)$ is given as

$$\{\widehat{L}_{\text{DST}_1}(\lambda), \widehat{L}_{\text{DST}_2}(\mu)\} = [\widehat{r}_{12}(\lambda, \mu), \widehat{L}_{\text{DST}_1}(\lambda)] - [\widehat{r}_{21}(\mu, \lambda), \widehat{L}_{\text{DST}_2}(\mu)],$$

where

$$\widehat{r}_{12}(\lambda, \mu) = \frac{1}{\mu - \lambda} \left(\mu \sum_{i \leq j} + \lambda \sum_{i > j} \right) E_{ij} \otimes E_{ji}.$$

Using the identity (4.1) once again, one finds that the gauge transformation (4.9) gives for $L_{\text{DST}}(\lambda)$ the r -matrix (4.4) associated with the cyclotomic $\mathfrak{gl}_T(\mathbb{C})$ -Gaudin model, as we would have anticipated. We then have the Poisson bracket

$$\{L_{\text{DST}_1}(\lambda), L_{\text{DST}_2}(\mu)\} = [r_{12}(\lambda, \mu), L_{\text{DST}_1}(\lambda)] - [r_{21}(\mu, \lambda), L_{\text{DST}_2}(\mu)].$$

Similar to the case of the periodic Toda chain, the Lax matrix $L_{\text{DST}}(\lambda)$ can be seen as a realisation of the cyclotomic Gaudin Lax matrix, this time with simple poles at the origin and all $\omega^k \zeta_1$, $k \in \{0, \dots, T-1\}$, for some $\zeta_1 \in \mathbb{C}^\times$, and a double pole at infinity, that is,

$$L_{\text{DST}}(\lambda) = \frac{K_0^{(0)}}{\lambda} + \frac{1}{T} \sum_{k=0}^{T-1} \frac{\sigma^k K_1}{\lambda - \omega^k \zeta_1} + K_\infty, \quad (4.10)$$

where

$$K_0^{(0)} = \sum_{i=1}^T c_i E_{ii}, \quad K_1 = \sum_{i,j=1}^T x_i X_j E_{ij}, \quad K_\infty = \sum_{i=1}^T E_{i,i+1}. \quad (4.11)$$

Orbit realisation. Set $D = \{0, 1, \infty\}$, and choose the non-dynamical element

$$\Lambda_{\text{DST}}(\lambda) = \frac{\Lambda_0^{(0)}}{\lambda} + \frac{1}{T} \sum_{k=0}^{T-1} \frac{\sigma^k \Lambda_1}{\lambda - \omega^k \zeta_1} + \Lambda_\infty \in \Omega_{D'}^\Gamma,$$

where

$$\Lambda_0^{(0)} = \sum_{i=1}^T c_i E_{ii} \in \mathfrak{g}^{(0)}, \quad \Lambda_1 = E_{11} \in \mathfrak{g}, \quad \Lambda_\infty = \sum_{i=1}^T E_{i,i+1} \in \mathfrak{g}^{(1)}.$$

The coadjoint action of the group elements $\varphi_+ = (\varphi_{0+}, \varphi_{1+}, \varphi_{\infty+})$, defined by (3.24), on $\Lambda_{\text{DST}}(\lambda)$ gives the orbit where $L_{\text{DST}}(\lambda)$ lives. From (3.27), we know that the components of $L_{\text{DST}}(\lambda)$ can then be expressed as

$$K_0^{(0)} = \phi_0^{(0)} \Lambda_0^{(0)} \phi_0^{(0)-1}, \quad K_1 = \phi_1 \Lambda_1 \phi_1^{-1}, \quad K_\infty = \Lambda_\infty, \quad (4.12)$$

where we have denoted $\phi_1^{(0)}$ by ϕ_1 for simplicity. This gives us a parameterisation of the Lax matrix $L_{\text{DST}}(\lambda)$ in (4.14) as an element of the coadjoint orbit $\mathcal{O}_\Lambda^{\text{DST}}$. We parameterise ϕ_1 as

$$\phi_1 = \sum_{i,j=1}^T s_{ij} E_{ij},$$

and denote by \hat{s}_{ij} , $i, j = 1, \dots, T$, the entries of its inverse, that is,

$$\phi_1^{-1} = \sum_{i,j=1}^T \hat{s}_{ij} E_{ij}.$$

From (4.12), we then have

$$K_0^{(0)} = \sum_{i=1}^T c_i E_{ii}, \quad K_1 = \sum_{i,j=1}^T s_{i1} \hat{s}_{1j} E_{ij}, \quad K_\infty = \sum_{i=1}^T E_{i,i+1}.$$

Defining

$$x_i = s_{i1} \quad \text{and} \quad X_i = \hat{s}_{1i} \quad \text{for} \quad i = 1, \dots, T,$$

gives us the desired realisation of the Lax matrix $L_{\text{DST}}(\lambda)$ as an element of the coadjoint orbit $\mathcal{O}_\Lambda^{\text{DST}}$. Notice that $\text{Tr} K_1 = \sum_{i=1}^T x_i X_i = 1$ is an orbit invariant and can be seen as being generated by the symmetry $x_i \rightarrow ax_i$, $X_i \rightarrow a^{-1}X_i$.

Lax equations. Let us choose invariant functions $H_{p,r}$ on $\tilde{\mathfrak{g}}_D^{(1)}$ as defined in (3.36), and treat $H_{p,0}$ and $H_{p,1}$ as the independent functions. The Lax equations for the DST model with respect to the elementary times t_p^r are then given by

$$\partial_{t_p^r} \iota_\lambda L_{\text{DST}} = [R_\pm \nabla H_{p,r}(\iota_\lambda L_{\text{DST}}), \iota_\lambda L_{\text{DST}}].$$

For $p = 1$, these take the form of the set of Lax equations in (3.34):

$$\partial_{t_1^r} L_{\text{DST}} = [M_{1,r}, L_{\text{DST}}] \quad \text{with} \quad M_{1,r} = \begin{cases} 0 & \text{for } r = 0, \\ -\frac{1}{T} \sum_{k=0}^{T-1} \frac{\omega^k \zeta_1 \sigma^k K_1}{\lambda - \omega^k \zeta_1} & \text{for } r = 1. \end{cases}$$

The t_1^0 equations are trivial. Similar to the case of the periodic Toda chain, the easiest way to get the t_1^1 equations of motion for x_i and X_i , for $i = 1, \dots, T$, is to undo the dressing. With $M = K_0^{(0)} + \zeta_1 K_\infty + \frac{1}{T} \sum_{k=1}^{T-1} \sigma^k K_1$, we find that $\phi_1^{-1}(\partial_{t_1^1} \phi_1 \phi_1^{-1} - M)\phi_1$ must commute with Λ_1 , and hence be block diagonal with a scalar ‘‘block’’, say ρ , and a $(T-1) \times (T-1)$ block which is irrelevant. Then we find, collecting x_i in the vector \mathbf{x} , and X_i in the vector \mathbf{X} ,

$$\partial_{t_1^1} \mathbf{x} - M\mathbf{x} = \rho\mathbf{x}, \quad \partial_{t_1^1} \mathbf{X}^T - \mathbf{X}^T M = \rho\mathbf{X}^T.$$

Using the freedom $x_i \rightarrow ax_i$, $X_i \rightarrow a^{-1}X_i$ mentioned above with $a = e^B$, $\partial_{t_1^1} B = \rho$, we can set $\rho = 0$. Explicitly, we obtain

$$\begin{aligned} \partial_{t_1^0} X_i &= 0, \\ \partial_{t_1^0} x_i &= 0, \\ \partial_{t_1^1} X_i &= -c_i X_i - \zeta_1 X_{i-1} - \frac{1}{T} \sum_{k=1}^{T-1} \sum_{j=1}^T \omega^{k(j-i)} x_j X_j X_i, \\ \partial_{t_1^1} x_i &= c_i x_i + \zeta_1 x_{i+1} + \frac{1}{T} \sum_{k=1}^{T-1} \sum_{j=1}^T \omega^{k(j-i)} X_j x_j x_i. \end{aligned} \tag{4.13}$$

Lagrangian description. We can now write a Lagrangian multiform for the DST hierarchy using Theorem 3.6 as

$$\mathcal{L}_{\text{DST}} = \sum_{p=1}^M \mathcal{L}_{p,0} dt_p^0 + \sum_{p=1}^M \mathcal{L}_{p,1} dt_p^1$$

with

$$\mathcal{L}_{p,r} = \sum_{i=1}^T X_i \partial_{t_p^r} x_i - H_{p,r}(\iota_\lambda L_{\text{DST}}), \quad r \in \{0, 1\},$$

where the potential term $H_{p,r}(\iota_\lambda L_{\text{DST}})$ is given by (3.36) for $r \in \{0, 1\}$. Notice that we have dropped the kinetic contribution to $\mathcal{L}_{p,r}$ from the pole at origin since being a total derivative it will not enter the Euler–Lagrange equations. For $p = 1$, the Lagrangian coefficients explicitly read

$$\begin{aligned} \mathcal{L}_{1,0} &= \sum_{i=1}^T X_i \partial_{t_1^0} x_i - \frac{1}{2} \sum_{i=1}^T c_i^2, \\ \mathcal{L}_{1,1} &= \sum_{i=1}^T X_i \partial_{t_1^1} x_i - \frac{1}{2T} \sum_{i,j=1}^T \sum_{k=0}^{T-1} \omega^{k(j-i)} x_i x_j X_i X_j - \sum_{i=1}^T c_i x_i X_i - \zeta_1 \sum_{i=1}^T x_{i+1} X_i. \end{aligned}$$

It can be checked that the Euler–Lagrange equations obtained from varying $\mathcal{L}_{1,0}$ and $\mathcal{L}_{1,1}$ with respect to X_i , x_i are exactly the equations in (4.13).

4.3 Coupled Toda-DST system

Finally, we turn to the task of coupling together the two hierarchies we have described variationally in this section. The Lax matrix of the coupled hierarchy can be expressed as

$$L_{\text{Toda-DST}}(\lambda) = L_{\text{Toda}}(\lambda) + \beta L_{\text{DST}}(\lambda), \quad (4.14)$$

where $L_{\text{Toda}}(\lambda)$ is the Lax matrix of the periodic Toda chain in (4.5), $L_{\text{DST}}(\lambda)$ is the DST Lax matrix (4.10), and the parameter β is a real-valued scalar parameter dictating the strength of coupling between the two hierarchies. Naturally, the Lax matrix $L_{\text{Toda-DST}}(\lambda)$ can be seen as a realisation of the cyclotomic Gaudin Lax matrix:

$$L_{\text{Toda-DST}}(\lambda) = \frac{J_0^{(0)} + \beta K_0^{(0)}}{\lambda} + \frac{J_0^{(1)}}{\lambda^2} + \frac{\beta}{T} \sum_{k=0}^{T-1} \frac{\sigma^k K_1}{\lambda - \omega^k \zeta_1} + J_\infty + \beta K_\infty$$

with the \mathfrak{g} -valued coefficients defined in (4.6) and (4.11).

As the number of finite (non-zero) poles in a cyclotomic Gaudin model is arbitrary, our construction allows us, in principle, to couple together an arbitrary number of copies of the DST model and a copy of the periodic Toda chain. Here we only illustrate it for a single copy each of the periodic Toda chain and the DST model.

Orbit realisation. Set $D = \{0, 1, \infty\}$. Since we already have a parameterisation of the Lax matrices $L_{\text{Toda}}(\lambda)$ and $L_{\text{DST}}(\lambda)$ as orbit elements, we only need to check that the action of a generic group element $\varphi_+ = (\varphi_{0+}, \varphi_{1+}, \varphi_{\infty+}) \in \tilde{G}_{D+}^{(0)}$ defined by (3.24) on the non-dynamical element

$$\Lambda_{\text{Toda-DST}}(\lambda) = \Lambda_{\text{Toda}}(\lambda) + \beta \Lambda_{\text{DST}}(\lambda) \in \Omega_{D'}^\Gamma,$$

where

$$\Lambda_{\text{Toda}}(\lambda) = \frac{\Lambda_0^{(1)}}{\lambda^2} + \Lambda_\infty, \quad \Lambda_{\text{DST}}(\lambda) = \frac{\Lambda_0^{(0)}}{\lambda} + \frac{1}{T} \sum_{k=0}^{T-1} \frac{\sigma^k \Lambda_1}{\lambda - \omega^k \zeta_1} + \Lambda_\infty$$

results in the Lax matrix $L_{\text{ Toda-DST}}(\lambda)$. Indeed, using the result in (3.27), we have that the components of $L_{\text{ Toda-DST}}(\lambda)$ take the form

$$\begin{aligned} J_0^{(0)} + \beta K_0^{(0)} &= [\phi_0^{(1)} \phi_0^{(0)-1}, J_0^{(1)}] + \beta \phi_0^{(0)} \Lambda_0^{(0)} \phi_0^{(0)-1}, \\ J_0^{(1)} &= \phi_0^{(0)} \Lambda_0^{(1)} \phi_0^{(0)-1}, \\ \beta K_1 &= \beta \phi_1 \Lambda_1 \phi_1^{-1}, \quad \text{and} \\ J_\infty + \beta K_\infty &= \Lambda_\infty + \beta \Lambda_\infty, \end{aligned}$$

as desired. This gives us a parameterisation of the Lax matrix $L_{\text{ Toda-DST}}(\lambda)$ in (4.14) as an element of the coadjoint orbit $\mathcal{O}_\Lambda^{\text{Toda}} \times \mathcal{O}_\Lambda^{\text{DST}}$.

Lax equations. As earlier, we will choose invariant functions $H_{p,r}$ on $\tilde{\mathfrak{g}}_D^{(1)}$ as defined in (3.36), for $r \in \{0, 1\}$. The Lax equations for the DST-Toda model with respect to the elementary times t_p^r are given by

$$\partial_{t_p^r} \iota_\lambda L_{\text{ Toda-DST}} = [R_\pm \nabla H_{p,r}(\iota_\lambda L_{\text{ Toda-DST}}), \iota_\lambda L_{\text{ Toda-DST}}].$$

For $p = 1$, these take the form of the set of Lax equations in (3.34):

$$\partial_{t_1^r} L_{\text{ Toda-DST}} = [M_{1,r}, L_{\text{ Toda-DST}}] \tag{4.15}$$

with

$$M_{1,r} = \begin{cases} -\frac{J_0^{(1)}}{\lambda} & \text{for } r = 0, \\ -\frac{\beta}{T} \sum_{k=0}^{T-1} \frac{\omega^k \zeta_1 \sigma^k K_1}{\lambda - \omega^k \zeta_1} & \text{for } r = 1. \end{cases}$$

Taking residues on both sides in (4.15) gives the equations of motion for p_i, q_i, X_i, x_i , for $i = 1, \dots, T$. Like in the cases of the periodic Toda chain and the DST model, to get the equations of motion for q_i, X_i, x_i , one can “undo” the dressing in the corresponding equations. Explicitly, one gets the following equations for p_i, q_i for $i = 1, \dots, T$:

$$\begin{aligned} \partial_{t_1^0} p_i &= (1 + \beta)(e^{q_i - q_{i+1}} - e^{q_i - 1 - q_i}) + \frac{\beta}{\zeta_1}(e^{q_i - 1 - q_i} x_{i-1} X_i - e^{q_i - q_{i+1}} x_i X_{i+1}), \\ \partial_{t_1^0} q_i &= -p_i - \beta c_i, \\ \partial_{t_1^1} p_i &= \frac{\beta}{\zeta_1}(e^{q_i - q_{i+1}} x_i X_{i+1} - e^{q_i - 1 - q_i} x_{i-1} X_i), \\ \partial_{t_1^1} q_i &= -\beta x_i X_i, \end{aligned} \tag{4.16}$$

and the following equations for X_i, x_i , for $i = 1, \dots, T$:

$$\begin{aligned} \partial_{t_1^0} X_i &= \frac{1}{\zeta_1} e^{q_i - q_{i+1}} X_{i+1}, \\ \partial_{t_1^0} x_i &= -\frac{1}{\zeta_1} e^{q_i - 1 - q_i} x_{i-1}, \\ \partial_{t_1^1} X_i &= -p_i X_i - \beta c_i X_i - \frac{1}{\zeta_1} e^{q_i - q_{i+1}} X_{i+1} - \frac{\beta}{T} \sum_{j=1}^T \sum_{k=0}^{T-1} \omega^{k(j-i)} x_j X_j X_i - (1 + \beta) \zeta_1 X_{i-1}, \\ \partial_{t_1^1} x_i &= p_i x_i + \beta c_i x_i + \frac{1}{\zeta_1} e^{q_i - 1 - q_i} x_{i-1} + \frac{\beta}{T} \sum_{j=1}^T \sum_{k=0}^{T-1} \omega^{k(j-i)} X_j x_j x_i + (1 + \beta) \zeta_1 x_{i+1}. \end{aligned} \tag{4.17}$$

Setting $\beta = 0$ in the above equations produces the equations of motion in (4.8) for the periodic Toda chain. In the limit $\beta \rightarrow \infty$, these reduce to the equations of motion we obtained for the DST model in (4.13). To see this, note that the Lax matrix L_{DST} comes as βL_{DST} in the coupled system. Therefore, in the limit $\beta \rightarrow \infty$, the time flow t_p^r is rescaled such that $\partial_{t_p^r}$ rescales to $\beta^p \partial_{t_p^r}$.

Lagrangian description. Using Theorem 3.6, we can now write a Lagrangian multiform for the Toda-DST hierarchy:

$$\mathcal{L}_{\text{Toda-DST}} = \sum_{p=1}^M \mathcal{L}_{p,0} dt_p^0 + \sum_{p=1}^M \mathcal{L}_{p,1} dt_p^1$$

with

$$\begin{aligned} \mathcal{L}_{p,r} = & \text{Tr}(J_0^{(0)} \partial_{t_p^r} \phi_0^{(0)} \phi_0^{(0)-1}) + \beta \text{Tr}(K_0^{(0)} \partial_{t_p^r} \phi_0^{(0)} \phi_0^{(0)-1}) \\ & + \beta \text{Tr}(K_1 \partial_{t_p^r} \phi_1 \phi_1^{-1}) - H_{p,r}(\iota_\lambda L_{\text{Toda-DST}}), \quad r \in \{0, 1\}, \end{aligned}$$

where the potential term $H_{p,r}(\iota_\lambda L_{\text{Toda-DST}})$ is given by (3.36) for $r \in \{0, 1\}$. Notice that $K_0^{(0)} = \sum_{i=1}^T c_i E_{ii}$. So, the second term on the right-hand side is, in fact, a total derivative and will not enter the Euler–Lagrange equations. So, we will simply drop it. In terms of the canonical coordinates, we then have

$$\mathcal{L}_{p,r} = - \sum_{i=1}^T p_i \partial_{t_p^r} q_i + \beta \sum_{i=1}^T X_i \partial_{t_p^r} x_i - H_{p,r}(\iota_\lambda L_{\text{Toda-DST}}), \quad r \in \{0, 1\}. \quad (4.18)$$

The decoupled limits of the periodic Toda and the DST hierarchies are easily recovered by setting $\beta = 0$ and taking the limit $\beta \rightarrow \infty$ in (4.18), respectively.

For $p = 1$, (4.18) gives the Lagrangian for the coupling of the periodic Toda chain and the DST model with the potential terms

$$\begin{aligned} H_{1,0} = & \frac{1}{2} \sum_{i=1}^T p_i^2 + \beta \sum_{i=1}^T c_i p_i + \frac{\beta^2}{2} \sum_{i=1}^T c_i^2 + (1 + \beta) \sum_{i=1}^T e^{q_i - q_{i+1}} - \frac{\beta}{\zeta_1} \sum_{i=1}^T e^{q_i - q_{i+1}} x_i X_{i+1}, \\ H_{1,1} = & \frac{\beta^2}{2T} \sum_{i,j=1}^T \sum_{k=0}^{T-1} \omega^{k(j-i)} x_i x_j X_i X_j + \beta \sum_{i=1}^T p_i x_i X_i + \beta^2 \sum_{i=1}^T c_i x_i X_i \\ & + \frac{\beta}{\zeta_1} \sum_{i=1}^T e^{q_i - q_{i+1}} x_i X_{i+1} + (\beta + \beta^2) \zeta_1 \sum_{i=1}^T x_{i+1} X_i. \end{aligned}$$

The variation $\delta \mathcal{L}_{1,r}$ gives the Euler–Lagrange equations for the coupled system with respect to the time flow t_1^r , for each $r \in \{0, 1\}$. It can be checked that these are exactly the equations we obtained in (4.16) and (4.17).

5 Conclusion

From the point of view of Lagrangian multiforms, our main result is the construction of a Lagrangian multiform for the class of cyclotomic Gaudin models which includes, as a particular example, the emblematic periodic Toda chain. The main technical aspect of this result lies in the pinning down of the necessary algebraic setup (in Section 3) associated with the classical r -matrix of the cyclotomic Gaudin model which is both non-skew-symmetric and depends on spectral parameters. This is the first time that a Lagrangian multiform has been constructed

for an r -matrix with both of these properties, as previous general constructions were either only presented for skew-symmetric ones [11, 15] or for non-skew-symmetric r -matrices but without spectral parameter [11].

The non-skew-symmetric r -matrix of the cyclotomic Gaudin model arises from a decomposition of a direct sum of (twisted) loop algebras into a pair of complementary subalgebras, which controls the corresponding Lie dialgebra structure entering its Lagrangian multiform. While finishing this work, the preprint [1] appeared on the construction of solutions of the general classical Yang–Baxter equation based on more general decompositions of such Lie algebras and to which one can associate generalised notions of Gaudin models. In particular, it appears that the cyclotomic setup considered here is a special case of their construction. It would be interesting to investigate if our Lagrangian multiform construction can be extended to the more general non-skew-symmetric r -matrices and associated generalised Gaudin models of [1].

Acknowledgements

The authors would like to thank the referees for their constructive feedback. A.A.S. is funded by the School of Mathematics EPSRC Doctoral Training Partnership Studentship (Project Reference Number 2704447). B.V. gratefully acknowledges the support of the Leverhulme Trust through a Leverhulme Research Project Grant (RPG-2021-154).

References

- [1] Abedin R., Maximov S., Stolin A., Generalized classical Yang–Baxter equation and regular decompositions, [arXiv:2405.04440](https://arxiv.org/abs/2405.04440).
- [2] Adler M., On a trace functional for formal pseudo differential operators and the symplectic structure of the Korteweg-de Vries type equations, *Invent. Math.* **50** (1978), 219–248.
- [3] Adler M., van Moerbeke P., Completely integrable systems, Euclidean Lie algebras, and curves, *Adv. Math.* **38** (1980), 267–317.
- [4] Avan J., Classical R -matrices with general automorphism groups, 1991, Preprint BROWN-HET-781, 15 pages.
- [5] Avan J., From rational to trigonometric R -matrices, *Phys. Lett. A* **156** (1991), 61–68.
- [6] Avan J., Talon M., Graded R -matrices for integrable systems, *Nuclear Phys. B* **352** (1991), 215–249.
- [7] Babelon O., Bernard D., Talon M., Introduction to classical integrable systems, *Cambridge Monogr. Math. Phys.*, Cambridge University Press, Cambridge, 2003.
- [8] Babelon O., Viallet C.-M., Hamiltonian structures and Lax equations, *Phys. Lett. B* **237** (1990), 411–416.
- [9] Bobenko A.I., Suris Yu.B., Integrable systems on quad-graphs, *Int. Math. Res. Not.* **2002** (2002), 573–611, [arXiv:nlin.SI/0110004](https://arxiv.org/abs/nlin.SI/0110004).
- [10] Caudrelier V., Crampé N., Classical N -reflection equation and Gaudin models, *Lett. Math. Phys.* **109** (2019), 843–856, [arXiv:1803.09931](https://arxiv.org/abs/1803.09931).
- [11] Caudrelier V., Dell’Atti M., Singh A.A., Lagrangian multiforms on coadjoint orbits for finite-dimensional integrable systems, *Lett. Math. Phys.* **114** (2024), 34, 52 pages, [arXiv:2307.07339](https://arxiv.org/abs/2307.07339).
- [12] Caudrelier V., Nijhoff F., Sleight D., Vermeeren M., Lagrangian multiforms on Lie groups and non-commuting flows, *J. Geom. Phys.* **187** (2023), 104807, 35 pages, [arXiv:2204.09663](https://arxiv.org/abs/2204.09663).
- [13] Caudrelier V., Stoppato M., Hamiltonian multiform description of an integrable hierarchy, *J. Math. Phys.* **61** (2020), 123506, 25 pages, [arXiv:2004.01164](https://arxiv.org/abs/2004.01164).
- [14] Caudrelier V., Stoppato M., Multiform description of the AKNS hierarchy and classical r -matrix, *J. Phys. A* **54** (2021), 235204, 43 pages, [arXiv:2010.07163](https://arxiv.org/abs/2010.07163).
- [15] Caudrelier V., Stoppato M., Vicedo B., Classical Yang–Baxter equation, Lagrangian multiforms and ultralocal integrable hierarchies, *Comm. Math. Phys.* **405** (2024), 12, 67 pages, [arXiv:2201.08286](https://arxiv.org/abs/2201.08286).
- [16] Christiansen P.L., Jørgensen M.F., Kuznetsov V.B., On integrable systems close to the Toda lattice, *Lett. Math. Phys.* **29** (1993), 165–173.

- [17] Eilbeck J.C., Lomdahl P.S., Scott A.C., The discrete self-trapping equation, *Phys. D* **16** (1985), 318–338.
- [18] Flaschka H., The Toda lattice. II. Existence of integrals, *Phys. Rev. B* **9** (1974), 1924–1925.
- [19] Gaudin M., Diagonalisation d’une classe d’Hamiltoniens de spin, *J. Physique* **37** (1976), 1089–1098.
- [20] Gaudin M., La fonction d’onde de Bethe, *Collect. Commissariat Énerg. Atom. Sér. Sci.*, Masson, Paris, 1983.
- [21] Kostant B., The solution to a generalized Toda lattice and representation theory, *Adv. Math.* **34** (1979), 195–338.
- [22] Kuznetsov V.B., Salerno M., Sklyanin E.K., Quantum Bäcklund transformation for the integrable DST model, *J. Phys. A* **33** (2000), 171–189, [arXiv:solv-int/9908002](https://arxiv.org/abs/solv-int/9908002).
- [23] Lobb S., Nijhoff F., Lagrangian multiforms and multidimensional consistency, *J. Phys. A* **42** (2009), 454013, 18 pages, [arXiv:0903.4086](https://arxiv.org/abs/0903.4086).
- [24] Mikhailov A.V., The reduction problem and the inverse scattering method, *Phys. D* **3** (1981), 73–117.
- [25] Nijhoff F.W., Lax pair for the Adler (lattice Krichever–Novikov) system, *Phys. Lett. A* **297** (2002), 49–58, [arXiv:nlin.SI/0110027](https://arxiv.org/abs/nlin.SI/0110027).
- [26] Nijhoff F.W., Lagrangian 3-form structure for the Darboux system and the KP hierarchy, *Lett. Math. Phys.* **113** (2023), 27, 19 pages, [arXiv:2206.14338](https://arxiv.org/abs/2206.14338).
- [27] Petrera M., Suris Yu.B., Variational symmetries and pluri-Lagrangian systems in classical mechanics, *J. Nonlinear Math. Phys.* **24** (2017), 121–145, [arXiv:1710.01526](https://arxiv.org/abs/1710.01526).
- [28] Petrera M., Vermeeren M., Variational symmetries and pluri-Lagrangian structures for integrable hierarchies of PDEs, *Eur. J. Math.* **7** (2021), 741–765, [arXiv:1906.04535](https://arxiv.org/abs/1906.04535).
- [29] Reshetikhin N.Yu., Faddeev L.D., Hamiltonian structures for integrable field theory models, *Theoret. and Math. Phys.* **56** (1983), 847–862.
- [30] Semenov-Tian-Shansky M., Integrable systems: the r -matrix approach, Preprint RIMS-1650, Kyoto University, 2008.
- [31] Skrypnyk T., New integrable Gaudin-type systems, classical r -matrices and quasigraded Lie algebras, *Phys. Lett. A* **334** (2005), 390–399.
- [32] Sleigh D., Nijhoff F., Caudrelier V., A variational approach to Lax representations, *J. Geom. Phys.* **142** (2019), 66–79, [arXiv:1812.08648](https://arxiv.org/abs/1812.08648).
- [33] Sleigh D., Nijhoff F., Caudrelier V., Variational symmetries and Lagrangian multiforms, *Lett. Math. Phys.* **110** (2020), 805–826, [arXiv:1906.05084](https://arxiv.org/abs/1906.05084).
- [34] Sleigh D., Nijhoff F.W., Caudrelier V., Lagrangian multiforms for Kadomtsev–Petviashvili (KP) and the Gelfand–Dickey hierarchy, *Int. Math. Res. Not.* **2023** (2023), 1420–1460, [arXiv:2011.04543](https://arxiv.org/abs/2011.04543).
- [35] Sleigh D., Vermeeren M., Semi-discrete Lagrangian 2-forms and the Toda hierarchy, *J. Phys. A* **55** (2022), 475204, 24 pages, [arXiv:2204.13063](https://arxiv.org/abs/2204.13063).
- [36] Suris Yu.B., Variational formulation of commuting Hamiltonian flows: multi-time Lagrangian 1-forms, *J. Geom. Mech.* **5** (2013), 365–379, [arXiv:1212.3314](https://arxiv.org/abs/1212.3314).
- [37] Suris Yu.B., Vermeeren M., On the Lagrangian structure of integrable hierarchies, in *Advances in Discrete Differential Geometry*, Springer, Berlin, 2016, 347–378, [arXiv:1510.03724](https://arxiv.org/abs/1510.03724).
- [38] Symes W.W., Systems of Toda type, inverse spectral problems, and representation theory, *Invent. Math.* **59** (1980), 13–51.
- [39] Toda T., Waves in nonlinear lattice, *Prog. Theor. Phys.* **45** (1970), 174–200.
- [40] Vermeeren M., Continuum limits of pluri-Lagrangian systems, *J. Integrable Syst.* **4** (2019), xyy020, 34 pages, [arXiv:1706.06830](https://arxiv.org/abs/1706.06830).
- [41] Vicedo B., Young C., Cyclotomic Gaudin models: construction and Bethe ansatz, *Comm. Math. Phys.* **343** (2016), 971–1024, [arXiv:1409.6937](https://arxiv.org/abs/1409.6937).
- [42] Vicedo B., Young C., Cyclotomic Gaudin models with irregular singularities, *J. Geom. Phys.* **121** (2017), 247–278, [arXiv:1611.09059](https://arxiv.org/abs/1611.09059).
- [43] Yoo-Kong S., Lobb S., Nijhoff F., Discrete-time Calogero–Moser system and Lagrangian 1-form structure, *J. Phys. A* **44** (2011), 365203, 39 pages, [arXiv:1102.0663](https://arxiv.org/abs/1102.0663).