



This is a repository copy of *First Indication of solar 8 B neutrinos via coherent elastic neutrino-nucleus scattering with XENONnT*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/220357/>

Version: Published Version

Article:

Aprile, E. orcid.org/0000-0001-6595-7098, Aalbers, J. orcid.org/0000-0003-0030-0030, Abe, K. orcid.org/0009-0000-9620-788X et al. (164 more authors) (2024) First Indication of solar 8 B neutrinos via coherent elastic neutrino-nucleus scattering with XENONnT. Physical Review Letters, 133. 191002. ISSN 0031-9007

<https://doi.org/10.1103/physrevlett.133.191002>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

First Indication of Solar ^8B Neutrinos via Coherent Elastic Neutrino-Nucleus Scattering with XENONnT

E. Aprile¹, J. Aalbers², K. Abe³, S. Ahmed Maouloud⁴, L. Althueser⁵, B. Andrieu⁴, E. Angelino^{6,7}, D. Antón Martín⁸, F. Arneodo⁹, L. Baudis¹⁰, M. Bazyk¹¹, L. Bellagamba¹², R. Biondi¹³, A. Bismark¹⁰, K. Boese¹³, A. Brown¹⁴, G. Bruno¹¹, R. Budnik¹⁵, C. Cai¹⁶, C. Capelli¹⁰, J. M. R. Cardoso¹⁷, A. P. Cimental Chávez¹⁰, A. P. Colijn¹⁸, J. Conrad¹⁹, J. J. Cuenca-García¹⁰, V. D'Andrea^{7,*}, L. C. Daniel Garcia⁴, M. P. Decowski¹⁸, A. Deisting²⁰, C. Di Donato^{21,7}, P. Di Gangi¹², S. Diglio¹¹, K. Eitel²², A. Elykov²², A. D. Ferella^{21,7}, C. Ferrari⁷, H. Fischer¹⁴, T. Flehmke¹⁹, M. Flierman¹⁸, W. Fulgione^{6,7}, C. Fuselli¹⁸, P. Gaemers¹⁸, R. Gaior⁴, M. Galloway¹⁰, F. Gao¹⁶, S. Ghosh²³, R. Giacomobono²⁴, R. Glade-Beucke¹⁴, L. Grandi⁸, J. Grigat¹⁴, H. Guan²³, M. Guida¹³, P. Gyorgy²⁰, R. Hammann¹³, A. Higuera²⁵, C. Hils²⁰, L. Hoetsch¹³, N. F. Hood²⁶, M. Iacovacci²⁴, Y. Itow²⁷, J. Jakob⁵, F. Joerg^{13,10}, Y. Kaminaga³, M. Kara²², P. Kavargin¹⁵, S. Kazama²⁷, M. Kobayashi²⁷, D. Koke⁵, A. Kopec^{26,†}, F. Kuger¹⁴, H. Landsman¹⁵, R. F. Lang²³, L. Levinson¹⁵, I. Li²⁵, S. Li²⁸, S. Liang²⁵, Y.-T. Lin¹³, S. Lindemann¹⁴, M. Lindner¹³, K. Liu^{16,‡}, M. Liu^{1,16}, J. Loizeau¹¹, F. Lombardi²⁰, J. Long⁸, J. A. M. Lopes^{17,§}, T. Luce¹⁴, Y. Ma²⁶, C. Macolino^{21,7}, J. Mahlstedt¹⁹, A. Mancuso¹², L. Manenti⁹, F. Marignetti²⁴, T. Marrodán Undagoitia¹³, K. Martens³, J. Masbou¹¹, E. Masson⁴, S. Mastroianni²⁴, A. Melchiorre^{21,7}, J. Merz²⁰, M. Messina⁷, A. Michael⁵, K. Miuchi²⁹, A. Molinaro⁶, S. Moriyama³, K. Morå¹, Y. Mosbacher¹⁵, M. Murra¹, J. Müller¹⁴, K. Ni²⁶, U. Oberlack²⁰, B. Paetsch¹⁵, Y. Pan⁴, Q. Pellegrini⁴, R. Peres¹⁰, C. Peters²⁵, J. Pienaar^{8,15}, M. Pierre¹⁸, G. Plante¹, T. R. Pollmann¹⁸, L. Principe¹¹, J. Qi²⁶, J. Qin²⁵, D. Ramírez García¹⁰, M. Rajado¹⁰, R. Singh²³, L. Sanchez²⁵, J. M. F. dos Santos¹⁷, I. Sarnoff⁹, G. Sartorelli¹², J. Schreiner¹³, P. Schulte⁵, H. Schulze Eiβing⁵, M. Schumann¹⁴, L. Scotto Lavina⁴, M. Selvi¹², F. Semeria¹², P. Shagin²⁰, S. Shi¹, J. Shi¹⁶, M. Silva¹⁷, H. Simgen¹³, A. Takeda³, P.-L. Tan¹⁹, D. Thers¹¹, F. Toschi²², G. Trinchero⁶, C. D. Tunnell²⁵, F. Tönnies¹⁴, K. Valerius²², S. Vecchi³⁰, S. Vetter²², F. I. Villazon Solar²⁰, G. Volta¹³, C. Weinheimer⁵, M. Weiss¹⁵, D. Wenz⁵, C. Wittweg¹⁰, V. H. S. Wu²², Y. Xing¹¹, D. Xu^{1,||}, Z. Xu¹, M. Yamashita³, L. Yang²⁶, J. Ye^{31,¶}, L. Yuan⁸, G. Zavattini³⁰ and M. Zhong²⁶

(XENON Collaboration)

¹Physics Department, Columbia University, New York, New York 10027, USA²Nikhef and the University of Groningen, Van Swinderen Institute, 9747AG Groningen, Netherlands³Kamioka Observatory, Institute for Cosmic Ray Research, and Kavli Institute for the Physics and Mathematics of the Universe (WPI), University of Tokyo, Higashi-Mozumi, Kamioka, Hida, Gifu 506-1205, Japan⁴LPNHE, Sorbonne Université, CNRS/IN2P3, 75005 Paris, France⁵Institut für Kernphysik, University of Münster, 48149 Münster, Germany⁶INAF-Astrophysical Observatory of Torino, Department of Physics, University of Torino and INFN-Torino, 10125 Torino, Italy⁷INFN-Laboratori Nazionali del Gran Sasso and Gran Sasso Science Institute, 67100 L'Aquila, Italy⁸Department of Physics, Enrico Fermi Institute and Kavli Institute for Cosmological Physics, University of Chicago, Chicago, Illinois 60637, USA⁹New York University Abu Dhabi - Center for Astro, Particle and Planetary Physics, Abu Dhabi, United Arab Emirates¹⁰Physik-Institut, University of Zürich, 8057 Zürich, Switzerland¹¹SUBATECH, IMT Atlantique, CNRS/IN2P3, Université de Nantes, Nantes 44307, France¹²Department of Physics and Astronomy, University of Bologna and INFN-Bologna, 40126 Bologna, Italy¹³Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany¹⁴Physikalisches Institut, Universität Freiburg, 79104 Freiburg, Germany¹⁵Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot 7610001, Israel¹⁶Department of Physics and Center for High Energy Physics, Tsinghua University, Beijing 100084, People's Republic of China¹⁷LIBPhys, Department of Physics, University of Coimbra, 3004-516 Coimbra, Portugal¹⁸Nikhef and the University of Amsterdam, Science Park, 1098XG Amsterdam, Netherlands¹⁹Oskar Klein Centre, Department of Physics, Stockholm University, AlbaNova, Stockholm SE-10691, Sweden²⁰Institut für Physik and Exzellenzcluster PRISMA⁺, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany

²¹*Department of Physics and Chemistry, University of L'Aquila, 67100 L'Aquila, Italy*²²*Institute for Astroparticle Physics, Karlsruhe Institute of Technology, 76021 Karlsruhe, Germany*²³*Department of Physics and Astronomy, Purdue University, West Lafayette, Indiana 47907, USA*²⁴*Department of Physics "Ettore Pancini", University of Napoli and INFN-Napoli, 80126 Napoli, Italy*²⁵*Department of Physics and Astronomy, Rice University, Houston, Texas 77005, USA*²⁶*Department of Physics, University of California San Diego, La Jolla, California 92093, USA*²⁷*Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, and Institute for Space-Earth Environmental Research, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan*²⁸*Department of Physics, School of Science, Westlake University, Hangzhou 310030, People's Republic of China*²⁹*Department of Physics, Kobe University, Kobe, Hyogo 657-8501, Japan*³⁰*INFN-Ferrara and Dipartimento di Fisica e Scienze della Terra, Università di Ferrara, 44122 Ferrara, Italy*³¹*School of Science and Engineering, The Chinese University of Hong Kong (Shenzhen), Shenzhen, Guangdong, 518172, People's Republic of China*

(Received 4 August 2024; revised 29 August 2024; accepted 25 September 2024; published 7 November 2024)

We present the first measurement of nuclear recoils from solar ^8B neutrinos via coherent elastic neutrino-nucleus scattering with the XENONnT dark matter experiment. The central detector of XENONnT is a low-background, two-phase time projection chamber with a 5.9 t sensitive liquid xenon target. A blind analysis with an exposure of $3.51 \text{ t} \times \text{yr}$ resulted in 37 observed events above 0.5 keV, with $(26.4_{-1.3}^{+1.4})$ events expected from backgrounds. The background-only hypothesis is rejected with a statistical significance of 2.73σ . The measured ^8B solar neutrino flux of $(4.7_{-2.3}^{+3.6}) \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$ is consistent with results from the Sudbury Neutrino Observatory. The measured neutrino flux-weighted CE ν NS cross section on Xe of $(1.1_{-0.5}^{+0.8}) \times 10^{-39} \text{ cm}^2$ is consistent with the Standard Model prediction. This is the first direct measurement of nuclear recoils from solar neutrinos with a dark matter detector.

DOI: [10.1103/PhysRevLett.133.191002](https://doi.org/10.1103/PhysRevLett.133.191002)

Introduction—Coherent elastic neutrino-nucleus scattering (CE ν NS) is a Standard Model (SM) process with low momentum transfer, which allows neutrinos to scatter coherently with nuclei [1–3]. This process has only recently been observed using an intense, pulsed spallation neutron source (SNS) [4,5]. The detection of CE ν NS events from solar neutrinos is more challenging due to the lower flux [6] and energy, as well as the lack of timing information. Therefore, it requires minimal backgrounds and maximizing the sensitive region of interest (ROI) with a low energy threshold. Liquid xenon (LXe) detectors searching for dark matter (DM) particles fulfill these requirements, but have not been able to reach the required sensitivity until now [7,8]. Solar ^8B neutrinos are expected to contribute the

largest detectable number of coherent neutrino-xenon scattering events, albeit at low nuclear recoil (NR) energies [9]. In this Letter, the first detection of CE ν NS induced by solar ^8B neutrinos with the XENONnT experiment is reported. This is a “first” in three different aspects: the first detection of elastic NRs from astrophysical neutrinos, the first measurement of the CE ν NS process with a Xe target, and the first step into the “neutrino fog” by a DM experiment [10,11].

Experiment—The XENONnT experiment [12], located at the INFN Laboratori Nazionali del Gran Sasso in Italy, is designed to search for weakly interacting massive particles (WIMPs) scattering off Xe nuclei, which has a similar NR signature as CE ν NS. The experiment consists of three nested detectors: a muon veto (MV), a neutron veto (NV), and an innermost LXe detector. The latter is a two-phase time projection chamber (TPC) housed in a double-walled cryostat filled with 8.5 t of LXe. The cylindrical TPC, 1.33 m in diameter and 1.49 m in height, is enclosed by polytetrafluoroethylene (PTFE) panels and viewed by 494 3-in. Hamamatsu R11410-21 photomultiplier tubes (PMTs) [13] arranged in a top and a bottom array. The active LXe mass in the TPC is 5.9 t.

Particle interactions in the TPC produce both scintillation photons and ionization electrons. The prompt scintillation photons are detected by the PMTs and are referred to as the S1 signal. The liberated electrons drift upward in the drift field to the liquid-gas interface, where they are

* Also at INFN-Roma Tre, 00146 Roma, Italy.

† Now at Department of Physics and Astronomy, Bucknell University, Lewisburg, Pennsylvania, USA.

‡ Contact author: lkx21@mails.tsinghua.edu.cn

§ Also at Coimbra Polytechnic—ISEC, 3030-199 Coimbra, Portugal.

|| Contact author: dacheng.xu@columbia.edu¶ Contact author: yejingqiang@cuhk.edu.cn

extracted into the gas and produce a secondary scintillation signal, called the $S2$ signal, via electroluminescence. The time difference between $S1$ and $S2$ signals is proportional to the interaction depth (Z). Event positions in the horizontal plane (X , Y) are reconstructed based on the hit patterns of $S2$ signals in the top PMT array.

The electric fields in the TPC are established by three parallel-wire electrodes made of stainless steel [14]. The cathode and gate electrodes establish a drift field at 23 V/cm, resulting in a maximum electron drift time of 2.2 ms. The extraction field in LXe is set to 2.9 kV/cm by the gate and anode electrodes, which are reinforced by two and four additional perpendicular wires, respectively, to minimize sagging [14]. Two additional parallel-wire electrodes shield the PMT arrays from electric fields [12].

Dataset—This search uses two datasets with a total live time of 316.5 days after accounting for dead time from data acquisition [15] and vetoes. The first dataset, taken between July 6, 2021 and November 28, 2021 is referred to as the SR0 dataset in this Letter with a live time of 108.0 days. The second dataset was collected between May 19, 2022 and August 8, 2023, a period referred to as SR1, with a live time of 208.5 days. During SR0 (SR1), the temperature and pressure in the detector are stable within (176.8 ± 0.4) $[(177.2 \pm 0.4)]$ K and (1.890 ± 0.004) $[(1.92 \pm 0.02)]$ bar, respectively. The liquid level in SR0 is stable within (5.02 ± 0.20) mm [16]. On July 15, 2022, the liquid level is lowered by 0.2 mm and the anode voltage is raised by 50 V to mitigate localized electron bursts and maintain a consistent extraction field strength. Before and after this adjustment, the liquid level in SR1 is maintained stable at 5.0 and 4.8 mm above the gate electrode, respectively. The systematic uncertainty of the liquid level measurement is 0.2 mm.

In addition to the 17 PMTs already excluded from the analyses during SR0 [17], three additional PMTs are removed in SR1 due to increased afterpulse or intermittent light emission. PMT gains are monitored weekly using pulsed LED signals and are found to be stable in SR0 (SR1) within 3% (3.5%). PMT hits are recorded on a per-PMT basis when crossing the digitization threshold, typically about 2.06 mV [15]. The mean single photoelectron (PE) acceptance in SR0 (SR1) is determined to be $(91.2 \pm 0.2\%)$ $[(92.1 \pm 0.7)\%]$. Clusters of PMT hits in time are divided into peaks, which are classified into $S1$ and $S2$ signals based on their waveforms and intensity distributions on PMT arrays [18,19].

A distortion of the drift field near the edges of the detector leads to a difference in positions between the interaction site and the extraction position. It also leads to a small charge-insensitive volume (CIV) [14] in the lower part of the TPC, from where the drifting electrons reach the PTFE wall instead of the liquid-gas interface. A data-driven correction for the radial coordinate is applied to reproduce the uniform distribution of $^{83\text{m}}\text{Kr}$ calibration events [20].

For SR0, the method from [21] is kept, where the CIV does not enter the position correction but is considered in the fiducial volume (FV) calculation. The FV mass uncertainty originated from field distortion and position reconstruction is less than 5%. In SR1, the event positions are corrected according to the boundary defined by the simulated drift field [14] to account for the CIV. After considering the field distortion correction and removing events with the interaction depth Z below -142 cm or above -13 cm due to an insufficient understanding of the detector and backgrounds, the FV mass for SR0 and SR1 are (3.97 ± 0.20) and (4.10 ± 0.19) t, respectively. The total exposure in this analysis is $3.51 \text{ t} \times \text{yr}$.

Light from $S1$ or $S2$ signals can create delayed electron signals via photoionization of impurities in the LXe [16]. The photoionization strength, defined as the ratio between the number of measured photoionization electrons within 2.2 ms after an $S2$ signal larger than 10 000 PE and the number of electrons in the $S2$ signal itself, increased tenfold after a long maintenance and upgrade phase between SR0 and SR1. One hypothesis of the increased photoionization is that components in the radon removal system [22] are releasing photoionizable impurities after the upgrade, which enabled high flow extraction from the LXe target. No impact is observed from these impurities on the electron lifetime, which is an attenuation coefficient for the attachment to electronegative impurities during the drift of ionization electrons.

Signal inhomogeneities due to position- and time-dependent effects are corrected as described in [21]. The increased and varying photoionization strength in SR1 requires further time-dependent corrections to the $S1$ and $S2$ signal areas. After all corrections, the stability of the corrected $S1$ and corrected $S2$ ($cS2$) signals in SR0 are within 1% and 1.9%, respectively, and 0.3% and 1.1% in SR1. The variations are propagated as uncertainties into the determination of the photon gain ($g1$) and electron gain ($g2$). Using the method described in [21], $g1$ and $g2$ in SR0 (SR1) are found to be (0.151 ± 0.001) $[(0.137 \pm 0.001)]$ PE/photon and (16.5 ± 0.6) $[(16.9 \pm 0.5)]$ PE/electron, respectively.

CE ν NS signal—The expected NR spectrum of ^8B CE ν NS in LXe, considering the solar ^8B neutrino flux measured by the Sudbury Neutrino Observatory (SNO) [23], the ^8B neutrino energy spectrum from [24], and the CE ν NS cross section on Xe predicted by the SM [25], is shown in Fig. 1, with 90% of detectable recoils between 0.7 and 2.1 keV. The main contribution is from neutrinos with energies between 8 and 15 MeV. The low-energy NR response in this search is calibrated with 152 keV neutrons from an external ^{88}YBe source [26], with the recoil spectrum also shown in Fig. 1. The uncertainty in signal acceptance arises from uncertainties in $S1$ reconstruction, classification acceptance, and event selection acceptance. A model for light yield (L_y) and charge yield (Q_y) is fitted




FIG. 1. Acceptance for detecting low-energy NRs in XENONnT (top) and ${}^8\text{B}$ CE ν NS energy spectrum (bottom). The light (dark) blue curve denotes the acceptance of detecting $S1$ ($S2$) signals, and the black curve represents the combined acceptance. The expected CE ν NS signal spectrum induced by solar ${}^8\text{B}$ neutrinos in XENONnT with (without) acceptance loss is shown by the dark (light) red line. The green line shows a scaled spectrum of all energy depositions from ${}^{88}\text{YBe}$ calibration.

[27] to calibration data using a method similar to that described in [28]. The uncertainties of yields are propagated into the final inference with two parameters, t_{L_y} and t_{Q_y} , which determine the relative shift of L_y and Q_y from their median toward the $\pm 1\sigma$ quantiles. This calibration will be presented in an upcoming publication [29]. L_y and Q_y below 0.5 keV are assumed to be zero, which has a negligible impact on the ${}^8\text{B}$ CE ν NS detection rate.

The expected ${}^8\text{B}$ CE ν NS rate in our previous WIMP search region [30] is 0.2 events/(t \times yr). To increase the rate of detected ${}^8\text{B}$ CE ν NS in this search, the signal acceptance is improved by lowering two thresholds. First, the $S2$ signal threshold is reduced from 200 PE in the WIMP search to 120 PE in this search. Second, the $S1$ coincidence requirement was lowered from threefold coincidence to twofold coincidence, now minimally requiring only two PMTs with hits within ± 50 ns around the maximal amplitude of the $S1$ waveform. The reduced thresholds lead to an expected ${}^8\text{B}$ CE ν NS detection rate of 3.7(3.3) events/(t \times yr) in SR0(1), a factor of ~ 17 larger than in the WIMP search.

The ROI in this analysis is defined to be two or three hits for $S1$ signals and (120, 500) PE for $S2$ signals. The upper bound of the $S2$ area range is set to retain most of CE ν NS signal and to remove electronic recoil (ER) backgrounds from β and γ radiation, which have higher

ratios of $S2$ to $S1$ than NRs [31]. $S1$ signals with more than three hits are rarely produced by ${}^8\text{B}$ CE ν NS and such events are therefore not included in this analysis. Events in the ROI are blinded except those with radial positions larger than 63.0 cm, which are used to model the surface events produced by ${}^{210}\text{Pb}$ plate out on the TPC wall [32] and are not part of the dataset for the search. Threefold events were unblinded in the SR0 WIMP search [14], which contributes to $\leq 3\%$ of total ${}^8\text{B}$ CE ν NS rate since twofold events dominate and SR1 has more exposure.

Cuts based on the features of $S1$ and $S2$ peaks, inherited from [17], are employed to ensure the quality of the reconstructed events. $S1$ signals composed of at least two hits are required to be larger than 1 PE. $S1$ signals up to 4 PE are accepted in size per PMT. $S2$ signals must be detected by both PMT arrays with a reasonable signal fraction of around 75% in the top array. $S2$ signals detected on the top array are also required to follow the expected pattern from the optical response of XENONnT. Events with multiple $S2$ signals are rejected to suppress the neutron background. As in [17], events found in coincidence with either MV or NV are rejected.

Backgrounds—This analysis considers accidental coincidence (AC), surface, neutron, and ER background components, as in the search for solar ${}^8\text{B}$ CE ν NS signals with the XENON1T detector [7,28]. The AC is the dominant background, formed by accidentally paired “isolated” $S1$ and $S2$ signals. The accidental pileup rate of these isolated $S1$ and $S2$ signals within the maximum drift time is significant, reaching several hundred events per day before mitigation measures are applied.

The primary source of the isolated $S1$ and $S2$ signals in the ${}^8\text{B}$ CE ν NS search ROI are delayed signals after high-energy (HE) interactions. These interactions, with characteristic $S2$ areas larger than 10000 PE induced predominantly by γ rays from the materials’ radioactivity, are known to contaminate their subsequent time interval with single photoelectron PMT hits and small $S2$ signals. This phenomenon has been observed in many LXe detectors [7,8,33]. While the physical mechanism is still under investigation [34,35], the AC background can be modeled by data-driven simulation, after applying dedicated cuts to remove the isolated peaks correlated with their preceding HE peaks.

The impact on an isolated signal by a preceding HE event is quantified by the ratio of $S2_{\text{pre}}$ to Δt_{pre} , where $S2_{\text{pre}}$ is the $S2$ area of the HE event and Δt_{pre} is the time between the HE event and the isolated signal. All the HE events 1 sec before the isolated signal are considered and the event with the largest ratio of $S2_{\text{pre}}$ to Δt_{pre} (defined as $S2_{\text{pre}}/\Delta t_{\text{pre}}$) is identified as the most influential one on the isolated signal rate. Cuts are then applied on $S2_{\text{pre}}/\Delta t_{\text{pre}}$ to minimize the isolated signal rate. A time window of 2.2 ms (one maximum drift time) is vetoed after any HE

interaction in SR0. In SR1, due to the increased photoionization rate, the veto window is extended to 4.4 ms. The cut on $S2_{\text{pre}}/\Delta t_{\text{pre}}$ for 2- (3-) hit $S1$ signals is less than 10.1(38.2) PE/ μs , effectively reducing isolated $S1$ rates by more than 80% (50%) while accepting 87% (96%) of ${}^8\text{B}$ CE ν NS signals. Localized bursts of intense single-electron (SE) emission observed in SR0 [30] appear more frequently in SR1, contributing also to the isolated $S2$ signals. For isolated $S2$ signals, correlations with preceding HE events and the localized SE burst in (X, Y) position are utilized, accounting for the uncertainty in position reconstruction. Two-dimensional cuts in time and position are developed, effectively rejecting over 50% of isolated $S2$ signals while accepting around 96% of ${}^8\text{B}$ CE ν NS signals.

After all the cuts, the average isolated $S1$ and $S2$ signal rates in SR0 (SR1) are 2.3 (2.2) Hz and 18 (26) mHz, respectively. By injecting simulated ${}^8\text{B}$ CE ν NS signals at random times and positions into the real data, the overall acceptance of these cuts is evaluated to be 75% (85%) for 2- (3-) hit signals. The isolated $S1$ and $S2$ waveforms are then sampled and assigned a random drift time before being merged into artificial AC events. Facilitated by [36], the simulation improved compared to [7] in preserving the $S2_{\text{pre}}/\Delta t_{\text{pre}}$ spectrum and modeling the time dependence to minimize the systematic uncertainties of the AC model.

Two boosted decision tree (BDT) classifiers are developed to distinguish between ${}^8\text{B}$ CE ν NS signals and the AC background events. The output scores from these classifiers are used as analysis dimensions in the final likelihood. The distributions of $S1$ photons of ${}^8\text{B}$ CE ν NS signals in time and across the PMT arrays differ from those of the isolated $S1$ signals induced by a random pileup of PMT hits. Features from these distributions are therefore combined in an $S1$ BDT score. Another BDT assesses the $S2$ signal shape and the time between the $S1$ and $S2$ signals, which in ${}^8\text{B}$ CE ν NS signals are correlated due to diffusion of the drifting electron cloud, but this correlation is absent for the AC background. A cut on the $S2$ BDT score is applied to reject about 90% of the AC background events while retaining more than 80% of the signal events.

The $S2$ pulse shape changes close to the perpendicular supporting wires [14,30], so applying the $S2$ BDT cut to those events would introduce systematic errors in signal acceptance. Consequently, events close to the perpendicular wires are excluded from the analysis. Because of the $S2$ -area-dependent position resolution, this leads to an $S2$ -area-dependent reduction in the $S2$ acceptance rather than a reduction of the fiducial mass. Simulated $S1$ and $S2$ waveforms [37] are used to assess the acceptance loss due to cuts. The difference between acceptances estimated by simulated events and calibration data is smaller than 10%, which is assigned as the uncertainty on the total acceptance.

Figure 1 shows the total acceptance for $S1$ - and $S2$ -based cuts as function of NR energy.

AC-rich datasets are selected to validate the AC background model, including events with unphysically long drift times, calibration datasets featuring a high rate of isolated peaks, and an AC sideband mainly made of events rejected by the $S2$ BDT cut. These validations are performed with a binned likelihood goodness of fit (GOF) test in all the same dimensions as used in the statistical inference to search for ${}^8\text{B}$ CE ν NS signals, including $cS2$, $S2_{\text{pre}}/\Delta t_{\text{pre}}$, $S1$ BDT score, and $S2$ BDT score. In all these validation datasets, good agreements between AC prediction and observation in the ${}^8\text{B}$ CE ν NS ROI is achieved, constraining systematic uncertainties on the AC rate to be below 5%. Conservatively, the systematic uncertainty of the AC background is solely estimated from the AC sideband, which is unblinded only after the AC prediction and event selections are both fixed. The AC model passed the binned likelihood GOF test with the sideband data at a p value of 0.16. The AC background uncertainty for SR0 (SR1) is 9.0% (5.8%), based on statistical uncertainties from the AC sideband data. The expected numbers of AC background in SR0 and SR1 are (7.5 ± 0.7) and (17.8 ± 1.0) , respectively. Details about the AC sideband unblinding are provided in Appendix A.

Surface events produced mainly by ${}^{210}\text{Pb}$ plate out on the TPC wall have reduced $S2$ signals [32], which could lead to leakage of events into the ROI. A data-driven approach is adopted to derive the radial distribution of this background. Because of the limited statistical data, deriving and validating the data-driven model across all four analysis dimensions is currently unfeasible. Consequently, the outer radius of the FV for SR0 (SR1) is set at 60.15 cm (59.60 cm), such that surface events are expected to be less than 0.12 (0.23), respectively. At this level, this background can be safely neglected without risk of signal-like mismodeling in the ${}^8\text{B}$ CE ν NS search according to a dedicated toy Monte Carlo (MC) study.

Radiogenic neutrons originating from the detector materials are modeled using the framework of [28] with neutron spectra from updated knowledge of the detector material radioactivity. The prediction for SR0 and SR1 are (0.13 ± 0.07) and (0.33 ± 0.19) events, respectively. The rate uncertainty of 58% is derived from neutron candidates in SR0 tagged by the NV. In the CE ν NS ROI, the NV and MV tagged one event each after a dedicated unblinding, which is in agreement with the expected number of events vetoed by accidental coincidence between the TPC and the veto detectors.

The ER background is composed mainly of β decays from radioactive impurities, such as ${}^{214}\text{Pb}$ and ${}^{85}\text{Kr}$, and electrons scattered by external γ rays and solar neutrinos [28]. The shape of the ER background in the ${}^8\text{B}$ CE ν NS ROI is generated by [27] with emission model

fit to the ^{220}Rn calibration data [28]. The rate of ER background events is derived by fitting the events with ER energy above 20 keV, assuming a flat ER spectrum. However, the emission model in low energy has large systematical uncertainty. If using the light and charge yields from the Noble Element Simulation Technique [38,39], the expected ER rate is 10 times lower. To account for this discrepancy, a 100% uncertainty is assigned to the ER rate. Consequently, the assumed ER background in SR0 and SR1 is taken to be at most 0.13 ± 0.13 and 0.56 ± 0.56 events, respectively. Measurement of the light and charge response in XENONnT with a 0.27 keV calibration using a ^{37}Ar electron capture (EC) source, which will be introduced in a future publication, also confirms the nominal rate of the ER background is a conservative choice.

Statistical inference— $S2_{\text{pre}}/\Delta t_{\text{pre}}$, $S1$ BDT score, $S2$ BDT score, and $cS2$ are the four dimensions used to discriminate between the ^8B CE ν NS signal and the dominating AC background. The background and signal models are coarsely binned, with three bins in each of the four analysis dimensions for a total of 81 bins. A four-dimensional binned likelihood analysis is performed. The bins are chosen to have the same expected number of AC background events in the projection of each dimension. The chance for mismodeling of the AC background due to the limited number of isolated $S1$ and $S2$ peaks is negligible, as validated via toy MC simulations.

The extended likelihood function is constructed as

$$\mathcal{L}(\mu, \vec{\theta}) = \prod_{i=0,1} \mathcal{L}_i(\mu, \vec{\theta}) \times \prod_m \mathcal{L}_m(\theta_m), \quad (1)$$

where the parameter of interest μ can either be the solar ^8B neutrino flux (Φ), or the flux-weighted CE ν NS cross section on Xe ($\sigma_{\text{CE}\nu\text{NS}}$). $\vec{\theta}$ are the nuisance parameters, i iterates through the two science runs, and m iterates through the nuisance terms: the constraints on t_{L_y} and t_{Q_y} , the signal acceptance uncertainty, and the uncertainties in the rates of the AC, neutron, and ER backgrounds. The nuisance parameters θ_m are constrained via external measurements, modeled by Gaussian pull terms $\mathcal{L}_m(\theta_m)$. The models of ^8B CE ν NS and neutrons change in shape and expectation value with t_{L_y} and t_{Q_y} . The AC background rates are independent between science runs, while all other parameters are coupled.

The ^8B CE ν NS discovery significance and the construction of a confidence interval for the ^8B neutrino flux are computed using a test statistic q_μ based on the profile log-likelihood ratio as in [28,40]. The critical region for the confidence interval construction and expected discovery significance are computed with toy MC simulations using [41]. Consistency between the model and data is evaluated by a combination of four binned likelihood GOF tests performed on the four one-dimensional projections, combining SR0 and SR1. The p values are computed based on the distribution of the binned likelihood GOF test

TABLE I. The expected and best-fit number of events from signal and background components in the ROI. The uncertainty in the expectation accounts for contributions from signal detection efficiency, L_y , and Q_y . The uncertainties of background expectations correspond to the width of the Gaussian constraints in the fit, the ^8B signal is not constrained.

Component	Expectation	Best fit
AC (SR0)	7.5 ± 0.7	7.4 ± 0.7
AC (SR1)	17.8 ± 1.0	17.9 ± 1.0
ER	0.7 ± 0.7	$0.5^{+0.7}_{-0.6}$
Neutron	$0.5^{+0.2}_{-0.3}$	0.5 ± 0.3
Total background	$26.4^{+1.4}_{-1.3}$	26.3 ± 1.4
^8B	$11.9^{+4.5}_{-4.2}$	$10.7^{+3.7}_{-4.2}$
Observed	37	

statistic obtained via toy MC simulations. A threshold of 0.013 is selected for each test to obtain a 95% confidence limit (CL) for the final combined test. The test is defined before unblinding and its suitability to reject mismodeling is assessed using toy MC simulations.

The strategy to report the result from the ^8B CE ν NS search is decided before unblinding. A Feldman-Cousins construction [42] is used to constrain the solar ^8B neutrino flux and the CE ν NS cross section $\sigma_{\text{CE}\nu\text{NS}}$ without setting a threshold on p values for reporting a two-sided measurement. The expected ^8B CE ν NS signal under the nominal emission model is $11.9^{+4.5}_{-4.2}$ events, with the uncertainty originating from $S1$ and $S2$ acceptances and detector response to low-energy NRs. The expected background is $26.4^{+1.4}_{-1.3}$ events, dominated by the AC background, as shown in Table I. With the final background prediction summarized in Table I, the probability of obtaining a $\geq 2\sigma$ (3σ) discovery significance with this dataset is estimated to be 80% (48%) using toy MC simulations.

Before unblinding the ^8B CE ν NS search data, the signal and background modeling are validated in the four-dimensional space by measuring the L_y of the ^{37}Ar L shell electron capture ER signal at 0.27 keV, where Q_y is constrained [43], but L_y has not yet been measured. The background in the ^{37}Ar data at this low-energy region is dominated by the AC background due to the high rate of isolated $S2$ signals. The L_y of ^{37}Ar L shell is measured by fitting the ^{37}Ar calibration data [21] with the ^{37}Ar signal and the AC background. This fitting is analogous to the search for ^8B CE ν NS signals in terms of the signal dependence on the light and charge yields, the dominant background, and the energy region. Using approaches on the signal and background modeling comparable to the ^8B CE ν NS search, the best fit of the ^{37}Ar signal model and the AC background is consistent with the data in all of the four analysis dimensions. More information about this validation is described in Appendix B.

Results—After unblinding, 9 and 28 events are observed in SR0 and SR1, respectively. The observed number of events is consistent with the expected ^8B CE ν NS signal on top of the background. The best-fit values of background components and ^8B CE ν NS signal from the unconstrained fit are also shown in Table I. The best-fit nuisance parameters $\bar{\theta}$ are all within $\pm 0.3\sigma$ constrained by the external measurements. The background-only hypothesis, with no ^8B CE ν NS signals, is disfavored with a p value of 0.003, corresponding to a statistical significance of 2.73σ .

The distributions of the observed 37 events and the best-fit model projected to each analysis dimension are shown in Fig. 2. A detailed plot showing the SR0 and SR1 results




FIG. 2. Distributions of best-fit signal and background, together with the data in the projected analysis dimensions, summing both science runs. The observed number of events with Poisson uncertainties in each bin is shown in black. The ^8B CE ν NS signal is represented by the light green histogram on top of the backgrounds, which are indicated by purple (AC), blue (ER), and yellow (neutron) histograms. As the bin edges on each analysis dimension vary from SR0 to SR1, the plot for $cS2$ is shown in double axis, and the other dimensions are shown in quantiles of the AC background for the summed results.

separately is presented in Appendix C. The p values in $cS2$, $S1$ BDT score, and $S2$ BDT score show a good match between the unconstrained best-fit model and observations. The p value in the $S2_{\text{pre}}/\Delta t_{\text{pre}}$ is 0.008, indicating a potential mismodeling. No other indication of possible mismodeling is found by inspecting the individual events in the dataset or the AC sideband data. Abandoning $S2_{\text{pre}}/\Delta t_{\text{pre}}$ in the statistical inference would lead to a larger best-fit ^8B CE ν NS signal of 13.1 events with a statistical significance of 3.22σ . In addition, two tests of overdensity in (X, Y) space were defined before unblinding, although not part of the analysis dimensions. One returned a p value below the threshold of 0.018, prompting checks including inspection of event distributions in all cut spaces that show no indication of mismodeling.

Assuming the flux-weighted CE ν NS cross section $\sigma_{\text{CE}\nu\text{NS}}$ predicted by the SM, Fig. 3 shows the XENONnT constraint on the solar ^8B neutrino flux of $(4.7^{+3.6}_{-2.3}) \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$ at 68% CL. With the solar ^8B neutrino flux being constrained by SNO [23], Fig. 4 shows the first measurement of the flux-weighted CE ν NS cross section $\sigma_{\text{CE}\nu\text{NS}}$ on Xe as $(1.1^{+0.8}_{-0.5}) \times 10^{-39} \text{ cm}^2$, consistent with the SM prediction of $1.2 \times 10^{-39} \text{ cm}^2$. Since the momentum transferred from a solar ^8B neutrino to a Xe nucleus is $\leq 20 \text{ MeV}/c$, this measurement is less sensitive to uncertainties in the nuclear form factor compared to CE ν NS measurements made by the COHERENT Collaboration with neutrinos produced by the SNS [44]. The measurements of the flux-weighted CE ν NS cross section on CsI [44], Ar [5], and Ge [45] nuclei by the COHERENT Collaboration are shown in Fig. 4 for




FIG. 3. Constraints on solar ^8B neutrino flux. Top: the 68% (90%) measurement of solar ^8B neutrino flux from this work is shown in black (gray). The 68% CL measurement from SNO [23] and 90% CL upper limits from XENON1T [7] and PandaX-4T [8] are also shown. Bottom: the solid red line shows the profile likelihood ratio test statistics q_μ as a function of solar ^8B neutrino flux. The constraints are derived with Feldman-Cousins construction at 68% (90%) CL, indicated by the black (gray) curve.




FIG. 4. Measurements of the flux-weighted $\text{CE}\nu\text{NS}$ cross section $\sigma_{\text{CE}\nu\text{NS}}$. The measurement using Xe nuclei solar ${}^8\text{B}$ neutrinos from this work is shown in black. The 90% CL upper limit from XENON1T [7] is shown in blue. The measurements with neutrinos from the SNS by the COHERENT Collaboration using CsI [44] (red), Ar [5] (green), and Ge [45] (orange) nuclei are also shown. For comparison, the SM predictions are shown by vertical dashed lines.

comparison. Because of the lower average energy, the solar ${}^8\text{B}$ neutrino flux-weighted $\text{CE}\nu\text{NS}$ cross section is the lowest one measured to date.

Summary—We performed a blind search for NR signals from solar ${}^8\text{B}$ neutrinos via $\text{CE}\nu\text{NS}$ with XENONnT using data from two science runs with a combined exposure of $3.51 \text{ t} \times \text{yr}$. By lowering the $S1$ and $S2$ thresholds, we are able to include NR signals as low as 0.5 keV. Various techniques are developed to reduce the dominant AC background. Various calibrations, including ${}^{88}\text{YBe}$ and ${}^{37}\text{Ar}$, are performed to understand the detector response, signal, and background modeling. The data disfavor the background-only hypothesis at 2.73σ . The unconstrained best-fit number of ${}^8\text{B}$ $\text{CE}\nu\text{NS}$ signals is $10.7^{+3.7}_{-4.2}$, consistent with the expectation of $11.9^{+4.5}_{-4.2}$ events, based on the measured solar ${}^8\text{B}$ neutrino flux from SNO [23], the theoretical $\text{CE}\nu\text{NS}$ cross section with Xe nuclei [25], and the calibrated detector response to low-energy NRs in XENONnT. Thus, the measured solar ${}^8\text{B}$ neutrino flux is $(4.7^{+3.6}_{-2.3}) \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$, consistent with SNO, and the measured neutrino flux-weighted $\text{CE}\nu\text{NS}$ cross section on Xe is $(1.1^{+0.8}_{-0.5}) \times 10^{-39} \text{ cm}^2$, consistent with the SM prediction. As XENONnT continues to take data, more precise measurements are expected in the future.

Note added—Recently, we noticed the results of the ${}^8\text{B}$ neutrino flux measurement from the PandaX Collaboration with a similar statistical significance in [46].

Acknowledgments—We would like to thank the COHERENT Collaboration for providing data points and predictions for the measurement of flux-weighted $\text{CE}\nu\text{NS}$ cross section $\sigma_{\text{CE}\nu\text{NS}}$ at the SNS. We gratefully acknowledge support from the National Science Foundation, Swiss National Science Foundation, German Ministry for Education and Research, Max Planck

Gesellschaft, Deutsche Forschungsgemeinschaft, Helmholtz Association, Dutch Research Council (NWO), Fundacao para a Ciencia e Tecnologia, Weizmann Institute of Science, Binational Science Foundation, Région des Pays de la Loire, Knut and Alice Wallenberg Foundation, Kavli Foundation, JSPS Kakenhi and JST FOREST Program ERAN in Japan, Tsinghua University Initiative Scientific Research Program, DIM-ACAV+ Région Ile-de-France, and Istituto Nazionale di Fisica Nucleare. This project has received funding and support from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 860881-HIDDeN. We gratefully acknowledge support for providing computing and data-processing resources of the Open Science Pool and the European Grid Initiative, at the following computing centers: the CNRS/IN2P3 (Lyon, France), the Dutch national e-infrastructure with the support of SURF Cooperative, the Nikhef Data-Processing Facility (Amsterdam, Netherlands), the INFN-CNAF (Bologna, Italy), the San Diego Supercomputer Center (San Diego, U.S.), and the Enrico Fermi Institute (Chicago, U.S.). We acknowledge the support of the Research Computing Center (RCC) at The University of Chicago for providing computing resources for data analysis. We thank INFN Laboratori Nazionali del Gran Sasso for hosting and supporting the XENON project.

- [1] D. Z. Freedman, Coherent effects of a weak neutral current, *Phys. Rev. D* **9**, 1389 (1974).
- [2] V. B. Kopeliovich and L. L. Frankfurt, Isotopic and chiral structure of neutral current, *JETP Lett.* **19**, 145 (1974), http://jetpletters.ru/ps/1776/article_27044.shtml.
- [3] A. Drukier and L. Stodolsky, Principles and applications of a neutral current detector for neutrino physics and astronomy, *Phys. Rev. D* **30**, 2295 (1984).
- [4] D. Akimov *et al.* (COHERENT Collaboration), Observation of coherent elastic neutrino-nucleus scattering, *Science* **357**, 1123 (2017).
- [5] D. Akimov *et al.* (COHERENT Collaboration), First measurement of coherent elastic neutrino-nucleus scattering on argon, *Phys. Rev. Lett.* **126**, 012002 (2021).
- [6] J. N. Bahcall and C. Pena-Garay, Solar models and solar neutrino oscillations, *New J. Phys.* **6**, 63 (2004).
- [7] E. Aprile *et al.* (XENON Collaboration), Search for coherent elastic scattering of solar ${}^8\text{B}$ neutrinos in the XENON1T dark matter experiment, *Phys. Rev. Lett.* **126**, 091301 (2021).
- [8] W. Ma *et al.* (PandaX Collaboration), Search for solar B8 neutrinos in the PandaX-4T experiment using neutrino-nucleus coherent scattering, *Phys. Rev. Lett.* **130**, 021802 (2023).
- [9] L. E. Strigari, Neutrino coherent scattering rates at direct dark matter detectors, *New J. Phys.* **11**, 105011 (2009).
- [10] J. Billard, L. Strigari, and E. Figueroa-Feliciano, Implication of neutrino backgrounds on the reach of next generation dark matter direct detection experiments, *Phys. Rev. D* **89**, 023524 (2014).
- [11] C. A. J. O’Hare, New definition of the neutrino floor for direct dark matter searches, *Phys. Rev. Lett.* **127**, 251802 (2021).

- [12] E. Aprile *et al.* (XENON Collaboration), The XENONnT dark matter experiment, *Eur. Phys. J. C* **84**, 784 (2024).
- [13] V. C. Antochi *et al.*, Improved quality tests of R11410-21 photomultiplier tubes for the XENONnT experiment, *J. Instrum.* **16**, P08033 (2021).
- [14] E. Aprile *et al.* (XENON Collaboration), Design and performance of the field cage for the XENONnT experiment, *Eur. Phys. J. C* **84**, 138 (2024).
- [15] E. Aprile *et al.* (XENON Collaboration), The triggerless data acquisition system of the XENONnT experiment, *J. Instrum.* **18**, P07054 (2023).
- [16] E. Aprile *et al.* (XENON Collaboration), Emission of single and few electrons in XENON1T and limits on light dark matter, *Phys. Rev. D* **106**, 022001 (2022).
- [17] E. Aprile *et al.* (XENON Collaboration), XENONnT analysis: Signal reconstruction, calibration and event selection, [arXiv:2409.08778](https://arxiv.org/abs/2409.08778).
- [18] J. Aalbers *et al.*, AxFoundation/strax: Stream analysis for xenon TPCs (2024), [10.5281/zenodo.11355772](https://doi.org/10.5281/zenodo.11355772).
- [19] XENON Collaboration, XENONnT/straxen: Streaming analysis for XENON (2024), [10.5281/zenodo.12608732](https://doi.org/10.5281/zenodo.12608732).
- [20] E. Aprile *et al.* (XENON Collaboration), XENON1T dark matter data analysis: Signal reconstruction, calibration and event selection, *Phys. Rev. D* **100**, 052014 (2019).
- [21] E. Aprile *et al.* (XENON Collaboration), Search for new physics in electronic recoil data from XENONnT, *Phys. Rev. Lett.* **129**, 161805 (2022).
- [22] M. Murra, D. Schulte, C. Huhmann, and C. Weinheimer, Design, construction and commissioning of a high-flow radon removal system for XENONnT, *Eur. Phys. J. C* **82**, 1104 (2022).
- [23] B. Aharmim *et al.* (SNO Collaboration), Combined analysis of all three phases of solar neutrino data from the Sudbury neutrino observatory, *Phys. Rev. C* **88**, 025501 (2013).
- [24] J. N. Bahcall, E. Lisi, D. E. Alburger, L. De Braeckelee, S. J. Freedman, and J. Napolitano, Standard neutrino spectrum from ^8B decay, *Phys. Rev. C* **54**, 411 (1996).
- [25] J. Barranco, O. G. Miranda, and T. I. Rashba, Probing new physics with coherent neutrino scattering off nuclei, *J. High Energy Phys.* **12** (2005) 021.
- [26] J. I. Collar, Applications of an $^{88}\text{Y}/\text{Be}$ photo-neutron calibration source to dark matter and neutrino experiments, *Phys. Rev. Lett.* **110**, 211101 (2013).
- [27] XENON Collaboration, XENONnT/appletree: A high-performance program simulates and fits response of xenon (2024), [10.5281/zenodo.12601629](https://doi.org/10.5281/zenodo.12601629).
- [28] E. Aprile *et al.* (XENON Collaboration), XENONnT WIMP search: Signal & background modeling and statistical inference, [arXiv:2406.13638](https://arxiv.org/abs/2406.13638).
- [29] E. Aprile *et al.* (XENON Collaboration) (to be published).
- [30] E. Aprile *et al.* (XENON Collaboration), First dark matter search with nuclear recoils from the XENONnT experiment, *Phys. Rev. Lett.* **131**, 041003 (2023).
- [31] E. Aprile, C. E. Dahl, L. DeViveiros, R. Gaitskell, K. L. Giboni, J. Kwong, P. Majewski, K. Ni, T. Shutt, and M. Yamashita, Simultaneous measurement of ionization and scintillation from nuclear recoils in liquid xenon as target for a dark matter experiment, *Phys. Rev. Lett.* **97**, 081302 (2006).
- [32] E. Aprile *et al.* (XENON Collaboration), Dark matter search results from a one ton-year exposure of XENON1T, *Phys. Rev. Lett.* **121**, 111302 (2018).
- [33] J. Aalbers *et al.* (LZ Collaboration), First dark matter search results from the LUX-ZEPLIN (LZ) experiment, *Phys. Rev. Lett.* **131**, 041002 (2023).
- [34] A. Kopec, A. L. Baxter, M. Clark, R. F. Lang, S. Li, J. Qin, and R. Singh, Correlated single- and few-electron backgrounds milliseconds after interactions in dual-phase liquid XENON time projection chambers, *J. Instrum.* **16**, P07014 (2021).
- [35] P. Sorensen, Electron train backgrounds in liquid xenon dark matter search detectors are indeed due to thermalization and trapping, [arXiv:1702.04805](https://arxiv.org/abs/1702.04805).
- [36] XENON Collaboration, XENONnT/axidence: Strax-based data-driven accidental coincidence background simulation and peak-level salting (2024), [10.5281/zenodo.12791105](https://doi.org/10.5281/zenodo.12791105).
- [37] XENON Collaboration, XENONnT/fuse: Refactor xenonnt epix and WFSim code (2024), [10.5281/zenodo.11551366](https://doi.org/10.5281/zenodo.11551366).
- [38] M. Szydagis *et al.*, A review of NEST models, and their application to improvement of particle identification in liquid xenon experiments, [arXiv:2211.10726](https://arxiv.org/abs/2211.10726).
- [39] M. Szydagis *et al.*, Noble element simulation technique is used to simulate noble-element energy deposition microphysics (2024), [10.5281/zenodo.6448408](https://doi.org/10.5281/zenodo.6448408).
- [40] S. S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses, *Ann. Math. Stat.* **9**, 60 (1938).
- [41] XENON Collaboration, XENONnT/alea: A tool to perform toymc-based inference constructions (2024), [10.5281/zenodo.10829030](https://doi.org/10.5281/zenodo.10829030).
- [42] G. J. Feldman and R. D. Cousins, A unified approach to the classical statistical analysis of small signals, *Phys. Rev. D* **57**, 3873 (1998).
- [43] E. Aprile *et al.* (XENON Collaboration), Low-energy calibration of XENON1T with an internal ^{37}Ar source, *Eur. Phys. J. C* **83**, 542 (2023).
- [44] D. Akimov *et al.* (COHERENT Collaboration), Measurement of the coherent elastic neutrino-nucleus scattering cross section on CsI by COHERENT, *Phys. Rev. Lett.* **129**, 081801 (2022).
- [45] S. Adamski *et al.*, First detection of coherent elastic neutrino-nucleus scattering on germanium, [arXiv:2406.13806](https://arxiv.org/abs/2406.13806).
- [46] Zihao Bo *et al.* (PandaX Collaboration), preceding Letter, First indication of solar ^8B neutrinos through coherent elastic neutrino-nucleus scattering in PandaX-4T, *Phys. Rev. Lett.* **133**, 191001 (2024).

End Matter

Appendix A: AC sideband validation—The AC sideband validation is also performed with a blind analysis, before unblinding the ^8B CE ν NS search data. After the AC event selection and prediction are both

fixed, the SR0 and SR1 AC sideband datasets are unblinded. With the initial S2 threshold of 100 PE, 133 (416) events are observed in SR0 (SR1) with an expectation of 135.9 (368.2). With the four-dimensional

TABLE II. AC sideband validation. The expected and observed numbers of events are for a 120 (100) PE S_2 threshold.

Science run	Expectation	Observation	p value
SR0	122.7 (135.9)	121 (133)	0.33 (0.74)
SR1	302.5 (368.2)	326 (416)	0.16 (0.03)

binned likelihood GOF test, the prediction and the observation in SR0 show an acceptable agreement. However, the test on SR1 showed a mismatch with a p value of 0.03. All the analysis dimensions are inspected and the mismatch is only present below 120 PE in S_2 suggesting that the mismatch in SR1 is most likely due to the increase in photoionization. The S_2 thresholds for both SR0 and SR1 are thus conservatively increased for the ${}^8\text{B}$ CE ν NS search data, with minor loss in the discovery potential of the solar ${}^8\text{B}$ CE ν NS signals. The

final prediction of the AC background and observations in the AC sideband are shown in Table II. The projection of the four analysis dimensions with the same binning used in the ${}^8\text{B}$ CE ν NS search in sideband data with the S_2 larger than 120 PE in both SR0 and SR1 data are shown in Fig. 5.

Appendix B: Modeling validation—The signal and background modeling is validated by the measurement of L_y of ${}^{37}\text{Ar}$ L shell EC, which is performed with a blind analysis. The AC background in this measurement is estimated to be 1062 ± 53 based on a similar modeling approach to that in the ${}^8\text{B}$ CE ν NS search. After unblinding, 1676 events are observed. The observed events above the expected AC background are strongly validated by a four-dimensional GOF test, yielding a p value of 0.92. Figure 6 shows the observed




FIG. 5. Distributions of expected AC background in AC sideband and the observed data in the projected analysis dimensions. Both expectation and observation have S_2 larger than 120 PE. The expected AC background is shown in the purple histogram. The observed number of events with Poisson uncertainties in each bin is shown in black.




FIG. 6. Distributions of the best-fit AC background, ${}^{37}\text{Ar}$ L shell EC signal, and the observed data in the ${}^8\text{B}$ CE ν NS analysis dimensions. The observed number of events with Poisson uncertainties in each bin is shown in black. The ${}^{37}\text{Ar}$ signal (AC background) is shown by the green (purple) histogram.




FIG. 7. Distributions of best-fit signal and background, together with the data in the projected analysis dimensions, with SR0 and SR1 shown in the left and right column, respectively. The observed number of events with Poisson uncertainties in each bin is shown in black. The ${}^8\text{B}$ CE ν NS signal is represented by the light green histogram on top of the backgrounds, which are indicated by purple (AC), blue (ER), and yellow (neutron) histograms.

events in the same analysis dimensions as the ${}^8\text{B}$ CE ν NS search along with the AC background and the best-fit ${}^{37}\text{Ar}$ L shell EC signal during the ${}^{37}\text{Ar}$ calibration. The measurement will be presented in a future publication.

Appendix C: Separate SR0/SR1 best-fit results—The distributions of the observed events in SR0 and SR1 and the corresponding best-fit model projected to each analysis dimension are shown individually in Fig. 7.