



This is a repository copy of *Great expectations? GPs' estimations of time required to deliver BMJ's '10 minute consultations'*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/222415/>

Version: Published Version

Article:

Bradley, S.H. orcid.org/0000-0002-2038-2056, Harper, A.M. orcid.org/0000-0001-5866-2128, Smith, L. orcid.org/0000-0002-4280-6323 et al. (6 more authors) (2024) Great expectations? GPs' estimations of time required to deliver BMJ's '10 minute consultations'. *BMJ Open*, 14 (2). e079578. ISSN 2044-6055

<https://doi.org/10.1136/bmjopen-2023-079578>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

BMJ Open Great expectations? GPs' estimations of time required to deliver BMJ's '10 minute consultations'

Stephen Henry Bradley ^{1,2}, Alice M Harper ³, Lesley Smith ⁴, Nigel Taylor,⁵ Harriet Delap,⁶ Helen Pyke,⁶ James Girkin,⁷ Carol Sinnott ^{8,9}, Jessica Watson ³

To cite: Bradley SH, Harper AM, Smith L, *et al*. Great expectations? GPs' estimations of time required to deliver BMJ's '10 minute consultations'. *BMJ Open* 2024;**14**:e079578. doi:10.1136/bmjopen-2023-079578

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2023-079578>).

Received 05 September 2023
Accepted 09 November 2023



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Academic Unit of Primary Care, University of Leeds, Leeds, UK

²York Street Practice, Leeds, UK

³Centre for Academic Primary Care, University of Bristol, Bristol, UK

⁴Leeds Institute of Health Sciences, University of Leeds, Leeds, UK

⁵Bradgate Surgery, Brentry, UK

⁶West Walk Surgery, Yate, UK

⁷Grove Medical Centre, Felixstowe, UK

⁸The Healthcare Improvement Studies (THIS) Institute, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

⁹Court St Medical Practice, Enniscorthy, Ireland

Correspondence to

Dr Stephen Henry Bradley; medsbra@leeds.ac.uk

ABSTRACT

Objectives

1. To estimate the time required to undertake consultations according to BMJ's 10-minute consultation articles.
2. To quantify the tasks recommended in 10-minute consultation articles.
3. To determine if, and to what extent, the time required and the number of tasks recommended have increased over the past 22 years.

Design Analysis of estimations made by four general practitioners (GPs) of the time required to undertake tasks recommended in *BMJ's 10-minute consultation* articles.

Setting Primary care in the UK.

Participants Four doctors with a combined total of 79 years of experience in the UK National Health Service following qualification as GPs.

Main outcome measures Median minimum estimated consultation length (the estimated time required to complete tasks recommended for all patients) and median maximum estimated consultation length (the estimated time required to complete tasks recommended for all patients *and* the additional tasks recommended in specific circumstances). Minimum, maximum and median consultation lengths reported for each year and for each 5-year period.

Results Data were extracted for 44 articles. The median minimum and median maximum estimated consultation durations were 15.7 minutes (IQR 12.6–20.9) and 28.4 minutes (IQR 22.4–33.8), respectively. A median of 17 tasks were included in each article. There was no change in durations required over the 22 years examined.

Conclusions The approximate times estimated by GPs to deliver care according to *10-minute consultations* exceed the time available in routine appointments. '10 minute consultations' is a misleading title that sets inappropriate expectations for what GPs can realistically deliver in their routine consultations. While maintaining aspirations for high-quality care is appropriate, practice recommendations need to take greater account of the limited time doctors have to deliver routine care.

INTRODUCTION

Across high-income countries, general practitioners (GPs) report excessive workloads with concerns that their ability to deliver

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ A novel design was employed whereby general practitioners (GPs) estimated the duration that would be required to deliver care as recommended in a long-standing article series.
- ⇒ Analysis was undertaken according to a pre-registered plan and all data have been made freely available.
- ⇒ There was substantial variation between estimates provided by GPs which could reflect differences in individual practice but may also be affected by difficulty in accurately estimating how long tasks would take to perform.

high-quality care is threatened by rising demand that exceeds the capacity of the medical workforce.^{1 2} In several countries the situation has worsened further since the beginning of the coronavirus pandemic.¹ Pressures are known to be particularly severe in the UK, where GPs have experienced simultaneous increases in patient contacts³ and declining numbers of full-time equivalent GPs⁴ both in absolute terms and in relation to the ratio of patients per GP,⁵ despite efforts to increase GP numbers.⁶ Appointments in general practice in the UK are among the shortest in Europe, with a standard appointment time of 10 minutes in most practices.⁷ While there are calls to lengthen appointment duration to at least 15 min,^{8–10} this could exacerbate difficulties in access to primary care where GPs report having an average of 37 patient contacts per day.¹¹ Prior research has demonstrated that delivering primary care, as specified by guidelines, is not feasible within the time that GPs have available.¹² Accordingly, it has been proposed that guideline bodies, such as the National Institute for Health and Care Excellence, should consider the time that would be taken to deliver them in routine practice.¹³



By contrast, *BMJ's 10-minute consultation* series are education articles orientated chiefly at primary care and explicitly aim to outline what a GP should cover in a time-limited consultation.¹⁴ While guidelines describe the recommended management of a limited number of specific diseases, *10-minute consultation* articles are generally themed around a presenting sign, symptom or concern that prompts a patient to seek medical assessment. Typically authored by GPs alongside specialists and spanning a broad range of topics, *10-minute consultations* constitute a valuable corpus of contemporaneously recommended practice by clinical peers.

In this study, we examined 22 years of *10-minute consultation* articles to determine if the recommendations they offer could be realistically implemented in 10 minutes. We also aimed to ascertain if expectation-inflation had occurred during that period, that is, if there were now greater expectations for how many tasks should be undertaken, and the time it would take to perform these, compared with 22 years ago.

METHODS

We obtained the full texts of 44 *10-minute consultation* articles (21% of the entire series) including the first published article in the series (October 2000), and thereafter the first and last articles for each year between 2001 and 2021, and the first article published in 2022. All included articles are listed in online supplemental table S1. From these, one author (AH) extracted all tasks that were recommended by the authors and categorised these by type (eg, history, examination, organising investigations, reviewing records, prescribing, shared decision-making, referral, etc). These categories are listed in table 1. Since some tasks were contingent on particular characteristics of the presentation or patient (eg, if a particular symptom was elicited, or patient sex), we categorised tasks according to whether or not they would be required for all patients or required only for specific circumstances.

A spreadsheet listing the tasks extracted from each article was created. Articles were listed in a random, not chronological order. Four qualified GPs (NT, HD, HP and JG), who cumulatively had more than 79 years of experience as GPs in the National Health Service (NHS), independently determined the amount of time, rounded to the nearest half minute, which they thought each task would take. The participating GPs were purposively recruited as clinicians who were not academics and who spend the majority of their working week delivering primary healthcare (online supplemental table S2).

Statistical analysis

The median total number of tasks and the median number of tasks for each category of task (eg, history, examination, etc) were determined for the entire dataset of 44 articles. For each article, the maximum estimated consultation length (or time required to complete all tasks including those recommended for only specific

Table 1 Categories of tasks and median number of tasks for each category for the 44 *10-minute consultation* articles examined

Task category (n)	Median number of tasks relating to this category across the 44 articles (range)
History (262)	5 (1–17)
Examination (76)	1 (0–8)
History and examination (10)	0 (0–2)
Organising investigations (56)	1 (0–5)
Review records (8)	0 (0–2)
Explanation/advice (136)	3 (0–9)
Obtaining consent (2)	0 (0–1)
Use clinical tool/calculator (13)	0 (0–2)
Shared decision-making (12)	0 (0–3)
Review medication (7)	0 (0–10)
Prescription (47)	1 (0–3)
Referral (83)	1 (0–10)
Signposting (7)	0 (0–2)
Arrange hospital admission (3)	0 (0–1)
Safety netting advice (5)	0 (0–1)
Follow-up (55)	1 (0–6)
Other (26)	0 (0–4)
The total number of tasks across all articles was 808.	

circumstances) and the minimum estimated consultation length (or time required to complete those tasks that applied to all patients) were calculated for each GP. Medians were calculated for the minimum and maximum estimated consultation lengths using the estimates from all 4 GPs

We also determined the median and range of red flag diagnoses mentioned in order to indicate the expectation placed on GPs to remain mindful of infrequent diagnoses. No additional time was added to account for such mention of red flag symptoms, other than specific tasks which were required to elicit these, for example, history taking or examination.

For each consultation, we added 2 minutes 38 seconds which was the median time calculated for the following three tasks in a time-motion observation study of 61 NHS GPs¹⁵:

- ▶ Preparing to see patients (eg, reading patient's notes) (median time=47 seconds).
- ▶ Calling patients into the consultation room (median time=36 s).
- ▶ Documentation of consultation in the electronic health record (median time=1 minute 15 seconds).

To assess the inter-rater reliability between the four GPs for the estimated minimum and maximum durations the intraclass correlation coefficient (ICC) and 95% CIs were calculated based on the mean absolute agreement two-way random effects model.

To ascertain if expectations had increased over time, median minimum and maximum estimated

consultation lengths were charted by year and grouped into 5-year periods. Trends for median estimated consultation length over time were assessed using linear regression modelling. For these models, we assessed the year-on-year trend assuming a linear trend and also assessing changes over 5-year time periods.

More information regarding the analysis is provided in the pre-registered protocol.¹⁶ There were two substantive deviations from the protocol. First, we included three articles prior to the stated starting point of 2002 in order to include articles from the beginning of the series in 2000. Second, the protocol stated we would add the mean, rather than the median, additional times added for standard tasks as median times were not available when the protocol was registered.¹⁵ Sensitivity analyses were undertaken as per original protocol.

Patient and public involvement

Patients and the public were not involved in this study.

RESULTS

A median of 17 tasks were outlined per article, 10 of which were required for all patients. The median minimum estimated consultation length was 15.7 min (IQR 12.6–20.9) and the median maximum estimated consultation length was 28.4 min (IQR 22.4–33.8). The ‘maximum estimated consultation length’ refers to the total duration of all tasks listed in the

article, although for several articles all tasks would not plausibly pertain to a single patient. The median minimum estimated consultation lengths for all articles exceeded 10 minutes. Figure 1 presents the estimated consultation lengths for 10-minute consultations by year of publication category, while figure 2 presents the estimated consultation lengths for each of the four GPs individually.

Articles included a median of one red flag diagnosis (range 0–8) that GPs should consider in their assessments. Data for the entire period 2000–2022 and in 5-year intervals (with the exception of the 3-year interval 2020–2022) are presented in table 2 and figure 1. The median number of tasks per task category across the 44 papers is presented in table 1, and online supplemental table S3 presents the estimated durations for each task category. The most frequent tasks related to history taking and providing explanations and advice. A small number of the tasks which were not required for all patients pertained to situations which were mutually exclusive, for example, two tasks outlining actions if red flag symptoms were or were not present. These are summarised in online supplemental table S4.

Linear regression demonstrated no significant changes over time in estimated consultation lengths, either when grouping the articles into 5-year (aside from 2020 to 2022) periods or when assessing for annual change (table 3).




Figure 1 Box plot of minimum and maximum estimated consultation lengths for 10-minute consultations by year of publication category.




Figure 2 Participating general practitioners' (GP) estimates of minimum estimated consultation lengths required for 10-minute consultation articles across the 44 included articles. The combined data are the median of all four GP estimates. Median (IQR) minimum durations are GP 1: 12.1 (9.4, 15.6), GP 2: 19.5 (15.6, 25.4), GP 3: 9.9 (8.6, 12.4), GP 4: 26.4 (18.9, 31.6), combined: 15.7 (12.6, 20.9).

ICC demonstrated consistency that was moderate between the GPs' estimates of minimum estimated consultation lengths (0.68, 95% CI 0.35 to 0.83) and poor for maximum estimated consultation lengths (0.46, 95% CI 0.07 to 0.70). There was substantial discordance between GPs' estimates of consultation lengths across all 44 articles. Figure 2 and online supplemental figure S1 show each GP's median and maximum estimated consultation lengths, respectively, across all 44 articles. The lowest median minimum estimated consultation length across the 44 articles was 9.9 min (IQR 8.6–12.4), while their corresponding median maximum estimated consultation length was 14.6 min (IQR 13.1, 18.4).

Sensitivity analyses were consistent with the findings outlined above (online supplemental tables S5 and S6 and figures S2–S4).

DISCUSSION

Summary of principal findings

Using estimates from four experienced GPs, this analysis indicates that the care recommended in 10-minute consultations exceeds the time available to GPs to deliver this care. However, estimates of time required did not increase according to publication year.

Interpretation and comparison with other studies

Previous work has demonstrated that the time required to deliver care recommended by clinical guidelines greatly exceeds the time which is available to clinicians in routine consultations.^{12 13} 10-minute consultations are not guidelines and they are explicitly written with respect to care that might reasonably be expected to be delivered within a single standard consultation. To our knowledge, this study is the first that has assessed the time required

Table 2 Median number of tasks in the 44 articles categorised according to whether they apply to all patients (median minimum estimated consultation length) or also included tasks relating to specific circumstances only (median maximum estimated consultation length)

Period	Median number of tasks recommended for all patients (IQR)	Median minimum estimated consultation length (IQR)	Median number of tasks, including those for specific circumstances only (IQR)	Median maximum estimated consultation length (IQR)
Overall (2000–2022)	10 (7, 12)	15.7 (12.6, 20.9)	17 (14, 22)	28.4 (22.4, 33.8)
2000–2004	6 (6, 8)	14.1 (11.4, 18.1)	15 (12, 15)	27.1 (22.4, 31.4)
2005–2009	10 (8, 12)	16.4 (14.1, 21.9)	18 (13, 22)	24.5 (21.9, 34.6)
2010–2014	14 (10, 17)	18.5 (15.3, 21.9)	21 (17, 28)	31.4 (29.6, 33.8)
2015–2019	10 (8, 12)	13.2 (11.9, 20.1)	17 (13, 24)	27.7 (21.6, 40.6)
2020–2022	9 (8, 12)	15.1 (14.9, 16.1)	15 (14, 15)	22.4 (22.4, 25.4)

Table 3 Results from linear regression models of change in estimated consultation lengths (minutes) both annually (2000–2022) assuming a linear trend and compared by grouping into categories

	Change per year/period for minimum estimated consultation lengths (95% CI)	Change per year/period for maximum estimated consultation lengths (95% CI)
Annual	−0.06 (−0.41 to 0.30)	0.01 (−0.38 to 0.39)
Publication year category		
2000–2004	Baseline	Baseline
2005–2009	3.79 (−3.28 to 10.86)	0.69 (−6.84 to 8.22)
2010–2014	3.13 (−3.94 to 10.20)	3.08 (−4.45 to 10.61)
2015–2019	1.61 (−5.45 to 8.68)	2.22 (−5.31 to 9.75)
2020–2022	−0.46 (−9.04 to 8.12)	−2.66 (−11.80 to 6.58)

to deliver care according to brief and accessible pragmatic recommendations pitched directly to clinicians, as opposed to formal clinical guidelines. Greater inclusion of, and consultation, with ‘front-line’ clinicians including GPs has been suggested to help ensure guidelines and recommendations are realistic. Estimates of the time required to implement *10-minute consultations* are lower than estimates relating to clinical guidelines, but meeting the standards recommended by the articles is still not realistic within a standard general practice appointment. *10-minute consultation* articles are typically authored by GPs, suggesting their inclusion in formulating recommendations should not be considered a panacea. It may be that authors tend to include more tasks than can be undertaken within 10 minutes to anticipate criticisms that important considerations had been overlooked or it may be that additional tasks were identified during the peer review process. Alternatively, it could be the case that there is a fundamental dissonance between presenting recommendations that are sufficiently comprehensive as to be deemed defensible practice in a prestigious journal with the limits of what can actually be achieved in so short a period. However, it may also be the case that *10-minute consultation* articles describe relatively more complicated problems than are typical and that such presentations may, to some extent, be counterbalanced by very straightforward presentations which GPs can conclude in less than 10 minutes.

Delivering *10-minute consultations* within the stated time frame could be even more challenging than our work indicates. Direct observation has demonstrated that GPs frequently contend with time and attention-consuming system failures and interruptions during consultations.¹⁵ In addition, GPs normally address several problems in a consultation—not just one, as this analysis is predicated on. An analysis of recorded consultations published in 2013 reported a mean of 2.5 problems discussed (95% CI 2.3 to 2.6, IQR 1–8).¹⁷

Strengths and limitations of the study

The study employed novel methodology to estimate how long recommended aspects of clinical care could take to implement in routine clinical appointments, by asking clinicians themselves to estimate how long instituting

these tasks could take, rather than drawing on estimates from literature or routinely collected data. The study was also novel in that it examined recommendations authored by clinical peers including GPs, rather than official guidelines from regulatory bodies or disease or organ-specific societies. Therefore, the recommendations that we examined might be supposed to be more sensitive to time constraints in general practice.

Enabling real-world implementation of guidelines requires guideline bodies to consider clinician time in a structured way. Approaches for measuring clinician time include time-motion methods, self-report through retrospective interview, self-report through time sheets and patient flow analysis.¹⁸ The time estimates generated by these different approaches can vary significantly, but this variation is most marked for clinician activities carried out when clients are not present. Time-motion studies, which are often considered the gold standard approach for measuring clinician time, have been criticised for variability in the specification and categorisation of observed tasks, the approach to multitasking and interruptions, and the rigour of training and reliability of results between observers.¹⁹ All of the approaches, by their very presence, might change the activity patterns and time estimates of the events under observation (eg, Hawthorne effect).

The approach we employed allowed us to generate estimated timings for the very specific tasks outlined in *10-minute consultations*, which would not have otherwise been possible. However, several limitations of relying on GPs’ own estimations of time required to deliver care should be acknowledged. The estimate of consultation durations might differ substantially if data was collected from observing consultations, both because of the multiplicity of patient, system and clinician influences on consultations and because of the risk that individual clinicians are not able to accurately estimate how long consultation tasks take. It is also possible that if GPs were observed and timed undertaking tasks they might employ pragmatic strategies, or ‘short cuts’ to help consult within allotted appointment times. However, observing enough consultations in undifferentiated clinic lists to ensure that all the situations outlined in the *10-minute consultation* could be observed would not be feasible, nor would it be possible



to ensure that GPs carried out all the tasks that were recommended so that these could all be timed.

While the total duration estimated for all the tasks in each article is described as the 'maximum estimated consultation length', for several articles it is not plausible that all described circumstances could apply to a single patient. Ten tasks out of 808 (1.2%) were identified which described circumstances which could not apply to a single patient, for example, separate actions to undertake if red flag symptoms were or were not present. These are reproduced in online supplemental table S4.

The clinical problems which are covered in *10-minute consultations* are not necessarily representative of those most commonly encountered in primary care. Indeed, as the topics chosen were deemed suitable for elaboration in an education article they might be considered more complex or difficult. It is important to recognise that the study aimed to examine the feasibility of performing tasks as outlined in *10-minute consultations* and cannot demonstrate whether the presentations which actually do typically present to primary care can be performed satisfactorily within 10 minutes. However, reference to the titles of the 44 articles examined in this study (online supplemental table S1) suggests the topics were not unusually complex and the majority could be considered typical problems presenting to primary care.

Our study is a pragmatic attempt to understand whether the volume of tasks described in *10-minute consultations* might reasonably be accomplished in that time frame and whether the burden of expectation has grown over the last 22 years. The study provides persuasive evidence that the consultations described in the series cannot be expected to be undertaken in 10 minutes. We did not find evidence that the time that would be required to deliver care according to authors' recommendations has increased, but there were only a relatively small number of articles within each time period so it is possible that any differences in terms of clinical topics between time periods could have overwhelmed any smaller differences, for example, any underlying tendency for expectations to have increased over time.

Inter-rater reliability was moderate or poor. This discordance may reflect different working habits of the GPs and variations in consulting speeds. It may also reflect difficulty that GPs had in estimating how long tasks would actually take with accuracy and that estimates were generated by only four GPs. It is also possible that GPs interpreted instructions, which were to estimate how long tasks would take 'in routine primary care' differently and their estimates may have been varying influenced by the real-world existence of time restraints.

An additional limitation is that the study did not involve patients. The project was not funded and we were reluctant to request that patients also contribute their time and expertise without reimbursement.

The study analysis plan was pre-registered, thereby mitigating the possibility of post hoc calibration of the analysis to deliver particular findings, a potential weakness of

many observational studies. Similarly, data were extracted from the *10-minute consultation* articles in a robust and reproducible manner and all study data and the analysis code has been shared, so that readers can explore the data and judge whether they feel the estimated timings reflect their own experience.

Policy implications

This study provides evidence supporting calls for increasing appointment length. There are important barriers to abandoning the *10-minute consultation* model, particularly the reduction in the availability of total number of appointments to accommodate longer consultations. While this situation cannot be readily remedied, clinicians and health service leaders could acknowledge explicitly that appointment durations are not long enough to deliver standards of care recommended by members of the profession.

Projecting the median minimum consultation time in this study (15.7 minutes) across 37 consultations per day¹¹ each with 2.5 problems would necessitate GPs spending over 24 hours/day consulting with patients, not including any additional time for breaks or administrative tasks. The sheer impossibility of delivering care as recommended at the volume of demand faced by GPs indicates the need for health service leaders to temper high expectations on behalf of patients with a sense of pragmatism and understanding for those working in primary care.

Recent policy, particularly in England, has sought to involve and broaden the general practice team to include several non-medically qualified clinicians, in order to allow GPs to focus particularly on patients with more complex problems.²⁰ It is not yet understood whether this approach could help GPs to offer longer appointments for patients with more complex needs or less straightforward presentations. Remote consultations have previously been hailed as a means to free up time for GPs, however, experience suggests that while the technology has important benefits in terms of convenience of access it does not necessarily reduce time required for GPs.²¹

As an immediate measure we recommend that the *10-minute consultation* article series is renamed. The articles provide useful exemplars of primary care consultations, but labelling these as *10-minute consultations* perpetuates unrealistic expectations of what can be achieved within a short appointment. Given the well-documented pressures on primary care,²² this risks contributing to moral distress of GPs who wish to provide high-quality care as outlined in *10-minute consultations*, but who would not routinely be able to do so within the time they have available. Authors and editors of *10-minute consultations* could also acknowledge the intense time pressures GPs work under and for each problem identify a core set of essential tasks that could realistically be undertaken within 10 minutes.

CONCLUSION

This research indicates that the primary care consultations outlined in *BMJ's* long-running education article series take longer than a standard appointment length and that the title *10-minute consultations* is a misnomer. *BMJ* should consider renaming the article series in order to avoid placing unrealistic expectations on GPs.

Twitter Carol Sinnott @cmsinnott and Jessica Watson @drjessicawatson

Acknowledgements The authors are grateful to a patient reviewer, Mr Mike Etkind who contributed additional insights in the absence of dedicated patient involvement for this study. In particular, Mr Etkind highlighted the need to address the broader policy context and pragmatic decisions clinicians need to make around implementing recommendations within time constraints.

Contributors SHB conceived the study in discussion with JW and authored the first draft of the manuscript. All authors contributed to subsequent revisions of the manuscript. SHB prepared the study protocol with JW, AH and LS contributing revisions. AH performed the data extraction. NT, HD, HP and JG provided the estimated timings for tasks. CS provided data on timings for standard tasks in GP consultations. LS undertook the statistical analyses.

Funding CS is supported by The Healthcare Improvement Studies (THIS) Institute. The article processing fee was funded by the University of Bristol Clinical Academic Training School; The Healthcare Improvement Studies Institute (THIS), University of Cambridge; and Leeds Institute for Health Sciences, University of Leeds. THIS Institute is funded by the Health Foundation (Grant/Award Number:RHZF/001 - RG88620), an independent charity committed to bringing about better health and health care for people in the UK.

Competing interests SHB, CS, NT, HD, HP, JG and JW are general practitioners (GP), while AH is a GP trainee.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Study protocol and data are available at: <https://osf.io/5shpv/>. Code is available at: <https://github.com/LFairleySmith/TenMinConsultation>

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Stephen Henry Bradley <http://orcid.org/0000-0002-2038-2056>

Alice M Harper <http://orcid.org/0000-0001-5866-2128>

Lesley Smith <http://orcid.org/0000-0002-4280-6323>

Carol Sinnott <http://orcid.org/0000-0002-8620-7461>

Jessica Watson <http://orcid.org/0000-0002-8177-6438>

REFERENCES

- 1 Beech J, Fraser C, Gardner T, *et al*. Stressed and overworked. *The Health Foundation* 2023. Available: <https://www.health.org.uk/publications/reports/stressed-and-overworked> [Accessed 23 Jun 2023].
- 2 Boniol M, Kunjumen T, Nair TS, *et al*. The global health workforce stock and distribution in 2020 and 2030: a threat to equity and “universal” health coverage? *BMJ Glob Health* 2022;7:e009316.
- 3 Performance Tracker 2022/23: spring update - general practice 2023. Institute for government. n.d. Available: <https://www.instituteforgovernment.org.uk/performance-tracker-2022-23/general-practice>
- 4 General Practice. NHS England. 2022. Available: <https://digital.nhs.uk/data-and-information/publications/statistical/general-and-personal-medical-services/31-december-2022> [Accessed 23 Jun 2023].
- 5 Office for National Statistics. Trends in patient-to-staff numbers at GP practices in England, . 2022 Available: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthcaresystem/articles/trendsinpatienttostaffnumbersatgppracticesinengland/2022> [Accessed 23 Jun 2023].
- 6 Conservative Party manifesto. The conservative party. 2019. Available: <https://www.conservatives.com/our-plan/conservative-party-manifesto-2019> [Accessed 23 Jun 2023].
- 7 Irving G, Neves AL, Dambha-Miller H, *et al*. International variations in primary care physician consultation time: a systematic review of 67 countries. *BMJ Open* 2017;7:e017902e017902.
- 8 Safe working in general practice: British Medical Association. Available: <https://www.bma.org.uk/advice-and-support/gp-practices/managing-workload/safe-working-in-general-practice> [Accessed 23 Jun 2023].
- 9 Spence D. Bad Medicine: It's time for 15-minute appointments. *Br J Gen Pract* 2019;69:306.
- 10 Iacobucci G. Scrap 10 minute consultations, RCGP urges. *BMJ* 2019;i2270.
- 11 Merrifield N. GPs working average 11-hour day, major survey reveals. *Pulse* Available: <https://www.pulsetoday.co.uk/news/workload/gps-working-average-11-hour-day-major-survey-reveals> [Accessed 23 Jun 2023].
- 12 Porter J, Boyd C, Skandari MR, *et al*. Revisiting the time needed to provide adult primary care. *J Gen Intern Med* 2023;38:147–55.
- 13 Johansson M, Guyatt G, Montori V. Guidelines should consider clinicians' time needed to treat. *BMJ* 2023;380:e072953.
- 14 Article types: BMJ, Available: <https://www.bmj.com/about-bmj/resources-authors/article-types> [Accessed 23 Jun 2023].
- 15 Sinnott C, Moxey JM, Marjanovic S, *et al*. Identifying how GPs spend their time and the obstacles they face: a mixed-methods study. *Br J Gen Pract* 2022;72:e148–60.
- 16 Gps estimations of duration that would be required to deliver BMJ's 'ten minute consultations': open science framework. Available: <https://osf.io/5shpv/> [Accessed 23 Jun 2023].
- 17 Salisbury C, Procter S, Stewart K, *et al*. The content of general practice consultations: cross-sectional study based on video recordings. *Br J Gen Pract* 2013;63:e751–9.
- 18 Bratt JH, Foreit J, Chen PL, *et al*. A comparison of four approaches for measuring clinician time use. *Health Policy Plan* 1999;14:374–81.
- 19 Lopetegui M, Yen P-Y, Lai A, *et al*. Time motion studies in healthcare: what are we talking about? *J Biomed Inform* 2014;49:292–9.
- 20 NHS England. Next steps for integrating primary care: fuller stocktake report. Available: <https://www.england.nhs.uk/publication/next-steps-for-integrating-primary-care-fuller-stocktake-report/> [Accessed 1 Aug 2023].
- 21 Greenhalgh T, Ladds E, Hughes G, *et al*. Why do GPs rarely do video consultations? Qualitative study in UK general practice. *Br J Gen Pract* 2022;72:e351–60.
- 22 Salisbury H. Helen Salisbury: fictive schedules and the unbearable load of general practice. *BMJ* 2023;p1038.