



This is a repository copy of *Modelling the effects of vegetation and urban form on air quality in real urban environments: A systematic review of measurements, methods, and predictions*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/222758/>

Version: Accepted Version

Article:

Yao, M., Smith, M. and Peng, C. orcid.org/0000-0001-8199-0955 (2025) Modelling the effects of vegetation and urban form on air quality in real urban environments: A systematic review of measurements, methods, and predictions. *Urban Forestry & Urban Greening*, 105. 128693. ISSN 1618-8667

<https://doi.org/10.1016/j.ufug.2025.128693>

© 2025 The Authors. Except as otherwise noted, this author-accepted version of a journal article published in *Urban Forestry & Urban Greening* is made available via the University of Sheffield Research Publications and Copyright Policy under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:


<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Graphical Abstract

Modelling the effects of vegetation and urban form on air quality in real urban environments: A systematic review of measurements, methods, and predictions

Mengxue Yao^a, Michael Smith^b, Chengzhi Peng^{a*}

^aSchool of Architecture and Landscape, The Arts Tower, University of Sheffield, S10 2TN, UK

^bDepartment of Computer Science, Regent Court, University of Sheffield, S1 4DP, UK

*Corresponding author

Email address: c.peng@sheffield.ac.uk (Dr Chengzhi Peng)

This is the manuscript was accepted for publication by *Urban Forestry & Urban Greening* on 23 January 2025. [Received 25 October 2024, Revised 22 January 2025, Accepted 23 January 2025, Available online 31 January 2025.]

Mengxue Yao, Michael Smith, Chengzhi Peng (2025), Modelling the effects of vegetation and urban form on air quality in real urban environments: A systematic review of measurements, methods, and predictions, *Urban Forestry & Urban Greening*, 2025, 128693, DOI:10.1016/j.ufug.2025.128693.

Highlights

Modelling the effects of vegetation and urban form on air quality in real urban environments: A systematic review of measurements, methods, and predictions

- *A systematic review of 79 studies on vegetation, urban form and air quality*
- *PM_{2.5} is most used among the nine groups of air quality indicators identified*
- *Increased uses of air quality data sourced from ground stations and satellites*
- *A variety of indices measuring vegetation and urban morphology attributes*
- *Predictive air quality models perform better helped by computational learning*

Modelling the effects of vegetation and urban form on air quality in real urban environments: A systematic review of measurements, methods, and predictions

Abstract

Air pollution poses a significant threat to public health and well-being. In recent decades, researchers have used direct measurements and predictive modelling to assess urban air quality. However, the impact of vegetation and urban form on air quality remains uncertain, particularly regarding their interconnected roles. This paper systematically reviews studies on real urban environments, focusing on how vegetation and urban form influence air quality assessment and prediction. It highlights key variables and their importance, as reported in the literature, and identifies areas needing further research to improve predictions of vegetation's effects on urban air quality in relation to urban morphology.

Keywords: Urban air quality, Air pollutants, Vegetation indices, Urban morphological indicators, Air quality predictions

1. Introduction

1.1. Background and importance

Air pollution has long been a by-product of energy extraction from carbon-based combustion, driving production and consumption at scales that significantly threaten human and ecological health (Perera, 2018, Myers et al., 2013, Tong et al., 2022). As social-economic developments evolve, the sources and interactions of air pollutants have become more diverse and complex. For instance, the primary air pollutants commonly observed in Europe include particulate matter (PM), black carbon (BC), sulphur oxides (SO_x), nitrogen oxides (NO_x), ammonia (NH₃), carbon monoxide (CO), methane (CH₄), non-methane volatile organic compounds and certain metals and poly-cyclic aromatic hydrocarbons (EEA). The main known sources of air pollution are associated with human activities, including industrial emissions (Azarov et al., 2017), traffic emissions (Bai et al., 2022), agricultural fires (Khanal et al., 2022), and household emissions (Apte and Salvi, 2016). Natural processes, such as volcanic eruptions, dust storms, atmospheric inversions, can also cause or exacerbate air pollution (Burhan and Mukminin, 2020).

According to the “*State of Global Air 2020*” report (hei, 2020), air pollution ranks as one of the leading causes of premature death and is closely linked to a variety of diseases. Exposure to air pollution has been strongly associated with specific health outcomes, including stroke, ischemic heart disease, chronic obstructive pulmonary disease, lung cancer, and pneumonia. Air quality has been a focus of global attention, and a number of air quality control standards, guidelines, laws, policies and agreements have been signed, such as the *WHO Global Air Quality Guidelines* (Organization, 2021), *the Clean Air Programme for Europe* (Amann et al., 2005), and *the UK Clean Air Act* (Act, 1970).

The relationship between urban form and air quality began to attract researchers' attention as early as the

1970s (Tolley and Cohen, 1976, Capannelli et al., 1977), including studies on urban vegetation (Nadel et al., 1977, Smith and Staskawicz, 1977). However, the relationships between vegetation and built urban form (i.e., urban morphological characteristics) affecting air quality are far from conclusive. In some studies, vegetation was considered a component of urban form, influencing aspects of vertical structures of an urban environment. On the other hand, vegetation was distinguished from urban form due to functional differences (e.g., natural vs. man-made and ecosystem services vs. structural functions) and the use of different metrics of measurement and analysis (e.g., biomass vs. construction density). Consequently, the impact of vegetation on air quality is often assessed under the influences of built urban form, intentionally or otherwise, making it challenging to isolate and quantify the effects of vegetation. Better understanding of the effects on air quality due to vegetation in real urban environments is required to inform urban greening planning and design decision-making.

1.2. Objectives and scope

This research aligns with the United Nations Sustainable Development Goals. Specifically, it supports Indicator 11.6.2 under Target 11.6, which specifies the annual mean levels of fine particulate matter (e.g. PM_{2.5} and PM₁₀) in cities (population weighted) (Division, 2023). By addressing critical gaps in understanding the relationships between vegetation and air quality in real urban environments, this review contributes to actionable insights that can guide urban planning and policy development to achieve this target. We focus on the studies of urban air quality that examine the effects of vegetation in relation to (built) urban form characteristics. In **Section 2**, we first summarise the questions and findings from the previous review articles published during 2015-2023, to identify the areas for a new systematic review. In **Section 3**, we explain how the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) principles and protocol were applied to identify a set of 79 papers meeting the inclusion and exclusion criteria. **Section 4** comprises five subsections that provide detailed analyses of the study methods and results under review: each corresponding to specific review questions. In Section 5, we discuss the key findings and implications of this systemic review in terms of the research transitions observed and the limitations of our review. We then conclude in Section 6 the significance of this review and the key pointers to further research.

2. Summary of previous reviews published in 2015-2023

We identified 23 review articles published during 2015-2023 that focused on vegetation, urban form and air quality. To inform our systematic review, we grouped and summarised these 23 review papers into four tables according to the keywords used by the authors (**Supplementary Tables S1.a – S1.d**). The first set of eight reviews (listed in **Table S1.a**) focused on vegetation's capacity to mitigate and remove air pollutants. The main questions discussed were the mitigation capacities of different vegetation strategies, such as green walls, green roofs, and green spaces/parks. Particulate matter (PM) was most discussed, along with O₃, NO₂, and poly-cyclic aromatic hydrocarbons (PAHs). These reviews also compared air cleaning effects according to vegetation species and traits.

The second set of six reviews in **Table S1.b** shows a common interest in the processes and effects of vegetation on air pollution removal through deposition and dispersion. The deposition effect is considered an air pollutant capture mechanism via plant surfaces, while the dispersion effect involves transporting air pollutants through air flows, changing pollutants concentrations at different locations within the urban environment. This group of reviews discusses the deposition and aerodynamic dispersion models of green

infrastructure at different scales, which include various research methods – on-site studies, wind tunnel research, and numerical simulations. Key parameters were discussed among different processes and models.

Table S1.c contains seven reviews concerning the effects of vegetation in urban street canyons or open streets with or without buildings or other structures on both sides. Here, traffic emissions were the main source of air pollutants; trees and hedges planted in these urban spaces could function as porous obstacles. Other green infrastructure, such as green walls and green roofs, were of interest. Considering the combined effects of the built forms, emission patterns, there were noticeable characteristics in the concentration of air pollutants in urban streets (enclosed or open). Vegetation porosity was suggested as a parameter influencing whether barriers or obstacles reduce or increase air pollution concentrations in urban street spaces.

Finally, there are two review papers (**Table S1.d**) summarising the studies that investigated the influence of vegetation not only on air pollution but also on other aspects of urban ecological systems, including runoff pollution removal and interactions with the intensity of urban heat and pollution islands. The reviews highlighted the increasingly multidisciplinary approaches to assessing urban vegetation as an ecosystem service.

In summary, the previous reviews have identified five focal areas of urban air quality research: (1) mitigation processes (deposition, dispersion); (2) models and parameters (numerical simulation, wind tunnel); (3) macro- and micro-structure of vegetation (e.g., leaf traits, porosity); (4) effects of plants in different urban forms and scales (street canyon, open road); and (5) ecological roles of plants in urban environments. However, these reviews also show some limitations. First, only five reviews were conducted following the PRISMA protocols (Diener and Mudu, 2021, Corada et al., 2021, Buccolieri et al., 2022, Chaudhuri and Kumar, 2022, Ernst et al., 2022). Second, the reviews reporting the effects of vegetation did not provide hierarchical correlation or regression accounts of the vegetation's effects on mitigating concentrations of air pollutants. Third, it is difficult to draw clear implications from reviews that mixed theoretical studies with studies of real urban environments. These limitations highlight key research gaps in previous reviews, including the limited focus on real urban environments and the lack of exploration into the hierarchical relationships between vegetation and air pollutants. To address these gaps, we have identified the following questions that necessitate a new systematic review:

1. *What urban air quality indicators and data sources were used in the studies examining the effects of vegetation on urban air quality?*
2. *What metrics or indices were used to quantify the morphological characteristics of urban vegetation in studies on air quality?*
3. *What metrics or indices were used to quantify urban forms of real cities for air quality studies?*
4. *What data sources and methods were used for developing predictive models for assessing urban air quality of real urban environments?*

3. Method and materials

Our systematic review follows the PRISMA framework, which provides a structured and transparent approach to reviewing the literature. This methodology addresses inconsistencies in prior reviews by ensuring that all included studies are systematically identified, screened, and evaluated based on predefined eligibility criteria. By focusing on quantitative studies conducted in real urban environments, we aim to overcome the lack of real-world validation highlighted in previous reviews. Additionally, the use of standardised metrics and comprehensive data extraction protocols ensures consistency across the studies

analysed, facilitating more robust comparisons and actionable insights for urban planning and policy development.

We first explain how the PRISMA guidelines (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) were applied in this systematic review (Section 3.1). We then introduce the core materials that appeared in the articles identified by the search and selection criteria (Section 3.2).

3.1. Method: The PRISMA guidelines applied

The PRISMA guidelines were applied in three stages: (1) identification, (2) two-step selection – initial selection by abstract and title search, followed by selection via full-text search, and (3) grouping the identified articles into two main categories: Correlation/Regression Studies, and Prediction Models.

The following keywords were used in the initial search: ‘air quality’ OR ‘air pollution’ OR ‘air pollutant’ OR ‘air pollutants’, AND ‘vegetation’ OR ‘green infrastructure’ OR ‘plant’, AND ‘urban’ OR ‘outdoor’. Publication dates were limited to 2012-2023. Scopus was selected as the database. The search focused on journal articles, including other relevant literature such as book chapters and conference papers. The initial search returned 3,456 papers.

The first selection was conducted based on the search for abstracts, followed by a more detailed full-text reading and extraction. The criteria for the first selection were as follows:

Inclusion Criteria:

1. Search for papers published only in English.
2. Include research articles published as open access full texts.
3. Include research articles focused on outdoor air quality and vegetation within urban areas.
4. Include research where air quality is the dependent variable and vegetation-related metrics or indices are part of the independent/explanatory variables.
5. Include studies that screen vegetation and air quality as interaction objects. For studies with additional objects, only review sections that address the influence of vegetation on air quality (e.g., interactions between vegetation, air pollution, and urban heat islands).
6. Focus on research examining the morphological characteristics of vegetation.
7. Include studies on air quality in real urban environments based on data-driven quantitative analyses or modeling.

Exclusion Criteria:

1. Exclude studies on indoor air quality and indoor vegetation.
2. Exclude studies on vegetation in large non-urban environments (e.g., ecological forests, peri-urban farmlands).
3. Exclude studies related to plant adaptation or tolerance to air pollution; non-airborne pollutants (e.g., those in the rain); and vegetation effects such as ecological, economic, or social influences (e.g., effects on the thermal environment, economic benefits of greening, resident satisfaction with urban green spaces).
4. Exclude studies on the biological structures of plants (e.g., leaf traits such as wax or chlorophyll).
5. Exclude qualitative studies (e.g., guidelines, policy interpretations), numerical simulations (e.g., wind tunnels, CFD), laboratory based measurements (e.g., leaf deposition measurements), and

studies of vegetation removal capacity based on biochemical processes (e.g., stomatal uptake).

6. Exclude articles primarily focused on population exposure, human health, and diseases.
7. Exclude all 23 previous review papers discussed in Section 2.

Our inclusion and exclusion criteria were intentionally designed to concentrate on specific research topics. To emphasize researches with real urban environments, we excluded numerical simulation studies. Furthermore, we did not categorise individual vegetation types—such as trees, hedges, green roofs, or green walls—because our primary objective was to summarise the vegetation indices employed in these studies. We assume that these indices are generally applicable to various types of vegetation.

Based on the above search and selection criteria, 301 papers were identified in the first round of the abstract search, of which 79 papers were retained after a more detailed extraction from the open access full texts.

Figure 1 shows the PRISMA process flowchart.




Figure 1: Systematic literature review searching and retrieving flowchart (PRISMA)

3.2. *Materials: Air quality, vegetation, urban morphology indicators/indices, and prediction models*

Our initial summarisation of the core materials used in the 79 original research articles suggests five headings: (1) air quality indicators, (2) vegetation-related indices, (3) non-vegetation-related urban morphology indicators, (4) interrelations between vegetation and non-vegetation indices, and (5) predictive urban air quality models, as introduced below.

(1) Air quality indicators

We identify nine groups of air quality indicators used in the reported urban air quality studies: particulate matter (PM), total suspended particulates (TSP), ultrafine particles (UFPs), nitrogen oxides (NO_x), ozone (O₃), carbon oxides (CO_x), sulphur dioxide (SO₂), black carbon (BC), and aerosol optical depth (AOD).

Particulate matter (PM) is a non-gaseous substance in the air composed of chemical compounds and materials. This review focuses on PM₁₀, PM_{2.5}, and PM₁, referring to particles smaller than 10, 2.5, and 1 micrometer in diameter, respectively.

Total suspended particulates (TSP) are airborne particulate matter (PM) with diameters of up to approximately 100 micrometers. These particulates originate from a variety of sources, encompassing both combustion and non-combustion activities. PM₁₀, PM_{2.5}, PM₁ are all components of TSP.

Ultrafine particles (UFPs) are particles characterized by an aerodynamic diameter of 0.1 μm (100 nm) or smaller. Due to their small size, UFPs can efficiently traverse the respiratory tract, reach and breach the alveolar-capillary barrier in the alveoli. Consequently, they can disseminate throughout the body via the circulatory system, posing a potential threat to human health (Kwon et al., 2020).

Nitrogen oxides (NO_x) are gases produced from natural sources, motor vehicles and other fuel-burning processes. They are primarily composed of nitric oxide (NO) and nitrogen dioxide (NO₂).

Ozone (O₃) is present throughout the atmosphere. Stratospheric ozone, found in the upper atmosphere, forms a protective layer that shields life from the sun's harmful ultraviolet rays. However, ozone at ground level can be a harmful air pollutant and is the main ingredient in "smog." Ground-level ozone (O₃) does not originate directly from anthropogenic sources; rather, it is a secondary pollutant formed through a complex series of chemical reactions in the presence of sunlight.

Carbon oxides (CO_x). Carbon monoxide (CO) is a colourless, odourless gas released when something is burned. The primary contributors of CO to the atmosphere are automobiles, trucks, and other vehicles or machinery powered by the combustion of fossil fuels. Carbon dioxide (CO₂), another gas released from burning fossil fuels, is also colourless and non-flammable. While CO₂ is not typically considered an air pollutant, it is a significant heat-trapping (greenhouse) gas.

Sulphur dioxide (SO₂) is a corrosive, acidic gas. Approximately 99% of atmospheric SO₂ comes from anthropogenic sources, primarily the combustion of fossil fuels such as coal, oil, and natural gas. SO₂ is a major air pollutant that is harmful to human lungs and can lead to serious respiratory diseases.

Black carbon (BC) is a component of PM_{2.5}, formed by the incomplete combustion of fossil fuels. BC can cause poor health and premature deaths, and it also warms the atmosphere by effectively absorbing sunlight.

Aerosol optical depth (AOD) is a dimensionless measurement that indicates how much sunlight is blocked due to the presence of fine solid particles or liquid droplets suspended in the air. As a measurement of the attenuation effects caused by atmospheric aerosols, AOD is increasingly used to evaluate the extent of ambient air pollution over large areas (Li et al., 2021a). AOD measurements can be obtained through satellite observations (Gupta et al., 2022), ground-based instruments (NOAA/ESRL, n.d.), or simulation and modelling (Boullisset et al., 2023).

(2) Vegetation-related indices

Various vegetation-related indices have been defined at macro, meso, and micro scales. Based on remote sensing or aerial imaging data, macro indices (e.g., vegetation land use, landscape metrics) address large green spaces or landscape structures in cities. Meso indices (e.g., vegetation structures, green view index) focus on plant communities within city neighbourhoods or districts, where street-view images can be systematically captured and analysed. Micro indices, such as the leaf area index, deal with individual plant

or vegetation characteristics.

(3) Non-vegetation-related urban morphology indices

Non-vegetation-related factors and indices identified in this review include meteorological data (e.g., temperature, wind speed), urban form and structure (e.g., road length), land cover and surface (e.g., water bodies, industrial land cover), and socioeconomic factors (e.g., population density).

(4) Interrelations of air quality indicators, vegetation and non-vegetation indices

We developed a novel data visualization scheme to highlight the interrelations of the air quality indicators, vegetation, and urban morphology indices (see **Figure 4**). The frequency of studies on different air pollutants and their associated variables was also calculated (see **Section 4.4**).

(5) Predictive modelling of urban air quality

Air quality prediction is an emerging and rapidly evolving research area, aiming to improve air quality prediction models. Combining multiple data sources, machine learning and artificial intelligence techniques are increasingly applied to enhance forecasting capabilities involving a large number of auxiliary variables.

In this review, 19 articles on air quality prediction were identified and summarised, including the modelling methods, the types of auxiliary variables used, and feature importance analysis.

4. Results

4.1. Air quality indicators and data sources

Based on the 79 papers reviewed, we identified three main sources of air quality data used in the studies: ground-level observations, satellite measurements, and online open datasets.

4.1.1. Ground-level observations

Ground-level observations are the most direct way to obtain data on air quality and various types of air pollution at the population level. The two most commonly used methods are ground-based station observations and on-site mobile measurements. The former typically involves an air quality monitoring network composed of fixed monitoring stations established or managed by government departments or specialised agencies. Generally, it features high temporal resolution but low spatial resolution, meaning the data often has temporal continuity but is geographically limited by the location and number of stations. In contrast, the latter method is more flexible, allowing researchers to use mobile tools to collect data with greater freedom in choosing sample locations. However, both are costly, requiring significant time, energy, equipment, and financial resources.

Among the 79 papers, 58 utilised ground-level observation data. Of these, 35 relied on ground-based station datasets, 22 employed on-site mobile measurement datasets, and one paper combined both methods. **Table S.2** summarises the sources of ground-level observation data.

4.1.2. Satellite measurements

Satellite measurement datasets are widely used as air quality data sources, employing remote sensing technologies to retrieve information via satellite imagery. This method is popular for obtaining global air quality grids due to its accessibility and availability, overcoming the location limitations of ground-based

measurements and the time- and energy-intensive nature of mobile measurements. Commonly used satellite sensing instruments include Moderate Resolution Imaging Spectroradiometer (MODIS), Ozone Monitoring Instrument (OMI), Thermal Infrared Sensor (TIRS), and Visible Infrared Imaging Radiometer Suite (VIIRS). However, these datasets cannot directly distinguish concentrations of different components; instead, they retrieve target pollutant data based on the satellite's observation of aerosol optical depth (AOD). Additionally, tropospheric NO₂ can be measured through satellite images. Due to the limitations of satellite orbits, the resolution of these datasets is typically constrained. **Table S.3** summarises the satellite measurement datasets and their spatial resolutions.

4.1.3. *Online open datasets*

The retrieval process is essential for satellite measurement data to obtain target air pollution concentrations and achieve higher resolution. This typically involves using other data sources, such as ground-based observations, for calibration. By combining large amounts of data from multiple sources and undergoing rigorous screening, cleansing, computation, and validation processes, this method produces high-resolution predictions for specific areas. As a result, substantial data and computing resources are required. Consequently, some of these retrieved and validated target air pollution datasets are published online. These datasets are generally considered highly accurate and are widely used due to their high resolution and accessibility. **Table S.4** summarises the online public datasets used in the reviewed research articles.

For the air pollution data collected, pre-processing is essential due to issues like missing data and resolution inconsistencies. **Table S.5** summarises the data pre-processing methods used in the literature. Data interpolation methods, such as Inverse Distance Weighted (IDW) and Kriging, are commonly used to predict data distribution over larger areas based on a limited set of known data. Some research also uses linear regression to predict and fill in missing data. Additionally, resampling is often employed to achieve uniform dataset resolution. Ensuring data integrity is crucial for the smooth progression of subsequent studies.

4.2. *Vegetation-related metrics and indices*

Among these 79 papers, vegetation-related metrics and indices can be summarised into three categories: (1) Individual characteristics, (2) Satellite or street view sensing measurements, and (3) Landscape pattern metrics/indices. The vegetation type cluster primarily describes characteristics of macroscopic plant communities, including vertical and horizontal vegetation structures. Individual characteristics focus mainly on mesoscopic plant morphology traits, such as canopy and porosity. With advances in remote sensing technology and increased accessibility to open-source data, satellite and street view images have become convenient tools for quantifying vegetation characteristics in three dimensions. Landscape metrics include indices describing vegetation patch types and arrangements, widely used in landscape research. Additionally, morphological spatial pattern analysis (MSPA) is used to describe geometry and connectivity through geometric concepts, though it is not widely utilised in this research area. **Table 1** presents selected vegetation-related metrics/indices in terms of definition, calculation, and unit of measurement.

Table 1. Summary of Vegetation-related metrics and indices

[1] Individual Characteristics

Term	Definition	Calculator	Unit [Value range]
Count	Vegetation/Trees Count: The total count of trees in certain areas.	$PVA = \frac{Vegetation\ Area}{Ground\ Area}$	Scalar [0, ∞]
LAI	Leaf Area Index: A dimensionless quantity characterizing plant canopies.	$LAI = \frac{Leaf\ Area}{Ground\ Area}$	Scalar [0, 10]
CC/CD	Canopy Cover or Density: a ratio between the area covered by tree crowns and a total area within an area.	$Canopy\ Cover = \frac{Tree\ Crowns\ Covered\ Area}{Ground\ Area}$	Scalar [0, ∞]
DBH	Diameter at Breast Height: The tree diameter measured at 4.5 feet above the ground.	Direct measurement	Meter (m)
PS	Proportion of Species: Proportion of specific species in a green space.	$PS = \frac{The\ number\ of\ species}{The\ total\ number\ of\ vegetation}$	Scalar [0, 1]
SR	Species Richness: The number of species in given samples.	$SR = \frac{The\ total\ number\ of\ species}{Area}$	Scalar [0, ∞]
Porosity	Tree Crowns/Belts Porosity: Ratio of area light penetrating trees in a planar or sectional area (<i>ha</i>).	Digital image processing	Scalar [0, 1]

[2] Satellite sensing and street view scanning measurement

Term	Definition	Calculator	Unit [Value range]
NDVI	Normalized Difference Vegetation Index: An index quantifies the vegetation by measuring the ratio of near-infrared (NIR) and visible red light (Red).	$NDVI = \frac{(NIR - Red)}{(NIR + Red)}$	Scalar [-1, 1]
SAVI	Soil Adjusted Vegetation Index: An index to correct NDVI for the influence of soil brightness in areas where vegetative cover (L [0,1]) is low.	$SAVI = \frac{(NIR - Red)}{(NIR + Red + L)} * (1 + L)$	Scalar [-1, 1]
EVI	Enhanced Vegetation Index: An index to correct NDVI for atmospheric resistance (C), canopy background (L), and values from Blue band in areas with dense vegetation.	$EVI = G * \frac{(NIR - Red)}{(NIR + C1 * Red - C2 * Blue + L)}$	Scalar [-1, 1]
GTCT	Greenness Tasseled Cap Transformation: An index to convert satellite data into three spectral indicators with the Greenness indicator for vegetation growth cycles in	GIS-based digital image processing	% [0, 100]

	particular.		
GVI	Green View Index: An objective measurement of urban green at the street level from a human-eye viewpoint.	$GVI = \frac{GreenPixelCount}{TotalPixelCount}$	% [0, 100]
VGVI	Viewshed Greenness Visibility Index: In a GIS framework, VGVI _j is the index value for the observer cell <i>j</i> ; G _j is the visible green cell, V _i is the visible non-green cell, and d _i is distance decay weight corresponding to visible cell <i>i</i> .	$VGVI_j = \frac{\sum_1^n G_i * d_i}{(\sum_1^n G_i * d_i) + (\sum_1^n V_i * d_i)}$ Where 0 = no green cells are visible, and 1 = all of the visible cells are green	Scalar [0, 1]

[3] Landscape pattern metrics and indices

Term	Definition	Calculator	Unit [Value range]
PLAND	Percentage of Landscape: Percentage of the total area of <i>j</i> _{th} patch of patch type <i>i</i> (<i>a_{ij}</i>) over the total area of the landscape (<i>A</i>).	$PLAND = \frac{\sum_{j=1}^n a_{ij}}{A} * 100$	% [0, 100]
PD	Patch Density: Density of a certain patch in the landscape.	$PD = \frac{NP}{A}$	Patches/km ² [0, ∞]
MPS	Mean Patch Size is the area of all patches of patch type <i>i</i> (<i>a_{ij}</i>) divided by the number of the patch of type <i>i</i> (<i>n_i</i>), divided by 10,000 (to convert to hectare).	$MPS = \frac{\sum_{j=1}^n a_{ij}}{n_i} * \left(\frac{1}{10000}\right)$	Hectares
LSI	Landscape Shape Index: The ratio between the actual landscape edge length (<i>E</i>) and the hypothetical minimum edge length <i>min E</i> .	$LSI = \frac{E}{\min E} \quad LSI = \frac{0.25P_{ij}}{\sqrt{a_{ij}}}$	Scalar [1, ∞]
AI	Aggregation Index: The degree of aggregation or clumping.	$AI = \left[\frac{g_{ii}}{\max(g_{ii})} \right] * 100$	% [0, 100]
ED	Edge Density: An landscape configuration description index which equals all edges in the landscape in relation to the landscape area.	$ED = \frac{E}{A}$	m/ha [0, ∞]
AWMSI	Area-Weighted Mean Shape Index: Averaging the shape index value of all landscape patches, with the perimeter of patch (<i>P_{ij}</i>), the area of patch (<i>a_{ij}</i>), the total area of the landscape (<i>A</i>), weighted by the patch areas.	$AWMSI = \sum_{i=1}^m \sum_{j=1}^n \left[\frac{2 \ln(0.25P_{ij})}{\ln(\sqrt{a_{ij}})} \right] \left(\frac{a_{ij}}{A} \right)$	Scalar [1, ∞]

Figure 2 shows the number of articles that used various types of vegetation-related indices. Apart from the three main categories and MSPA, there is another set of metrics used in the studies, including vegetation structure (VS), and Land Use/Land Cover (LULC). NDVI is the most frequently used index, appearing in 36 papers, followed by various types of vegetation land types used in 27 papers. Landscape pattern indices constitute a broad category that includes several indices. Although different landscape pattern indices are sometimes combined in a single study, the total number of articles using them is relatively low. Overall, the application rate of indices in the macro-horizontal dimension is much higher than the usage rate of multi-dimensional indicators in the vertical dimension. For each of the vegetation-related indices/metrics used in the studies, a list of the literature reviewed is presented in **Table S.6**.




Figure 2: Vegetation-related indices and numbers of associated research articles reviewed

4.3. Summary and classification of non-vegetation related indices

The non-vegetation related indicators or indices used in the literature can be classified into four clusters: (1) meteorological data, (2) urban form and structure, (3) non-vegetation land cover and surface, and (4) economic and social data. **Figure 3** shows the number of articles reviewed in terms of the non-vegetation-related indices used.

The meteorological data cluster contains 16 variables, with *wind speed*, *temperature*, and *humidity* being the top three used. The urban form and structure cluster includes 26 indicators or indices, consisting of the natural topographical characteristics of urban areas, such as *elevation* and *slope*, as well as the structural

characteristics of the urban environment, such as *road length* and *road density*. Among these, *elevation* and *DEM* were the most used variables. The non-vegetation land cover and surface cluster comprises variables that describe different land use types in 1- or 2-dimensional spaces. The economic and social data cluster covers population related, economic, traffic, social, and industrial activities.




Figure 3: Non-vegetation related indices and numbers of associated articles reviewed

4.4. Interrelations of vegetation, air quality indicators, and non-vegetation indices

The air quality indicators and associated variables of interest across the 79 papers were analysed. Among

these variables, only data from more than two articles on the same air pollutant were retained. **Figure 4** shows the interrelations between air quality indicators, vegetation-related indices, and non-vegetation related indices, with the air quality indicators forming the central spine. The mapping shows that PM_{2.5} is the pollutant of highest concern linked to most variables, followed by PM₁₀ and NO₂. Among the vegetation related variables, those from satellite or street view imagery measurements and vegetation land type clusters were linked to more air quality indicators, while the landscape metrics cluster had a lower utilization rate. In contrast, the non-vegetation related variables/indices, as grouped in four sectors, were more evenly connected to air pollutants, with a slightly higher connection in the meteorological data.




Figure 4: Mapping the connections between Vegetation-Related Indices, Air Quality Indicators (Air Pollutants), and Non-Vegetation Related Indices used in the 79 original research articles reviewed.

4.5. Predictive modelling of urban air quality

Recently, identifying and evaluating variables for air quality prediction has become a focus of research. Of the 79 reviewed papers, 19 addressed urban air quality predictions. These studies outline a three-step process: (1) auxiliary variable selection, (2) predictive model development and validation, and (3) feature importance analysis. Step one involves data pre-processing, such as handling missing values and standardising dataset resolutions (see **Table S.5**). Step two, the core stage, encompasses algorithm selection, parameter/hyperparameter tuning, model training/testing, and validation (details in **Table S.7**). While the first two steps are essential, feature importance analysis is not always included. We identified at four types

of prediction models and summarised their feature importance analyses.

4.5.1. Prediction methods and models

Predictive urban air quality models can be categorised into four types: (1) spatial estimation models, (2) traditional land use regression (LUR) models, (3) machine learning (ML) and deep learning (DL) models, and (4) ML/DL LUR hybrid models.

(1) Spatial estimation models

These are divided into two sub-types: interpolation methods and spatial regression models.

- Interpolation methods, such as Inverse Distance Weighting (IDW) and kriging (Ramos et al., 2016), estimate air quality in unmonitored areas using mathematical and geostatistical approaches. They rely on spatial relationships based on distances to known points, which often overlook other influential variables, leading to large margins of error. As a result, these methods are now primarily used for preprocessing, such as filling data gaps.
- Spatial regression models, particularly the Geographically Weighted Regression (GWR) model (Li et al., 2017), account for variable autocorrelation and heterogeneity, making them less dependent on ground-based stations.

(2) Traditional land use regression (LUR) models

LUR models derive air quality (dependent variable) from station monitoring data and extract auxiliary variables (independent variables) from buffer zones around the stations.

- Their accuracy is limited by station location and density, with most located in urban areas. Auxiliary variables extracted from multiple buffer zones can lead to data redundancy, collinearity and other issues, making variable selection critical. Common methods include correlation analysis and the Variance Inflation Factor (VIF).
- Although typically reliant on Multiple Linear Regression or stepwise (forward/backward) methods, LUR models struggle with non-linear relationships and handling large datasets effectively. **Figure 5** illustrates traditional LUR workflows (Han et al., 2022a; Kong and Tian, 2020; Guo et al., 2020; Van Ryswyk et al., 2019; Liu et al., 2019; Masri et al., 2019; Wu et al., 2017; Wu et al., 2015a; Meng et al., 2015; Rao et al., 2014).




Figure 5: Land use regression (LUR) based air quality models using multiple linear regression (left), stepwise (forward/backward) linear regression (right)




Figure 6: ML/DL prediction modelling (left) and ML/DL LUR mixed modelling (right)

(3) Machine/Deep learning prediction models

As machine learning (ML) and deep learning (DL) algorithms advance, ML/DL models are becoming as popular tools for predicting and analysing large, multidimensional datasets. These models excel at identifying non-linear relationships between air quality and auxiliary variables, surpassing traditional

models by incorporating diverse data types (e.g., 3D variable indices) and sources (e.g., satellite AOD datasets) without reliance on ground monitoring stations. In the era of big data, this adaptability is invaluable. However, effective data cleaning and preprocessing are essential to address challenges like missing values and varying dataset resolutions. A key limitation of ML/DL models is their “black boxes” nature, which hinders interpretability. Techniques like feature important analysis are necessary to enhance understanding. **Figure 6 (left)** summarises the ML/DL predictive modelling process (Tella and Balogun, 2021; Shogrkhodaei et al., 2021; Liu et al., 2020; Li et al., 2020b; Zhang and Hu, 2017).

(4) ML/DL LUR hybrid models

To address the limitations of individual methods, a combined approach integrating LUR with ML/DL techniques has been developed. Based on ground monitoring data, LUR extracts buffer data, while ML/DL algorithms process large multidimensional datasets and analyse non-linear relationships. **Figure 6 (right)** presents a flowchart of the mixed ML/DL LUR predictive air quality model (Qi et al., 2022; Babu Saheer et al., 2022; Han et al., 2022a).

4.5.2. Feature variable importance ranking

Across 19 reviewed articles on predictive air quality models, 55 feature variables were utilised. Five studies included feature variable importance analysis during model development. **Table 2** summarises the results, with two studies focused on PM_{2.5} prediction (Shogrkhodaei et al., 2021; Li et al., 2020b) and three on NO₂, O₃, and PM₁₀, respectively (Qi et al., 2022; Han et al., 2022a; Tella and Balogun, 2021). Despite variations in feature selection and ranking, three common variables—humidity, maximum temperature, and elevation—were consistently used in modelling.

Table 2. Feature variable importance ranking in predictive air quality modelling

[1] NO₂ prediction: LUR using street view imagery (SVI) and satellite sensing data (Qi et al., 2022)

Feature variable	Ranking	Feature Variable	Ranking
Built Environment (SVI)	3 [9]	Transport Vehicles (SVI)	1 [9]
People count (SVI)	8 [9]	Vegetation (SVI)	5 [9]
Natural scenery (SVI)	6 [9]	Water (SVI)	7 [9]
Ozone Monitoring Instrument (SVI)	4 [9]	Year	9 [9]
Transport Network (SVI)	2 [9]		

[2] O₃ prediction: Ozone and UHI using a ML modified LUR method (Han et al., 2022a)

Feature variable	Ranking	Feature Variable	Ranking
Average Temperature	5 [20]	Industrial and Mining Land	13 [20]
Average Wind Speed	4 [20]	Major Road	12 [20]
Distance to Cultivated Land	9 [20]	Maximum Temperature	2 [20]
Distance to Major Road	18 [20]	Maximum Wind Speed	3 [20]
Distance to Printing Factory	10 [20]	PM ₁₀	16 [20]
Distance to Ridge Line	11 [20]	PM _{2.5}	19 [20]
Elevation	6 [20]	Point of Interest (POI) - Factory	15 [20]
Green Space	8 [20]	POI - Gas Station	20 [20]
Gross Domestic Product (GDP)	7 [20]	Population	14 [20]
Humidity	1 [20]	Water Body (5000m buffer)	17 [20]

[3] PM10 prediction: KNN, XGBoost, RF, and NB model (Tella and Balogun, 2021)

K-Nearest Neighbor (KNN) model, Selangor State, Malaysia

Feature variable	Ranking	Feature Variable	Ranking
Average Wind Speed	2 [8]	Normalized Difference Vegetation Index (NDVI)	6 [8]
Build-up Index	7 [8]	Road Density	4 [8]
Elevation	1 [8]	Soil Adjusted Vegetation Index (SAVI)	5 [8]
Land Surface Temperature	3 [8]	Slope	8 [8]

XGBoost model, Selangor State, Malaysia

Feature variable	Ranking	Feature Variable	Ranking
Average Wind Speed	2 [8]	Normalized Difference Vegetation Index (NDVI)	6 [8]
Build-up Index	8 [8]	Road Density	4 [8]
Elevation	1 [8]	Soil Adjusted Vegetation Index (SAVI)	7 [8]
Land Surface Temperature	3 [8]	Slope	5 [8]

Random Forest (RF) model, Selangor State, Malaysia

Feature variable	Ranking	Feature Variable	Ranking
Average Wind Speed	2 [8]	Normalized Difference Vegetation Index (NDVI)	7 [8]
Build-up Index	8 [8]	Road Density	5 [8]
Elevation	1 [8]	Soil Adjusted Vegetation Index (SAVI)	6 [8]
Land Surface Temperature	3 [8]	Slope	4 [8]

Naive Bayes (NB) model, Selangor State, Malaysia

Feature variable	Ranking	Feature Variable	Ranking
Average Wind Speed	2 [8]	Normalized Difference Vegetation Index (NDVI)	7 [8]
Build-up Index	6 [8]	Road Density	4 [8]
Elevation	1 [8]	Soil Adjusted Vegetation Index (SAVI)	5 [8]
Land Surface Temperature	3 [8]	Slope	8 [8]

[4] PM2.5 prediction: Seasonal models using three ML algorithms (Shogrkhodaei et al., 2021)

Spring model, Tehran metropolis

Feature variable	Ranking	Feature Variable	Ranking
Average Temperature	5 [10]	Maximum Wind Speed	6 [10]
Distance to Industrial	2 [10]	Normalized Difference Vegetation Index (NDVI)	1 [10]
Humidity	8 [10]	Population Density	4 [10]
Maximum Temperature	10 [10]	Rainfall	7 [10]
Minimum Temperature	9 [10]	Road Density	3 [10]

Summer model, Tehran metropolis

Feature variable	Ranking	Feature Variable	Ranking
Average Temperature	1 [10]	Maximum Wind Speed	10 [10]
Distance to Industrial	7 [10]	Normalized Difference Vegetation Index (NDVI)	6 [10]
Humidity	9 [10]	Population Density	8 [10]
Maximum Temperature	2 [10]	Rainfall	5 [10]
Minimum Temperature	4 [10]	Road Density	3 [10]

Autumn model, Tehran metropolis

Feature variable	Ranking	Feature Variable	Ranking
Average Temperature	6 [10]	Maximum Wind Speed	7 [10]
Distance to Industrial	4 [10]	Normalized Difference Vegetation Index (NDVI)	1 [10]
Humidity	5 [10]	Population Density	3 [10]
Maximum Temperature	9 [10]	Rainfall	8 [10]
Minimum Temperature	10 [10]	Road Density	2 [10]

Winter model, Tehran metropolis

Feature variable	Ranking	Feature Variable	Ranking
Average Temperature	8 [10]	Maximum Wind Speed	7 [10]
Distance to Industrial	1 [10]	Normalized Difference Vegetation Index (NDVI)	2 [10]
Humidity	5 [10]	Population Density	3 [10]
Maximum Temperature	9 [10]	Rainfall	10 [10]
Minimum Temperature	6 [10]	Road Density	4 [10]

[5] PM2.5 prediction: Ensemble-based deep learning over California (Li et al., 2020b)

Feature variable	Ranking	Feature Variable	Ranking
10-meter Northward Wind	20 [20]	Maximum Temperature	4 [20]
CO	1 [20]	Pressure	9 [20]
Daily Mean Downward Shortwave Radiation	15 [20]	Product of Latitude and Longitude	3 [20]
Dry Deposition of Ox	19 [20]	Sea Salt Concentrations in PM2.5	10 [20]
Elevation	18 [20]	Square of Latitude	7 [20]
Humidity	16 [20]	Square of Longitude	11 [20]
Impervious Layer	12 [20]	Temporal Basis Function 1	2 [20]
Latitude	5 [20]	Temporal Basis Function 2	14 [20]
Longitude	6 [20]	Temporal Basis Function 3	17 [20]
MAIAC AOD	8 [20]	Year	13 [20]

5. Discussion

5.1. Transition of the indices' measurements and prediction models

The initial search for this systemic review, conducted in early 2023, targeted studies from the past decade, a period marked by: (1) the proliferation of data sources through advancements in remote sensing and digital imaging, and (2) improved computational performance driven by machine learning and deep learning innovations.

5.1.1. The transition of air quality measurements and vegetation-related indices

Ground-based station monitoring (**Tables S.2–S.4**) remains widely used for direct and accurate air pollutant measurements. Since 2017, satellite sensing and online datasets have gained prominence, offering broader geographic coverage and higher resolution. Despite these advancements, ground-based monitoring remains essential for retrieving target air pollutant concentrations and validating satellite-derived data. Its role has evolved from a primary data source to a verification tool, improving research efficiency by addressing the spatial and cost limitations of ground monitoring networks and expanding air quality datasets.

Vegetation-related indices quantify horizontal and vertical characteristics. Horizontal indices, such as satellite-derived NDVI and land use metrics (e.g., PLAND), are preferred for their simplicity and ease of use, whereas vertical indices like Leaf Area Index (LAI) and porosity require specialised instruments and

manual labour, resulting in smaller datasets. Landscape pattern metrics are less common due to their complexity.

Since 2020, street view images have provided an accessible, low-cost method for capturing vertical vegetation characteristics in urban areas. Advances in machine learning and deep learning have enhanced image analysis, significantly improving the efficiency and applicability of street-view images in urban greening research.

5.1.2. The evolution of air quality prediction models

Air quality prediction models have advanced from relatively simple spatial estimation techniques to land use regression (LUR) models using linear regression, and, after 2020, to machine learning and deep learning models. By 2022, hybrid models combining machine learning, deep learning, and LUR emerged, driven by high-performance computational algorithms capable of processing high-dimensional data and capturing complex non-linear relationships. Earlier methods like inverse distance weighting (IDW) and Kriging are now primarily used as preprocessing tool to fill missing data and harmonise dataset resolutions.

5.1.3. Some observations of the latest research trends (2023-24)

Since concluding our initial search for this systematic review, we have noted ongoing trends reported in recent research (post-2023). Traditional studies continue to focus on the impact of vegetation on air pollutants through localised sampling in specific areas such as schools, elderly care centres, and parks (Pan et al., 2024; Wang et al., 2024b; Wu et al., 2024; Ta and Promchan, 2024). Meanwhile, the growing availability of large datasets, including remote sensing data, has enabled broader-scale analyses of spatiotemporal air quality and vegetation distribution (Liu et al., 2024; Saha et al., 2024; Naboureh et al., 2024; Sheng et al., 2023; Kan et al., 2023; Mansourmoghaddam et al., 2023).

Recent advancements in remote sensing and street-view imaging have enhanced the quantification of urban vegetation. Machine learning (ML) and deep learning (DL) algorithms are increasingly applied to image-based tasks such as vegetation detection, canopy identification, and species classification. High-resolution datasets (e.g., WorldView-2/3 series, GeoEye-1, Planet Labs' SkySat, and Pleiades) enable detailed vegetation mapping (Guo et al., 2023; Sicard et al., 2023). Furthermore, integrating 3D vegetation analysis from street-view imagery marks a shift from traditional 2D studies. AI-driven computer vision now extracts 3D structural attributes, including vertical dimensions, offering deeper insights into urban greenery (Gupta et al., 2024; Xu et al., 2023). ML and DL algorithms are also pivotal in air quality prediction, improving analysis of high-dimensional datasets and assessing interactions among multiple variables (Wang et al., 2024a; Gündoğdu and Elbir, 2024). However, isolating the effects of individual factors remains a challenge, limiting the robustness of these analyses.

5.2. Limitations of this systematic review

We used the PRISMA framework to systematically review literature on the relationships between vegetation morphology, non-vegetation factors, and air quality in real urban environments. The inclusion and exclusion criteria defined the search and review scope, though some limitations remain.

5.2.1. Short of a meta-analysis

In reviewing prior research, we initially considered conducting a meta-analysis to evaluate the impact of vegetation on air quality improvement. However, significant variations in research scales, indices,

methodologies, and metrics rendered a consistent meta-analysis infeasible. For instance, studies using the NDVI index employed different air quality indicators (e.g., AOD in Yang et al., 2022, and PM_{2.5} in Llaguno-Munitxa et al., 2021) or distinct data processing methods (e.g., correlation analysis in Yang et al., 2022, and random forest modelling in Shogrkhodaei et al., 2021). As a result, we adopted a qualitative approach, organising findings into two methodological themes: correlation/regression studies and predictive models. This enabled us to map the range of air quality indicators, vegetation indices, urban form indices, and meteorological data, as well as the predictive models developed over the past decade.

While a meta-analysis in this field is feasible, it was not suitable at the time of this review. Future systematic reviews could enable meta-analyses by focusing on specific research questions addressed by multiple studies, ensuring consistency in metrics such as vegetation indices (e.g., NDVI) or types of green infrastructure (e.g., urban trees or green roofs/walls) within defined climatic regions and/or seasons. Such meta-analyses could adopt methodologies similar to those in evidence-based medicine, which is well developed and widely applied.

5.2.2. Limited green infrastructure types

Urban green infrastructure (GI) encompasses diverse forms beyond traditional plantings, including green walls and roofs, which play key roles in urban ecology. This review focuses on indices for quantifying vegetation characteristics, assuming their broad applicability across vegetation types. For example, NDVI captures horizontal greenery like green roofs, while GVI characterises vertical greenery such as street trees and green walls. Although, different GI types uniquely impact air quality—a topic warranting further review—recent studies (Vashishta et al., 2024; Barriuso and Urbano, 2021) highlight their potential to complement traditional urban vegetation.

5.2.3. Real urban environments and numerical simulations

This review focuses on studies conducted in real-world settings, excluding numerical simulations. Our aim is to summarise effective quantitative methods for complex environmental contexts in urban areas. While CFD-based numerical simulations can isolate and quantify vegetation's effects on air quality by simplifying urban scenarios, they often depend on researcher-assumed parameters, potentially introducing biases. These assumptions may overlook the complexity of real-world processes involving numerous interacting variables. By concentrating on non-simulation research, we better capture insights from analysing large-scale high-resolution field measurements. However, validated simulations can contribute to enhanced ML/DL models where comprehensive and reliable field measurements are limited. In particular, these combined capabilities are essential for evaluating planning and design proposals which do not exist in reality. Comparing and combining empirical field studies with simulation-based findings offers a promising direction for future research.

5.2.4. Selection of searchable databases, languages, and accessibility

We conducted a literature search on Scopus, a widely recognised database, yielding 3,456 initial results. Scopus was chosen over alternatives like Web of Science due to its broader journal coverage across disciplines relevant to vegetation, urban form, and air quality, such as environmental sciences, urban studies, and engineering. Its robust citation metrics also facilitated the identification of influential studies and authors, enhancing the systematic review process. While articles not indexed in Scopus were excluded, its comprehensive coverage and alignment with PRISIMA guidelines supported transparent screening and

selection. Limiting the review to English-language and open-access papers may have further excluded some relevant studies.

5.3. *Future Research priorities*

Air quality significantly impacts human health and the environment, prompting global policy actions. Assessing and predicting air quality remains challenging due to the transient nature of urban air pollution (e.g., emission, dispersion, deposition, resuspension, interaction with urban lights). This review highlights the complexity of quantifying vegetation's effects on air quality in dynamic urban environments, where numerous variables interact. Advances in large-scale urban datasets from satellite and street-view imaging now enable the transition from 2D (planar) to 3D (volumetric) vegetation quantification. Integrating 3D vegetation indices with machine learning and deep learning shows new revenues for predictive air quality modelling. These technologies also enable automated plant species recognition, supporting evidence-based urban greening interventions to improve air quality.

However, a key challenge lies in the “black-box” nature of AI-based predictions, which obscures the interactions among variables affecting urban air quality. Enhancing the explainability of these models is crucial to inform urban planning and design decisions effectively.

5.4. *Significant of this review*

This review summarises advancements in vegetation-related indices, urban form, and their impact on urban air quality over the past decade, emphasizing their relevance to urban and landscape design. It also reviews air quality prediction models and algorithms, highlighting data-driven approaches and future research directions. These insights provide urban planners, designers, and policymakers with a foundation for informed science-based decisions for creating sustainable, air-purifying environments.

6. Conclusion

Informed by previous review articles, we set out a new systematic review to provide an up-to-date summary of 79 studies identified that focus on effects of vegetation and urban morphological characteristics on air quality. We identify four key questions to be addressed in the new systematic review concerning the range of air quality indicators and data sources used in the studies, the indices defined for quantifying vegetation morphology and urban form, and predictive models for assessing air quality of real urban environments.

There are nine groups of air quality indicators used ranging from particulate matter (PM) to aerosol optical depth (AOD) which is also increasingly used to derive air pollutant concentrations of various kinds covering large urban areas. Among them, PM_{2.5} is the AQ indicator attracting the largest number of studies due to its significant impact on public health. In terms of data sources, we observe increasing utilisation of combined ground-based air quality monitoring measurements and satellite measurements to obtain the spatial-temporal resolutions required in the studies

In vegetation indices, two-dimensional indices are more commonly used to quantify vegetation morphological attributes. This is due to the fact that remote sensing technology has enabled quantification of top-view features over large areas, such as NDVI. In contrast, three-dimensional indices that capture vertical vegetation characteristics are less developed due to the technical challenge of semantic segmentation of vegetation in large volumes of urban scenes or street views.

Traditional on-site measurements can only provide small-scale vertical characteristics such as canopy

features. With the advancement of large urban image datasets, including street view scans, advanced machine-learning techniques are being developed and applied in quantifying vertical dimension of urban vegetation in real cities. However, due to limited open-source urban image datasets and reliable image processing techniques, this remains a research topic to be further addressed in future research. More importantly, how vegetation indices defined and measured in both horizontal and vertical dimensions may be combined to form new three-dimensional vegetation indices remain to be further developed.

This systematic review identifies and summarises several air pollution prediction models. It shows that the traditional Land Use Regression (LUR) model is relatively well-established. LUR was frequently used pre-2020, but since then, the advancement of machine learning (ML) and deep learning (DL) algorithms has led to the rapid development of ML/DL models, which exhibited better prediction performance due to the computational power and intelligence unavailable before.

Finally, our attempt at summarising how vegetation and urban form variables may interact with different air pollutants can be explained to some extent by a feature importance analysis. We find that conducting a thorough meta-analysis of the effects of vegetation an urban form on air quality is not without substantial difficulties. There is a need for establishing a comprehensive air pollution research data repository, linking related studies to enable classifying and summarising data by pollutant types, research locations, spatial-temporal scales, and modelling methods. This could lead to identification of the key variables impacting specific air pollutants as the basis for bringing forward evidence-based guidelines applicable to cleaner air landscaping planning and urban design.

In conclusion, this systematic review offers a comprehensive summary of advancements in vegetation-related indices and other influencing factors, such as urban form, over the past decade, focusing on their impact on urban air quality. These elements are deeply intertwined with urban design, making our findings highly relevant for urban planners and designers aiming to create environments that promote nature-based solutions to air purification. Furthermore, we summarise the air quality prediction models and algorithms developed over the past ten years, prompting the emerging research trends of developing large-scale data-driven approaches. Advance in quantifying effects of vegetation on air quality in real urban environments can improve the proficiency of evidence-based planning and design decision-making.

CRedit authorship contribution statement

Conceptualization: MY, CP.

Methodology: MY, CP, MS.

Data collection, curation: MY.

Data analysis, visualization: MY.

Writing – original draft: MY, CP.

Writing – review & editing: CP, MS, MY.

Supervision: CP, MS.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Supplementary Tables

Table S1.a. The previous eight reviews on mitigation and removal capacity of vegetation (Keywords: mitigation, phytoremediation, vegetation traits, air pollutants capture)

Article	Review Questions	Summary of Review Findings
(Hellebaut et al., 2022)	Current understanding and knowledge gaps in air quality and plants traits on green walls, evidence of knowledge application in design practice	Hairiness, roughness and leaf size are the traits that affect particulate matter (PM) capture. Review six green wall designs in Belgium & showing knowledge of plants traits and air pollutants removal in green wall design remained partial.
(Han et al., 2022b)	Phytoremediation of indoor air pollutants; removal efficiency of plants on different air pollutants under different environmental settings	Absorption and purification of different pollutants (formaldehyde, aromatic compounds and inorganic pollutants) are affected by different trait variables (leaf characteristics, planting patterns, species). The effectiveness of pollutant removal differs under different environmental settings (lab-scale studies, real-world site-specific indoor/outdoor conditions).
(Wróblewska and Jeong, 2021)	Effectiveness of plants on removing particulate matter (PM)	1) Deposition on leaf surfaces; 2) Factors affecting PM capture efficiency: leaf area index, morphological characteristics of leaf surfaces, environment (e.g., wind), and 3D geometry of city design; 3) different types of green infrastructure - green roofs, living walls, water reservoirs, urban farming; 4) PM removal capacity could be improved by species selection and increasing biodiversity.
(Diener and Mudu, 2021)	Effects and mechanisms of green spaces on reducing PM exposure to protect public health	1) Three mechanisms to reduce PM exposure: deposition, dispersion and modification; 2) Public health interventions to reduce PM exposure should consider sensitivity of green spaces in mitigating PM exposure: location of green spaces at a regional scale, porosity of green spaces at a local scale.
(Corada et al., 2021)	Effective leaf traits for enhancing PM capture	1) Coniferous needle leaves, 2) Small, rough and textured broad-leaves, 3) Extended oval shapes, 4) Waxy coatings and high-density trichomes; 5) Ancillary factors and the context of plantings should also be considered to improve PM removal, e.g., plant species, wind conditions, and locations.
(Sicard et al.,	Quantification of O ₃	For O ₃ removal, urban trees are more efficient and

2018)	removal capacity of trees, shrubs, and green roofs	cost effective than green roofs; broad-leaf tree species perform better than conifers, while evergreen are better than deciduous broadleaf.
(Gourdji, 2018)	Effects of green roofs on mitigating air pollution; plant species for PM, O ₃ , NO ₂ reduction; effects of green roofs on air quality in the Montreal region, Canada	Air pollutants removal processes: 1) Deposition of PM, 2) Deposition of O ₃ on plant or soil surfaces via stomatal conductance and non-stomatal uptake, 3) NO ₂ removed by stomatal absorption; 4) Small Zone 5 hardiness tolerant plants on intensive green roofs was recommended for Montreal.
(Huang et al., 2018)	PAHs removal; PAHs accumulating capability of pine needles, Holm oak leaves, and moss	1) PAHs (Polycyclic aromatic hydrocarbons) uptake via absorption and adsorption; 2) Moss perform better in PAHs capture than oak leaves and pine needles; 3) Environment factors (temperature, seasonality, photolysis) could affect the transfer process of PHAs from atmosphere to vegetation.

Table S1.b. The previous reviews of deposition and dispersion effects and processes of vegetation (Keywords: deposition, dispersion, aerodynamic, CFD, green walls, green infrastructure)

Article	Review Questions	Summary of Review Findings
(Li et al., 2022)	Removal of NO ₂ by dry deposition of plants	Plant structure, chemical composition of leaves, nitrogen content of leaves, meteorological conditions, and other related factors affect the deposition mechanism and the efficiency of NO ₂ removal.
(Ysebaert et al., 2021)	Effectiveness of green walls in removing PM	Species, pollution level and residence time affecting PM deposition on green walls. More field, wind tunnel and model validation studies are needed to eliminate discrepancies about the key parameters affecting PM capture by green walls.
(Badach et al., 2020)	Effects of urban greenery on mitigating air pollution in Polish cities	Urban greenery can have combined deposition and aerodynamic effects on air quality. Critical evaluation of local urban planning practice in Gdańsk, Warsaw, and Poznań found limited applicability of the known effects due to lack of accurate models and tools.
(Tiwari et al., 2019)	Green infrastructure (GI) impact on air pollution and health risk assessment	Ten studies that have quantified the linkage between GI, air pollution reduction and health benefits were identified and summarized. Simplified deposition schemes may lead to uncertainties in removal estimation. Future dispersion models need to account for wind speed based GI porosity as well as GI at different spatial scales (microscale and macroscale).

(Buccolieri et al., 2019)	The effects of urban trees on air quality and thermal conditions learned from Computational Fluid Dynamics (CFD) studies	Parameterizations of urban vegetation (trees) are appropriate to account for aerodynamic and deposition effects; resuspension and thermal effects of different types of trees need more works in CFD; “the right tree in the right street” is a better approach.
(Janhäll, 2015)	Vegetation as ecosystem services for air quality improvements – effects of vegetation choice on air pollution from different sources and particle sizes	Urban vegetation effects on air quality summarised as 1) deposition process (particle properties and vegetation properties) at different scales (parks, regional); 2) dispersion effect of vegetation barriers. The studies reviewed include on-site measurements, wind tunnel studies and CFD modelling.

Table S1.c. Previous reviews of the influence of vegetation in urban streets/roads (Keywords: barriers, obstacles, street canyons, open roads, street greening)

Article	Review Questions	Summary of Review Findings
(Buccolieri et al., 2022)	Influence of obstacles (porous & non-porous) on urban canyon ventilation including air pollutant dispersion	The isothermal flow dispersion effects of porous (trees, hedgerows) and non-porous obstacles (parked cars, low boundary walls or baffles, noise/roadside barriers, wind catchers, solar chimneys), and the efficacy, costs, as well as pros and cons.
(Chaudhuri and Kumar, 2022)	Strategic urban greening for long-term air pollution prevention and control measures	PRISMA-based review of global literature (post-2005) and a meta-analysis to be considered by air quality regulatory authorities with particular references to Indian cities to enhance tree species selection, removal strategies in street canyon and open road environment.
(Tomson et al., 2021)	Optimal form and arrangement of Green Infrastructure (GI) for air quality in street canyons	Deposition and dispersion are the main impact pathways for vegetation on air pollution. The effectiveness of different GI forms (trees, hedges, green roofs, green walls and green screens) in the street canyon environment and the methods for assessing effectiveness.
(Barwise and Kumar, 2020)	Vegetation barriers in open-road environment, optimal configuration as barriers between traffic emissions and adjacent spaces	Effective barriers design principles in different spatial scales (city scale, local scale) and plant selection recommendations (e.g., ecophysiological and morphological characteristics, species emissions) for open-road environment and street canyons in the UK.
(Mori et al., 2018)	The effect of air pollutants on human health, and the vegetation characteristics help to optimise air	Species selection and planting schemes (density of vegetation, disposition of plants, global dimensions of GI) should be considered according to different planting site characteristics (open areas vs. street

	pollutants interception	canyons).
(Abhijith et al., 2017)	Aerodynamic effects and reduction potentials of vegetation in street canyon and open road; vegetation types and characteristics help air pollution reduction	1) In street canyons, hedges improve air quality while trees led to deterioration; 2) In open road, low porosity and tall vegetation helps to downwind pollutant reductions; 3) Green walls and roofs on building envelopes can be effective ways to improve air quality.
(Gallagher et al., 2015)	Passive methods for improving air quality and reducing personal exposure – porous and solid barriers	The strengths and limitations and modelling approaches of porous barriers (trees and vegetation) and solid barriers (noise barriers, low boundary walls, parked cars).

Table S1.d. The previous reviews of plants as urban ecosystem services where air quality is a part (Keywords: urban ecosystems, nature-based, pollutants removal, urban greening, microclimate)

Article	Review Questions	Summary of Review Findings
(Biswal et al., 2022)	Nature-based systems for reducing pollutants in storm water, rainwater and urban air	1) Physico-chemical removal through filtration, adsorption, precipitation, and complexation; 2) Biological removal via air phytoremediation plants; 3) Roadside removal via vegetation characteristics of height, thickness, coverage, porosity.
(Ernst et al., 2022)	The relationships between urban greening, canopy layer urban heat island (UHI) and urban pollution island (UPI), air quality, and urban microclimate	1) The links between microclimate and air quality studies were weak; 2) tools for assessing greening's impacts on both microclimate and air quality with good accuracy at the city scale were not well developed; 3) interactions between plant functioning, microclimate and atmospheric composition may hold the key to modelling the links between urban greening, UHI, UPI.

Table S.2. Summary of air quality indicators/indices and measurements from ground stations

Article	Source/Agency	Air Pollutants	No. of Stations
(O'Regan et al., 2022)	Purple Air Network for Cork City, Ireland	PM _{2.5} , PM ₁₀ , PM ₁	12
(Tella and Balogun, 2021, Halim et al., 2020, Shahrin et al., 2019)	the Malaysian Department of Environment (DOE)	PM ₁₀ , CO, O ₃ , NO, NO ₂ , NO _x , SO ₂	8
(Wang et al., 2022b, Zhao et al., 2022, Li et al., 2021b, Wang et al., 2021, Han et al., 2020, Luan et al., 2020, Liu et	China Environmental Monitoring Station (http://www.cnemc.cn/)	PM _{2.5} , PM ₁₀ , SO ₂ , NO ₂ , O ₃ , CO, AQI,	1589

al., 2019, Tian et al., 2019, Wang et al., 2018b, Zhang and Hu, 2017, Chen et al., 2016, Wu et al., 2015b, Wu et al., 2015a)			
(Khan et al., 2022)	Environmental Protection Apartment, Punjab, Lahore	NO _x , CO, SO ₂ , PM ₁₀	20
(Zeng et al., 2022)	Shenzhen Municipal Ecological Environment Bureau, China (http://meeb.sz.gov.cn/)	PM _{2.5}	74
(Babu Saheer et al., 2022)	UK Air, Department for Environment Food & Rural Affairs (uk-air.defra.gov.uk/)	PM _{2.5} , PM ₁₀ , NO ₂	1500+
(Han et al., 2022a)	Xi'an Air Quality Monitoring Stations, China	O ₃	139
(Shogrkhodaei et al., 2021)	Tehran Air pollution control stations, Iran	PM _{2.5}	23
(Li et al., 2020c)	Shenyang Environment Monitoring Center, China	PM _{2.5} , PM ₁₀ , NO ₂ , SO ₂	11
(Tian et al., 2020)	Georgia Department of Natural Resources, Environmental Protection Division (EPD)	PM _{2.5}	9
(Li et al., 2020a)	Weifang PM _{2.5} Monitoring Stations, China	PM _{2.5}	38
(Kong and Tian, 2020)	Beijing Municipal Environmental Monitoring Center, China (BMEMC 2018)	PM _{2.5}	35
(Guo et al., 2020)	Xi'an Air Quality Daily Reporting System, China (http://www.xianemc.gov.cn/)	PM _{2.5}	13
(Guo et al., 2019)	Tianjin air pollution monitoring stations, China	NO ₂	23
(Li et al., 2017)	US Environmental Protection Agency (EPA)'s Air Quality System (AQS) (www.epa.gov/aqs)	PM _{2.5}	55
(Wu et al., 2017)	Taipei metropolis air pollutant monitoring database, Taiwan Air Quality Monitoring Network (airtw.moenv.gov.tw/eng/)	PM _{2.5}	17
(Ramos et al., 2016)	National Air Pollution Surveillance (NAPS) network of	PM _{2.5}	10

	Environment Canada		
(Meng et al., 2015)	Shanghai Environmental Monitoring Centre (SEMC), China	NO ₂	38

Table S.3. Summary of satellite sensing air quality datasets

Article	Dataset	Resolution
(Rahman and Haque, 2022)	Landsat data to retrieve AOD (earthexplorer.usgs.gov)	30m
(Islam et al., 2012)	Sentinel-5P Level-3 NO ₂ Daily Product (V1)	0.01 arc-degree
(Qi et al., 2022)	OMI/Aura NO ₂ Tropospheric, Stratospheric & Total Columns MINDS Daily L3 Global Gridded (DOI: 10.5067/MEASURES/MINDS/DATA304)	0.25° × 0.25°
(Sun et al., 2022, Xie and Sun, 2021, Li et al., 2020b)	MODIS MAIAC remote sensing AOD data (MCD19A2) (lpdaac.usgs.gov/products/mcd19a2v006)	1km
(Zhang and Hu, 2017)	MODIS Collection 6 Level 2 aerosol products (ladsweb.modaps.eosdis.nasa.gov/archive/allData/6)	3km
(Li et al., 2017)	MODIS AOD Level 2 product (Collection 5.1)	10km
(Li and Myint, 2021)	Landsat 5 satellite images to retrieve AOD	60m
(Syafei et al., 2019)	GOME-2 MetOP-A satellite datasets for NO ₂	80*40km ² or 80*10km ²
(He et al., 2019)	Aqua and Terra MODIS Collection 6 Level 2 aerosol products	3km
(Wang ChengHao et al., 2017)	MODIS Terra Atmosphere Aerosol Level 2 Product	3km
(Ye et al., 2016)	HJ-1B satellite images for AOD retrieving	30m

Table S.4. Summary of publicly available online datasets

Article	Dataset	Resolution
(Hassan et al., 2022)	Socioeconomic Data and Applications Center (SEDAC) and Sentinel-5p data of the European Space Agency for the last 18 years (2002–2020)	1km
(Lin and Jiang, 2022)	Ground-level air pollutants for China (ChinaHighAirPollutants, CHAP) PM _{2.5}	1km
(Wei et al., 2021)	Socioeconomic Data and Applications Center (sedac.ciesin.columbia.edu/search/data?contains=PM2.5)	1km
(Li et al., 2021b)	Gridded global surface PM _{2.5} concentration dataset	0.01°
(van Oorschot et al., 2021)	Annual mean PM ₁₀ concentrations Map, Hague (www.atlasleefomgeving.nl/kaarten)	25m

(Wang et al., 2019)	Open-Data Inventory for Anthropogenic Carbon dioxide (ODIAC) fossil fuel emission dataset from the Center for Global Environmental Research (db.cger.nies.go.jp/dataset/ODIAC/), National Institute for Environment Studies	1km
(Wang et al., 2020a, Lu et al., 2019)	The global annual average surface PM _{2.5} concentrations grids provided by Atmospheric Composition Analysis Group (ACAG) at Dalhousie University	0.01°
(Bechle et al., 2017)	Publicly available global estimates of gridded annual surface NO ₂ concentrations.	0.1° × 0.1°

Table S.5. Summary of air quality data pre-processing methods

Article	Method	Description
(Hassan et al., 2022, Sun et al., 2022, Halim et al., 2020, Arista et al., 2020, Li et al., 2020a, Shahrin et al., 2019, Cui et al., 2019, Chen et al., 2016, Ramos et al., 2016)	Inverse Distance Weighted (IDW)	IDW is one of the commonly used methods of spatial interpolation in air pollution prediction areas. It makes predictions on the concentration of unknown points based on a function of inverse distance from a known point, assuming that the closer they are to the known point, the greater the influence.
(Sun et al., 2022, Llaguno-Munitxa et al., 2021, Shogrkhodaei et al., 2021, Xie and Sun, 2021)	Kriging Interpolation	A set of geostatistical interpolation techniques wherein the value at an unobserved location is estimated through a linear combination of values from neighbouring locations. The weights assigned to these values are determined by a semivariogram which considers the spatial correlation. It has an effective performance in the data points having spatial autocorrelation. Ordinary Kriging and Universal kriging are widely used in air pollution prediction areas.
(Li et al., 2020b, Zhang and Hu, 2017)	Linear Regression	A commonly employed technique for replacing missing values in a dataset. Typically, it involves predicting the missing data in the target dataset by establishing a linear relationship with a reference dataset.
(Qi et al., 2022, Yang et al., 2022, Hassan et al., 2022, Wei et al., 2021, Xie and Sun, 2021, Wang et al., 2019, Lu et al., 2019, Wang et al., 2018b, Wang ChengHao	Resampling	Common data processing methods for uniform resolution of a different data set. It changes the dataset's spatial resolution by aggregating or interpolating values. Common types include Nearest Neighbor and Bilinear Interpolation.

et al., 2017, Ye et al.,
2016)

Table S.6. Summary of vegetation-related indices papers

Vegetation-related indices	Articles
Vegetation Structure	(Niu et al., 2022, Jiang et al., 2021, Qiu et al., 2019, Qiu et al., 2018, Chen et al., 2015)
Land Use/Land Cover (LULC)	(Sun et al., 2022, Zeng et al., 2022, Han et al., 2022a, Llaguno-Munitxa et al., 2021, Li and Myint, 2021, Xie and Sun, 2021, Liu et al., 2020, Li et al., 2020b, Halim et al., 2020, Li et al., 2020c, Li et al., 2020a, Tian et al., 2020, Luan et al., 2020, Kong and Tian, 2020, Guo et al., 2020, Qiu et al., 2019, Van Ryswyk et al., 2019, Shahrin et al., 2019, Liu et al., 2019, Guo et al., 2019, Fan et al., 2019, Chen et al., 2016, Ye et al., 2016, Wu et al., 2015b, Wu et al., 2015a, Meng et al., 2015, Rao et al., 2014)
Count	(Babu Saheer et al., 2022, Llaguno-Munitxa et al., 2021, Yli-Pelkonen et al., 2017)
Leaf Area Index (LAI)	(Niu et al., 2022, van Oorschot et al., 2021, Wang et al., 2020b)
Canopy Cover/Density (CC/CD)	(Niu et al., 2022, Jiang et al., 2021, Wang et al., 2020b, Qiu et al., 2018, Chen et al., 2015, Islam et al., 2012)
Diameter at Breast Height (DBH)	(Niu et al., 2022, Yli-Pelkonen et al., 2017)
Vegetation Height (VH)	(Jiang et al., 2021, Wang et al., 2020b, Hart et al., 2020)
Species Related (Proportion of Species and Species Richness)	(Niu et al., 2022, Wang et al., 2020b, Grzędzicka, 2019, Desyana et al., 2017, Yli-Pelkonen et al., 2017)
Porosity	(Grzędzicka, 2019, Chen et al., 2015, Islam et al., 2012)
Percentage of Vegetation Area / Vegetation Coverage (PAV/VC)	(Grzędzicka, 2019, Masri et al., 2019, Lu et al., 2019, Syafei et al., 2019, Bechle et al., 2017)
NDVI (Normalized Difference Vegetation Index)	(Sun et al., 2022, Zeng et al., 2022, O'Regan et al., 2022, Yang et al., 2022, Hassan et al., 2022, Lin and Jiang, 2022, Zhao et al., 2022, Deb et al., 2022, Llaguno-Munitxa et al., 2021, Li and Myint, 2021, Tella and Balogun, 2021, Li et al., 2021b, Wang et al., 2021, Wei et al., 2021, Shogrkhodaei et al., 2021, Li et al., 2020b, Kong and Tian, 2020, Hart et al., 2020, Van Ryswyk et al., 2019, Han et al., 2020, Wang et al., 2020a, Arista et al., 2020, Wang et al., 2019, Masri et al., 2019, Cui et al., 2019, Tian et al., 2019, He et al., 2019, Wang et al., 2018a, Wang et al., 2018b, Wang ChengHao et al., 2017, Zhang and Hu, 2017, Li et al., 2017, Wu et al., 2017, Farrell et al., 2015, Wu et al., 2015a, Dadvand et al., 2015)
SAVI (Soil Adjusted Vegetation Index)	(Tella and Balogun, 2021)
EVI (Enhanced Vegetation Index)	(Islam et al., 2022)
GTCT (Greenness Tasseled Cap Transformation)	(Ramos et al., 2016)

GVI/GVA (Green View Index / Green View Amount)	(Zeng et al., 2022, O'Regan et al., 2022, Wang et al., 2022b, Qi et al., 2022, Liu et al., 2020)
VGVI ()	(Labib et al., 2021)
Percentage of Landscape Types (PLAND)	(Wang et al., 2022a, Fan et al., 2019, Ye et al., 2016, Wu et al., 2015b)
Patch Density (PD)	(Wang et al., 2022a, Tian and Yao, 2022, Li et al., 2021b, Tian et al., 2020, Fan et al., 2019, Ye et al., 2016, Wu et al., 2015b)
Landscape Shape Index (LSI)	(Wang et al., 2022a, Tian and Yao, 2022, Li et al., 2021b, Tian et al., 2020, Ye et al., 2016)
Class Area (CA)	(Tian and Yao, 2022, Tian et al., 2020)
Largest Patch Index (LPI)	(Tian and Yao, 2022, Li et al., 2021b, Tian et al., 2020, Ye et al., 2016)
Aggregation Index (AI)	(Tian and Yao, 2022, Li et al., 2021b, Tian et al., 2020, Fan et al., 2019)
Area-Weighted Mean Shape Index (AWMSI)	(Li et al., 2021b)
Patch Cohesion Index (COHESION)	(Li et al., 2021b)
Edge Density (ED)	(Tian et al., 2020, Ye et al., 2016, Wu et al., 2015b)
Contagion (CONTAG)	(Ye et al., 2016, Wu et al., 2015b)
Shannon's Evenness Index (SHEI)	(Wu et al., 2015b)
Mean Patch Size (MPS)	(Fan et al., 2019)
MSPA	(Li et al., 2021b)

Table S.7. Types of air quality prediction models, algorithms, output resolutions, and validation

Model	Article	Algorithm	Resolution	Model Validation and Prediction Accuracy
Spatial Estimation Models	(Liu et al., 2019)	OK; IDW	500m	LOOCV
	(Li et al., 2017)	GWR	10km	10-fold CV; RMSE/MAE/RRMSE/RMAE-
	(Ramos et al., 2016)	KED; IDW; KED-IDW	100m	LOOCV
	(Meng et al., 2015)	IDW; OK	1km	LOOCV; R ² /RMSE
Traditional Land Use Regression (LUR) Models	(Han et al., 2022a)	/	500m grid	Splitting 80% training and 20% test; R ² /MSE/RMSE/MAE
	(Kong and Tian, 2020)	SMR	10km grid	LOOCV
	(Guo et al., 2020)	SMR	100m grid	CV; R ² /RMSE/MPE
	(Van Ryswyk et al., 2019)	BFSR	10m grid	LOOCV
	(Liu et al., 2019)	/	500m grid	LOOCV
	(Masri et al., 2019)	BFSR	1km grid	LOOCV
	(Wu et al., 2017)	SLR	250m grid	10-fold CV; External Verification (out-of-sample observations from 2013)
(Wu et al., 2015a)	SLR	30m grid	LOOCV; RMSE/NMSE	

	(Meng et al., 2015)	SFR	1km grid	LOOCV; R ² /RMSE
	(Rao et al., 2014)	/	200m	CV
Machine/ Deep Learning Prediction Models	(Tella and Balogun, 2021)	XGBoost; RF; KNN; NB	60m grid	Confusion matrix; Statistical Measures; ROC-AUC
	(Shogrkhodaei et al., 2021)	RF; AdaBoost; SGD	30m grid	Splitting 70% training and 30% test; RMSE/MAE; ROC-AUC
	(Liu et al., 2020)	RF; SVM; MLR	15 m distance along the route	10-fold CV (Splitting 70% training and 30% test); R ² /RMSE/MAE/IA
	(Li et al., 2020b)	Full Residual Deep Network	1km grid	63.3% samples for training and validation (80% training/20% validation), 36.7% for independent test; 4 monitoring sites for independent tests; R ² /RMSE
	(Zhang and Hu, 2017)	LEM	3km grid	10-fold CV; R ² /MPE/RMSE
ML/DL LUR Models	(Qi et al., 2022)	RF	100m grid (within 500m of each monitor)	10-fold CV (splitting training and testing set randomly, temporally, or spatially); R ² /MAE/RMSE
	(Babu Saheer et al., 2022)	LR; SVR; LSTM	Within 1km of monitor	MAE/MSE/RMSE/R ² /MAPE
	(Han et al., 2022a)	RF, MLR	500m grid	Splitting 80% training and 20% test; R ² /MSE/RMSE/MAE

OK: Ordinary Kriging, **IDW:** Inverse Distance Weighted, **KED:** Kriging with external drift, **KED-IDW:** A Hybridization of KED and IDW, **SMR:** Stepwise Multiple Regression, **BFSR:** Backwards and Forwards Stepwise Regression, **SLR:** Stepwise Linear Regression, **SFR:** Supervised Forward Regression, **LR:** Linear Regression, **MLR:** Multiple Linear Regression, **LEM:** Linear Mixed-effects Model, **XGBoost:** eXtreme Gradient Boosting algorithms, **RF:** Random Forest, **KNN:** K-Nearest Neighbour, **NB:** Naive Bayes, **SGD:** The Stochastic Gradient Descent algorithm, **SVR:** Support Vector Regression, **SVM:** Support Vector Machine, **LSTM:** Long Short Term Memory, **CV:** Cross-Validation, **LOOCV:** Leave-one-out cross-validation.

Statistical Measures includes Recall (REC), Precision (PREC), Specificity, Kappa Index (KI), F-measure, Accuracy, Fitting Index (IA). ROC-AUC: The area under the ROC curve, Error Metrics includes mean average error (MAE), mean squared error (MSE), root mean squared error (RMSE), mean absolute percentage error (MAPE), R²score, mean percentage error (MPE), relative root mean squared error (RRMSE), relative mean absolute error (RMAE), normalized mean squared error (NMSE).

References

ABHIJITH, K., KUMAR, P., GALLAGHER, J., MCNABOLA, A., BALDAUF, R., PILLA, F., BRODERICK, B., DI SABATINO, S. & PULVIRENTI, B. 2017. Air pollution abatement performances of green infrastructure in open road and built-up street canyon environments—A review. *Atmospheric environment*, 162, 71-86. DOI: 10.1016/j.atmosenv.2017.05.014

- ACT, C. A. 1970. Clean Air Act. *United States*. [Google Scholar].
- AMANN, M., BERTOK, I., COFALA, J., GYARFAS, F., HEYES, C., KLIMONT, Z., SCHÖPP, W. & WINIWARTER, W. 2005. Baseline scenarios for the clean air for Europe (CAFE) programme. *Final report*, 79.
- APTE, K. & SALVI, S. 2016. Household air pollution and its effects on health. *F1000Research*, 5. DOI: 10.12688/f1000research.7552.1
- ARISTA, F., SARASWATI, R. & WIBOWO, A. Spatial modelling of particular matter 10 distribution in Bandung City. *IOP Conference Series: Earth and Environmental Science*, 2020. IOP Publishing, 012047.
- AZAROV, V., SERGINA, N. & KONDRATENKO, T. Problems of protection of urban ambient air pollution from industrial dust emissions. *MATEC Web of Conferences*, 2017. EDP Sciences, 07017.
- BABU SAHEER, L., BHASY, A., MAKTABDAR, M. & ZARRIN, J. 2022. Data-driven framework for understanding and predicting air quality in urban areas. *Frontiers in big Data*, 5, 822573. DOI: 10.3389/fdata.2022.822573
- BADACH, J., DYMNICKA, M. & BARANOWSKI, A. 2020. Urban vegetation in air quality management: A review and policy framework. *Sustainability*, 12, 1258. DOI: 10.3390/su12031258
- BAI, R., XUE, Y., JIANG, X., JIN, C. & SUN, Y. 2022. Retrieval of high-resolution aerosol optical depth for urban air pollution monitoring. *Atmosphere*, 13, 756. DOI: 10.3390/atmos13050756
- BARRIUSO, F. & URBANO, B. 2021. Green roofs and walls design intended to mitigate climate change in urban areas across all continents. *Sustainability*, 13, 2245. DOI: 10.3390/su13042245
- BARWISE, Y. & KUMAR, P. 2020. Designing vegetation barriers for urban air pollution abatement: A practical review for appropriate plant species selection. *Npj Climate and Atmospheric Science*, 3, 12. DOI: 10.1038/s41612-020-0115-3
- BECHLE, M. J., MILLET, D. B. & MARSHALL, J. D. 2017. Does urban form affect urban NO₂? Satellite-based evidence for more than 1200 cities. *Environmental Science & Technology*, 51, 12707-12716. DOI: 10.1021/acs.est.7b01194
- BISWAL, B. K., BOLAN, N., ZHU, Y.-G. & BALASUBRAMANIAN, R. 2022. Nature-based Systems (NbS) for mitigation of stormwater and air pollution in urban areas: A review. *Resources, Conservation and Recycling*, 186, 106578. DOI: 10.1016/j.resconrec.2022.106578
- BOULISSET, V., ATTIÉ, J.-L., TOURNIER, R., CEAMANOS, X., ANDREY, J., PEQUIGNOT, E., LAURET, N. & GASTELLU-ETCHEGORRY, J.-P. 2023. Aerosol Optical Depth Measurements from a Simulated Low-Cost Multi-Wavelength Ground-Based Camera: A Clear Case over a Peri-Urban Area. *Remote Sensing*, 16, 140. DOI: 10.3390/rs16010140
- BUCCOLIERI, R., CARLO, O. S., RIVAS, E., SANTIAGO, J. L., SALIZZONI, P. & SIDDIQUI, M. S. 2022. Obstacles influence on existing urban canyon ventilation and air pollutant concentration: A

- review of potential measures. *Building and Environment*, 214, 108905. DOI: 10.1016/j.buildenv.2022.108905
- BUCCOLIERI, R., SANTIAGO, J.-L., RIVAS, E. & SÁANCHEZ, B. 2019. Reprint of: Review on urban tree modelling in CFD simulations: Aerodynamic, deposition and thermal effects. *Urban Forestry & Urban Greening*, 37, 56-64. DOI: 10.1016/j.ufug.2018.07.004
- BURHAN, E. & MUKMININ, U. 2020. A systematic review of respiratory infection due to air pollution during natural disasters. *Medical Journal of Indonesia*, 29, 11-8. DOI: 10.13181/mji.oa.204390
- CAPANNELLI, G., GOLLO, E., MUNARI, S. & RATTO, G. 1977. Nitrogen oxides: analysis of urban pollution in the city of Genoa. *Atmospheric Environment*, 11, 719-727. DOI: 10.1016/0004-6981(77)90180-9
- CHAUDHURI, S. & KUMAR, A. 2022. Urban greenery for air pollution control: a meta-analysis of current practice, progress, and challenges. *Environmental Monitoring and Assessment*, 194, 235. DOI: 10.1007/s10661-022-09808-w
- CHEN, J., ZHU, L., FAN, P., TIAN, L. & LAFORTEZZA, R. 2016. Do green spaces affect the spatiotemporal changes of PM 2.5 in Nanjing? *Ecological processes*, 5, 1-13. DOI: 10.1186/s13717-016-0052-6
- CHEN, X., PEI, T., ZHOU, Z., TENG, M., HE, L., LUO, M. & LIU, X. 2015. Efficiency differences of roadside greenbelts with three configurations in removing coarse particles (PM10): A street scale investigation in Wuhan, China. *Urban Forestry & Urban Greening*, 14, 354-360. DOI: 10.1016/j.ufug.2015.02.013
- CORADA, K., WOODWARD, H., ALARAJ, H., COLLINS, C. M. & DE NAZELLE, A. 2021. A systematic review of the leaf traits considered to contribute to removal of airborne particulate matter pollution in urban areas. *Environmental Pollution*, 269, 116104. DOI: 10.1016/j.envpol.2020.116104
- CUI, Y., JIANG, L., ZHANG, W., BAO, H., GENG, B., HE, Q., ZHANG, L. & STREETS, D. G. 2019. Evaluation of China's environmental pressures based on satellite NO2 observation and the extended STIRPAT model. *International Journal of Environmental Research and Public Health*, 16, 1487. DOI: 10.3390/ijerph16091487
- DADVAND, P., RIVAS, I., BASAGAÑA, X., ALVAREZ-PEDREROL, M., SU, J., PASCUAL, M. D. C., AMATO, F., JERRET, M., QUEROL, X. & SUNYER, J. 2015. The association between greenness and traffic-related air pollution at schools. *Science of the Total Environment*, 523, 59-63. DOI: 10.1016/j.scitotenv.2015.03.103
- DEB, S., DWIVEDI, A. K., CHAUDHARI, S. & RAJAN, K. S. Spatial Factor Analysis of Mobile IoT Data: A Case Study on PM Across India. IGARSS 2022-2022 IEEE International Geoscience and Remote Sensing Symposium, 2022. IEEE, 1272-1275.
- DESYANA, R., SULISTYANTARA, B., NASRULLAH, N. & FATIMAH, I. Study of the effectiveness of several tree canopy types on roadside green belt in influencing the distribution of NO2 gas emitted from transportation. IOP Conference Series: Earth and Environmental Science, 2017.

IOP Publishing, 012045.

- DIENER, A. & MUDU, P. 2021. How can vegetation protect us from air pollution? A critical review on green spaces' mitigation abilities for air-borne particles from a public health perspective-with implications for urban planning. *Science of the Total Environment*, 796, 148605. DOI: 10.1016/j.scitotenv.2021.148605
- DIVISION, U. N. S. 2023. Metadata for SDG Indicator 11.6.2: Annual mean levels of fine particulate matter (e.g. PM_{2.5} and PM₁₀) in cities (population weighted). <https://unstats.un.org/sdgs/metadata/> (Accessed 21/01/2025)
- ERNST, M., LE MENTEC, S., LOUVRIER, M., LOUBET, B., PERSONNE, E. & STELLA, P. 2022. Impact of urban greening on microclimate and air quality in the urban canopy layer: Identification of knowledge gaps and challenges. *Frontiers in Environmental Science*, 10, 924742. 10.3389/fenvs.2022.924742
- FAN, S., LI, X. & DONG, L. 2019. Field assessment of the effects of land-cover type and pattern on PM₁₀ and PM_{2.5} concentrations in a microscale environment. *Environmental Science and Pollution Research*, 26, 2314-2327. DOI: 10.1007/s11356-018-3697-0
- FARRELL, W. J., CAVELLIN, L. D., WEICHENTHAL, S., GOLDBERG, M. & HATZOPOULOU, M. 2015. Capturing the urban canyon effect on particle number concentrations across a large road network using spatial analysis tools. *Building and environment*, 92, 328-334. DOI: 10.1016/j.buildenv.2015.05.004
- GALLAGHER, J., BALDAUF, R., FULLER, C. H., KUMAR, P., GILL, L. W. & MCNABOLA, A. 2015. Passive methods for improving air quality in the built environment: A review of porous and solid barriers. *Atmospheric environment*, 120, 61-70. DOI: 10.1016/j.atmosenv.2015.08.075
- GOURDJI, S. 2018. Review of plants to mitigate particulate matter, ozone as well as nitrogen dioxide air pollutants and applicable recommendations for green roofs in Montreal, Quebec. *Environmental pollution*, 241, 378-387. DOI: 10.1016/j.envpol.2018.05.053
- GRZEDZICKA, E. 2019. Is the existing urban greenery enough to cope with current concentrations of PM_{2.5}, PM₁₀ and CO₂? *Atmospheric Pollution Research*, 10, 219-233. DOI: 10.1016/j.apr.2018.08.002
- GÜNDOĞDU, S. & ELBİR, T. 2024. Elevating hourly PM_{2.5} forecasting in Istanbul, Türkiye: Leveraging ERA5 reanalysis and genetic algorithms in a comparative machine learning model analysis. *Chemosphere*, 364, 143096. DOI: 10.1016/j.chemosphere.2024.143096
- GUO, B., WANG, X., ZHANG, D., PEI, L., ZHANG, D. & WANG, X. 2020. A land use regression application into simulating spatial distribution characteristics of particulate matter (PM_{2.5}) concentration in city of Xi'an, China. *Polish Journal of Environmental Studies*, 29. DOI: 10.15244/pjoes/118426
- GUO, J., XU, Q., ZENG, Y., LIU, Z. & ZHU, X. X. 2023. Nationwide urban tree canopy mapping and coverage assessment in Brazil from high-resolution remote sensing images using deep learning. *ISPRS Journal of Photogrammetry and Remote Sensing*, 198, 1-15. DOI:

10.1016/j.isprsjprs.2023.02.007

GUO, Y., SU, J. G., DONG, Y. & WOLCH, J. 2019. Application of land use regression techniques for urban greening: An analysis of Tianjin, China. *Urban Forestry & Urban Greening*, 38, 11-21. DOI: 10.1016/j.ufug.2018.10.013

GUPTA, A., MORA, S., ZHANG, F., RUTTEN, M., PRASAD, R. V. & RATTI, C. 2024. GreenScan: Towards large-scale terrestrial monitoring the health of urban trees using mobile sensing. *IEEE Sensors Journal*. DOI: 10.1109/JSEN.2024.3397490

GUPTA, G., RATNAM, M. V., MADHAVAN, B. & NARAYANAMURTHY, C. 2022. Long-term trends in Aerosol Optical Depth obtained across the globe using multi-satellite measurements. *Atmospheric Environment*, 273, 118953. DOI: 10.1016/j.atmosenv.2022.118953

HALIM, N. D. A., LATIF, M. T., MOHAMED, A. F., MAULUD, K. N. A., IDRUS, S., AZHARI, A., OTHMAN, M. & SOFWAN, N. M. 2020. Spatial assessment of land use impact on air quality in mega urban regions, Malaysia. *Sustainable Cities and Society*, 63, 102436. DOI: 10.1016/j.scs.2020.102436

HAN, L., TAN, X., ZHOU, W., LI, W. & QIAN, Y. 2020. Better urban vegetation planning for maximum utility in air pollutant reduction: A theoretical perspective and preliminary analysis in Chinese cities. *Sustainable Cities and Society*, 62, 102377. DOI: 10.1016/j.scs.2020.102377

HAN, L., ZHAO, J., GAO, Y. & GU, Z. 2022a. Prediction and evaluation of spatial distributions of ozone and urban heat island using a machine learning modified land use regression method. *Sustainable Cities and Society*, 78, 103643. DOI: 10.1016/j.scs.2021.103643

HAN, Y., LEE, J., HAIPING, G., KIM, K.-H., WANXI, P., BHARDWAJ, N., OH, J.-M. & BROWN, R. J. 2022b. Plant-based remediation of air pollution: a review. *Journal of Environmental Management*, 301, 113860. DOI: 10.1016/j.jenvman.2021.113860

HART, R., LIANG, L. & DONG, P. 2020. Monitoring, mapping, and modeling spatial-temporal patterns of PM_{2.5} for improved understanding of air pollution dynamics using portable sensing technologies. *International journal of environmental research and public health*, 17, 4914. DOI: 10.3390/ijerph17144914

HASSAN, S., ISLAM, T. & BHUIYAN, M. A. H. 2022. Effects of economic and environmental factors on particulate matter (PM_{2.5}) in the middle parts of Bangladesh. *Water, Air, & Soil Pollution*, 233, 328. DOI: 10.1007/s11270-022-05819-y

HE, Q., GU, Y. & ZHANG, M. 2019. Spatiotemporal patterns of aerosol optical depth throughout China from 2003 to 2016. *Science of The Total Environment*, 653, 23-35. DOI: 10.1016/j.scitotenv.2018.10.307

HEI 2020. State of Global Air 2020. Health Effects Institute Boston, MA, USA.

HELLEBAUT, A., BOISSON, S. & MAHY, G. 2022. Do plant traits help to design green walls for urban air pollution control? A short review of scientific evidences and knowledge gaps. *Environmental Science and Pollution Research*, 29, 81210-81221. DOI: 10.1007/s11356-022-23439-1

- HUANG, S., DAI, C., ZHOU, Y., PENG, H., YI, K., QIN, P., LUO, S. & ZHANG, X. 2018. Comparisons of three plant species in accumulating polycyclic aromatic hydrocarbons (PAHs) from the atmosphere: a review. *Environmental Science and Pollution Research*, 25, 16548-16566. DOI: 10.1007/s11356-018-2167-z
- ISLAM, K. A., ADNAN, M. S. G., ZANNAT, K. E. & DEWAN, A. 2022. Spatiotemporal dynamics of NO₂ concentration with linear mixed models: a Bangladesh case study. *Physics and Chemistry of the Earth, Parts A/B/C*, 126, 103119. DOI: 10.1016/j.pce.2022.103119
- ISLAM, M. N., RAHMAN, K.-S., BAHAR, M. M., HABIB, M. A., ANDO, K. & HATTORI, N. 2012. Pollution attenuation by roadside greenbelt in and around urban areas. *Urban forestry & urban greening*, 11, 460-464. DOI: 10.1016/j.ufug.2012.06.004
- JANHÄLL, S. 2015. Review on urban vegetation and particle air pollution—Deposition and dispersion. *Atmospheric environment*, 105, 130-137. DOI: 10.1016/j.atmosenv.2015.01.052
- JIANG, B., SUN, C., MU, S., ZHAO, Z., CHEN, Y., LIN, Y., QIU, L. & GAO, T. 2021. Differences in airborne particulate matter concentration in urban green spaces with different spatial structures in Xi'an, China. *Forests*, 13, 14. DOI: 10.3390/f13010014
- KAN, X., LIU, X., ZHOU, Z., ZHANG, Y., ZHU, L., SIAN, K. T. C. L. K. & LIU, Q. 2023. Analysis of spatiotemporal variation and influencing factors of PM_{2.5} in China based on multisource data. *Sustainability*, 15, 14656. DOI: 10.3390/su151914656
- KHAN, M. K., NAEEM, K., HUO, C. & HUSSAIN, Z. 2022. The nexus between vegetation, urban air quality, and public health: an empirical study of Lahore. *Frontiers in Public Health*, 10, 842125. DOI: 10.3389/fpubh.2022.842125
- KHANAL, S., POKHREL, R. P., POKHAREL, B., BECKER, S., GIRI, B., ADHIKARI, L. & LAPLANTE, M. D. 2022. An episode of transboundary air pollution in the central Himalayas during agricultural residue burning season in North India. *Atmospheric Pollution Research*, 13, 101270. DOI: 10.1016/j.apr.2021.101270
- KONG, L. & TIAN, G. 2020. Assessment of the spatio-temporal pattern of PM_{2.5} and its driving factors using a land use regression model in Beijing, China. *Environmental monitoring and assessment*, 192, 1-19. DOI: 10.1007/s10661-019-7943-9
- KWON, H.-S., RYU, M. H. & CARLSTEN, C. 2020. Ultrafine particles: unique physicochemical properties relevant to health and disease. *Experimental & molecular medicine*, 52, 318-328. DOI: 10.1038/s12276-020-0405-1
- LABIB, S., HUCK, J. J. & LINDLEY, S. 2021. Modelling and mapping eye-level greenness visibility exposure using multi-source data at high spatial resolutions. *Science of the Total Environment*, 755, 143050. DOI: 10.1016/j.scitotenv.2020.143050
- LI, C., ZHANG, K., DAI, Z., MA, Z. & LIU, X. 2020a. Investigation of the impact of land-use distribution on PM_{2.5} in Weifang: seasonal variations. *International Journal of Environmental Research and Public Health*, 17, 5135. DOI: 10.3390/ijerph17145135
- LI, J., GE, X., HE, Q. & ABBAS, A. 2021a. Aerosol optical depth (AOD): spatial and temporal variations

- and association with meteorological covariates in Taklimakan desert, China. *PeerJ*, 9, e10542. DOI: 10.7717/peerj.10542
- LI, K., LI, C., LIU, M., HU, Y., WANG, H. & WU, W. 2021b. Multiscale analysis of the effects of urban green infrastructure landscape patterns on PM_{2.5} concentrations in an area of rapid urbanization. *Journal of Cleaner Production*, 325, 129324. DOI: 10.1016/j.jclepro.2021.129324
- LI, L., GIRGUIS, M., LURMANN, F., PAVLOVIC, N., MCCLURE, C., FRANKLIN, M., WU, J., OMAN, L. D., BRETON, C. & GILLILAND, F. 2020b. Ensemble-based deep learning for estimating PM_{2.5} over California with multisource big data including wildfire smoke. *Environment international*, 145, 106143. DOI: 10.1016/j.envint.2020.106143
- LI, L., ZHAO, Z., WANG, H., WANG, Y., LIU, N., LI, X. & MA, Y. 2020c. Concentrations of four major air pollutants among ecological functional zones in Shenyang, Northeast China. *Atmosphere*, 11, 1070. DOI: 10.3390/atmos11101070
- LI, M., GU, H., LAM, S. S., SONNE, C. & PENG, W. 2022. Deposition-mediated phytoremediation of nitrogen oxide emissions. *Environmental Pollution*, 308, 119706. DOI: 10.1016/j.envpol.2022.119706
- LI, X., ZHANG, C., LI, W. & LIU, K. 2017. Evaluating the use of DMSP/OLS nighttime light imagery in predicting PM_{2.5} concentrations in the northeastern United States. *Remote Sensing*, 9, 620. DOI: 10.3390/rs9060620
- LI, Y. & MYINT, S. W. 2021. Fine resolution air quality dynamics related to socioeconomic and land use factors in the most polluted desert metropolitan in the American Southwest. *Science of the Total Environment*, 788, 147713. DOI: 10.1016/j.scitotenv.2021.147713
- LIN, H. & JIANG, P. 2022. Analyzing the phased changes of socioeconomic drivers to carbon dioxide and particulate matter emissions in the Yangtze River Delta. *Ecological Indicators*, 140, 109044. DOI: 10.1016/j.ecolind.2022.109044
- LIU, T., YAO, J., CAO, Y., QIN, T., WU, Q., MO, F., ZHAI, H., GONG, H. & LIU, Z. 2024. Spatial-temporal heterogeneity of vegetation reduces concentration of atmospheric pollution particles in the East China Metropolitan Area. *Ecological Indicators*, 169, 112902. DOI: 10.1016/j.ecolind.2024.112902
- LIU, X., ZHANG, X., SCHNELLE-KREIS, J. R., JAKOBI, G., CAO, X., CYRYS, J., YANG, L., SCHLOTTER-HAI, B., ABBASZADE, G. L. & ORASCHE, J. R. 2020. Spatiotemporal characteristics and driving factors of black carbon in Augsburg, Germany: combination of mobile monitoring and street view images. *Environmental Science & Technology*, 55, 160-168. DOI: 10.1021/acs.est.0c04776
- LIU, Z., GUAN, Q., LUO, H., WANG, N., PAN, N., YANG, L., XIAO, S. & LIN, J. 2019. Development of land use regression model and health risk assessment for NO₂ in different functional areas: A case study of Xi'an, China. *Atmospheric Environment*, 213, 515-525. DOI: 10.1016/j.atmosenv.2019.06.044
- LLAGUNO-MUNITXA, M., SHU, X. & MISTRY, B. Multivariate analysis of the influence between

- building design and energy performance, socio-demographic metrics, and the intra-urban environment. *Journal of Physics: Conference Series*, 2021. IOP Publishing, 012056.
- LU, L., WENG, Q., GUO, H., FENG, S. & LI, Q. 2019. Assessment of urban environmental change using multi-source remote sensing time series (2000–2016): A comparative analysis in selected megacities in Eurasia. *Science of the Total Environment*, 684, 567-577. DOI: 10.1016/j.scitotenv.2019.05.344
- LUAN, Q., JIANG, W., LIU, S. & GUO, H. 2020. Impact of Urban 3D Morphology on Particulate Matter 2.5 (PM 2.5) Concentrations: Case Study of Beijing, China. *Chinese geographical science*, 30, 294-308. DOI: 10.1007/s11769-020-1112-5
- MANSOURMOGHADDAM, M., ROUSTA, I., OLAFSSON, H., TKACZYK, P., CHMIEL, S., BARANOWSKI, P. & KRZYSZCZAK, J. 2023. Monitoring of Carbon Monoxide (CO) changes in the atmosphere and urban environmental indices extracted from remote sensing images for 932 Iran cities from 2019 to 2021. *International Journal of Digital Earth*, 16, 1205-1223. DOI: 10.1080/17538947.2023.2196445
- MASRI, S., HOU, H., DANG, A., YAO, T., ZHANG, L., WANG, T., QIN, Z., WU, S., HAN, B. & CHEN, J.-C. J. 2019. Development of spatiotemporal models to predict ambient ozone and NOx concentrations in Tianjin, China. *Atmospheric environment*, 213, 37-46. DOI: 10.1016/j.atmosenv.2019.05.060
- MENG, X., CHEN, L., CAI, J., ZOU, B., WU, C.-F., FU, Q., ZHANG, Y., LIU, Y. & KAN, H. 2015. A land use regression model for estimating the NO2 concentration in Shanghai, China. *Environmental research*, 137, 308-315. DOI: 10.1016/j.envres.2015.01.003
- MORI, J., FERRINI, F. & SAEBO, A. 2018. Air pollution mitigation by urban greening. *Italus Hortus*, 25, 13-22. DOI: 10.26353/j.itahort/2018.1.1322
- MYERS, S. S., GAFFIKIN, L., GOLDEN, C. D., OSTFELD, R. S., H. REDFORD, K., H. RICKETTS, T., TURNER, W. R. & OSOFSKY, S. A. 2013. Human health impacts of ecosystem alteration. *Proceedings of the National Academy of Sciences*, 110, 18753-18760. DOI: 10.1073/pnas.1218656110
- NABOUREH, A., LI, A., BIAN, J., LEI, G., NAN, X., ZHANG, Z., SHAMI, S. & LIN, X. 2024. Green space coverage versus air pollution: a cloud-based remote sensing data analysis in Sichuan, Western China. *International Journal of Digital Earth*, 17, 2383454. DOI: 10.1080/17538947.2024.2383454
- NADEL, I. B., OBERLANDER, C. H. & BOHM, L. R. 1977. *Trees in the City*, Elsevier Science & Technology.
- NIU, X., LI, Y., LI, M., ZHANG, T., MENG, H., ZHANG, Z., WANG, B. & ZHANG, W. 2022. Understanding vegetation structures in green spaces to regulate atmospheric particulate matter and negative air ions. *Atmospheric Pollution Research*, 13, 101534. DOI: 10.1016/j.apr.2022.101534
- O'REGAN, A. C., BYRNE, R., HELLEBUST, S. & NYHAN, M. M. 2022. Associations between Google Street View-derived urban greenspace metrics and air pollution measured using a distributed

- sensor network. *Sustainable Cities and Society*, 87, 104221. DOI: 10.1016/j.scs.2022.104221
- ORGANIZATION, W. H. 2021. *WHO global air quality guidelines: particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide*, World Health Organization.
- PAN, J., CHEN, S., XU, N., CHENG, M., WANG, X., LAN, J., WANG, R. & WANG, Y. 2024. Effect of spatial differentiation of plant communities on PM2.5 and O3 in urban green spaces in Beijing, China. *Journal of Environmental Engineering and Landscape Management*, 32, 372–380-372–380. DOI: 10.3846/jeelm.2024.22359
- PERERA, F. 2018. Pollution from fossil-fuel combustion is the leading environmental threat to global pediatric health and equity: Solutions exist. *International journal of environmental research and public health*, 15, 16. DOI: 10.3390/ijerph15010016
- QI, M., DIXIT, K., MARSHALL, J. D., ZHANG, W. & HANKEY, S. 2022. National land use regression model for NO2 using street view imagery and satellite observations. *Environmental Science & Technology*, 56, 13499-13509. DOI: 10.1021/acs.est.2c03581
- QIU, L., LIU, F., ZHANG, X. & GAO, T. 2018. The reducing effect of green spaces with different vegetation structure on atmospheric particulate matter concentration in Baoji City, China. *Atmosphere*, 9, 332. DOI: 10.3390/atmos9090332
- QIU, L., LIU, F., ZHANG, X. & GAO, T. 2019. Difference of airborne particulate matter concentration in urban space with different green coverage rates in Baoji, China. *International journal of environmental research and public health*, 16, 1465. DOI: 10.3390/ijerph16081465
- RAHMAN, M. M. & HAQUE, S. 2022. Retrieving spatial variation of aerosol level over urban mixed land surfaces using Landsat imageries: Degree of air pollution in Dhaka Metropolitan Area. *Physics and Chemistry of the Earth, Parts a/b/c*, 126, 103074. DOI: 10.1016/j.pce.2021.103074
- RAMOS, Y., ST-ONGE, B., BLANCHET, J.-P. & SMARGIASSI, A. 2016. Spatio-temporal models to estimate daily concentrations of fine particulate matter in Montreal: Kriging with external drift and inverse distance-weighted approaches. *Journal of exposure science & environmental epidemiology*, 26, 405-414. DOI: 10.1038/jes.2015.79
- RAO, M., GEORGE, L. A., ROSENSTIEL, T. N., SHANDAS, V. & DINNO, A. 2014. Assessing the relationship among urban trees, nitrogen dioxide, and respiratory health. *Environmental pollution*, 194, 96-104. DOI: 10.1016/j.envpol.2014.07.011
- SAHA, M., AL KAFY, A., BAKSHI, A., NATH, H., ALSULAMY, S., RAHAMAN, Z. A. & SAROAR, M. 2024. The urban air quality nexus: Assessing the interplay of land cover change and air pollution in emerging South Asian cities. *Environmental Pollution*, 361, 124877. DOI: 10.1016/j.envpol.2024.124877
- SHAHRIN, N. N., ASMAT, A., HAZALI, N. A. & SAHAK, N. Land use and land cover (LULC) modification on the climate and air quality variations. IOP Conference Series: Earth and Environmental Science, 2019. IOP Publishing, 012009.
- SHENG, Q., JI, Y., ZHOU, C., ZHANG, H. & ZHU, Z. 2023. Spatiotemporal Variation and Pattern Analysis of Air Pollution and Its Correlation with NDVI in Nanjing City, China: A Landsat-Based

- Study. *Forests*, 14, 2106. DOI: 10.3390/f14102106
- SHOGRKHODAEI, S. Z., RAZAVI-TERMEH, S. V. & FATHNIA, A. 2021. Spatio-temporal modeling of PM_{2.5} risk mapping using three machine learning algorithms. *Environmental Pollution*, 289, 117859. DOI: 10.1016/j.envpol.2021.117859
- SICARD, P., AGATHOKLEOUS, E., ARAMINIENE, V., CARRARI, E., HOSHIKA, Y., DE MARCO, A. & PAOLETTI, E. 2018. Should we see urban trees as effective solutions to reduce increasing ozone levels in cities? *Environmental pollution*, 243, 163-176. DOI: 10.1016/j.envpol.2018.08.049
- SICARD, P., COULIBALY, F., LAMEIRO, M., ARAMINIENE, V., DE MARCO, A., SORRENTINO, B., ANAV, A., MANZINI, J., HOSHIKA, Y. & MOURA, B. B. 2023. Object-based classification of urban plant species from very high-resolution satellite imagery. *Urban Forestry & Urban Greening*, 81, 127866. DOI: 10.1016/j.ufug.2023.127866
- SMITH, W. H. & STASKAWICZ, B. J. 1977. Removal of atmospheric particles by leaves and twigs of urban trees: Some preliminary observations and assessment of research needs. *Environmental management*, 1, 317-330. DOI: 10.1007/BF01865859
- SUN, Y., ZENG, J. & NAMAITI, A. 2022. Research on the spatial heterogeneity and influencing factors of air pollution: a case study in Shijiazhuang, China. *Atmosphere*, 13, 670. DOI: 10.3390/atmos13050670
- SYAFEI, A., IRAWANDANI, T., BOEDISANTOSO, R., ASSOMADI, A., SLAMET, A. & HERMANA, J. The influence of environmental conditions (vegetation, temperature, equator, and elevation) on tropospheric nitrogen dioxide in urban areas in Indonesia. IOP Conference Series: Earth and Environmental Science, 2019. IOP Publishing, 012034.
- TA, A. T. & PROMCHAN, N. A relationship between urban green space and PM_{2.5} levels in urban and suburban areas of Bangkok. IOP Conference Series: Earth and Environmental Science, 2024. IOP Publishing, 012004.
- TELLA, A. & BALOGUN, A.-L. 2021. GIS-based air quality modelling: Spatial prediction of PM₁₀ for Selangor State, Malaysia using machine learning algorithms. *Environmental Science and Pollution Research*, 1-17. DOI: 10.1007/s11356-021-16150-0
- TIAN, Y. & YAO, X. 2022. Urban form, traffic volume, and air quality: A spatiotemporal stratified approach. *Environment and Planning B: Urban Analytics and City Science*, 49, 92-113. DOI: 10.1177/2399808321995822
- TIAN, Y., YAO, X. & CHEN, L. 2019. Analysis of spatial and seasonal distributions of air pollutants by incorporating urban morphological characteristics. *Computers, Environment and Urban Systems*, 75, 35-48. DOI: 10.1016/j.compenvurbsys.2019.01.003
- TIAN, Y., YAO, X. A., MU, L., FAN, Q. & LIU, Y. 2020. Integrating meteorological factors for better understanding of the urban form-air quality relationship. *Landscape Ecology*, 35, 2357-2373. DOI: 10.1007/s10980-020-01094-6
- TIWARI, A., KUMAR, P., BALDAUF, R., ZHANG, K. M., PILLA, F., DI SABATINO, S., BRATTICH,

- E. & PULVIRENTI, B. 2019. Considerations for evaluating green infrastructure impacts in microscale and macroscale air pollution dispersion models. *Science of The Total Environment*, 672, 410-426. DOI: 10.1016/j.scitotenv.2019.03.350
- TOLLEY, G. S. & COHEN, A. S. 1976. Air pollution and urban land-use policy. *Journal of Environmental Economics and Management*, 2, 247-254. DOI: 10.1016/S0095-0696(76)80002-2
- TOMSON, M., KUMAR, P., BARWISE, Y., PEREZ, P., FOREHEAD, H., FRENCH, K., MORAWSKA, L. & WATTS, J. F. 2021. Green infrastructure for air quality improvement in street canyons. *Environment international*, 146, 106288. DOI: 10.1016/j.envint.2020.106288
- TONG, S., BAMBRICK, H., BEGGS, P. J., CHEN, L., HU, Y., MA, W., STEFFEN, W. & TAN, J. 2022. Current and future threats to human health in the Anthropocene. *Environment international*, 158, 106892. DOI: 10.1016/j.envint.2021.106892
- VAN OORSCHOT, J., SPRECHER, B., VAN'T ZELFDE, M., VAN BODEGOM, P. M. & VAN OUDENHOVEN, A. P. 2021. Assessing urban ecosystem services in support of spatial planning in the Hague, the Netherlands. *Landscape and Urban Planning*, 214, 104195. DOI: 10.1016/j.landurbplan.2021.104195
- VAN RYSWYK, K., PRINCE, N., AHMED, M., BRISSON, E., MILLER, J. D. & VILLENEUVE, P. J. 2019. Does urban vegetation reduce temperature and air pollution concentrations? Findings from an environmental monitoring study of the Central Experimental Farm in Ottawa, Canada. *Atmospheric Environment*, 218, 116886. DOI: 10.1016/j.atmosenv.2019.116886
- VASHISHTA, N., GUPTA, H. & BHARGAVA, B. 2024. Green Walls, Green Roofs, and Urban City Landscapes as Nature-Based Ecological Solutions. *Ornamental Horticulture: Latest Cultivation Practices and Breeding Technologies*. Springer.
- WANG, C., GUO, M., JIN, J., YANG, Y., REN, Y., WANG, Y. & CAO, J. 2022a. Does the spatial pattern of plants and green space affect air pollutant concentrations? Evidence from 37 garden cities in China. *Plants*, 11, 2847. DOI: 10.3390/plants11212847
- WANG CHENGHAO, W. C., WANG CHUYUAN, W. C., MYINT, S. & WANG ZHIHUA, W. Z. 2017. Landscape determinants of spatio-temporal patterns of aerosol optical depth in the two most polluted metropolitans in the United States. DOI: 10.1016/j.scitotenv.2017.07.273
- WANG, H., LIU, G. & SHI, K. 2019. What are the driving forces of urban CO₂ emissions in China? A refined scale analysis between national and urban agglomeration levels. *International Journal of Environmental Research and Public Health*, 16, 3692. DOI: 10.3390/ijerph16193692
- WANG, J., DU, Y., CUI, T., YUE, J., SUN, L., SI, H., DONG, P., LIU, H., LI, Y. & SHI, L. 2024a. A multi-perspective assessment of satellite surface Ozone products in China: Spatiotemporal variability, land cover impacts and pollution monitoring capability. *Remote Sensing Applications: Society and Environment*, 36, 101359. DOI: 10.1016/j.rsase.2024.101359
- WANG, Q., AO, R., CHEN, H., LI, J., WEI, L. & WANG, Z. 2024b. Characteristics of PM_{2.5} and CO₂ Concentrations in Typical Functional Areas of a University Campus in Beijing Based on Low-Cost Sensor Monitoring. *Atmosphere*, 15, 1044. DOI: 10.3390/atmos15091044

- WANG, S., LIU, X., YANG, X., ZOU, B. & WANG, J. 2018a. Spatial variations of PM_{2.5} in Chinese cities for the joint impacts of human activities and natural conditions: A global and local regression perspective. *Journal of cleaner production*, 203, 143-152. DOI: 10.1016/j.jclepro.2018.08.249
- WANG, W., SAMAT, A., ABUDUWAILI, J. & GE, Y. 2020a. Spatio-temporal variations of satellite-based PM_{2.5} concentrations and its determinants in Xinjiang, northwest of China. *International Journal of Environmental Research and Public Health*, 17, 2157. DOI: 10.3390/ijerph17062157
- WANG, W., TIAN, P., ZHANG, J., AGATHOKLEOUS, E., XIAO, L., KOIKE, T., WANG, H. & HE, X. 2022b. Big data-based urban greenness in Chinese megalopolises and possible contribution to air quality control. *Science of The Total Environment*, 824, 153834. DOI: 10.1016/j.scitotenv.2022.153834
- WANG, X., TENG, M., HUANG, C., ZHOU, Z., CHEN, X. & XIANG, Y. 2020b. Canopy density effects on particulate matter attenuation coefficients in street canyons during summer in the Wuhan metropolitan area. *Atmospheric Environment*, 240, 117739. DOI: 10.1016/j.atmosenv.2020.117739
- WANG, Y., DU, H., XU, Y., LU, D., WANG, X. & GUO, Z. 2018b. Temporal and spatial variation relationship and influence factors on surface urban heat island and ozone pollution in the Yangtze River Delta, China. *Science of the Total Environment*, 631, 921-933. DOI: 10.1016/j.scitotenv.2018.03.050
- WANG, Y., GUO, Z. & HAN, J. 2021. The relationship between urban heat island and air pollutants and them with influencing factors in the Yangtze River Delta, China. *Ecological Indicators*, 129, 107976. DOI: 10.1016/j.ecolind.2021.107976
- WEI, F., LI, S., LIANG, Z., HUANG, A., WANG, Z., SHEN, J., SUN, F., WANG, Y., WANG, H. & LI, S. 2021. Analysis of Spatial Heterogeneity and the Scale of the Impact of Changes in PM_{2.5} Concentrations in Major Chinese Cities between 2005 and 2015. *Energies*, 14, 3232. DOI: 10.3390/en14113232
- WRÓBLEWSKA, K. & JEONG, B. R. 2021. Effectiveness of plants and green infrastructure utilization in ambient particulate matter removal. *Environmental Sciences Europe*, 33, 110. DOI: 10.1186/s12302-021-00547-2
- WU, C.-D., CHEN, Y.-C., PAN, W.-C., ZENG, Y.-T., CHEN, M.-J., GUO, Y. L. & LUNG, S.-C. C. 2017. Land-use regression with long-term satellite-based greenness index and culture-specific sources to model PM_{2.5} spatial-temporal variability. *Environmental pollution*, 224, 148-157. DOI: 10.1016/j.envpol.2017.01.074
- WU, H.-W., KUMAR, P. & CAO, S.-J. 2024. The role of roadside green infrastructure in improving air quality in and around elderly care centres in Nanjing, China. *Atmospheric Environment*, 332, 120607. DOI: 10.1016/j.atmosenv.2024.120607
- WU, J., LI, J., PENG, J., LI, W., XU, G. & DONG, C. 2015a. Applying land use regression model to estimate spatial variation of PM_{2.5} in Beijing, China. *Environmental Science and Pollution Research*, 22, 7045-7061. DOI: 10.1007/s11356-014-3893-5

- WU, J., XIE, W., LI, W. & LI, J. 2015b. Effects of urban landscape pattern on PM_{2.5} pollution—a Beijing case study. *PLoS one*, 10, e0142449. DOI: 10.1371/journal.pone.0142449
- XIE, Q. & SUN, Q. 2021. Monitoring the spatial variation of aerosol optical depth and its correlation with land use/land cover in Wuhan, China: a perspective of urban planning. *International Journal of Environmental Research and Public Health*, 18, 1132. DOI: 10.3390/ijerph18031132
- XU, J., LIU, M., CHEN, H. & LUO, M. 2023. Spatially heterogeneous influence of street greenery on street-level PM_{2.5} pollution using mobile monitoring from a three-dimensional perspective. *Urban Climate*, 48, 101414. DOI: 10.1016/j.uclim.2023.101414
- YANG, Y., CERMAK, J., YANG, K., PAULI, E. & CHEN, Y. 2022. Land use and land cover influence on Sentinel-2 aerosol optical depth below city scales over Beijing. *Remote Sensing*, 14, 4677. DOI: 10.3390/rs14184677
- YE, L., FANG, L., TAN, W., WANG, Y. & HUANG, Y. 2016. Exploring the effects of landscape structure on aerosol optical depth (AOD) patterns using GIS and HJ-1B images. *Environmental Science: Processes & Impacts*, 18, 265-276. DOI: 10.1039/c5em00538h
- YLI-PELKONEN, V., VIIPPOLA, V., KOTZE, D. J. & SETälä, H. 2017. Greenbelts do not reduce NO₂ concentrations in near-road environments. *Urban Climate*, 21, 306-317. DOI: 10.1016/j.uclim.2017.08.005
- YSEBAERT, T., KOCH, K., SAMSON, R. & DENYS, S. 2021. Green walls for mitigating urban particulate matter pollution—A review. *Urban Forestry & Urban Greening*, 59, 127014. DOI: 10.1016/j.ufug.2021.127014
- ZENG, L., HANG, J., WANG, X. & SHAO, M. 2022. Influence of urban spatial and socioeconomic parameters on PM_{2.5} at subdistrict level: A land use regression study in Shenzhen, China. *Journal of Environmental Sciences*, 114, 485-502. DOI: 10.1016/j.jes.2021.12.002
- ZHANG, X. & HU, H. 2017. Improving satellite-driven PM_{2.5} models with VIIRS nighttime light data in the Beijing–Tianjin–Hebei region, China. *Remote Sensing*, 9, 908. DOI: 10.3390/rs9090908
- ZHAO, Y., ZHANG, X., CHEN, M., GAO, S. & LI, R. 2022. Regional variation of urban air quality in China and its dominant factors. *Journal of Geographical Sciences*, 32, 853-872. DOI: 10.1007/s11442-022-1975-8