



UNIVERSITY OF LEEDS

This is a repository copy of *Osteoarthritis After Anterior Cruciate Ligament Reconstruction: A Systematic Review of Contributing Factors and Potential Treatments*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/224152/>

Version: Accepted Version

Article:

Charles, S. and Preston, N. orcid.org/0000-0001-8429-7320 (2024) Osteoarthritis After Anterior Cruciate Ligament Reconstruction: A Systematic Review of Contributing Factors and Potential Treatments. *Cureus*, 16 (10). e71188. ISSN 2168-8184

<https://doi.org/10.7759/cureus.71188>

This is an author produced version of an article published in *Cureus: Journal of Medical Science*, made available under the terms of the Creative Commons Attribution License (CC-BY), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

ABSTRACT

Anterior cruciate ligament injuries result in abnormal knee motion and long-term degradation of the joint. ACL reconstruction (ACLR) is done with the aim of restoring normal knee kinematics and slowing the joint degradation process. It does appear that this inevitably happens and can be impacted by a multitude of factors. The aim of this review was to examine the factors that influence the progression of osteoarthritis (OA) after ACLR and examine possible treatments that can aid in slowing that progression. A systematic review was conducted by searching all levels of evidence, for all studies in English that assessed risk factors for developing OA after ACL reconstruction, had a minimum follow-up of 10 years, and used radiographical outcomes to measure the presence of OA. Studies on trial treatments to reduce osteoarthritis after ACL reconstruction were also included. It was found that among the factors associated with an increased risk of post-ACLR OA include meniscal lesions, meniscectomy, increased age at the time of ACLR, increased time from injury to surgery, male sex, loss of knee flexion and extension, smaller thigh girth, graft rupture and residual laxity of the graft. Additionally, in performing the ACLR, anteromedial femoral tunnel placement, higher graft tension and following guidelines for performing anatomic ACLR have been shown to reduce the risk of OA as well. Patients should be adequately counselled on their risk pre-operatively for informed decision-making. Surgeons should also be aware of potential risk factors and how they can be mitigated.

TITLE

Osteoarthritis after ACL Reconstruction - a systematic review of contributing factors and potential treatments

Aims/Hypothesis

The aims of this review were to determine the factors that impact the development of tibiofemoral osteoarthritis after surgical reconstruction of the anterior cruciate ligament injuries in adults, and to investigate what interventions there are to mitigate the development of osteoarthritis after reconstruction surgery for anterior cruciate ligament injuries in adults.

Background/Introduction

The anterior cruciate ligament (ACL) is one of the most commonly injured ligaments in the knee. It accounts for approximately half of all knee ligament injuries [1]. Suffering an ACL tear can be severely damaging to an athlete's career, as it can have them side-lined for several months and even limit their future ability to participate in cutting sports, though approximately 83% of high-level athletes do return to sport after injury. [2]. The impact of this injury on the general population, with persons aiming to maintain activity for their well-being, is hard to quantify. ACL tears can be treated conservatively or via surgical methods, including repair and reconstruction. Surgical reconstruction aims to restore the knee's natural biomechanics and kinematics, which become impaired in the ACL-deficient knee. [3] However, it has been reported that approximately 50% of patients who undergo reconstruction still develop osteoarthritis (OA) in the medium to long term [4], with 1.1% progressing to knee arthroplasty

[5]. The impact of osteoarthritis cannot be overstated. Patients suffer from pain, limitation in ability to conduct daily activities, and functional impairment. Given that ACL injuries mainly occur in a younger patient group [3], this means that a significant group of young people suffer from an increased risk of this post-traumatic arthritis, even if they have reconstruction done. Bearing this in mind, patients choosing to undergo ACL reconstruction (ACLR) should be knowledgeable of their risk of progression to OA. This knowledge will also help in future determinations of which patients are most likely to benefit from ACLR and the development of an optimal standardized pathway for the management of ACL injuries, as well as in determining how best personally tailored treatments can be provided. While previous studies have looked at the development of post-traumatic arthritis in patients with conservatively managed ACL injuries, less data is looking specifically at OA development in patients that have already had ACLR, specifically the factors that influence its progression. Therefore, this study aims to determine the factors that impact the development of tibiofemoral OA after ACLR and to look at the treatments available to mitigate the progression of osteoarthritis after ACLR in adults.

METHODOLOGY

Study Search and Selection

The study search process began with searching the EMBASE, CINAHL, DARE, and Cochrane databases. Additionally, Google Scholar and the University of South Wales Library were searched for additional studies to identify any possible grey literature relevant to the topic. The search was conducted using the terms “Arthritis” OR “post-traumatic arthritis” OR

“Osteoarthritis” OR “post-surgical arthritis” OR “OA” AND “anterior cruciate ligament” OR “ACL” AND “Reconstruction” OR “Surgery” OR “repair” OR “ACLR”. Study titles were examined for relevance and progressed to abstract screening once suitable, where they were screened according to the inclusion/exclusion criteria. The Covidence Systematic Review Software/Program was used to manage the workflow and automate the removal of duplicates. Due to the nature of this intervention, there was some limitation in finding studies that assess the outcomes prospectively and have suitable control groups, or participants without other confounding factors. This is so because ACL injuries are commonly associated with other lesions within the knee joint, all of which can impact outcomes.

The papers included had to be systematic reviews, meta-analyses, randomized control trials, case-control, cohort, or high-quality retrospective studies, published between the years 2005 and 2024, had to assess risk factors for developing OA after ACL reconstruction, must have had a minimum follow-up of 10 years, must have been in English, and must have used a radiographical outcome measure to assess the presence of OA.

Studies on the impact of knee biomechanics and kinematics on the development of arthritis after ACL reconstruction, studies assessing radiographical impact on cartilage after ACL Reconstruction, and studies on available treatments and trial treatments to reduce osteoarthritis after ACL reconstruction were included to answer the review questions adequately. Studies on revision ACL surgery, those done on animal models, or those with follow-ups of less than ten years were excluded. Those studies solely on conservatively managed (no surgery) ACL injuries were excluded. Studies on patellofemoral OA, those that defined OA symptoms using questionnaires/surveys without radiographic evidence, were excluded. Studies that met the eligibility criteria for the review had their entire manuscripts analysed and assessed according to

the Critical Appraisal Skills Programme (CASP) protocols to assess their extrinsic and intrinsic quality, and bias risk. Issues that arose regarding study selection and data extraction were handled by consultation with a specialist in the field, as well as the tutor for this project.

Data Management

Data extracted from the studies included general information (e.g., Author, Article title), study characteristics (e.g., Aim/objectives of the study, Study design, Study inclusion, and exclusion criteria), participant characteristics (e.g., demographics, co-morbidities), data on the Intervention, the radiological outcome measure used, statistical techniques used, length of follow-up, etc. A systematic analysis was conducted on the assessed studies. This involved an analysis of the commonly emerging themes seen between the studies to determine which risk factors may contribute to the development of osteoarthritis after ACLR, with any odds/hazard ratios reported. The studies and their characteristics and findings were tabulated for visual representation and ease of comparison. Due to the significant heterogeneity between the studies, this precluded us from doing a meta-analysis of the reviews included, and a systematic review was therefore done.

Ethical Approval

Ethical Approval was sought and granted by the Ethics Review Board of the University of South Wales. This was deemed to be a low-risk study as it involved no patient-identifiable data and did not influence the course of patient treatment.

Disclaimer

This work was completed in partial fulfilment of an MSc in Sports and Exercise Medicine from the University of South Wales.

RESULTS

The above-mentioned search criteria yielded over 8000 titles from several databases. 762 titles were highlighted for formal title screening, of which 279 were selected for abstract screening. 81 of those studies progressed to full-text screening, of which 29 were included in the final analysis and had their data extracted. This is highlighted in Figure 1. Data extraction was done according to the fields highlighted in the methodology and put into a templated data extraction spreadsheet (Table 1).

In total, 29 studies were included in this analysis. There was a significant level of heterogeneity among the studies. Their structures ranged from case series to retrospective and prospective cohorts, to meta-analyses. There was also significant heterogeneity among the studies in terms of numbers of participants, methods used, and units of assessments of outcomes used.

The factors influencing the progression of osteoarthritis after ACLR were examined. It is best to interpret the results in a systematic manner. Therefore, we will look at pre-surgical, surgical, and post-surgical factors.

Pre-Surgical Factors

Meniscal Lesions

Meniscal lesions found in the preoperative MRI were found to impact the development of OA post-ACLR in our review. This was seen by Barenius et al [6] who in a study of 164 patients with long-term follow-up, found an increased risk of OA development in persons who had medial meniscus lesions at inclusion of the study ((OR 2.5 (1.23-5.24) p = 0.011). Notably,

Filbay et al [7] in a study that included 234 patients at very long-term (32-37 years post-ACLR) follow-up, found that a baseline meniscal lesion increased the odds of developing osteoarthritis by three times (OR: 3.2 (95% CI: 1.3 to 8.3)). Costa-Paz et al. [8] found that patients with meniscal lesions had nearly four times (OR 3.96, $p = 0.039$) increased odds of OA developing after ACLR. Even though the study by Seon et al. [9] had a small sample size of 58 patients, a meniscal lesion was associated with a greatly increased risk (OR 9.19), $p < 0.001$). Furthermore, in a study by Øiestad et al (2010) that included 181 participants, it was seen that the ACL rupture combined with meniscal injury group had a higher prevalence of knee OA compared with the isolated ACL injury group. [10]

Sex

Generally, as seen in a multicentre study by Curado et al. [11] with 182 patients (67 of them being male), male sex appears to confer an increased risk ($p = 0.00018$). This was confirmed by a study by Shelbourne et al. [12], where it was seen that male sex resulted in an increased odds (OR 2.38 (1.44-3.94), $P = .0007$) of developing moderate to severe OA.

Age at Surgery

In our review, it was found that increased age at the time of ACLR was associated with an increased risk of OA. This was seen in several studies including Shelbourne et al. [12] where increased age conferred an odds ratio of 1.06 (1.03-1.09), $P < .0001$ for increasing OA risk. Curado et al. [11] showed that age >30 was associated with increased risk ($p = 0.0026$). Age >25 at ACLR was shown to have increased risk in the study by Seon et al. [9], with a nearly four times increased risk (OR 3.365, $p = 0.034$). Lindanger et al. [13] found that patients with a

Kellgren-Lawrence rating greater than 2 were on average, slightly older as well (OR 1.046 (1.005-1.088), P value .023).

BMI

Curado et al. [11] found that BMI was not associated with an increased risk of post-ACLR OA. However, only 2% of the patients in the study were obese, no comment on the statistical significance of the result was made, and thus it would be difficult to form a conclusion on the relationship from the study. Barenius et al. [6] however, did find BMI to be a risk factor. In this study, 164 patients were randomized to receive either a PT graft or an ST graft, and outcomes were assessed using the Kellgren-Lawrence classification. After adjusting for medial meniscal resection, it was found that BMI resulted in an increased risk of OA for the medial compartment (OR 3.1 (1.22-7.89)) (P = .004), as well as for the patellofemoral joint (OR 3.5 (1.53-7.84) (P = .004).

Time from Injury to Surgery

In several studies in this review, it was seen that the time from initial injury to having ACLR impacted OA outcomes. Curado et al. [11] found that those who waited more than 16 months to have surgery after injury had a higher risk of OA (p= 0.0041). In a study by Sanders et al. [13], early surgery was defined as ACLR within 1 year of injury, and delayed surgery as more than one year after injury. It was found that those who delayed surgery had a four times increased risk of developing a meniscal tear (HR of 3.9 (95% CI, 2.2-6.9)), and a six times higher risk of developing OA (HR 6.2 (95% CI, 3.4-11.4)). Of note, the study by Seon et al. [9] found that even delaying ACLR by 6 months after injury increased the chances of developing OA (OR of

4.767, p-value 0.021). Only the study by Barenius et al. [6] reported that time from injury to surgery was not a risk factor, in groups defined as surgery within 6 months or after, or within 12 months or after (OR 1.0 (0.99-1.02), p-value 0.45).

Surgical Factors

Meniscectomy

Meniscectomy has been shown to be independently associated with increased risk ($p < 0.05$) of OA post-ACLR [12], and this association has been seen in several studies. For example, in a meta-analysis done by Claes et al. [16] that included 1554 patients across all studies, meniscectomy was likely to increase the risk of developing OA by 3.54 times ((95 % CI 2.56–4.91). Even though there was significant heterogeneity between the studies, the combined estimate for the prevalence of OA in the group without meniscectomy was 16.4% (95 % CI 7.0–33.9 %), compared to 50.4% (95 % CI 27.4–73.1 %) in the group with meniscectomy. A study by Ruano et al. (2017) also showed the same odds ratio of 3.54 for meniscectomy. [15] Nakagawa et al. [17] showed that meniscectomy increased the odds (OR 34.1, 95% CI 2.2-522.4, $p = 0.01$), with no significant difference if it was medial or lateral meniscectomy ($p=0.17$). Lindanger et al. [13] also showed the same. Medial meniscectomy increased the risk (OR 1.876 (1.026-3.431), p-value 0.041), as well as lateral meniscectomy (OR 1.960 (1.048-3.669), p-value 0.035). Shelbourne et al. [12] had similar results, with patients with medial meniscectomy having an increased odds ratio of 1.8 (95% CI 1.4-2.5, p-value 0.0012), and OR 1.1 (95% CI 0.85-1.7, p-value .294) for lateral meniscectomy. Interestingly, Barenius et al. [6] showed that meniscus

repair (OR 0.8) reduced the risk of OA compared to meniscectomy (OR 4.2), making a case for meniscus preservation over removal.

Open vs. Arthroscopically Assisted Surgery

Holm et al. [18] conducted a randomised control trial with a sample size of 67 patients and found that there was no difference ($P = 1.000$) in radiographic osteoarthritis outcomes between open and arthroscopically assisted surgery. This is, however, quite a small sample size.

Graft Type

D'Ambrosi et al. [19] found, in a study including over 1500 operated knees with 20 years of follow-up, that patients with ITB grafts had the worst OA outcomes, with 71% of patients developing moderate to severe OA. Patients with BPTB grafts had more radiographic OA than those with HT grafts (29% compared to 13%). This, however, was not proven to be statistically significant. Holm et al. [20] showed that BPTB grafts had a higher rate of OA compared to HT grafts at 10 years follow-up (55% with mild to moderate OA compared to 64%, ($P = 0.27$)). Furthermore, Sanders et al. [14] found in a study that included 964 patients, that the use of allograft tissue in ACLR was associated with an increased risk of developing arthritis. (HR 4.9 (2.05-11.65, $P < .001$)).

Tunnel Placement

A systematic review and meta-analysis by Cinque et al. [21] found that using the transtibial approach to bone tunnel placement was associated with a higher risk of developing OA when compared to the anteromedial approach. When compared to the anteromedial approach, the rates

of PTOA on long-term follow-up of 10 years for the transtibial and anteromedial approaches were 45.6 % and 31.2% respectively ($p < 0.0001$).

Graft Tension

In a randomised controlled trial conducted by Costa et al. [22], patients 10-12 years after surgery were allotted to either have their grafts inserted under high or low tension at the time of surgery. The level of tension was determined by either inserting the graft with the knee in thirty degrees of flexion for the high-tension group or with the knee fully extended (zero degrees of flexion) for the low-tension group. OA outcomes were then assessed using the MRI WORM and radiographic OARSI score. Generally, it was found that there was no significant difference between the groups for the WORM ($P = 0.374$) and OARSI ($P = 0.179$) scores. However, it was seen that males appeared to have relatively poorer outcomes for the OARSI scores in the low-tension group ($P = 0.006$). This seems to suggest that males with low-tension grafts may be predisposed to having OA after ACLR. It is important to note, however, that this study was conducted with a relatively small group of patients ($n=85$), and there was a variety of grafts used, introducing heterogeneity and bias.

Anatomic vs. Non-anatomic Reconstruction

In a study by Rothrauff et al. [23], it was assessed whether doing an anatomic reconstruction would result in less risk of OA compared to a non-anatomic one. It was found that the anatomic group had an OA prevalence of 23.2% and the non-anatomic group had an OA prevalence of 43.9%. Notably, the anatomic group on average checked 9.2 +/- 1.3 of the anatomic ACL reconstruction checklist (AARSC) criteria as defined by van Eck et al. [24], and the non-

anatomic group 5.1 +/- 1.1. An anatomic ACLR was therefore associated with a lower risk of post-ACLR OA.

Post-Surgical Factors

Range of Motion Loss

This review showed that ROM after surgery was a prognostic factor in the progression of OA. Shelbourne et al. [25] found that loss of any knee extension at long-term follow-up was associated with an increased risk of OA development. Loss of knee extension (3.86 (2.38-6.26), $P < 0.0001$) and loss of knee flexion (OR 3.36 (2.14-5.27), $P < 0.0001$) increased the likelihood of having OA at 20-year follow-up. It is therefore necessary to aim to achieve full knee ROM at discharge from physiotherapy, as not achieving this goal increases the risk of knee extension loss (OR 19.7 (95% CI, 10.59-36.65; $P < 0.001$)) and flexion loss (7.97 (95% CI, 4.96-12.86; $P < 0.001$)).

Quadriceps Strength

Øiestad et al. [26] found that loss of quadriceps strength over time was associated with an increased risk of OA (OR 1.00, 95% CI 1.00-1.01). This, however, is a very minuscule increase in risk as seen by the OR of 1. Zandiyeh et al. [27] however, in a study that assessed the differences in symmetry of muscle activation more than 10 years after ACLR, found that on MRI assessment via the WORMS score, patients with smaller thigh muscle girth had more signs of degenerative OA on long term follow up ($p = 0.009$). This was, however, a quite small study of 11 cases and 12 controls for comparison.

Return to Activity

Øiestad et al., [28] assessed the impact of returning to pivoting sport on the progression of OA after ACLR. 71% of patients in that study returned to sport post-ACLR. Of that population, 51% were able to return to their previous sport, with 18.5% of them having radiographic OA at long-term follow-up. Despite this, only 5.5% of the return to sport cohort had symptomatic OA as determined by KOOS scales. In the group that did not return to pivoting sport, 42% had radiographic OA and 25% symptomatic OA. It was seen therefore, that return to pivoting sport post-ACLR was associated with a reduction in OA risk (OR 0.40, 95% CI 0.17 to 0.98) at 15-year long-term follow-up. A similar result was also seen in a study by Lindanger et al. [13], where a return to preinjury level of sports was associated with a reduced risk of developing OA (P = 0.062).

Graft Rupture

In a study by Söderman et al. [29], KL classification was used to determine OA, and MRI was used to determine the integrity of the graft at long-term follow-up of 30 years post ACLR. Patients with ruptured and missing grafts were found to have worse outcome scores and increased signs of OA in the tibiofemoral and patellofemoral compartments (p = 0.0003). This displays the impact of the ACLR on restoring stable ligamentous structures in the joint and its ability to reduce the risk of progression to OA.

Residual Laxity

Struewer et al. [30] conducted a retrospective analysis of hamstring grafts with long-term follow-up. The study did suffer from significant loss to follow-up, however, it did show that OA progression was associated with decreased stability of the graft and joint over time ($p < 0.05$). This was also seen by Curado et al. [11] who demonstrated that residual laxity of the joint $>5\text{mm}$ after ACLR was associated with an increased risk of OA ($p < 0.05$).

Possible Treatments

Options for the treatment of OA post-ACLR are a new region of medicine being discovered. There are, however, some available options on the market. For example, in a randomised controlled trial with 75 patients, Blackburn et al. [31] allocated patients to either receive whole body (WBV) or local muscle (LMV) vibratory stimuli with the aim of improving neuromuscular control post-ACLR. For the whole-body group this involved using a specific machine to apply a vibratory stimulus to the entire body, to cause repeated flexion and extension stimuli to the joints of the lower extremities. The local muscle device caused a similar response but was attached instead to one muscle (e.g. the quadriceps muscle). It was found that WBV increased the peak internal knee extension and that LMV improved the loading rates of the joint in gait. Both these tenets of the gait cycle were associated with PTOA development after ACLR. [31]

Further to this, Andriolo et al. [32] conducted a systematic review of the use of platelet-rich plasma (PRP) in the ACLR population. In that study, it was seen that PRP can be used at many steps along the ACLR process, including in the harvest site for healing, intraarticularly at the end of the surgery and in the bone tunnels. In the final analysis, the use of PRP did have mixed results but did show itself to be a safe option, with potential benefits for graft harvest site

healing and graft maturation. No significant benefit was seen for graft integration. Additionally, Kon et al. [33] also conducted a systematic review of treatments for post-ACLR OA including uses of PRP and stem cells. In that review, two studies showed no benefit in the use of either Adipose-Derived Stem Cells (ADSCs) or Bone Marrow Concentrate (BMC) respectively when they were integrated into the ACLR procedure. Regarding the use of PRP, several papers in the review reported positive benefits of PRP for graft maturation, reducing post-op stiffness and improving graft stability.

DISCUSSION

ACL injuries are common. They affect approximately 1 in 3500 people in the United States, and there are around 400,000 ACLRs done yearly [3]. ACL tears account for approximately half of all knee ligament injuries. These injuries can happen to anyone, but athletes participating in pivoting sports, as well as female athletes, are at greater risk of suffering an ACL tear. [34]. An ACL injury can have an athlete sidelined for months. Approximately 83% of high-level athletes are able to return to sport after injury. [2]. Female athletes are also especially susceptible to ACL injury [34]. This is thought to be due to anatomic factors like their smaller femoral intercondylar notch and their pelvic Q angle, as well as hormonal factors that seem to impact ligamentous laxity. [34, 35]. The average age at which one suffers an ACL injury is 33.9 years. [36]

There are several ways in which an ACL tear can be managed. Conservative management mandates aggressive physiotherapy to regain similar pre-morbid function, and the patient-reported outcomes have been shown to be similar to that of surgical management [37]. Patients selecting this treatment option must be aware, however, of the implications of the ACL deficient knee and its impact on the meniscus [34], which works to prevent aberrant force transmission in the joint and plays a pivotal role in the prevention of osteoarthritis.

It was seen that the status of the meniscus is one of the greatest prognostic indicators. Loss of meniscal substance and meniscal tears results in abnormal force distribution and degenerative changes in the joint over time, accelerating the OA process. Acute tears, those smaller than 2cm and vertical longitudinal tears increase the chances of a meniscal repair healing, as well as tears in the vascularised outer zone [36]. As much as possible, surgeons

should attempt to preserve the meniscus, as without it, as seen in this review, patients are predisposed to pain and early-onset OA [38, 39].

As mentioned earlier, females are at higher risk of developing ACL tears due to structural and hormonal factors. From the review, however, it was males who seemed to have a higher risk of OA progression after ACLR. The results have been mixed though. Against this background, it is important to note that males were more likely to return to sport and have higher activity levels than females after ACLR [40].

It was seen that increased age at the time of ACLR significantly increased the risk of OA post-ACLR. Due to the high heterogeneity between the studies, however, it is difficult to quantify how the level of risk increases with age at the time of ACLR. This therefore makes it difficult to possibly determine a cut-off age after which the risk of doing an ACLR to slow the progression of OA, may outweigh the benefits. What is partially unclear as well, is whether patients being older at the time of surgery is also associated with more chronic ACL tears, and there may therefore be higher levels of OA already present in the knee, thereby acting as a confounding factor. Older patients must therefore be adequately counselled before undergoing ACLR on the risks, including their long-term risk of OA progression.

The time from injury to surgery also significantly influenced disease progression. The impact of an ACL-deficient knee on OA progression has been highlighted above. The question then becomes, what is ideal timing, how early is too early and how late is too late? In a paper by Evans et al. [41], several systematic reviews on the topic were investigated to determine an answer. It was found that surgery too early (before 4-6 weeks) can be complicated by arthrofibrosis, and surgery after 5 months but before 24 months was associated with worse subjective outcome scores than those who had early surgery. Furthermore, it was seen in our

study that surgery later than 6 months after injury increased the risk of developing OA. It would therefore appear that the optimal timing would be somewhere within that window. While the heterogeneity of the studies may have made it difficult to ascertain a golden or specific time frame within which the risk of OA progression is minimized, the general trend was that earlier surgery was better.

The influence of BMI remains contentious. Further studies will have to be done to truly determine if it is a significant contributor. It would seem though, that as with regular osteoarthritis, a higher BMI would result in a higher OA risk [42]. This association remains to be determined clearly in the post-ACLR population, however.

Regarding surgical technique, it was seen that an anatomical reconstruction was superior to a non-anatomical reconstruction for reducing OA risk. The definition of an anatomic ACLR was based on the Anatomic ACL Reconstruction Checklist (AARSC) as developed by van Eck et al. [24]. The AARSC was also shown in a study by Samuelsson et al. [43] to have a lower risk of revision surgery compared with trans-portal drilling with anatomic tunnel placement. Moreover, as seen by Rothrauff et al. [23], the use of the AARSC also reduced the risk of developing OA post-ACLR. The AARSC is therefore a suitable guide for surgeons doing ACLR to follow.

As seen in this review, there was no additional benefit of arthroscopic surgery compared to open mini arthrotomy when looking at OA outcomes [16]. Arthroscopic surgery does, however, have the benefits of smaller incisions, less post-operative pain and shorter length of stay [44], and it would be understood why patients would opt for that method over open surgery.

Regarding OA outcomes, it was found in this review that there was no significant difference in the risk of developing OA post-ACLR between the different types of autografts, with the main ones used including PT and HT grafts. The choice of graft therefore comes down

to surgeon and patient preference and which pros and cons they are willing to accept. On the other hand, however, autografts generally have been shown to have better outcomes than allografts as allografts increase the risk of OA as well as of graft rejection, have a higher rupture risk, and have less favorable patient-reported outcomes [45].

During an ACLR, once the type of graft has been selected, one has to determine how they are going to place the graft. The main options for femoral tunnel drilling and placement are the transtibial (TT), anteromedial (AM) and outside-in (OI) techniques. There are advantages and disadvantages of each technique. The TT technique, however, has been associated with longer and more oblique femoral tunnels, possibly resulting in the AM technique giving better stability and postoperative outcomes compared to the TT technique [46]. In a two-year follow-up study by Carllee et al. [47], it was found that there was no significant difference in graft rupture, outcome scores and function between the OI and AM techniques. Furthermore, in our review, it was found that the TT approach was associated with a higher risk of developing OA post-ACLR than the AM approach [19]. It has been suggested that the TT approach may be less anatomic and thus may allow for more abnormal rotation and anterior translation of the joint compared to the anteromedial approach [48].

Lastly, early and aggressive physiotherapy is the cornerstone of recovery from ACLR. It is just as important as the surgery itself in getting patients back on their feet and improving their outcomes. [49]. Persons who suffer ACL injuries are often young, active individuals who participate in pivoting/cutting sports. These injuries are big blows to their usual levels of activity, and they often want to return to sport as soon as possible. In this review, it was seen that a return to pivoting sport was associated with a reduction in the risk of developing OA post-ACLR [28]. It can be that those patients with less symptomatic and radiographic OA progressed well to

return to sport, while the others didn't, thereby skewing the result. However, as it stands, it appears that a return to pivoting sport may be a protective factor against OA.

Patients with ACL injury have abnormal gait. Analysis of their gait displays reduced peak knee flexion angles and sagittal plane moments. These have been shown to be associated with poor quadriceps function [31]. As seen in the study by Blackburn et al. [31], a vibratory stimulus was shown to improve quadricep muscle function and improve gait biomechanics associated with post-traumatic osteoarthritis. This was done by either the use of a whole body or a local muscle vibration machine. Both of these stimuli were shown to improve aspects of the gait cycle. This is therefore a method that can be integrated into the physiotherapy regimen. Furthermore, PRP was shown to have some benefits when integrated into ACLR. In the systematic reviews by Andriolo et al. [32] and Kon et al. [33], it was shown that PRP can be used at many of the ACLR steps, including in the harvest site for healing, intraarticularly at the end of the surgery and in the bone tunnels. Integration of PRP was shown to have potential benefits for graft harvest site healing, graft maturation, reducing post-op stiffness and improving graft stability.

There were several limitations in this study. As early ACLR surgery was routinely done using BPTB grafts, many of the studies assess outcomes related to those grafts. There remains to be a significant body of long-term data on OA outcomes in HT grafts and allografts. No studies assessed the impact of ethnicity or race on the progression of OA after ACLR. A lot of the studies included were based on majority Caucasian/European populations. It would be interesting to see how this factor can impact, especially given that some African and Indian populations have greater articular mobility [50]. Furthermore, this review was an assessment of radiological OA, and not symptomatic OA. It was seen that not all patients with radiographic features of OA are symptomatic [18]. There was significant heterogeneity between the methods used in the

studies. This included the types of grafts used, the surgical methods, the units of measurement of radiographic OA, and the time from injury to surgery. Additionally, there was also a significant difference between the study methods, as some studies were prospective, while others were retrospective.

Though there was significant heterogeneity, it also contributed to the benefits of this analysis, as this review was able to capture a scoping view of the topic from many perspectives, and through many methods, over a long period of time.

Bias

Due to the heterogeneity between the study methods, participants and outcome measures used etc, there was certainly some elevated risk of bias in this review. Most of the studies were also found on online databases, resulting in some measure of publication bias as well. Additionally, only English studies were included, introducing language bias into the review. Finally, some of the studies did not report on the size of the effects measured with confidence intervals or p values. This made it difficult to know whether some results were truly significant or not. To mitigate the risks highlighted, the search for studies was significantly robust to ensure that all appropriate studies on the topic were included. Furthermore, significant attention was paid to ensure the studies selected for analysis strictly fit the eligibility criteria and were of high quality.

RECOMMENDATIONS

In the study by Rothrauff et al. [23], it was seen that an anatomic ACLR was associated with a decreased risk of OA. It also guides adequate documentation of the procedure and has been shown to have benefits for patients. It was also seen that deficiencies in quadriceps strength as

well as limited range of motion after ACLR can be deleterious to patients. All patients should therefore undergo early and tailored physiotherapy with the aim of achieving full range of motion in flexion and extension and developing adequate quadriceps muscle mass by the point of discharge from physiotherapy. Though only the study by Costa et al. [22] explicitly investigated the impact of tension of the ACL graft, there was some evidence that low-tension grafts should be avoided. Due to the potential benefit of return to sport for OA outcomes, it can be encouraged, but should be guided by professionals in the field. Early surgery is better. Several studies showed that delay of surgery was associated with poorer OA outcomes. ACLR should therefore be done as early as safely possible, as the risk of OA was shown to increase if surgery was done even at 6 months post-injury. Surgeons should consider persevering the meniscus as much as possible to reduce OA risk. This would involve attempting repair rather than heading straight to removal for meniscal injuries seen intra-operatively. [51, 52] Surgeons can also give thought to the use of PRP during ACLR as positive results have been shown thus far for graft maturation, stability and donor site healing.

FURTHER RESEARCH

Though some studies did partially assess it, the outcomes of meniscectomy versus repair of meniscal tears, and that impact on the development of OA deserve a dedicated study with a sufficiently large sample size. Further to this, biologics are being investigated as a potential mediator of PTOA after ACLR. However, these outcomes have to be assessed with radiographic outcomes after long-term follow-up > 10 years.

CONCLUSION

This review showed that there are a multitude of factors that can impact on the progression of OA after ACLR. Those studies with the sizes of their effects stated have been summarised in Figure 2. Among the factors associated with an increased risk of post-ACLR OA include preoperatively diagnosed meniscal lesions, meniscectomy, increased age at the time of ACLR, increased time from injury to surgery, male sex, loss of knee flexion and extension, smaller thigh girth, graft rupture and residual laxity of the graft. Additionally, in performing the ACLR, anteromedial femoral tunnel placement, higher graft tension and following guidelines for performing anatomic ACLR have been shown to reduce the risk of OA as well. Finally, the importance of aggressive physiotherapy was seen as any deficit in the range of motion in flexion and extension was shown to impact on OA outcomes. Deficits in quadriceps strength were also shown to contribute to abnormal knee kinematics, which can lead to increased risk. Further to this, it was seen that vibratory stimuli were able to reduce some of the quadriceps dysfunction associated with ACL injury and work as a useful adjunct in rehabilitation. PRP was also shown to be a useful aid, with its ability to be integrated and show benefit at multiple steps in ACLR. Patients should be adequately counselled on their risk pre-operatively for informed decision-making. Surgeons should also be aware of potential risk factors and the steps they can take for those risks to be mitigated.

1. Kaeding CC, Léger-St-Jean B, Magnussen RA. Epidemiology and diagnosis of anterior cruciate ligament injuries. *Clinics in Sports Medicine*. 2017 Jan;36(1):1–8. doi:10.1016/j.csm.2016.08.001
2. Lai CC, Ardern CL, Feller JA, Webster KE. Eighty-three per cent of elite athletes return to preinjury sport after Anterior Cruciate Ligament Reconstruction: A systematic review with meta-analysis of return to sport rates, graft rupture rates and performance outcomes. *British Journal of Sports Medicine*. 2017 Feb 21;52(2):128–38. doi:10.1136/bjsports-2016-096836
3. Evans J. Anterior cruciate ligament knee injury [Internet]. U.S. National Library of Medicine; 2023 [cited 2024 Jun 22]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK499848/>
4. Barenus B, Ponzer S, Shalabi A, Bujak R, Norlén L, Eriksson K. Increased risk of osteoarthritis after anterior cruciate ligament reconstruction. *The American Journal of Sports Medicine*. 2014 Mar 18;42(5):1049–57. doi:10.1177/0363546514526139
5. Visnes H, Gifstad T, Persson A, Lygre SH, Engebretsen L, Drogset JO, et al. ACL reconstruction patients have increased risk of knee arthroplasty at 15 years of follow-up. *JBJS Open Access*. 2022;7(2). doi:10.2106/jbjs.oa.22.00023
6. Barenus B, Ponzer S, Shalabi A, Bujak R, Norlén L, Eriksson K. Increased risk of osteoarthritis after ACL reconstruction – a 14-year follow-up study of a randomized controlled trial. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2013 Oct;29(10). doi:10.1016/j.arthro.2013.07.080
7. Filbay S, Gauffin H, Andersson C, Kvist J. Prognostic factors for tibiofemoral and patellofemoral osteoarthritis 32–37 years after anterior cruciate ligament injury managed with early surgical repair or rehabilitation alone. *Osteoarthritis and Cartilage*. 2021 Dec;29(12):1682–90. doi:10.1016/j.joca.2021.08.009
8. Costa-Paz M, Garcia-Mansilla I, Marciano S, Ayerza MA, Muscolo DL. Knee-related quality of life, functional results and osteoarthritis at a minimum of 20 years' follow-up after Anterior Cruciate Ligament Reconstruction. *The Knee*. 2019 Jun;26(3):666–72. doi:10.1016/j.knee.2019.04.010
9. Seon JK, Song EK, Park SJ. Osteoarthritis after anterior cruciate ligament reconstruction using a patellar tendon autograft. *International Orthopaedics*. 2006 Jan 25;30(2):94–8. doi:10.1007/s00264-005-0036-0
10. Øiestad BE, Holm I, Aune AK, Gunderson R, Myklebust G, Engebretsen L, et al. Knee function and prevalence of knee osteoarthritis after anterior cruciate ligament reconstruction. *The American Journal of Sports Medicine*. 2010 Aug 16;38(11):2201–10. doi:10.1177/0363546510373876
11. Curado J, Hulet C, Hardy P, Jenny J-Y, Rousseau R, Lucet A, et al. Very long-term osteoarthritis rate after anterior cruciate ligament reconstruction: 182 cases with 22-year' follow-up. *Orthopaedics & Traumatology: Surgery & Research*. 2020 May;106(3):459–63. doi:10.1016/j.otsr.2019.09.034
12. Shelbourne KD, Benner RW, Gray T. Results of anterior cruciate ligament reconstruction with patellar tendon autografts: Objective factors associated with the development of osteoarthritis at 20 to 33 years after surgery. *The American Journal of Sports Medicine*. 2017 Aug 14;45(12):2730–8. doi:10.1177/0363546517718827
13. Lindanger L, Strand T, Mølster AO, Solheim E, Fischer-Bredenbeck C, Ousdal OT, et al. Predictors of osteoarthritis development at a median 25 years after anterior cruciate

- ligament reconstruction using a patellar tendon autograft. *The American Journal of Sports Medicine*. 2022 Mar 2;50(5):1195–204. doi:10.1177/03635465221079327
14. Sanders TL, Kremers HM, Bryan AJ, Fruth KM, Larson DR, Pareek A, et al. Is anterior cruciate ligament reconstruction effective in preventing secondary meniscal tears and osteoarthritis? *The American Journal of Sports Medicine*. 2016 Mar 8;44(7):1699–707. doi:10.1177/0363546516634325
 15. Ruano JS, Sitler MR, Driban JB. Prevalence of radiographic knee osteoarthritis after anterior cruciate ligament reconstruction, with or without meniscectomy: An evidence-based practice article. *Journal of Athletic Training*. 2017 Jun 1;52(6):606–9. doi:10.4085/1062-6050-51.2.14
 16. Claes S, Hermie L, Verdonk R, Bellemans J, Verdonk P. Is osteoarthritis an inevitable consequence of Anterior Cruciate Ligament Reconstruction? A meta-analysis. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2012 Oct 26;21(9):1967–76. doi:10.1007/s00167-012-2251-8
 17. Nakagawa T, Kimura M, Ogoshi A, Yanagisawa S, Yorifuji H, Takagishi K. Risk factors for the development of osteoarthritis after Anterior Cruciate Ligament Reconstruction. *The Kitakanto Medical Journal*. 2017;67(2):129–33. doi:10.2974/kmj.67.129
 18. Holm I, Øiestad BE, Risberg MA, Gunderson R, Aune AK. No differences in prevalence of osteoarthritis or function after open versus endoscopic technique for Anterior Cruciate Ligament Reconstruction. *The American Journal of Sports Medicine*. 2012 Sept 13;40(11):2492–8. doi:10.1177/0363546512458766
 19. D’Ambrosi R, Carrozzo A, Meena A, Corona K, Yadav AK, Annibaldi A, et al. A slight degree of osteoarthritis appears to be present after anterior cruciate ligament reconstruction compared with contralateral healthy knees at a minimum of 20 years: A systematic review of the literature. *Journal of Experimental Orthopaedics*. 2024 Apr;11(2). doi:10.1002/jeo2.12017
 20. Holm I, Øiestad BE, Risberg MA, Aune AK. No difference in knee function or prevalence of osteoarthritis after reconstruction of the anterior cruciate ligament with 4-strand hamstring autograft versus patellar tendon—bone autograft. *The American Journal of Sports Medicine*. 2010 Jan 23;38(3):448–54. doi:10.1177/0363546509350301
 21. Cinque ME, Kunze KN, Williams BT, Moatshe G, LaPrade RF, Chahla J. Higher Incidence of Radiographic Posttraumatic Osteoarthritis With Transtibial Femoral Tunnel Positioning Compared With Anteromedial Femoral Tunnel Positioning During Anterior Cruciate Ligament Reconstruction: A Systematic Review and Meta-analysis. *The American Journal of Sports Medicine*. 2021 Mar 26;50(1):255–63. doi:10.1177/0363546521993818
 22. Costa MQ, Badger GJ, Chrostek CA, Carvalho OD, Faiola SL, Fadale PD, et al. Effects of initial graft tension and patient sex on knee osteoarthritis outcomes after ACL reconstruction: A randomized controlled clinical trial with 10- to 12-year follow-up. *The American Journal of Sports Medicine*. 2022 Oct 19;50(13):3510–21. doi:10.1177/03635465221124917
 23. Rothrauff BB, Jorge A, de Sa D, Kay J, Fu FH, Musahl V. Anatomic ACL reconstruction reduces risk of post-traumatic osteoarthritis: A systematic review with minimum 10-year follow-up. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2019 Aug 30;28(4):1072–84. doi:10.1007/s00167-019-05665-2


24. van Eck CF, Gravare-Silbernagel K, Samuelsson K, Musahl V, van Dijk CN, Karlsson J, et al. Evidence to support the interpretation and use of the anatomic anterior cruciate ligament reconstruction checklist. *The Journal of Bone & Joint Surgery*. 2013 Oct;95(20). doi:10.2106/jbjs.l.01437
25. Shelbourne KD, Urch SE, Gray T, Freeman H. Loss of normal knee motion after anterior cruciate ligament reconstruction is associated with radiographic arthritic changes after surgery. *The American Journal of Sports Medicine*. 2011 Oct 11;40(1):108–13. doi:10.1177/0363546511423639
26. Øiestad BE, Holm I, Gunderson R, Myklebust G, Risberg MA. Quadriceps muscle weakness after anterior cruciate ligament reconstruction: A risk factor for knee osteoarthritis? *Arthritis Care & Research*. 2010 Nov 30;62(12):1706–14. doi:10.1002/acr.20299
27. Zandiyeh P, Parola LR, Costa MQ, Hague MJ, Molino J, Fleming BC, et al. Long-term bilateral neuromuscular function and knee osteoarthritis after anterior cruciate ligament reconstruction. *Bioengineering*. 2023 Jul 6;10(7):812. doi:10.3390/bioengineering10070812
28. Øiestad BE, Holm I, Risberg MA. Return to pivoting sport after ACL reconstruction: Association with osteoarthritis and knee function at the 15-year follow-up. *British Journal of Sports Medicine*. 2018 Mar 17;52(18):1199–204. doi:10.1136/bjsports-2017-097718
29. Söderman T, Werner S, Wretling M-L, Hänni M, Mikkelsen C, Sundin A, et al. Knee function 30 years after ACL reconstruction: A case series of 60 patients. *Acta Orthopaedica*. 2021 Jul 28;92(6):716–21. doi:10.1080/17453674.2021.1959709
30. Struwer J, Ziring E, Frangen TM, Efe T, Meißner S, Buecking B, et al. Clinical outcome and prevalence of osteoarthritis after isolated anterior cruciate ligament reconstruction using hamstring graft: Follow-up after two and ten years. *International Orthopaedics*. 2012 Sept 2;37(2):271–7. doi:10.1007/s00264-012-1653-z
31. Blackburn T, Padua DA, Pietrosimone B, Schwartz TA, Spang JT, Goodwin JS, et al. Vibration improves gait biomechanics linked to posttraumatic knee osteoarthritis following anterior cruciate ligament injury. *Journal of Orthopaedic Research*. 2020 Aug 12;39(5):1113–22. doi:10.1002/jor.24821
32. Andriolo L, Di Matteo B, Kon E, Filardo G, Venieri G, Marcacci M. PRP augmentation for ACL reconstruction. *BioMed Research International*. 2015;2015:1–15. doi:10.1155/2015/371746
33. Kon E, Di Matteo B, Altomare D, Iacono F, Kurpyakov A, Lychagin A, et al. Biologic agents to optimize outcomes following ACL repair and Reconstruction: A systematic review of clinical evidence. *Journal of Orthopaedic Research*. 2021 Mar 8;40(1):10–28. doi:10.1002/jor.25011
34. Azar FM, Beaty JH, Canale ST, Campbell WC. *Campbell's Operative Orthopaedics*. Philadelphia, PA: Elsevier; 2021.
35. Ireland ML. The female ACL: Why is it more prone to injury? *Orthopedic Clinics of North America*. 2002 Oct;33(4):637–51. doi:10.1016/s0030-5898(02)00028-7
36. Nicholls M, Aspelund T, Ingvarsson T, Briem K. Nationwide study highlights a second peak in ACL Tears for women in their early forties. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2017 Nov 21;26(2):648–54. doi:10.1007/s00167-017-4807-0
37. Migliorini F, Oliva F, Eschweiler J, Torsiello E, Hildebrand F, Maffulli N. Knee osteoarthritis, joint laxity and Proms following conservative management versus surgical

- reconstruction for ACL Rupture: A meta-analysis. *British Medical Bulletin*. 2022 Nov 21;145(1):72–87. doi:10.1093/bmb/ldac029
38. Raj MA. Knee Meniscal tears [Internet]. U.S. National Library of Medicine; 2023 [cited 2024 Jun 30]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK431067/>
 39. Beaufils P, Becker R, Kopf S, Matthieu O, Pujol N. The knee meniscus: Management of traumatic tears and degenerative lesions. *EFORT Open Reviews*. 2017 May;2(5):195–203. doi:10.1302/2058-5241.2.160056
 40. Bruder AM, Culvenor AG, King MG, Haberfield M, Roughead EA, Mastwyk J, et al. Let's talk about sex (and gender) after ACL injury: A systematic review and meta-analysis of self-reported activity and knee-related outcomes. *British Journal of Sports Medicine*. 2023 Mar 8;57(10):602–10. doi:10.1136/bjsports-2022-106099
 41. Evans S, Shaginaw J, Bartolozzi A. ACL reconstruction - it's all about timing [Internet]. U.S. National Library of Medicine; 2014 [cited 2024 Jun 30]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4004131/>
 42. Coggon D, Reading I, Croft P, McLaren M, Barrett D, Cooper C. Knee osteoarthritis and obesity. *International Journal of Obesity*. 2001 May;25(5):622–7. doi:10.1038/sj.ijo.0801585
 43. Samuelsson K, Desai N, Andernord D, Sundemo DS, Alentorn-Geli E, Musahl V, et al. Revision surgery in Anterior Cruciate Ligament Reconstruction - A cohort study of 17,682 patients using the anatomic anterior cruciate ligament reconstruction scoring checklist applied to the Swedish National Knee Ligament Register. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2017 Oct;33(10). doi:10.1016/j.arthro.2017.08.171
 44. Weale AE, Ackroyd CE, Mani GV, Winson IG. Day-case or short-stay admission for arthroscopic knee surgery: A randomised controlled trial [Internet]. U.S. National Library of Medicine; 1998 [cited 2024 Jun 23]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2503003/>
 45. Baawa-Ameyaw J, Plastow R, Begum FA, Kayani B, Jeddy H, Haddad F. Current concepts in graft selection for Anterior Cruciate Ligament Reconstruction. *EFORT Open Reviews*. 2021 Sept;6(9):808–15. doi:10.1302/2058-5241.6.210023
 46. Mao Y, Zhang K, Li J, Fu W. Transtibial versus anteromedial portal technique for femoral tunnel drilling in primary single-bundle Anterior Cruciate Ligament Reconstruction: A meta-analysis of level 1 and 2 evidence of clinical, revision, and radiological outcomes. *The American Journal of Sports Medicine*. 2021 Oct 15;51(1):250–62. doi:10.1177/03635465211044476
 47. Carllee T, Ries Z, Wolf B, Amendola A, Hettrich C, Bollier M. Two year follow-up comparing 2-incision vs anteromedial portal techniques for femoral drilling during primary ACL reconstruction. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2016 Jun;32(6). doi:10.1016/j.arthro.2016.03.046
 48. Alentorn-Geli E, Samitier G, Álvarez P, Steinbacher G, Cugat R. Anteromedial portal versus transtibial drilling techniques in ACL reconstruction: A blinded cross-sectional study at two- to five-year follow-up. *International Orthopaedics*. 2010 Apr 20;34(5):747–54. doi:10.1007/s00264-010-1000-1
 49. Kochman M, Kasprzak M, Kielar A. ACL reconstruction: Which additional physiotherapy interventions improve early-stage rehabilitation? A systematic review.

International Journal of Environmental Research and Public Health. 2022 Nov
29;19(23):15893. doi:10.3390/ijerph192315893

50. Beighton P, Solomon L, Soskolne CL. Articular mobility in an African population. *Annals of the Rheumatic Diseases*. 1973 Sept 1;32(5):413–8. doi:10.1136/ard.32.5.413
51. Hart AJ, Buscombe J, Malone A, Dowd GS. Assessment of osteoarthritis after reconstruction of the Anterior Cruciate Ligament. *The Journal of Bone and Joint Surgery British volume*. 2005 Nov;87-B(11):1483–7. doi:10.1302/0301-620x.87b11.16138
52. Gerhard P, Bolt R, Dück K, Mayer R, Friederich NF, Hirschmann MT. Long-term results of arthroscopically assisted anatomical single-bundle anterior cruciate ligament reconstruction using patellar tendon autograft: Are there any predictors for the development of osteoarthritis? *Knee Surgery, Sports Traumatology, Arthroscopy*. 2012 Apr 10;21(4):957–64. doi:10.1007/s00167-012-2001-y

APPENDIX 1 –PRISMA Chart



APPENDIX 2 – Summary of Studies

Study ID	Methods: Design	Sample size at follow-up	Length of Follow Up	Risk Factor/Treatment/Intervention Assessed	Outcome Measures Used	Main Result
Barenius 2013 [6]	Randomized controlled trial	135	14 years	Patellar tendon vs Semitendinosus grafts	KL. Patellar height: Insall Salvati index	No significant difference was seen between the graft types (P = .073). Risk Factors for Osteoarthritis were: BMI 25 kg/m ² at 2-year follow-up (OR 3.1 (1.22-7.89) p = 0.004) and Medial meniscus resection (OR 3.6 (1.42-9.29) P= 0.001)
Filbay 2021 [7]	Prospective cohort study	251	32 to 37 years	early surgical repair vs non-augmented repair plus physiotherapist-supervised rehabilitation vs physiotherapy only	Kellgren and Lawrence (KL) classification system	A preoperative meniscal injury increased the odds of OA (OR: 3.2 (95% CI: 1.3 to 8.3)), compared to no meniscal injury. Meniscal surgery resulted in a higher risk of OA (OR 3.0 (95% CI 1.2, 7.8)) compared to no meniscal injury or surgery.
Costa-Paz 2019 [8]	Retrospective cohort study	72	median follow-up of 22 (IQR 21–25)	Long term ACLR follow up	IKDC criteria	Meniscal lesions were associated with an increased risk of developing OA (OR 3.96, 95% CI 1.07-14.65)
Seon 2006 [9]	Prospective cohort study	58	11.2 years (range 8.6–13.8 years)	Long term ACLR follow up, Patellar tendon autograft	Kellgren and Lawrence's classification	Factors associated with increased OA risk: Age >25 years at the time of reconstruction - OR 3.37, p=0.034. Time from injury to reconstruction >6 months OR 4.77, p=0.021. Meniscal injury OR 9.19, p<0.001. Medial meniscectomy OR 3.71, p=0.03, and lateral meniscectomy OR 1.20, p>0.05.
Øiestad 2010 [10]	Prospective cohort study	181	10-15-years	Bone-patellar tendon-bone (BPTB) autograft, either with miniopen or arthroscopic procedure. Isolated vs combined ACLR and meniscal lesions	Kellgren and Lawrence (KL) classification system	Combined injury group resulted in significantly higher prevalence of radiographic knee OA compared with the isolated injury group (80% and 62%, P< 0.008)
Curado 2020 [11]	Retrospective cohort study	182	22 ± 1 years	Long term ACLR follow up	IKDC criteria	Risk factors for OA after ACLR: Age > 30 years at surgery (p=0.0026), Male sex (p=0.00018), Injury-to-surgery time > 16 months (p=0.0041), Pivot sport (p< 0.05), Moderate or strenuous physical activity (p< 0.05), Meniscectomy (p< 0.05), Residual laxity >5 mm (p< 0.05).
Shelbourne 2017 [12]	Retrospective cohort study	423	22.5 years (range, 20.0-33.1 years)	Long term ACLR follow up, patella tendon autografts	IKDC criteria	Factors associated with increased OA risk after regression model include medial meniscectomy OR 2.9 (1.8-4.7) p<.0001, loss of knee extension OR 2.1 (1.1-3.9) p= 0.024, older age at surgery OR 1.04 (1.0-1.07) p= 0.011.
Lindanger 2022 [13]	Prospective cohort study	235	median 25 years (range, 20-31)	Long term ACLR follow up	Kellgren and Lawrence (KL) classification system	Medial and lateral meniscal surgery were found to be independent risk factors for OA development after ACLR (OR 1.88 (95% CI, 1.03-3.43; P = .041 for medial meniscal surgery and OR 1.96 (95% CI, 1.05-3.67; P = .035) for lateral meniscal surgery. Return to sports was associated with a decreased odds of developing OA (P = .06)

Sanders 2016 [14]	Retrospective cohort study	964	13.7 years	Timing of ACLR, Long term ACLR follow up	Physician-diagnosed arthritis	Factors associated with increased risk of OA were: delayed reconstruction HR of 6.2 (95% CI, 3.4-11.4), age older than 21 years at the time of injury (P = .02), presence of both medial and lateral meniscal tears at the time of ACL injury (P < .001), treatment with meniscectomy (P = .07), an articular cartilage injury at the time of the ACL tear (P = .006) and use of allograft tissue during ACLR (P < 0 .001).
Ruano 2017 [15]	Systematic review	1554	10 to 24.5 years	ACLR with and without meniscectomy	Ahlbäck, modified Fairbanks grades, KL and IKDC	Meniscectomy significantly increased the risk of developing OA by 3.54 times.
Claes 2013 [16]	Meta analysis	1554	10 to 24.5 years	Long term ACLR follow up	IKDC criteria	Meniscectomy was associated with a 3.54 times higher risk (95 % CI 2.56–4.91) than ACLR without meniscectomy.
Nakagawa 2017 [17]	Retrospective cohort study	40	Minimum 15 years	Long term ACLR follow up	Kellgren and Lawrence (KL) classification system	Meniscectomy was identified as the only risk factor for OA progression (p=0.01)
Holm 2012 [18]	Randomized controlled trial	67	12 years	endoscopic patellar tendon–bone (PTB) reconstruction (ENDO) vs open PTB reconstruction (OPEN)	Kellgren and Lawrence (KL) classification system	There were no significant differences between the OPEN and ENDO groups for OA outcomes, with 80% and 79% of the respective groups developing grade 2 or higher OA. (P = 1.000)
D'Ambrosi 2024 [19]	Systematic review	1552	Mean 23.34 ± 2.56 years.	Long term follow up of various ACLR grafts	Kellgren–Lawrence, IKDC and Ahlbäck classification.	71% of patients that received ITB graft were classified as IKDC C or D at radiographic follow-up. For PT grafts (IKDC grades C and D—29% vs. 13%). No significant difference between the PT and HS grafts in terms of OA were seen. (19.3% for PT vs. 19.6% for HT, p = 0.94).
Holm 2010 [20]	Randomized controlled trial	57	10 years	Patellar tendon vs 4 bundle Hamstring graft	Kellgren and Lawrence (KL) classification system	No significant differences were found between the two types of grafts. 55% and 64% of the participants had significant OA in the hamstring and the PT groups, respectively (P=0.27).
Cinque 2022 [21]	Systematic Review	1546	10.9 years (5.4-17.8)	Anteromedial vs Transtibial approach to femoral tunnel positioning	Kellgren-Lawrence (KL) classification, Ahlbäck, International Knee Documentation Committee (IKDC), and Osteoarthritis Research Society International (OARSI).	TT approach associated with increased risk of PTOA (45.6%) compared with the AM approach (31.2%) at long-term follow-up (Z = 8.58; P < .0001)
Costa 2022 [22]	Randomized controlled trial	95	10 to 12 years	Graft tension	OARSI, Whole-Organ Magnetic Resonance Imaging Score [WORMS]	Males with low tension grafts had significantly worse outcomes compared to females (P=.006).
Rothrauff 2020 [23]	Systematic review	2224	Mean 15.3-15.9 years	Anatomic vs non-anatomic ACLR	OARSI, WORMS, Anatomic ACL Reconstruction Checklist (AARSC)	23.2% of patients with anatomic ACLR developed OA compared to 43.9% that had non-anatomic ACLR.

Shelbourne 2012 [25]	Prospective cohort study	780	mean of 10.5 ± 4.2 years	Knee motion after ACLR	IKDC criteria	At final follow-up, loss of range of motion, those who had partial medial meniscectomy, and those who suffered articular cartilage damage each had about 2 times the odds of having OA.
Øiestad 2010 [26]	Prospective cohort study	258	10-15 years	Quadriceps muscle weakness	muscle strength - cybex 6000. radiologic examination - Kellgren/Lawrence classification	Loss of quadriceps muscle strength showed no significant association with developing OA post-ACLR. OR 1.00, 95% CI 1.0-1.01. Increased age and meniscal injury were associated with higher OA risk (OR 1.06 and 2.05 respectively).
Zandiyeh 2023 [27]	Case-control study	11	11.9 ± 1.3 years	ACLR and Neuromuscular function impact on OA	WORMS	Smaller muscle girth of the thigh was significantly associated with inferior WORM scores ($B_0 = -8.3$, $\beta = -20.8$, $p = 0.009$)
Øiestad 2018 [28]	Prospective cohort study	210	15 years	Return to pivoting sport	Kellgren and Lawrence (KL) classification system	Participants that returned to pivoting sports were likely to develop radiographic OA (OR 0.40, 95%CI 0.17 to 0.98) compared with those who did not.
Söderman 2021 [29]	Retrospective cohort study	60	31 years (range 28-33 years)	Graft rupture	KL, MRI was used to assess the menisci and the structural integrity of the ACL graft.	Ruptured ACL graft was associated with increased risk of medial tibiofemoral OA ($p=0.0003$)
Struwer 2013 [30]	Retrospective cohort study	52	Average 10.2 years (8-13 years)	Hamstring tendon grafts/anterior laxity	Jäger- Wirth classification	There was a significant correlation between instrumental anterior laxity and radiological OA ($p<0.05$)
Blackburn 2021 [31]	Randomized controlled trial	75	Not stated	Whole body and local muscle vibration	Gait speed, peak knee flexion angle, peak knee virus angle, peak vGRF, Linear vGFR loading rate, Instantaneous vGFR loading rate, KAM - internal knee abduction moment, KEM - internal knee extension moment	Vibration reduced pain, improved quadriceps function, and enhanced CNS excitability.
Andriolo 2015 [32]	Systematic Review	Not stated	Not stated	PRP integration in ACLR	Variable	PRP showed positive effects on graft maturation, bone healing
Kon 2022 [33]	Systematic Review	Not stated	Not stated	Biologic Agent use in ACLR	Variable	No benefit seen of ADSCs, or BMC. PRP shown to have less swelling post op, better function post op, better graft maturation
Hart 2005 [51]	Case-control study	31	Mean 10 years (9 to 13)	arthroscopic bone-patellar tendon-bone reconstruction without notch-plasty	Ahlback system	ACLR with partial meniscectomy resulted in greater degenerative changes than ACLR without meniscectomy. ($P < 0.05$)
Gerhard 2013 [52]	Retrospective cohort study	63	mean 16 ± 1 years	Long term ACLR follow up	Kellgren and Lawrence (KL) classification system	Those with a meniscal lesion at the time of ACL tear showed less favourable outcomes than those with isolated tears ($p < 0.05$)

APPENDIX 3 – Stated Odds Ratios of Risk Factors

