

This is a repository copy of *Stochastic Modified Flows for Riemannian Stochastic Gradient Descent*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/224353/>

Version: Accepted Version

Article:

Gess, Benjamin, Kassing, Sebastian and Rana, Nimit (2024) Stochastic Modified Flows for Riemannian Stochastic Gradient Descent. *SIAM Journal on Control and Optimization*. pp. 3288-3314. ISSN 1095-7138

<https://doi.org/10.48550/arXiv.2402.03467>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

STOCHASTIC MODIFIED FLOWS FOR RIEMANNIAN STOCHASTIC GRADIENT DESCENT

BENJAMIN GESS, SEBASTIAN KASSING, AND NIMIT RANA

ABSTRACT. We give quantitative estimates for the rate of convergence of Riemannian stochastic gradient descent (RSGD) to Riemannian gradient flow and to a diffusion process, the so-called Riemannian stochastic modified flow (RSMF). Using tools from stochastic differential geometry we show that, in the small learning rate regime, RSGD can be approximated by the solution to the RSMF driven by an infinite-dimensional Wiener process. The RSMF accounts for the random fluctuations of RSGD and, thereby, increases the order of approximation compared to the deterministic Riemannian gradient flow. The RSGD is built using the concept of a retraction map, that is, a cost efficient approximation of the exponential map, and we prove quantitative bounds for the weak error of the diffusion approximation under assumptions on the retraction map, the geometry of the manifold, and the random estimators of the gradient.

1. INTRODUCTION

Many optimization problems of the form

$$(1) \quad \text{Find } x^* \in \arg \min_{y \in M} f(y)$$

are posed on non-linear sets M , e.g. M being a subset of a Euclidean space due to non-linear constraints. For example, principal component analysis (PCA) leads to optimization problems on the Grassmann manifold or the Stiefel manifold, see Section 6.1. In supervised learning problems with positive homogeneous activation function, such as ReLU, one can restrict to networks with normalized weights, leading to optimization on the sphere [SK16, HSK⁺12, DK22b], see Section 6.2. Other examples include learning hierarchical representations [CCD17, NK17, WL18], e.g. in language models, where the optimization is often performed on hyperbolic space, see Section 6.3, and optimization over a parametrized family of probability measures, e.g. in the training of generative adversarial networks [GPAM⁺14, SZRM18, SWRH20], see Section 6.4.

In all of these examples, the search space M forms a Riemannian manifold. Therefore, popular optimization schemes for numerically solving (1), such as stochastic gradient descent (SGD), have been transferred to the Riemannian setting [Bon13, ZS16, ZRS16].

The analysis of the dynamics of SGD, including its algorithmic, implicit bias, and the empirically observed good generalization properties of artificial neural networks trained by SGD, is highly involved. Since there exists a large toolbox from optimal control theory and stochastic analysis for the investigation of continuous time processes that is difficult to apply in discrete time, one is led to the derivation of continuous time limits of SGD. In this work, we present continuum limits for Riemannian SGD on manifolds in the small learning rate regime and provide quantitative estimates on the rate of convergence.

2020 *Mathematics Subject Classification.* Primary 62L20; Secondary 58J65, 60J20, 65K05.

Key words and phrases. Riemannian stochastic gradient descent; diffusion approximation; supervised learning; weak error; Riemannian gradient flow.

In [Sha21, KHK22] the ODE method for Riemannian SGD is introduced proving that, as the learning rate tends to zero, the dynamics of SGD can be approximated by the solution to the Riemannian gradient flow ODE. As the first main result, we give the first quantified bounds on the weak error of this approximation, see Theorem 1.1.

Note that the deterministic Riemannian gradient flow describes the typical behavior of SGD in the small learning rate regime without taking into account the randomness of the gradient estimators. For the second main result, we introduce a class of stochastic differential equations (SDEs) on M , which we call Riemannian stochastic modified flow (RSMF), that capture both the mean behavior of the dynamical system as well as its random fluctuations. This carefully chosen limiting object is proved to capture the dynamics of SGD more precisely, giving a higher order approximation in the weak sense compared to the limiting ODE. In the Euclidean setting similar SDEs have been introduced in [LTE17, LTE19] with extensions to the overparametrized, infinite-particle limit given in [GKK24].

Let us introduce the central objects of the present work. See Appendix A for an overview of the notation used in this work. Let M be a complete, connected C^∞ -Riemannian manifold. With an eye on the applications detailed in Section 6, note that we do not assume M to be compact. Let $(\Xi, \mathcal{G}, \vartheta)$ be a probability space such that $L^2((\Xi, \vartheta); \mathbb{R})$ is separable and $\text{grad } \tilde{f} : M \times \Xi \rightarrow TM$ be a function that satisfies for all $x \in M$ that $\mathbb{E}_\vartheta[\|\text{grad } \tilde{f}(x, \xi)\|_x^2] < \infty$ and

$$\mathbb{E}_\vartheta[\text{grad } \tilde{f}(x, \xi)] = \text{grad } f(x),$$

where $\|\cdot\|_x$ denotes the norm on $T_x M$ given by the Riemannian metric. E.g., if ϑ is a rotation invariant probability measure on $\mathbb{R}^d$ with bounded second moment, where $d = \dim(M)$, the choice $\text{grad } \tilde{f}(x, \xi) = \text{grad } f(x) + \xi$ satisfies the conditions above (identifying $\mathbb{R}^d$ with $T_x M$). Formally, $\text{grad } \tilde{f}(\cdot, \xi)$ is not required to be a gradient vector field for every $\xi \in \Xi$. We use this notation to highlight the fact that $\text{grad } \tilde{f}$ is a random estimator of $\text{grad } f$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $(\xi_n)_{n \in \mathbb{N}}$ be an i.i.d. sequence of ϑ -distributed random variables on $(\Omega, \mathcal{F}, \mathbb{P})$ and set $(\mathcal{F}_n)_{n \in \mathbb{N}_0} = (\sigma(\xi_1, \dots, \xi_n))_{n \in \mathbb{N}_0}$. For $\eta > 0$ and $x \in M$ we consider an $(\mathcal{F}_n)_{n \in \mathbb{N}_0}$ -adapted, M -valued, process $(Z_n^\eta(x))_{n \in \mathbb{N}_0}$ satisfying $Z_0^\eta(x) = x$ and

$$(2) \quad Z_n^\eta(x) = \text{retr}_{Z_{n-1}^\eta(x)}(-\eta \text{grad } \tilde{f}(Z_{n-1}^\eta(x), \xi_n)), \quad \text{for all } n \in \mathbb{N},$$

where $\text{retr}_z : T_z M \rightarrow M$ denotes the Riemannian exponential map $\exp_z : T_z M \rightarrow M$ at $z \in M$ or a computationally efficient approximation of the exponential map, see Definition 2.1. We call $(Z_n^\eta(x))_{n \in \mathbb{N}_0}$ the *Riemannian SGD scheme* or *RSGD scheme* with *learning rate* η , started at x . Note that, since M is (geodesically) complete, $\exp_z$ is defined on the whole tangent space $T_z M$. Thus, the expression on the right-hand side of (2) is well-defined.

For small learning rates $\eta > 0$ the dynamics of $(Z_n^\eta(x))_{n \in \mathbb{N}_0}$ can be compared to continuous time processes using the numerical time-scale $t_n = n\eta$. Since, in expectation, SGD performs an Euler-step for the gradient flow ODE

$$(3) \quad \dot{z}_t = -\text{grad } f(z_t),$$

it seems natural to compare the dynamics of RSGD in the small learning rate regime with those of a solution $(z_t(x))_{t \geq 0}$ of (3) with initial condition $z_0(x) = x$. We quantify the quality of the approximation in the following theorem.

Theorem 1.1 (See Theorem 3.1). *Assume that M has bounded curvature, $\text{retr} : TM \rightarrow M$ is an appropriate approximation of $\exp$ and $\text{grad } \tilde{f} : M \times \Xi \rightarrow TM$ is sufficiently regular. Then, for all $T > 0$ and sufficiently regular test functions $g : M \rightarrow \mathbb{R}$ there exists a constant $C > 0$*

such that for all $\eta > 0$

$$\sup_{x \in M} \max_{n=0, \dots, \lfloor T/\eta \rfloor} |\mathbb{E}[g(Z_n^\eta(x))] - g(z_{n\eta}(x))| \leq C\eta.$$

In light of Theorem 1.1, we say that the solution to the gradient flow is a weak order 1 approximation of RSGD.

In order to get a continuous-in-time approximation of SGD that is of order $\mathcal{O}(\eta^2)$ one has to introduce a diffusion term that takes into account the random fluctuations of RSGD. We introduce a class of diffusion processes on a different probability space and compare their marginal distributions to those of RSGD. Let $(\tilde{\Omega}, (\tilde{\mathcal{F}}_t)_{t \geq 0}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ be a complete, filtered probability space with right-continuous filtration $(\tilde{\mathcal{F}}_t)_{t \geq 0}$. We consider the solution to the SDE taking values in M

$$(4) \quad dX_t^\eta(x) = B^\eta(X_t^\eta(x)) dt + G^\eta(X_t^\eta(x), \cdot) \circ dW_t$$

with initial condition $X_0^\eta(x) = x$, where $(W_t)_{t \geq 0}$ denotes a cylindrical Wiener process on $L^2((\Xi, \vartheta); \mathbb{R})$ defined on $(\tilde{\Omega}, (\tilde{\mathcal{F}}_t)_{t \geq 0}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, see Section 4. As coefficients we choose

$$(5) \quad G(x, \xi) := G^\eta(x, \xi) := \sqrt{\eta}(\text{grad } f(x) - \text{grad } \tilde{f}(x, \xi))$$

and

$$(6) \quad B(x) := B^\eta(x) := -\text{grad } f(x) - \frac{1}{2}\eta \left(\nabla_{\text{grad } f(x)}(\text{grad } f) + \int_{\Xi} \nabla_{\bar{G}(x, \xi)} \bar{G}(\cdot, \xi) \vartheta(d\xi) \right),$$

where ∇ denotes the Riemannian connection on M and $\bar{G} = \frac{1}{\sqrt{\eta}}G$. We call the solution to (4) *Riemannian stochastic modified flow* (RSMF).

Additionally to the diffusion term (5), we have to introduce two correction terms in (6) in order to get a weak order 2 approximation of RSGD: a bias correction term that compensates the second order term in the Euler discretization of the gradient flow and a term that accounts for using the Stratonovich formulation in (4), which is the more natural choice for defining SDEs on manifolds.

Let us state an informal version of the second main result of this article.

Theorem 1.2 (see Theorem 5.1). *Assume that M and $\text{grad } \tilde{f} : M \times \Xi \rightarrow TM$ are sufficiently regular and $\text{retr} : TM \rightarrow M$ is an appropriate approximation of $\exp$. Then, for all $T > 0$ and sufficiently regular test functions $g : M \rightarrow \mathbb{R}$ there exists a constant $C > 0$ such that for all $\eta > 0$*

$$\sup_{x \in M} \max_{n=0, \dots, \lfloor T/\eta \rfloor} |\mathbb{E}[g(Z_n^\eta(x))] - \tilde{\mathbb{E}}[g(X_{n\eta}^\eta(x))]| \leq C\eta^2,$$

where $(X_t^\eta(x))_{t \geq 0}$ denotes the unique solution to (4).

If M is compact then the assumptions on the geometry of M are satisfied. Moreover, if M is the Euclidean space one can choose $\text{retr}_x(v) = x + v$ for every $x \in M$ and $v \in T_x M \simeq \mathbb{R}^d$ and recover the results in [LTE19, Theorem 9] and [GKK24, Corollary 14] for the Stratonovich formulation of the Euclidean stochastic modified flow. However, Theorem 1.2 includes unbounded manifolds that satisfy uniform boundedness conditions on the geometry, see Definition 4.4. We verify the assumptions on the geometry of the manifold and define appropriate approximations of the exponential map for principle component analysis, see Section 6.1, and for weight normalization in artificial neural networks, see Section 6.2.

The proofs of Theorem 1.1 and Theorem 1.2 proceed by a precise analysis of the (Markov) semigroups corresponding to the ODE (3) and the SDE (4) and their flow maps $(x, t) \mapsto z_t(x)$

and $(x, t) \mapsto X_t^\eta(x)$. We show regularity results for the flow maps and give quantitative bounds on their derivatives with respect to the initial condition. These bounds depend on the regularity of the random vector field $\text{grad } \tilde{f}$ as well as on the curvature of M . Compared to the Euclidean case, the non-explosion of the solution to (4) does not follow from the Lipschitz-continuity of the coefficients alone. Therefore, on non-compact manifolds one has to be especially careful to ensure that the solution to (4) does not explode in finite time and that there exists a global flow map, see e.g. [Elw82, Li94a, Li94b].

Note that our proofs do not use the fact that $\text{grad } f$ is a gradient vector field and $\text{grad } f$ can be replaced by any sufficiently regular vector field. In that sense, Theorem 1.1 and Theorem 1.2 naturally extend to Riemannian stochastic approximation schemes for non-gradient vector fields, as well as accelerated optimization methods defined on the tangent bundle TM (see also Theorem 14 and 16 in [LTE19]).

The remainder of this article is organized as follows. In Section 1.1, we give an overview of the existing literature on Riemannian stochastic gradient descent and continuous time approximations for SGD in the Euclidean and Riemannian setting. In Section 2, we define uniform retraction maps, see Definition 2.1. In Section 3, we prove the first main result, Theorem 1.1, by analyzing the flow of a vector field as well as the dynamics of RSGD. In Section 4, we introduce SDEs on manifolds driven by a cylindrical Wiener process. We give an existence and uniqueness result for locally Lipschitz continuous coefficients. Moreover, under additional regularity assumptions on the manifold we prove the strong completeness of the SDE (4), i.e. the existence of a global flow map, and give bounds for the derivatives of the corresponding Markov semigroup. In Section 5, we prove the second main result, Theorem 1.2. Finally, in Section 6 we comment on the geometry in principle component analysis, weight normalization, hyperbolic space and statistical manifolds and give appropriate retraction maps in the respective optimization tasks. See Appendix A for the notation used throughout the article and Appendix B for an introduction into derivatives of higher order for scalar-valued functions and vector fields.

We refer the reader to [DC92] for a more detailed introduction into the general theory of Riemannian manifolds, to [Udr94, AMS08, Bou23] for an introduction into optimization on Riemannian manifolds and to [Elw82, Hsu02, IW89] for an introduction into SDEs on manifolds.

1.1. Overview of the literature. Diffusion approximations of Euclidean SGD in the small learning rates regime have been introduced by Li, Tai and E in [LTE17] and [LTE19]. Following these original papers several results were derived for SDE approximations of SGD, e.g. generator based proofs [FLL18, HLLL19], approximations for SGD without reshuffling [AP22] and uniform-in-time estimates for strongly convex objective functions [FGL⁺20, LW22]. Feng et al. presented a diffusion approximation for SGD performed on the sphere [FLL18]. In [GKK24], a Euclidean analog of the stochastic modified flow (4) has been proposed in order to approximate the multi-point motion of SGD. This work also presents an approximation for the dynamics of SGD in the small learning rate - infinite width scaling regime for overparametrized neural networks. A diffusion approximation result for SGD with time-dependent learning rate has been derived in [FDBD21]. In [SHSW23] the concept of second order retractions is used in order to approximate the Brownian motion on a Riemannian manifold in a cost-efficient way. This result generalizes the classical Donsker's theorem in the Riemannian setting, see [Jør75]. For a discussion on the validity of the diffusion approximation for finite (non-infinitesimal) learning rate see [LMA21].

The derivation of stochastic continuum limits of SGD has proven instrumental in the analysis of optimization dynamics in several regards. For example, one of the motivations for the diffusion

approximation of SGD is to simplify the derivation of optimal hyperparameter schedules, e.g. for the learning rate [LTE17] or the batch-size [ZLP⁺22, Per23], using optimal control theory. Regarding the asymptotic behavior of SGD, diffusion approximations can be used for finding a Lyapunov function [GK23, MTB23], developing dynamical systems arguments [FGJ20], investigating the critical noise decay rate for the convergence property [DK22a] and analyzing the implicit bias of SGD [Woj24, LWA21].

Regarding SGD on a Riemannian manifold, the convergence of the objective function $f : M \rightarrow \mathbb{R}$ and its corresponding Riemannian gradient under the classical Robbins-Monro conditions has been shown by Bonnabel [Bon13] using either the exponential map or a retraction map. This analysis has been refined for Hadamard manifolds [SI23] and Riemannian stochastic approximation schemes, where the practitioner is only able to simulate a biased estimator of the vector field [DJM⁺21]. In [Sha21] and [KHKM22], the ODE method for SGD with decreasing learning rates is transferred to the Riemannian setting. For SGD with constant learning rate $\eta > 0$, [DJMS21] considered the invariant measure of the Markov-chain $(Z_n^\eta(x))_{n \in \mathbb{N}_0}$ and its asymptotic behavior as $\eta \rightarrow 0$. Tripuraneni et al. [TFBJ18] introduced a version of the Ruppert-Polyak averaging technique for the Riemannian setting to improve convergence rates for the approximation of an isolated stable minimum. [CB19, SFF19, HKKM23] considered escaping saddle points for perturbed gradient descent methods. See also [ZS16, ZRS16] regarding stochastic optimization results for geodesically convex target functions.

2. UNIFORM RETRACTIONS

In this section, we present the assumptions on the mapping $\text{retr} : TM \rightarrow M$ used in the definition of the RSGD scheme, see (2). In many applications, the exponential map is difficult to compute and it is more cost efficient to work with an approximation, a so-called retraction map. We introduce the required assumptions on the retraction map in the following definition.

Definition 2.1. (i) A $\mathcal{C}^1$ -map $\text{retr} : TM \rightarrow M$ is called a retraction map if for all $x \in M$ the restriction $\text{retr}_x : T_x M \rightarrow M$ satisfies $\text{retr}_x(0) = x$ and $D_0 \text{retr}_x : T_0(T_x M) \simeq T_x M \rightarrow T_x M$ is the identity map. A $\mathcal{C}^2$ -retraction map is called uniform first order retraction if there exists a constant $C \geq 0$ such that for all $x \in M$, $t \geq 0$ and $v \in T_x M$ with $\|v\| = 1$ one has

$$\|\dot{\gamma}_t\| \vee \left\| \frac{\nabla}{dt} \dot{\gamma}_t \right\| \leq C,$$

where $(\gamma_s)_{s \geq 0} = (\text{retr}_x(sv))_{s \geq 0}$.

(ii) A $\mathcal{C}^2$ -retraction map is called second order retraction if for all $x \in M$ and $v \in T_x M$ one has

$$\left. \frac{\nabla}{dt} \right|_{t=0} \dot{\gamma}_t = 0,$$

where $(\gamma_s)_{s \geq 0} = (\text{retr}_x(sv))_{s \geq 0}$. Moreover, a $\mathcal{C}^3$ -second order retraction is called uniform second order retraction if there exists a constant $C \geq 0$ such that for all $x \in M$, $t \geq 0$ and $v \in T_x M$ with $\|v\| = 1$ one has

$$\|\dot{\gamma}_t\| \vee \left\| \frac{\nabla}{dt} \dot{\gamma}_t \right\| \vee \left\| \frac{\nabla^2}{dt^2} \dot{\gamma}_t \right\| \leq C.$$

Example 2.2. (i) The exponential map is a uniform second order retraction. In fact, for all $x \in M$, $t \geq 0$ and $v \in T_x M$ one has $\|\dot{\gamma}_t\| = \|v\|$ and $\frac{\nabla}{dt} \dot{\gamma}_t = 0$, where $(\gamma_t)_{t \geq 0} = (\exp_x(tv))_{t \geq 0}$.

- (ii) The stereographic projection is a uniform second order retraction for the unit sphere $\mathbb{S}^2 \subset \mathbb{R}^3$. For $p = (0, 0, -1)$ we define $\text{retr}_p : T_p M \simeq \mathbb{R}^2 \rightarrow \mathbb{S}^2$ via

$$\text{retr}_p(x, y) = \left(\frac{x}{1 + \frac{1}{4}x^2 + \frac{1}{4}y^2}, \frac{y}{1 + \frac{1}{4}x^2 + \frac{1}{4}y^2}, \frac{-1 + \frac{1}{4}x^2 + \frac{1}{4}y^2}{1 + \frac{1}{4}x^2 + \frac{1}{4}y^2} \right).$$

Then, $\text{retr}_p(0, 0) = p$ and

$$D_0 \text{retr}_p = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

To show that retr is a uniform second order retraction, by symmetry, it is sufficient to consider $\frac{\nabla}{dt} \gamma_t$, where $\gamma_t = \text{retr}_p(tv)$ and $v = (1, 0)$. We get

$$\frac{d^2}{dt^2} \gamma_t = \left(\frac{8t(t^2 - 12)}{(t^2 + 4)^3}, 0, \frac{64 - 48t^2}{(t^2 + 4)^3} \right),$$

so that $\frac{\nabla}{dt}|_{t=0} \dot{\gamma} = P_p(\frac{d^2}{dt^2}|_{t=0} \gamma) = 0$. Here, P_p denotes the orthogonal projection onto $T_p M \simeq \mathbb{R}^2 \times \{0\}$. Moreover, $\|\dot{\gamma}_t\|$, $\|\frac{\nabla}{dt} \dot{\gamma}_t\|$ and $\|\frac{\nabla^2}{dt^2} \dot{\gamma}_t\|$ are uniformly bounded.

In general, let $M \subset \mathbb{R}^N$ be a smooth submanifold and set

$$(7) \quad \text{retr}_x(v) = \text{proj}(x + v), \quad x \in M, v \in T_x M,$$

where $\text{proj} : \mathbb{R}^N \rightarrow M$ denotes a metric projection, i.e.

$$(8) \quad \text{proj}(z) \in \arg \min_{y \in M} d(y, z), \quad z \in \mathbb{R}^N.$$

By [LS21, Theorem 1], there exists an open set $U \supset \mathbb{R}^N$ containing M such that for every $z \in U$ the minimizer in (8) is unique and $\text{proj}|_U : U \rightarrow \mathbb{R}^N$ is C^∞ . Moreover, Theorem 4.9 in [AM12] shows that for every $x \in M$ and $v \in T_x M$ it holds that $D_0 \text{retr}_x v = v$ and $\frac{\nabla}{dt}|_{t=0} \dot{\gamma}_t = 0$, where $(\gamma_s)_{s \geq 0} = (\text{retr}_x(sv))_{s \geq 0}$. We apply a cutoff function to the mapping retr defined in (7) in order to construct a uniform second order retraction for compact submanifolds.

Lemma 2.3. *Assume that M is compact and let $K \subset \mathbb{R}^N$ be a compact set such that $M \subset K \subset U$, where U is as above. Let $c : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ be a smooth, bounded function such that $c(x, 0) = 0$, $D_v c(x, 0) = \text{id}_{\mathbb{R}^N}$ and $D_v^2 c(x, 0) = 0$ for all $x \in M$, $x + c(x, v) \in K$ for all $x \in M$ and $v \in T_x M \subset \mathbb{R}^N$, and $D_v^\alpha c$ is bounded on $M \times \mathbb{R}^N$ for all $\alpha = 1, 2, 3$. Then the function $\text{retr} : TM \rightarrow M$ given by*

$$\text{retr}_x(v) = \text{proj}(x + c(x, v)), \quad x \in M, v \in T_x M,$$

is a uniform second order retraction.

Proof. For all $x \in M$ we denote by $P_x \in \mathbb{R}^{N \times N}$ the matrix that corresponds to the orthogonal projection from $\mathbb{R}^N$ onto the tangent space $T_{\text{proj}(x)} M$ of M at $\text{proj}(x)$ which is a C^∞ -smooth function on U .

For $x \in M$ and $v \in T_x M$ with $\|v\| = 1$ let $(\gamma_t)_{t \geq 0} = (\text{proj}(x + c(x, tv)))_{t \geq 0}$. Then, for all $t \geq 0$

$$\dot{\gamma}_t = D \text{proj}(x + c(x, tv)) D_v c(x, tv) v,$$

where D_v denotes the Jacobi matrix w.r.t. the second argument in c . Hence, $\dot{\gamma}_0 = D \text{proj}(x) v$. Moreover,

$$\begin{aligned} \frac{\nabla}{dt} \dot{\gamma}_t &= P_{\gamma_t} \left((D^2 \text{proj}(x + c(x, tv))) (D_v c(x, tv) v, D_v c(x, tv) v) \right) \\ &\quad + P_{\gamma_t} \left(D \text{proj}(x + c(x, tv)) D_v^2 c(x, tv) (v, v) \right), \end{aligned}$$

so that $\frac{\nabla}{dt}\dot{\gamma}_0 = P_x((D^2 \text{proj}(x))(v, v))$. Here we used the fact that for a vector field V along γ one has $\frac{\nabla}{dt}V(t) = P_{\gamma_t}(\frac{d}{dt}V(t))$, where P_{γ_t} denotes the orthogonal projection onto $T_{\gamma_t}M \subset \mathbb{R}^N$ and, for all t , $V(t)$ is identified as an element of $\mathbb{R}^N$ (see e.g. Proposition 5.3.2 in [AMS08]). Thus, using [AM12, Theorem 4.9], retr is a second order retraction. Lastly, using the boundedness of DP on K as well as, for $\alpha = 1, 2, 3$, the boundedness of $D^\alpha \text{proj}$ on K and $D_v^\alpha c$ on $M \times \mathbb{R}^N$ we have that $\|\dot{\gamma}_t\|$, $\|\frac{\nabla}{dt}\dot{\gamma}_t\|$ and $\|\frac{\nabla^2}{dt^2}\dot{\gamma}_t\|$ are uniformly bounded in $x \in M$, $t \geq 0$ and $v \in T_x M$ with $\|v\| = 1$. $\square$

Remark 2.4. One can choose $c(x, v) = v$ for all $x \in M$ and $v \in T_x M$ that satisfy $x + v \in K'$, where $K' \subset \mathbb{R}^N$ is a compact set such that $M \subset K' \subset U' \subset K$ for an open set U' . Therefore, if for ϑ -almost all $\xi \in \Xi$ we have $x + \text{grad } \tilde{f}(x, \xi) \in K'$ for all $x \in M$, we can without loss of generality assume that the second order retraction (7) is a uniform second order retraction.

3. ORDER 1 APPROXIMATION

In this section, we quantify the weak approximation error for comparing the dynamics of Riemannian SGD with the solution to the gradient flow ODE

$$(9) \quad \dot{z}_t(x) = -\text{grad } f(z_t(x))$$

with initial condition $z_0(x) = x$. Recall that, for $x \in M$, $R(x) : (T_x M)^3 \rightarrow T_x M$ denotes the curvature of M at x given by

$$R(x)(u, v)w = (R(U, V)W)(x),$$

where $U, V, W \in \mathfrak{X}^\infty(M)$ with $U(x) = u$, $V(x) = v$ and $W(x) = w$, see Appendix A. Moreover,

$$\|R(x)\| = \sup_{v, w, u \in T_x M} \frac{\|R(x)(u, v)w\|}{\|u\| \|v\| \|w\|}.$$

We next state the main result of this section, proving that the ODE (9) is an order 1 approximation of RSGD if M has bounded curvature and $f : M \rightarrow \mathbb{R}$ is sufficiently regular.

Theorem 3.1. *Let $\text{retr} : TM \rightarrow M$ be a uniform first order retraction and $\text{grad } \tilde{f} : M \times \Xi \rightarrow TM$ be a function that satisfies for all $x \in M$*

$$\mathbb{E}_\vartheta[\text{grad } \tilde{f}(x, \xi)] = \text{grad } f(x) \quad \text{and} \quad \sup_{y \in M} \mathbb{E}_\vartheta[\|\text{grad } \tilde{f}(y, \xi)\|^2] < \infty.$$

Assume that $\text{grad } f \in \mathfrak{X}_b^2(M)$ and $\sup_{x \in M} \|R(x)\| < \infty$. Then for all $T \geq 0$ and $g \in \mathcal{C}_b^2(M)$ there exists a constant $C \geq 0$ such that for all $\eta > 0$

$$\sup_{x \in M} \sup_{n=0, \dots, \lfloor T/\eta \rfloor} |\mathbb{E}[g(Z_n^\eta(x))] - g(z_{n\eta}(x))| \leq C\eta.$$

For the proof of Theorem 3.1, we first show regularity results for the flow of a vector field, in Section 3.1, and, afterwards, compare a single iteration step of RSGD with running the ODE for time η , in Section 3.2.

3.1. Flow of a vector field. Let $V \in \mathfrak{X}_b^1(M)$ and consider the ODE

$$(10) \quad \dot{z}_t(x) = V(z_t(x))$$

with initial condition $z_0(x) = x$. Then, for every $x \in M$ there exists a unique solution $(z_t(x))_{t \in \mathbb{R}}$ of (10) that does not explode in finite time. Moreover the mapping $M \times \mathbb{R} \ni (x, t) \mapsto z_t(x) \in M$ is C^1 , see e.g. [DK00, Theorem B.3]. We give a quantitative bound on the first derivative with respect to the initial condition. This follows immediately from [DC92, Lemma 3.4] and Gronwall's inequality.

Lemma 3.2. *For $x \in M$ and $v \in T_x M$, $(D_x z_t(v))_{t \geq 0}$ satisfies the differential equation*

$$\frac{\nabla}{dt}(D_x z_t(v)) = \nabla_{D_x z_t(v)} V.$$

Moreover, for all $t \geq 0$ we have $\|D_x z_t(v)\| \leq \|v\| \exp(\|V(x)\|_{\mathfrak{X}_b^1(M)} t)$.

Next, we consider quantitative bounds on the second derivative of the flow. For this estimate we need a bound on the Riemannian curvature.

Lemma 3.3. *Assume that $V \in \mathfrak{X}_b^2(M)$ and $\sup_{x \in M} \|R(x)\| < \infty$. Then the mapping $(x, t) \mapsto z_t(x)$ is $\mathcal{C}^2$ and for all $T \geq 0$ there exists a constant $C \geq 0$ that only depends on T , $\|V\|_{\mathfrak{X}_b^2(M)}$ and $\sup_{x \in M} \|R(x)\|$ such that for all $0 \leq t \leq T$, $s \geq 0$, $x \in M$ and $v, w \in T_x M$ with $\|v\| = \|w\| = 1$ we have*

$$\left\| \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) \right\| \leq C,$$

where $(\gamma_s)_{s \in \mathbb{R}} = (\exp_x(sw))_{s \in \mathbb{R}}$ and $(v_s)_{s \in \mathbb{R}}$ is given by $v_s = \text{Par}_{\gamma|_{[0,s]}} v$ for $s \geq 0$ and $v_s = (\text{Par}_{\gamma|_{[s,0]}})^{-1} v$ for $s < 0$.

Proof. For the statement that $(x, t) \mapsto z_t(x)$ is $\mathcal{C}^2$ see [DK00, Theorem B.3]. Let $x \in M$, $v, w \in T_x M$ with $\|v\| = \|w\| = 1$, $(\gamma_s)_{s \in \mathbb{R}} = (\exp_x(sw))_{s \in \mathbb{R}}$ and $(v_s)_{s \in \mathbb{R}}$ be given by $v_s = \text{Par}_{\gamma|_{[0,s]}} v$ for $s \geq 0$ and $v_s = (\text{Par}_{\gamma|_{[s,0]}})^{-1} v$ for $s < 0$. Then $\mathbb{R}^2 \ni (s, t) \mapsto z_t(\gamma_s) \in M$ is a parametrized surface and $(s, t) \mapsto D_{\gamma_s} z_t(v_s)$ is a vector field along this surface. Using [DC92, Lemma 4.1] and Lemma 3.2, we get

$$\begin{aligned} \frac{\nabla}{dt} \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) &= \frac{\nabla}{ds} (\nabla_{D_{\gamma_s} z_t(v_s)} V) + R(z_t(\gamma_s))(D_{\gamma_s} z_t(\dot{\gamma}_s), V(z_t(\gamma_s))) D_{\gamma_s} z_t(v_s) \\ &= \nabla_{\frac{\nabla}{ds} D_{\gamma_s} z_t(v_s)} V + (\nabla^2 V)(D_{\gamma_s} z_t(\dot{\gamma}_s), D_{\gamma_s} z_t(v_s)) \\ &\quad + R(z_t(\gamma_s))(D_{\gamma_s} z_t(\dot{\gamma}_s), V(z_t(\gamma_s))) D_{\gamma_s} z_t(v_s), \end{aligned}$$

see also Appendix B. Now, $\|\frac{\nabla}{ds} D_{\gamma_s} z_0(v_s)\|^2 = \|\frac{\nabla}{ds} v_s\|^2 = 0$ and

$$\begin{aligned} \frac{d}{dt} \left\| \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) \right\|^2 &= 2 \left\langle \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s), \frac{\nabla}{dt} \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) \right\rangle \\ &\leq 2 \left\| \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) \right\| \|V\|_{\mathfrak{X}_b^2(M)} \left(\left\| \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) \right\| + (1 + \|R\|_\infty) \|D_{\gamma_s} z_t\|^2 \right) \\ &\leq 2 \left(\left\| \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) \right\|^2 + 1 \right) \|V\|_{\mathfrak{X}_b^2(M)} \left(1 + (1 + \|R\|_\infty) \|D_{\gamma_s} z_t\|^2 \right), \end{aligned}$$

where $\|R\|_\infty := \sup_{x \in M} \|R(x)\|$. Using Lemma 3.2 and Gronwall's inequality we get

$$\left\| \frac{\nabla}{ds} D_{\gamma_s} z_t(v_s) \right\|^2 \leq \left(\int_0^t \alpha_s ds \right) e^{\int_0^t \alpha_s ds},$$

where $\alpha_s = 2\|V\|_{\mathfrak{X}_b^2(M)} (1 + (1 + \|R\|_\infty) \exp(2\|V\|_{\mathfrak{X}_b^2(M)} t))$. $\square$

With Lemma 3.2 and Lemma 3.3 at hand, we analyze the mapping $x \mapsto g(z_t(x))$ for a $g \in \mathcal{C}_b^2$.

Proposition 3.4. *Assume that $\sup_{x \in M} \|R(x)\| < \infty$ and let $g \in \mathcal{C}_b^2(M)$ and $V \in \mathfrak{X}_b^2(M)$. Then for all $T \geq 0$ there exists a constant $C \geq 0$ that only depends on $\|g\|_{\mathcal{C}_b^2(M)}$, $\|V\|_{\mathfrak{X}_b^2(M)}$ and $\sup_{x \in M} \|R(x)\|$ such that for all $0 \leq t \leq T$ the function*

$$\varphi_t : M \rightarrow \mathbb{R} \quad ; \quad x \mapsto g(z_t(x))$$

satisfies $\|\varphi_t\|_{\mathcal{C}_b^2(M)} \leq C$.

Proof. For all $t \in T$ we clearly have $\varphi_t \in \mathcal{C}^2$ and $\|\varphi_t\|_{\mathcal{C}_b^0} \leq \|g\|_{\mathcal{C}_b^0}$. For $x \in M$ and $v \in T_x M$,

$$(11) \quad \langle \text{grad } \varphi_t(x), v \rangle = D_x \varphi_t(v) = (D_{z_t(x)} g)(D_x z_t(v)) = \langle \text{grad } g(z_t(x)), D_x z_t(v) \rangle$$

so that, using Lemma 3.2, we get

$$\|\text{grad } \varphi_t(x)\| \leq \sup_{x \in M} \|\text{grad } g(x)\| \sup_{x \in M} \|D_x z_t\| \leq \|g\|_{\mathcal{C}_b^1(M)} \exp(\|V\|_{\mathfrak{X}_b^1(M)} t).$$

By polarization, it suffices to derive a bound for $\langle \text{Hess } \varphi_t(x) v, v \rangle$ for all $x \in M$ and $v \in T_x M$ with $\|v\| = 1$. Write $(\gamma_s)_{s \in \mathbb{R}} = (\exp_x(sv))_{s \in \mathbb{R}}$ and note that $\dot{\gamma}_s = \text{Par}_{\gamma|_{[0,s]}} v$ for all $s \geq 0$. Using [AMS08, Proposition 5.5.4] and (11), we get

$$\begin{aligned} \langle \text{Hess } \varphi_t(x) v, v \rangle &= \frac{d^2}{ds^2} \big|_{s=0} \varphi_t(\exp_x(sv)) = \frac{d}{ds} \big|_{s=0} \langle \text{grad } g(z_t(\gamma_s)), D_{\gamma_s} z_t(\dot{\gamma}_s) \rangle \\ &= \langle \nabla_{D_x z_t(v)} \text{grad } g, D_x z_t(v) \rangle + \langle \text{grad } g(z_t(x)), \frac{\nabla}{ds} \big|_{s=0} D_{\gamma_s} z_t(\dot{\gamma}_s) \rangle, \end{aligned}$$

so that the statement follows from applying Lemma 3.2 and Lemma 3.3. $\square$

3.2. Proof of Theorem 3.1. Fix $T \geq 0$ and for $0 \leq t \leq T$ let $\varphi_t : M \rightarrow \mathbb{R}$ be given by $\varphi_t(x) = g(z_t(x))$. By Proposition 3.4, there exists a constant C such that $\sup_{0 \leq t \leq T} \|\varphi_t\|_{\mathcal{C}_b^2(M)} \leq C$.

For $x \in M$ and $n \in \mathbb{N}$ we get, using the triangle inequality and the Markov property for $(Z_n^\eta(x))_{n \in \mathbb{N}_0}$,

$$\begin{aligned} (12) \quad |\mathbb{E}[g(Z_n^\eta(x))] - g(z_{n\eta}(x))| &= \left| \sum_{i=1}^n \mathbb{E}[\varphi_{(n-i)\eta}(Z_i^\eta(x)) - \varphi_{(n-i)\eta}(z_\eta(Z_{i-1}^\eta(x)))] \right| \\ &\leq \sum_{i=1}^n |\mathbb{E}[\mathbb{E}[\varphi_{(n-i)\eta}(Z_i^\eta(x)) | \mathcal{F}_{i-1}] - \varphi_{(n-i)\eta}(z_\eta(Z_{i-1}^\eta(x)))]| \\ &\leq \sum_{i=1}^n \sup_{y \in M} |\mathbb{E}[\varphi_{(n-i)\eta}(Z_1^\eta(y))] - \varphi_{(n-i)\eta}(z_\eta(y))|. \end{aligned}$$

We next derive a uniform bound that holds for each individual summand in the latter sum. For $t \geq 0$ and $\varphi \in \mathcal{C}_b^2(M)$ we have

$$\begin{aligned} \varphi(z_t(x)) &= \varphi(x) - \int_0^t \langle \text{grad } \varphi(z_s(x)), \text{grad } f(z_s(x)) \rangle ds \\ &= \varphi(x) - t \langle \text{grad } \varphi(x), \text{grad } f(x) \rangle - \int_0^t \int_0^s \frac{d}{du} \langle \text{grad } \varphi(z_u(x)), \text{grad } f(z_u(x)) \rangle du ds. \end{aligned}$$

Using the fact that for all $u \geq 0$

$$\begin{aligned} \frac{d}{du} \langle \text{grad } \varphi(z_u(x)), \text{grad } f(z_u(x)) \rangle &= - \langle \text{Hess } \varphi(z_u(x)) \text{grad } f(z_u(x)), \text{grad } f(z_u(x)) \rangle \\ &\quad - \langle \text{grad } \varphi(z_u(x)), \nabla_{\text{grad } f(z_u(x))} \text{grad } f \rangle, \end{aligned}$$

as well as $\varphi \in \mathcal{C}_b^2(M)$ and $\text{grad } f \in \mathfrak{X}_b^2(M)$, there exists a constant $C \geq 0$ that only depends on $\|\varphi\|_{\mathcal{C}_b^2(M)}$ and $\|\text{grad } f\|_{\mathfrak{X}_b^1(M)}$ such that

$$(13) \quad \sup_{x \in M} |\varphi(z_t(x)) - \varphi(x) + t \langle \text{grad } \varphi(x), \text{grad } f(x) \rangle| \leq Ct^2.$$

Next, we turn to the SGD scheme defined in (2). For $x \in M$ and $\xi \in \Xi$ we let $(\gamma_t)_{t \in [0,1]} = (\text{retr}_x(-\eta t \text{grad } \tilde{f}(x, \xi)))_{t \in [0,1]}$ and note that

$$\varphi(\gamma_1) = \varphi(x) + \frac{d}{ds}|_{s=0} \varphi(\gamma_s) + \int_0^t \int_0^s \frac{d^2}{du^2} \varphi(\gamma_u) du ds.$$

Since $\text{retr} : TM \rightarrow M$ is a retraction, see Definition 2.1, we get

$$\frac{d}{ds}|_{s=0} \varphi(\gamma_s) = \langle \text{grad } \varphi(x), D_0 \text{retr}_x(-\eta \text{grad } \tilde{f}(x, \xi)) \rangle = -\eta \langle \text{grad } \varphi(x), \text{grad } \tilde{f}(x, \xi) \rangle.$$

For the remainder, we note that for all $u \geq 0$

$$\frac{d^2}{du^2} \varphi(\gamma_u) = \langle \text{Hess } \varphi(\gamma_u) \dot{\gamma}_u, \dot{\gamma}_u \rangle + \langle \text{grad } \varphi(\gamma_u), \frac{\nabla}{du} \dot{\gamma}_u \rangle.$$

Using that retr is a uniform first order retraction, there exists a constant $C \geq 0$ such that $\|\dot{\gamma}_u\| \leq C\eta \|\text{grad } \tilde{f}(x, \xi)\|$ and $\|\frac{\nabla}{du} \dot{\gamma}_u\| \leq C\eta^2 \|\text{grad } \tilde{f}(x, \xi)\|^2$ for all $u \geq 0$. Thus, there exists a constant $C \geq 0$ that only depends on φ via $\|\varphi\|_{\mathcal{C}_b^2(M)}$ such that

$$(14) \quad \sup_{x \in M} |\mathbb{E}[\varphi(Z_1^\eta(x))] - \varphi(x) + \eta \langle \text{grad } \varphi(x), \text{grad } f(x) \rangle| \leq C\eta^2 \left(\sup_{x \in M} \mathbb{E}[\|\text{grad } \tilde{f}(x, \xi_1)\|^2] \right).$$

Combining (13) and (14), there exists a constant $C \geq 0$ that only depends on $\|\varphi\|_{\mathcal{C}_b^2(M)}$, $\|\text{grad } f\|_{\mathfrak{X}_b^1(M)}$ and $\sup_{x \in M} \mathbb{E}_\vartheta[\|\text{grad } \tilde{f}(x, \xi)\|^2]$ such that

$$(15) \quad \sup_{x \in M} |\mathbb{E}_\vartheta[\varphi(Z_1^\eta(x))] - \varphi(z_\eta(x))| \leq C\eta^2.$$

Using that $\sup_{0 \leq t \leq T} \|\varphi_t\|_{\mathcal{C}_b^2(M)} < \infty$ and applying (15) to each summand on the right-hand side of (12), there exists a constant $C \geq 0$ such that for all $n \in \{0, \dots, \lfloor T/\eta \rfloor\}$

$$\sup_{x \in M} |\mathbb{E}[g(Z_n^\eta(x))] - g(z_{n\eta}(x))| \leq nC\eta^2 \leq CT\eta.$$

4. SDES ON MANIFOLDS DRIVEN BY CYLINDRICAL WIENER PROCESSES

In this section, we give well-posedness and regularity results for stochastic differential equations (SDEs) on a Riemannian manifold driven by a cylindrical Wiener process. For a more detailed introduction into stochastic analysis on Riemannian manifolds we refer the reader to [Elw82, IW89, Hsu02]. Regarding SDEs on Euclidean space driven by a cylindrical Wiener process we refer the reader to [DPZ14, LR15, Rie11].

Let $(\Xi, \mathcal{G}, \vartheta)$ be a measure space such that ϑ is a finite measure and the space $L^2((\Xi, \vartheta); \mathbb{R})$ is separable. We denote by $\langle \cdot, \cdot \rangle_\vartheta$, resp. $\|\cdot\|_\vartheta$, the usual inner product, resp. its associated norm, on the space $L^2((\Xi, \vartheta); V)$, where V denotes a finite-dimensional Hilbert space.

Let $(W_t)_{t \geq 0}$ be a cylindrical Wiener process on $L^2((\Xi, \vartheta); \mathbb{R})$ defined on a filtered, complete probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \geq 0}, \tilde{\mathbb{P}})$ with right-continuous filtration $(\tilde{\mathcal{F}}_t)_{t \geq 0}$, that is,

- (i) for every $t \geq 0$, the map $W_t : L^2((\Xi, \vartheta); \mathbb{R}) \rightarrow L^2((\tilde{\Omega}, \tilde{\mathbb{P}}); \mathbb{R})$ is linear;
- (ii) for every $h \in L^2((\Xi, \vartheta); \mathbb{R})$, $(W_t(h))_{t \geq 0}$ is an $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ -Brownian motion with $\text{Var}(W_t(h)) = \|h\|_\vartheta^2 t$ for every $t \geq 0$.

For an $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ -progressively measurable $L^2((\Xi, \vartheta); \mathbb{R})$ -valued process $(G(t, \cdot))_{t \geq 0}$ that almost surely satisfies $G \in L_{\text{loc}}^2([0, \infty); L^2((\Xi, \vartheta); \mathbb{R}))$ we define

$$\int_0^t \int_\Xi G(s, \xi) W(d\xi, ds) := \int_0^t \Upsilon(s) dW_s,$$

where $\Upsilon(s)$ is given by $\Upsilon(s)h = \langle G(s, \cdot), h \rangle_{\vartheta}$ for all $h \in L^2((\Xi, \vartheta); \mathbb{R})$. For the definition of the integral with respect to a cylindrical Wiener process see, e.g., [GM10, Section 2.2.4]. It is known that there exist an orthonormal basis of $L^2((\Xi, \vartheta); \mathbb{R})$, $(e_i)_{i \in \mathbb{N}}$, and independent $\mathbb{R}$ -valued $(\tilde{\mathcal{J}}_t)_{t \geq 0}$ -Brownian motions, $(W_t^{(1)})_{t \geq 0}, (W_t^{(2)})_{t \geq 0}, \dots$, such that

$$(16) \quad \int_0^t \int_{\Xi} G(s, \xi) W(d\xi, ds) = \sum_{i=1}^{\infty} \int_0^t \langle G(s, \cdot), e_i \rangle_{\vartheta} dW_s^{(i)},$$

where the integrals on the right-hand side of (16) are classical Itô-integrals with respect to $\mathbb{R}$ -valued Brownian motions and the sum is almost surely finite, see e.g. [DPZ14, Section 4.2.2]. Moreover,

$$\|\Upsilon(s)\|_{\text{HS}}^2 = \sum_{i=1}^{\infty} |\Upsilon(s)e_i|^2 = \sum_{i=1}^{\infty} |\langle G(s, \cdot), e_i \rangle_{\vartheta}|^2 = \|G(s, \cdot)\|_{\vartheta}^2,$$

where the left-hand side denotes the Hilbert-Schmidt norm of the operator $\Upsilon(s) : L^2((\Xi, \vartheta); \mathbb{R}) \rightarrow \mathbb{R}$.

Next, we introduce the notion of a solution to an SDE on a manifold driven by a cylindrical Wiener process. This generalizes the approach in [Hsu02] for SDEs on manifolds driven by a finite-dimensional Brownian motion. We need some additional notation.

Let $\tilde{L}^2(M \times \Xi; TM)$ be the space of all functions $G : M \times \Xi \rightarrow TM$ such that, for all $x \in M$ and $\xi \in \Xi$, $G(x, \xi) \in T_x M$ and

$$(17) \quad \|G(x, \cdot)\|_{\vartheta}^2 := \int_{\Xi} \|G(x, \xi')\|_x^2 \vartheta(d\xi') < \infty.$$

We denote by $\tilde{\mathfrak{X}}^0(M)$ the space of functions $G \in \tilde{L}^2(M \times \Xi; TM)$ such that for ϑ -a.e. $\xi \in \Xi$ we have $G(\cdot, \xi) \in \mathfrak{X}^0(M)$ and for all $x \in M$ there exists a neighborhood $U \subset M$ such that $x \in U$ and

$$(18) \quad \int_{\Xi} \sup_{y \in U} \|G(y, \xi)\|_y^2 \vartheta(d\xi) < \infty.$$

Note that (18) together with the dominated convergence theorem implies that for all $g \in \mathcal{C}^1(M)$ the mapping $x \mapsto Gg(x, \cdot) = \langle G(x, \cdot), \text{grad } g \rangle \in L^2((\Xi, \vartheta); \mathbb{R})$ is continuous w.r.t. the norm $\|\cdot\|_{\vartheta}$. Moreover, we denote by $\tilde{\mathfrak{X}}_b^0(M)$ the space of functions $G \in \tilde{\mathfrak{X}}^0(M)$ with

$$\|G\|_{\tilde{\mathfrak{X}}_b^0(M)} := \sup_{x \in M} \|G(x, \cdot)\|_{\vartheta}^2 < \infty.$$

Similarly, for $\alpha \in \mathbb{N}$ we denote by $\tilde{\mathfrak{X}}^{\alpha}(M)$ the space of functions $G \in \tilde{\mathfrak{X}}^0(M)$ such that for ϑ -a.e. $\xi \in \Xi$ we have $G(\cdot, \xi) \in \mathfrak{X}^{\alpha}(M)$ and for all $1 \leq \beta \leq \alpha$ and $x \in M$ there exists a neighborhood $U \subset M$ such that $x \in U$ and

$$\int_{\Xi} \sup_{y \in U} \|\nabla^{\beta} G(y, \xi)\|^2 \vartheta(d\xi) < \infty,$$

see Definition B.1. Again, the dominated convergence theorem implies that for all $1 \leq \beta \leq \alpha$, $V_1, \dots, V_{\beta} \in \mathfrak{X}^{\beta-1}(M)$ and $g \in \mathcal{C}^1(M)$ we have that

$$x \mapsto (\nabla^{\beta} G(x, \cdot))(V_1, \dots, V_{\beta})g \in L^2((\Xi, \vartheta); \mathbb{R})$$

is continuous w.r.t. the norm $\|\cdot\|_{\vartheta}$. We analogously define the space $\tilde{\mathfrak{X}}_b^{\alpha}(M)$ and its respective norm.

Let $B \in \mathfrak{X}^1(M)$ and $G \in \tilde{\mathfrak{X}}^2(M)$ and consider the following formal SDE

$$(19) \quad dX_t(x) = B(X_t(x)) dt + \int_{\Xi} G(X_t(x), \xi) \circ W(d\xi, ds)$$

with initial condition $X_0(x) = x \in M$. To make sense of this expression we consider the one-point compactification $\hat{M} = M \cup \{\partial_M\}$ of M and apply test functions. We interpret the second summand in (19) as the sum of an Itô-integral and a corresponding Stratonovich correction term.

Definition 4.1. (i) Let $(K_n)_{n \in \mathbb{N}}$ be an increasing sequence of compact subsets of M satisfying $M = \bigcup_{n \in \mathbb{N}} K_n$. Such a sequence exists since differentiable manifolds are, by definition, second countable. For a continuous, $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ -adapted process $(X_t)_{t \geq 0}$ taking values in $\hat{M}$ we define

$$e := \lim_{n \rightarrow \infty} \tau_n \quad \text{with} \quad \tau_n := \inf\{t \geq 0 : X_t \notin K_n\},$$

which is an $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ -stopping time and independent of the choice for $(K_n)_{n \in \mathbb{N}}$. We call e the explosion time of $(X_t)_{t \geq 0}$.

- (ii) A continuous $\hat{M}$ -valued, $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ -adapted process $(X_t(x))_{t \geq 0}$ with $X_0(x) = x$ and explosion time $e(x)$ is called solution to the SDE (19) started in x if for all $g \in C^\infty(M)$ one has almost surely that for all $t \in [0, e(x))$

$$(20) \quad \begin{aligned} g(X_t(x)) = & g(x) + \int_0^t Bg(X_s(x)) ds + \frac{1}{2} \int_0^t \int_{\Xi} GGg(X_s(x), \xi) \vartheta(d\xi) ds \\ & + \int_0^t \int_{\Xi} Gg(X_s(x), \xi) W(ds, d\xi). \end{aligned}$$

We next present an existence and uniqueness statement for the SDE (19). This result can be obtained as in the proof of Theorem 1.2.9 in [Hsu02], where SDEs on manifolds driven by a finite dimensional noise were considered. We take special attention to the required smoothness of the coefficients in (19).

Proposition 4.2. Let $B \in \mathfrak{X}^1(M)$ and $G \in \tilde{\mathfrak{X}}^2(M)$. Then for every $x \in M$ there exists a solution $(X_t(x))_{t \geq 0}$ to the SDE (19) with initial condition $X_0(x) = x$ and explosion time $e(x)$ that is unique up to its explosion time. Moreover, $(X_t(x))_{t \geq 0}$ satisfies (20) for all $g \in C^2(M)$.

In the second main result, Theorem 5.1, we will work with solutions to SDEs that do not explode in finite time and under the following assumption on the regularity of the corresponding Markov semigroup.

Definition 4.3. Let $\alpha \in \{2, 3, \dots\}$. We say that M has regularity α if for all $B \in \mathfrak{X}_b^\alpha(M)$ and $G \in \tilde{\mathfrak{X}}_b^{\alpha+1}(M)$ we have that

- (i) the SDE (19) is complete, i.e. for all $x \in M$ we have $e(x) = \infty$ almost surely, where $e(x)$ is the explosion time for the SDE (19) started in x , and
- (ii) for all $g \in C_b^\alpha(M)$ and $t \geq 0$ we have that $\Psi_t(x) := \mathbb{E}[g(X_t(x))]$ $\in C_b^\alpha(M)$. Moreover, for all $T \geq 0$ there exists a constant $C \geq 0$ that only depends on T , $\|B\|_{\mathfrak{X}_b^\alpha(M)}$, $\|G\|_{\tilde{\mathfrak{X}}_b^{\alpha+1}(M)}$ and $\|g\|_{C_b^\alpha(M)}$ such that

$$\sup_{t \in [0, T]} \|\Psi_t\|_{C_b^\alpha(M)} \leq C.$$

Definition 4.3 summarizes the necessary assumptions on the manifold in order to derive the order 2 approximation result in Theorem 5.1. Verifying the completeness for SDEs on Riemannian manifolds is more involved than simply using the Lipschitz-continuity of the coefficients as in the Euclidean case. It is known that the existence of a uniform cover [Elw82, Corollary 6.1] or a weak uniform cover [Li94a, Theorem 2.4] implies completeness of the corresponding SDE. In the next section, we introduce a large class of manifolds, including all compact manifolds, that satisfy the assumptions in Definition 4.3.

4.1. Embedding of M with uniform boundedness conditions on the geometry. In this section, we introduce a class of manifolds that admit an embedding into a Euclidean space with certain boundedness conditions on the metric projection. This concept allows us to extend vector fields on M to the ambient space and control the Euclidean derivatives of these extensions. The definition is inspired by the notion of manifolds of bounded geometry, see [Eld13, Definition 2.1]. Here, we additionally assume the existence of a uniform tubular neighborhood of the normal bundle.

Let $N \in \mathbb{N}$ and $\iota : M \rightarrow \mathbb{R}^N$ be an isometric embedding of M into $\mathbb{R}^N$. By [LS21, Theorem 1], there exists an open set $U \supset \iota(M)$ containing $\iota(M)$ such that for every $x \in U$ there exists a unique minimizer

$$(21) \quad \text{proj}(x) := \arg \min_{y \in \iota(M)} d(x, y)$$

that satisfies $x - \text{proj}(x) \in (T_{\text{proj}(x)}\iota(M))^\perp$ and $\text{proj}|_U : U \rightarrow \mathbb{R}^N$ is C^∞ . Clearly, we have $\text{proj}(x) = x$ for all $x \in \iota(M)$. Moreover, for $x \in U$ let $P_x \in \mathbb{R}^{N \times N}$ be the matrix that corresponds to the orthogonal projection from $\mathbb{R}^N$ onto the tangent space $T_{\text{proj}(x)}\iota(M)$ of $\iota(M)$ at $\text{proj}(x)$.

Definition 4.4. Let $\alpha \in \mathbb{N}_0$. We say that a complete and connected C^∞ -manifold M is a $\text{BG}(\alpha)$ -manifold if there exist $N \in \mathbb{N}$, $r > 0$, an isometric embedding $\iota : M \rightarrow \mathbb{R}^N$ and an open set $U \subset \mathbb{R}^N$ such that $x+v \in U$ for all $x \in \iota(M)$ and $v \in (T_x\iota(M))^\perp$ with $|v| \leq r$ and $\text{proj} : U \rightarrow \mathbb{R}^N$ and $P : U \rightarrow \mathbb{R}^{N \times N}$ exist and their derivatives up to order α exist and are uniformly bounded.

In the Euclidean setting, i.e. $M = \mathbb{R}^N$, one can set $\iota = \text{id}$ and note that $(T_x\iota(M))^\perp = \emptyset$. Therefore, the Euclidean space is a $\text{BG}(\alpha)$ -manifold for all $\alpha \in \mathbb{N}_0$. Moreover, every compact C^∞ -manifold is a $\text{BG}(\alpha)$ -manifold for all $\alpha \in \mathbb{N}_0$, see e.g. [LS21].

The motivation for introducing $\text{BG}(\alpha)$ -manifolds is that we can extend functions on M to functions defined on the ambient space $\mathbb{R}^N$ such that uniform boundedness of the Riemannian derivatives is equivalent to uniform boundedness of the Euclidean derivatives for the extended functions. For the definition of the spaces $\mathcal{C}_b^\alpha(N)$ and $\mathfrak{X}_b^\alpha(N)$ for a manifold N and $\alpha \in \mathbb{N}_0$ as well as the respective norms $\|\cdot\|_{\mathcal{C}_b^\alpha(N)}$ and $\|\cdot\|_{\mathfrak{X}_b^\alpha(N)}$, see Definition B.1.

Lemma 4.5. Let $\alpha \in \mathbb{N}_0$ and M be a $\text{BG}(\alpha)$ -manifold with corresponding embedding $\iota : M \rightarrow \mathbb{R}^N$, open set U and constant $r > 0$. Then

- (i) there exists a constant $C_1 \geq 0$ such that for every function $g \in \mathcal{C}_b^{\alpha+1}(U)$ one has

$$\|g|_{\iota(M)}\|_{\mathcal{C}_b^{\alpha+1}(\iota(M))} \leq C_1 \|g\|_{\mathcal{C}_b^{\alpha+1}(U)},$$

- (ii) there exists a constant $C_2 \geq 0$ such that for every $g \in \mathcal{C}_b^\alpha(\iota(M))$ there exists an extension $\hat{g} \in \mathcal{C}_b^\alpha(U)$ of g with

$$\|\hat{g}\|_{\mathcal{C}_b^\alpha(U)} \leq C_2 \|g\|_{\mathcal{C}_b^\alpha(\iota(M))}$$

and

- (iii) *there exists a constant $C_3 \geq 0$ such that for every $V \in \mathfrak{X}_b^\alpha(\iota(M))$ there exists an extension $\hat{V} \in \mathfrak{X}_b^\alpha(\mathbb{R}^N)$ of V with $\hat{V}(x) = 0$ for all $x \in \mathbb{R}^N$ with $d(x, \iota(M)) \geq r/2$ and*

$$\|\hat{V}\|_{\mathfrak{X}_b^\alpha(\mathbb{R}^N)} \leq C_3 \|V\|_{\mathfrak{X}_b^\alpha(\iota(M))}.$$

Proof. (i): Denote $\tilde{g} = g|_{\iota(M)}$. Then $\text{grad } \tilde{g}(x) = P_x \text{grad } g(x)$ and, if $\alpha \geq 1$, for $v \in T_x \iota(M)$ we have

$$\begin{aligned} \text{Hess } \tilde{g}(x)v &= \nabla_v \text{grad } \tilde{g} = P_x \left(\frac{d}{dt} \Big|_{t=0} P_{\gamma_t} \text{grad } g(\gamma_t) \right) \\ &= P_x \left(\text{Hess } g(x)v + (DP_x v) \text{grad } g(x) \right). \end{aligned}$$

where $\gamma : \mathbb{R} \rightarrow \iota(M)$ is a smooth curve satisfying $\gamma_0 = x$ and $\dot{\gamma}_0 = v$. If $\alpha \geq 2$, we get for vector fields $V_1, V_2, V_3 \in \mathfrak{X}^2(\iota(M))$ that

$$\nabla^3 \tilde{g}(V_1, V_2, V_3) = V_1((\nabla^2 \tilde{g})(V_2, V_3)) - (\nabla^2 \tilde{g})(\nabla_{V_1} V_2, V_3) - (\nabla^2 \tilde{g})(V_2, \nabla_{V_1} V_3)$$

as well as $V_1((\nabla^2 \tilde{g})(V_2, V_3)) = \langle \nabla_{V_1}(\text{Hess } \tilde{g} V_2), V_3 \rangle + (\nabla^2 \tilde{g})(V_2, \nabla_{V_1} V_3)$. Now, for all $x \in \iota(M)$ and $v \in T_x \iota(M)$ we have $P_x \text{Hess } g(x)v = P_x \text{Hess } g(x)P_x v$ so that

$$\begin{aligned} \nabla_{V_1(x)}(\text{Hess } \tilde{g} V_2) &= P_x(DP_x(V_1(x)))\text{Hess } g(x)V_2(x) + P_x D^3 g(x)(V_1(x), V_2(x)) \\ &\quad + P_x \text{Hess } g(x)(DP_x(V_1(x)))V_2(x) + P_x \text{Hess } g(x)P_x(DV_2(x)(V_1(x))) \\ &\quad + P_x(DP_x(V_1(x)))(DP_x(V_2(x)))\text{grad } g(x) + P_x D^2 P_x(V_1(x), V_2(x))\text{grad } g(x) \\ &\quad + P_x(DP_x(DV_2(x)(V_1(x))))\text{grad } g(x) + P_x(DP_x(V_2(x)))\text{Hess } g(x)V_1(x). \end{aligned}$$

Using that

$$\begin{aligned} (\nabla^2 \tilde{g})(\nabla_{V_1} V_2, V_3)(x) &= \langle P_x \text{Hess } g(x)P_x(DV_2(x)(V_1(x))) \\ &\quad + P_x(DP_x(P_x(DV_2(x)(V_1(x))))\text{grad } g(x), V_3(x) \rangle \end{aligned}$$

and $DP_x v = 0$ for all $v \in (T_x \iota(M))^\perp$, we get

$$\begin{aligned} \nabla^3 \tilde{g}(V_1, V_2, V_3)(x) &= \langle P_x(DP_x(V_1(x)))\text{Hess } g(x)V_2(x) + P_x D^3 g(x)(V_1(x), V_2(x)) \\ &\quad + P_x \text{Hess } g(x)(DP_x(V_1(x)))V_2(x) + P_x(DP_x(V_1(x)))(DP_x(V_2(x)))\text{grad } g(x) \\ &\quad + P_x D^2 P_x(V_1(x), V_2(x))\text{grad } g(x) + P_x(DP_x(V_2(x)))\text{Hess } g(x)V_1(x), V_3(x) \rangle. \end{aligned}$$

Using $\sup_{x \in \iota(M)} \max_{\beta=0, \dots, \alpha} \|D^\beta P_x\| < \infty$, we proved (i) for $\alpha \in \{0, 1, 2\}$. The proof for higher derivatives is analogous.

(ii): For $x \in U$ define $\hat{g}(x) := g(\text{proj}(x))$. Then $\hat{g} \in \mathcal{C}^\alpha(U)$ and $\|\hat{g}(x)\|_{\mathcal{C}_b^0(U)} = \|g(x)\|_{\mathcal{C}_b^0(\iota(M))}$. If $\alpha \geq 1$, we have for $x \in U$ and $i = 1, \dots, N$

$$\frac{d}{dx_i} \hat{g}(x) = \langle \text{grad } g(\text{proj}(x)), D \text{proj}(x) e_i \rangle,$$

where $(e_1, \dots, e_N)$ denotes the standard orthonormal basis of $\mathbb{R}^N$. If $\alpha \geq 2$, note that for all $x \in \iota(M)$ we have $\text{grad } \hat{g}(x) = \text{grad } g(x) \in T_x \iota(M)$. Thus, for all $i, j = 1, \dots, N$ and $x \in U$

$$\begin{aligned} \frac{d}{dx_j} \frac{d}{dx_i} \hat{g}(x) &= \frac{d}{dx_j} \langle \text{grad } \hat{g}(\text{proj}(x)), D \text{proj}(x) e_i \rangle \\ &= \langle P_x \frac{d}{dx_j} \text{grad } \hat{g}(\text{proj}(x)), D \text{proj}(x) e_i \rangle + \langle \text{grad } \hat{g}(\text{proj}(x)), \frac{d}{dx_j} D \text{proj}(x) e_i \rangle \\ &= \langle \nabla_v \text{grad } g, D \text{proj}(x) e_i \rangle + \langle \text{grad } g(\text{proj}(x)), D^2 \text{proj}(x)(e_i, e_j) \rangle, \end{aligned}$$

where $v := D \text{proj}(x)e_j \in T_{\text{proj}(x)}\iota(M)$. Using $\sup_{x \in \iota(M)} \max_{\beta=0,\dots,\alpha} \|D^\beta \text{proj}(x)\| < \infty$, we proved (i) for $\alpha \in \{0, 1, 2\}$. The proof for higher derivatives is analogous.

(iii): Let $\psi : [0, \infty) \rightarrow [0, 1]$ be a C^∞ -cutoff function that satisfies $\psi(0) = 1$, $\psi(y) = 0$ for all $y \geq r^2/4$ and such that all derivative of ψ at 0 and $r^2/4$ vanish. For $x \in \mathbb{R}^N$ define

$$\hat{V}(x) := \begin{cases} \psi(|\text{proj}(x) - x|^2)V(\text{proj}(x)), & \text{if } x \in U \\ 0, & \text{otherwise.} \end{cases}$$

Then, $\hat{V} \in \mathfrak{X}^\alpha(U)$, $\|\hat{V}(x)\|_{\mathfrak{X}_b^0(M)} = \|V(x)\|_{\mathfrak{X}_b^0(\iota(M))}$ and $\hat{V} = 0$ for all $x \in \mathbb{R}^N$ with $d(x, \iota(M)) \geq r/2$. If $\alpha = 1$, we use $v = P_x v$ for $x \in \iota(M)$ and $v \in T_x \iota(M)$ and get for $i = 1, \dots, N$ and $x \in U$

$$\begin{aligned} \frac{d}{dx_i} \hat{V}(x) &= 2\psi'(|\text{proj}(x) - x|^2)(\text{proj}(x) - x)^\dagger \left(\frac{d}{dx_i} \text{proj}(x) - e_i \right) V(\text{proj}(x)) \\ &\quad + \psi(|\text{proj}(x) - x|^2) \left(\frac{d}{dx_i} P_x \right) V(\text{proj}(x)) \\ &\quad + \psi(|\text{proj}(x) - x|^2) P_x (DV(\text{proj}(x))) (D \text{proj}(x) e_i), \end{aligned}$$

where $a^\dagger$ denotes the transpose of a vector $a \in \mathbb{R}^N$ and $(e_1, \dots, e_N)$ denotes the standard orthonormal basis of $\mathbb{R}^N$. Using $P_x(DV(\text{proj}(x)))(D \text{proj}(x) e_i) = \nabla_{D \text{proj}(x) e_i} V$ and the bounds for DP_x and $D \text{proj}(x)$ we get the statement for $\alpha = 1$. The proof for higher derivatives is analogous. $\square$

Using the extensions of the vector fields and the test functions constructed in Lemma 4.5, the assumptions in Definition 4.3 can be verified for M replaced by $\mathbb{R}^N$. A completeness result for the SDE (19) on $\mathbb{R}^N$ with Lipschitz continuous coefficients can be found in [DPZ14, Theorem 7.5]. Boundedness of the first two derivatives of $x \mapsto \tilde{E}[g(X_t(x))]$ can be found in [DPZ14, Theorem 9.23] and [DPZ14, Remark 9.4].

Proposition 4.6. *Let $\alpha \in \{2, 3, \dots\}$ and M be a $\text{BG}(\alpha + 1)$ -manifold. Then M has regularity α .*

5. ORDER 2 APPROXIMATION

In this section, we prove the second main result which quantifies the order of approximation of RSGD by the solution to the RSMF (4) with coefficients G and B as in (5) and (6), respectively.

Theorem 5.1. *Let $\text{grad } \tilde{f} \in \tilde{\mathfrak{X}}_b^5(M)$, $\text{retr} : TM \rightarrow M$ be a uniform second order retraction and assume that M has regularity 4, see Definition 4.3. Moreover, assume that for all $x \in M$ one has $\mathbb{E}_\vartheta[\text{grad } \tilde{f}(x, \xi)] = \text{grad } f(x)$ and*

$$(22) \quad \mathfrak{f} := \sup_{x \in M} \int_{\Xi} \|\text{grad } \tilde{f}(x, \xi)\|^3 \vartheta(d\xi) < \infty.$$

Then, for every $g \in \mathcal{C}_b^4(M)$ and $T \geq 0$, there exists a constant $C \geq 0$ such that for all $\eta \geq 0$

$$\sup_{x \in M} \sup_{n=0, \dots, \lfloor T/\eta \rfloor} |\mathbb{E}[g(Z_n^\eta(x))] - \tilde{\mathbb{E}}[g(X_{n\eta}^\eta(x))]| \leq C\eta^2.$$

The proof consists in comparing an evolution step of RSGD with the solution of the RSMF for time η on test functions of the form Ψ_t , for $t \geq 0$, defined in Definition 4.3. First, we show that the coefficients of the RSMF are sufficiently regular.

Lemma 5.2. *Let $T > 0$, $\alpha \in \mathbb{N}$, $\text{grad } \tilde{f} \in \tilde{\mathfrak{X}}_b^\alpha(M)$ and B^η, G^η as in (5) and (6). Then, for all $0 \leq \eta \leq T$ it holds that $B^\eta \in \mathfrak{X}_b^{\alpha-1}(M)$ and $G^\eta \in \tilde{\mathfrak{X}}_b^\alpha(M)$ and*

$$\sup_{0 \leq \eta \leq T} \|B^\eta\|_{\mathfrak{X}_b^{\alpha-1}(M)} \vee \sup_{0 \leq \eta \leq T} \|G^\eta\|_{\tilde{\mathfrak{X}}_b^\alpha(M)} < \infty.$$

Proof. First, we will show that $\text{grad } f \in \mathfrak{X}_b^\alpha(M)$. Let $\iota : M \rightarrow \mathbb{R}^N$ be an isometric embedding and $\hat{f} \in \tilde{\mathfrak{X}}^\alpha(\mathbb{R}^N)$ be an extension of $\text{grad } \tilde{f}$. Then, for all compact sets $K \subset \mathbb{R}^N$ and $0 \leq \beta \leq \alpha$ we have that

$$\int_{\Xi} \sup_{y \in K} \|D^\beta \hat{f}(y, \xi)\|^2 \vartheta(d\xi) < \infty.$$

Using the dominated convergence theorem, we get $\mathbb{E}_\vartheta[\hat{f}(\cdot, \xi)] \in \mathfrak{X}^\alpha(\mathbb{R}^N)$ and, thus, $\text{grad } f \in \mathfrak{X}^\alpha(M)$. Moreover, for all $x \in M$ and $0 \leq \beta \leq \alpha$,

$$(23) \quad \|\nabla^\beta \text{grad } f(x)\|^2 = \|\mathbb{E}_\vartheta[\nabla^\beta \text{grad } \tilde{f}(x, \xi)]\|^2 \leq \|\nabla^\beta \text{grad } \tilde{f}(x, \cdot)\|_\vartheta^2$$

where the right-hand side of (23) is uniformly bounded in x , since $\text{grad } \tilde{f} \in \tilde{\mathfrak{X}}_b^\alpha(M)$. Thus, $\text{grad } f \in \mathfrak{X}_b^\alpha(M)$ which immediately implies that $\nabla_{\text{grad } f(\cdot)}(\text{grad } f) \in \mathfrak{X}_b^{\alpha-1}(M)$ and $G^\eta \in \tilde{\mathfrak{X}}_b^\alpha(M)$. Similarly, we have for an extension $\hat{G}^\eta \in \tilde{\mathfrak{X}}^\alpha(\mathbb{R}^N)$ of G^η , $0 \leq \beta \leq \alpha - 1$ and all compact sets $K \subset \mathbb{R}^N$ that

$$\int_{\Xi} \sup_{y \in K} \|D^\beta (D\hat{G}^\eta(y, \xi)\hat{G}^\eta(y, \xi))\| \vartheta(d\xi) < \infty.$$

Moreover, $z \mapsto \mathbb{E}_\vartheta[\|\nabla^\beta (\nabla_{G^\eta(z, \xi)} G^\eta(\cdot, \xi))\|]$ is uniformly bounded over $z \in M$ and $0 \leq \eta \leq T$ for all $0 \leq \beta \leq \alpha - 1$, so that $z \mapsto \mathbb{E}_\vartheta[\nabla_{G^\eta(z, \xi)} G^\eta(\cdot, \xi)] \in \mathfrak{X}_b^{\alpha-1}(M)$. $\square$

We next control the weak error of the diffusion approximation for one evolution step of RSGD.

Lemma 5.3. *Let retr be a uniform second order retraction, $\text{grad } \tilde{f} \in \tilde{\mathfrak{X}}_b^5(M)$ and $g \in \mathcal{C}_b^4(M)$ and $T \geq 0$. Moreover, assume that M has regularity 4 and (22) holds. Then, there exists a constant $C \geq 0$ that only depends on T , $\|g\|_{\mathcal{C}_b^4(M)}$, $\|\text{grad } \tilde{f}\|_{\tilde{\mathfrak{X}}_b^5(M)}$ and $\mathfrak{f}$ such that for all $0 \leq \eta \leq T$*

$$\sup_{x \in M} |\mathbb{E}_\vartheta[g(Z_1^\eta(x))] - \tilde{\mathbb{E}}[g(X_\eta^\eta(x))]| \leq C\eta^3.$$

Proof. Throughout this proof, C denotes a constant that only depends on T , $\|g\|_{\mathcal{C}_b^4(M)}$, $\|\text{grad } \tilde{f}\|_{\tilde{\mathfrak{X}}_b^5(M)}$ and $\mathfrak{f}$. Note that, using Lemma 5.2, the norms $\|B^\eta\|_{\mathfrak{X}_b^4(M)}$ and $\|G^\eta\|_{\tilde{\mathfrak{X}}_b^5(M)}$ can be uniformly bounded over $0 \leq \eta \leq T$.

Let $x \in M$ and $0 \leq \eta \leq T$. First, we consider the solution to the RSMF (4) and briefly write $(X_t)_{0 \leq t \leq \eta} := (X_t^\eta(x))_{0 \leq t \leq \eta}$. By definition, we have for all $0 \leq t \leq \eta$

$$(24) \quad \begin{aligned} g(X_t) - g(x) &= \int_0^t Bg(X_s) ds + \sqrt{\eta} \int_0^t \int_{\Xi} \bar{G}g(X_s, \xi) W(ds, d\xi) \\ &\quad + \frac{\eta}{2} \int_0^t \int_{\Xi} \bar{G}\bar{G}g(X_s, \xi) \vartheta(d\xi) ds. \end{aligned}$$

Now, for all $0 \leq t \leq \eta$ we get by Itô's isometry that

$$\left[\int_0^\cdot \int_{\Xi} \bar{G}g(X_s, \xi) W(ds, d\xi) \right]_t = \int_0^t \|\bar{G}g(X_s, \cdot)\|_\vartheta^2 ds,$$

so that

$$(M_t)_{0 \leq t \leq \eta} := \left(\int_0^t \int_{\Xi} \bar{G}g(X_s, \xi) W(ds, d\xi) \right)_{0 \leq t \leq \eta}$$

is a martingale and we have $\tilde{\mathbb{E}}[M_\eta] = 0$. Moreover, for the first term on the right-hand side of (24) we get, using Itô's formula,

$$\begin{aligned} Bg(X_s) &= Bg(x) + \int_0^s BBg(X_u) du + \int_0^s \int_{\Xi} \sqrt{\eta} \bar{G}Bg(X_u, \xi) W(du, d\xi) \\ &\quad + \frac{\eta}{2} \int_0^s \int_{\Xi} \bar{G}\bar{G}Bg(X_u, \xi) \vartheta(d\xi) du. \end{aligned}$$

Using Fubini's theorem, we get

$$\begin{aligned} \tilde{\mathbb{E}} \left[\int_0^\eta \int_0^s \int_{\Xi} \sqrt{\eta} \bar{G}Bg(X_u, \xi) W(du, d\xi) ds \right] \\ = \int_0^\eta \tilde{\mathbb{E}} \left[\int_0^s \int_{\Xi} \sqrt{\eta} \bar{G}Bg(X_u, \xi) W(du, d\xi) \right] ds = 0. \end{aligned}$$

We can bound

$$\tilde{\mathbb{E}} \left[\left| \int_0^\eta \frac{\eta}{2} \int_0^s \int_{\Xi} \bar{G}\bar{G}Bg(X_u, \xi) \vartheta(d\xi) du \right| \right] \leq \frac{\eta^3}{4} \sup_{x \in M} \mathbb{E}_\vartheta[|\bar{G}\bar{G}Bg(x, \xi)|] \leq C\eta^3,$$

for a constant $C \geq 0$. Again, using Itô's formula

$$\begin{aligned} BBg(X_s) &= BBg(x) + \int_0^s BBBg(X_u) du + \int_0^s \int_{\Xi} \sqrt{\eta} \bar{G}BBg(X_u, \xi) W(du, d\xi) \\ &\quad + \frac{\eta}{2} \int_0^s \int_{\Xi} \bar{G}\bar{G}BBg(X_u, \xi) \vartheta(d\xi) du, \end{aligned}$$

where for all $s \in [0, \eta]$ one has $\tilde{\mathbb{E}} \left[\int_0^s \int_{\Xi} \sqrt{\eta} \bar{G}BBg(X_u, \xi) W(du, d\xi) \right] = 0$.

We bound the contribution of the last term on the right-hand side of (24). An application of the dominated convergence theorem gives $\varphi(x) = \int_{\Xi} \bar{G}\bar{G}g(x, \xi) \vartheta(d\xi) \in \mathcal{C}_b^2(M)$. Thus,

$$\begin{aligned} \varphi(X_s) - \varphi(x) &= \int_0^t B\varphi(X_s) ds + \sqrt{\eta} \int_0^t \int_{\Xi} \bar{G}\varphi(X_s, \xi) W(ds, d\xi) \\ &\quad + \frac{\eta}{2} \int_0^t \int_{\Xi} \bar{G}\bar{G}\varphi(X_s, \xi) \vartheta(d\xi) ds, \end{aligned}$$

where, for a constant $C \geq 0$, for all $0 \leq u \leq \eta$ one has $\tilde{\mathbb{E}} \left[\int_0^u \int_{\Xi} \bar{G}\varphi(X_s, \xi) W(ds, d\xi) \right] = 0$.

Altogether, we can bound the terms of order $\mathcal{O}(\eta^3)$ in the equations above and get a constant $C \geq 0$ such that

$$\sup_{x \in M} \left| \tilde{\mathbb{E}}[g(X_\eta^\eta(x))] - g(x) - \eta Bg(x) - \frac{1}{2} \eta^2 \left(BBg(x) + \int_{\Xi} \bar{G}\bar{G}g(x, \xi) \vartheta(d\xi) \right) \right| \leq C\eta^3.$$

Recall that, for all $x \in M$ and $\xi \in \Xi$, $BBg(x) = \langle (\text{Hess } g(x))B(x), B(x) \rangle + (\nabla_{B(x)} B)g$ and $\bar{G}\bar{G}g(x, \xi) = \langle \text{Hess } g(x)\bar{G}(x, \xi), \bar{G}(x, \xi) \rangle + (\nabla_{\bar{G}(x, \xi)} \bar{G}(\cdot, \xi))g$. Hence, using the definition of B in (6) there exists a constant $C \geq 0$ such that

$$\begin{aligned} (25) \quad & \left| \tilde{\mathbb{E}}[g(X_\eta)] - g(x) + \eta \langle \text{grad } f(x), \text{grad } g(x) \rangle - \frac{1}{2} \eta^2 \left(\langle (\text{Hess } g(x)) \text{grad } f(x), \text{grad } f(x) \rangle \right. \right. \\ & \quad \left. \left. - \int_{\Xi} \langle \text{Hess } g(x) \bar{G}(x, \xi), \bar{G}(x, \xi) \rangle \vartheta(d\xi) \right) \right| \leq C\eta^3. \end{aligned}$$

Next, we turn to the RSGD scheme. For $x \in M$ and $\xi \in \Xi$, let $(\gamma_t)_{t \in [0,1]} = (\text{retr}_x(-\eta t \text{grad } \tilde{f}(x, \xi)))_{t \in [0,1]}$ and note that

$$g(\gamma_1) = g(x) + \frac{d}{ds}|_{s=0}g(\gamma_s) + \frac{1}{2} \frac{d^2}{ds^2}|_{s=0}g(\gamma_s) + \int_0^1 \int_0^s \int_0^u \frac{d^3}{d\ell^3}g(\gamma_\ell) d\ell du ds.$$

Since $\text{retr} : TM \rightarrow M$ is a second order retraction, see Definition 2.1, we get $\frac{d}{ds}|_{s=0}g(\gamma_s) = -\eta \langle \text{grad } g(x), \text{grad } \tilde{f}(x, \xi) \rangle$ as well as

$$\begin{aligned} \frac{d^2}{ds^2}|_{s=0}g(\gamma_s) &= \langle \nabla_{\dot{\gamma}_0} \text{grad } g, \dot{\gamma}_0 \rangle + \langle \text{grad } g(\gamma_0), \frac{\nabla}{ds}|_{s=0} \dot{\gamma}_s \rangle \\ &= \eta^2 \langle (\text{Hess } g(x))(\text{grad } \tilde{f}(x, \xi)), \text{grad } \tilde{f}(x, \xi) \rangle. \end{aligned}$$

For the remainder, note that for all $\ell \in [0, 1]$

$$\frac{d^3}{d\ell^3}g(\gamma_\ell) = \nabla^3 g(\dot{\gamma}_\ell, \dot{\gamma}_\ell, \dot{\gamma}_\ell) + 3\nabla^2 g\left(\frac{\nabla}{d\ell} \dot{\gamma}_\ell, \dot{\gamma}_\ell\right) + \langle \text{grad } g(\gamma_\ell), \frac{\nabla^2}{d\ell^2} \dot{\gamma}_\ell \rangle,$$

where we used that the Riemannian Hessian is symmetric. Since retr is a uniform second order retraction, there exists a constant $C \geq 0$ that only depends on $\|g\|_{\mathcal{C}_b^3(M)}$ such that $|\frac{d^3}{d\ell^3}g(\gamma_\ell)| \leq C\eta^3 \|\text{grad } \tilde{f}(x, \xi)\|^3$. Taking expectation,

$$\begin{aligned} (26) \quad & |\mathbb{E}_\vartheta[g(\text{retr}_x(-\eta \text{grad } \tilde{f}(x, \xi)))] - g(x) + \eta \langle \text{grad } f(x), \text{grad } g(x) \rangle \\ & - \frac{1}{2} \eta^2 \mathbb{E}_\vartheta[\langle (\text{Hess } g(x))(\text{grad } \tilde{f}(x, \xi)), \text{grad } \tilde{f}(x, \xi) \rangle]| \leq C\eta^3 \mathbb{E}_\vartheta[\|\text{grad } \tilde{f}(x, \xi)\|^3], \end{aligned}$$

for a constant $C \geq 0$, where

$$\begin{aligned} \mathbb{E}_\vartheta[\langle (\text{Hess } g(x)) \text{grad } \tilde{f}(x, \xi), (\text{grad } \tilde{f}(x, \xi)) \rangle] &= \langle (\text{Hess } g(x)) \text{grad } f(x), (\text{grad } f(x)) \rangle \\ &+ \mathbb{E}_\vartheta[\langle (\text{Hess } g(x)) \bar{G}(x, \xi), \bar{G}(x, \xi) \rangle]. \end{aligned}$$

Using that $\mathbb{E}_\vartheta[g(\text{retr}_x(-\eta \text{grad } \tilde{f}(x, \xi)))] = \mathbb{E}[g(Z_1^\eta(x))]$ and comparing (26) with (25), we get for a constant $C \geq 0$ that

$$\sup_{x \in M} |\mathbb{E}[g(Z_1^\eta(x))] - \tilde{\mathbb{E}}[g(X_\eta^\eta(x))]| \leq C\eta^3.$$

□

Proof of Theorem 5.1. Fix $T \geq 0$. By Lemma 5.2, we have $B^\eta \in \mathfrak{X}_b^4(M)$ and $G^\eta \in \tilde{\mathfrak{X}}_b^5(M)$ uniformly over $0 \leq \eta \leq T$. By Definition 4.3, for all $0 \leq \eta \leq T$ there exists a unique solution to the SDE (4) that does not explode in finite time and for all $g \in \mathcal{C}_b^4$ there exists a constant $C \geq 0$ such that for all $0 \leq t \leq T$ and $0 \leq \eta \leq T$ the function $\Psi_t^\eta : M \rightarrow \mathbb{R}$ given by $\Psi_t^\eta(x) = \tilde{\mathbb{E}}[g(X_t^\eta(x))]$ satisfies $\|\Psi_t^\eta\|_{\mathcal{C}_b^4(M)} \leq C$.

We conceive the probability spaces $(\Omega, \mathcal{F}, \mathbb{P})$ and $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ as the projections of the product space $(\Omega \times \tilde{\Omega}, \mathcal{F} \otimes \tilde{\mathcal{F}}, \mathbb{P} \times \tilde{\mathbb{P}})$ so that $(Z_n^\eta(x))_{n \in \mathbb{N}_0}$ and $(X_t^\eta(x))_{t \geq 0}$ are independent processes. For $x \in M$ and $n \in \mathbb{N}$ we get, using the triangle inequality and the tower property of the conditional expectation,

$$\begin{aligned} |\mathbb{E}[g(Z_n^\eta(x))] - \tilde{\mathbb{E}}[g(X_{n\eta}^\eta(x))]| &= \left| \sum_{i=1}^n \mathbb{E} \left[\Psi_{(n-i)\eta}^\eta(Z_i^\eta(x)) - \tilde{\mathbb{E}}[\Psi_{(n-i)\eta}^\eta(X_\eta^\eta(Z_{i-1}^\eta(x)))] \right] \right| \\ &\leq \sum_{i=1}^n \sup_{z \in M} |\mathbb{E}[\Psi_{(n-i)\eta}^\eta(Z_1^\eta(z))] - \tilde{\mathbb{E}}[\Psi_{(n-i)\eta}^\eta(X_\eta^\eta(z))]|. \end{aligned}$$

Applying Lemma 5.3 to each summand on the right-hand side of the inequality above, there exists a constant $C \geq 0$ such that for all $n \in \{0, \dots, \lfloor T/\eta \rfloor\}$

$$|\mathbb{E}[g(Z_n^\eta(x))] - \tilde{\mathbb{E}}[g(X_{n\eta}^\eta(x))]| \leq nC\eta^3 \leq CT\eta^2.$$

□

6. EXAMPLES

6.1. Principal component analysis. In principal component analysis (PCA), the aim is to find the r principal eigenvectors of a matrix $A := \mathbb{E}_\vartheta[z(\xi)z(\xi)^T]$, where $z : \Xi \rightarrow \mathbb{R}^n$ is a random data vector and v^T denotes the transpose of a vector or a matrix v . For this problem, a natural choice of the search space is the Stiefel manifold $\text{St}(r, n)$ [Oja92, MHM96] or the Grassmann manifold $G(r, n)$ [HHT07]. The Stiefel manifold $\text{St}(r, n)$ given by $\text{St}(r, n) := \{B \in \mathbb{R}^{n \times r} : B^T B = \mathbb{I}_r\}$ is a compact, smooth manifold of dimension $nr - \frac{1}{2}r(r+1)$, see [AMS08, Section 3.3.2]. The Grassmann manifold $G(r, n)$ consisting of all r -dimensional subspaces of $\mathbb{R}^n$ is a compact, smooth manifold of dimension $r(n-r)$, see [AMS08, Section 3.4.4]. By compactness, $\text{St}(r, n)$ and $G(r, n)$ are $\text{BG}(\alpha)$ -manifolds for all $\alpha \in \mathbb{N}$, see Definition 4.4.

On $\text{St}(r, n)$ we can define a loss function

$$f(B) = -\frac{1}{2} \text{tr}(B^T A B),$$

which is minimal if B consists of eigenvectors that correspond to the r largest eigenvalues of A . We can regard f as a function on the Euclidean space $\mathbb{R}^{n \times r}$ with gradient $Df(B) = -AB$. If $\text{St}(r, n)$ is equipped with the Riemannian submanifold metric inherited from $\mathbb{R}^{n \times r}$ then the Riemannian gradient of f at $B \in \text{St}(r, n)$ is given by $\text{grad } f(B) = P_B(Df(B))$, where

$$P_B(Z) = \frac{1}{2}Z(B^T Z - Z^T B) + (\mathbb{I}_n - BB^T)Z$$

is the orthogonal projection of $Z \in \mathbb{R}^{n \times r}$ onto the tangent space $T_B \text{St}(r, n) = \{\Delta \in \mathbb{R}^{n \times r} : \Delta^T B + B^T \Delta = 0\}$, see [EAS99], and $\mathbb{I}_n$ denotes the identity matrix in $\mathbb{R}^{n \times n}$. By [AM12, Proposition 3.4], for all $B \in \text{St}(r, n)$ and $\bar{B} \in \mathbb{R}^{n \times r}$ with $\|B - \bar{B}\| < 1$ the projection (8) exists, is unique and has the expression

$$\text{proj}(\bar{B}) = UV^T.$$

Here, for a matrix $X \in \mathbb{R}^{n \times r}$, $\|X\|$ denotes the Frobenius norm, given by $\|X\|^2 = \text{tr}(X^T X)$ and $U \in \mathbb{R}^{n \times n}$, $V \in \mathbb{R}^{r \times r}$ are orthogonal matrices given by the singular value decomposition $B = U\Sigma V^T$, where $\Sigma \in \mathbb{R}^{n \times r}$ denotes the diagonal matrix consisting of the singular values of B in decreasing order. Using Lemma 2.3, we get a uniform second order retraction via $\text{retr}_B(\Delta) = \text{proj}(B + c(B, \Delta))$, where $c : \mathbb{R}^{n \times r} \times \mathbb{R}^{n \times r} \rightarrow \mathbb{R}^{n \times r}$ is chosen according to Lemma 2.3.

See also [EAS99, LSW19, LFH⁺20] for different retractions and Riemannian metrics on $\text{St}(r, n)$ as well as an analysis of the geometry of $G(r, n)$. For other common optimization tasks on matrix manifolds we refer the reader to Section 2 in [AMS08].

6.2. Normalizing the weights of a neural network. We consider optimizing a neural network with a positive homogeneous activation function. For simplicity, we restrict the section to neural networks with one-hidden layer although the arguments remain true for deep neural networks with multiple hidden layers.

Let $d_0, d_1 \in \mathbb{N}$ and $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ be positive homogeneous, i.e. $\sigma(\lambda x) = \lambda \sigma(x)$ for all $x \in \mathbb{R}$ and $\lambda \geq 0$. This property is satisfied by e.g. the ReLU activation function or any linear activation. The networks configuration can be described by the weights $W^1 = (w_{j,i}^1)_{j=1, \dots, d_1, i=1, \dots, d_0} \in \mathbb{R}^{d_1 \times d_0}$ and

$W^2 = (w_1^2, \dots, w_{d_1}^2) \in \mathbb{R}^{d_1}$, and the biases $b^1 = (b_i^1)_{i=1, \dots, d_1} \in \mathbb{R}^{d_1}$ and $b^2 \in \mathbb{R}$. For $j = 1, \dots, d$, we write $w_j^1 = (w_{j,1}^1, \dots, w_{j,d_{\text{in}}}^1)^T$. The search space is given by

$$\mathbb{W} = (W^1, b^1, W^2, b^2) \in \mathbb{R}^{d_1 \times d_0} \times \mathbb{R}^{d_1} \times \mathbb{R}^{d_1} \times \mathbb{R} =: \mathcal{W}_{d_0, d_1}$$

and for $\mathbb{W} \in \mathcal{W}_{d_0, d_1}$ we define $\mathfrak{N}^{\mathbb{W}} : \mathbb{R}^{d_0} \rightarrow \mathbb{R}$ via

$$\mathfrak{N}^{\mathbb{W}}(x) = \sum_{j=1}^{d_1} w_j^2 \sigma(x^T w_j^1 + b_j^1) + b^2,$$

which is the *response* of the neural network to the input x for the configuration $\mathbb{W}$. Using the positive homogeneity of σ we have

$$\mathfrak{N}^{\mathbb{W}}(x) = \sum_{j=1}^{d_1} w_j^2 \|w_j^1\| \sigma\left(x^T \frac{w_j^1}{\|w_j^1\|} + \frac{b_j^1}{\|w_j^1\|}\right) + b^2$$

so that the optimization can be restricted to the Riemannian manifold

$$M := \{\mathbb{W} \in \mathcal{W}_{d_0, d_1} : \|w_j^1\| = 1 \forall j = 1, \dots, d_1\} \cong (\mathbb{S}^{d_0-1})^{d_1} \times \mathbb{R}^{d_1} \times \mathbb{R}^{d_1} \times \mathbb{R}.$$

Although M is non-compact, it is clearly a $\text{BG}(\alpha)$ -manifold for all $\alpha \in \mathbb{N}$. A uniform second order retraction is given by a componentwise stereographic projection, see Example 2.2, or componentwise metric projection, see (8). The idea of decoupling the length and the direction of the weight vectors in neural networks was popularized by Salimans and Kingma with their weight normalization algorithm [SK16]. The corresponding gradient flow was shown to have a beneficial implicit bias while being less sensitive to the initialization [PLB20, MR22, CRW23].

6.3. Hyperbolic space. In recent years, one of the most popular applications of machine learning methods are natural language processing tasks, e.g. learning hierarchical representations of words through unsupervised learning. The aim of embedding methods is to position the words in the ambient space such that the distance reflects their semantic and functional similarity. The numerical experiments in [CCD17, NK17] suggest that the hyperbolic space $\mathbb{H}^d$ for $d \in \mathbb{N}$ is particularly well-suited as an ambient space. As a heuristic argument for the empirical findings we note that the volume of a ball in the hyperbolic space increases exponentially with respect its radius. This makes $\mathbb{H}^d$ a natural choice for embedding tree-like structures [SDSGR18], which appear naturally e.g. in many real-world information networks [ASM13].

Using the hyperboloid model of the hyperbolic space, there exist simple expressions for the Riemannian gradient and exponential map defined in the ambient Minkowski space, see [WL18]. Since $\mathbb{H}^d$ has constant sectional curvature, the Riemannian curvature tensor is bounded, see e.g. [DC92, Lemma 3.4]. Thus, Theorem 3.1 can be applied to the hyperbolic space.

The exponential growth of the volume of a ball w.r.t. its radius makes it impossible to find an isometric embedding with uniform tubular neighborhood in the sense of Definition 4.4. Hence, in order to apply the second order approximation result, Theorem 5.1, it remains to verify Definition 4.3 for the hyperbolic space, which is left for future research.

6.4. Statistical manifolds. A key task in generative AI is the inference of a probability distribution in a parametrized family $(\nu_\theta)_{\theta \in \Theta}$ of probability measures on $\mathbb{R}^{d_{\text{data}}}$ that is a good approximation to a given distribution ν_{data} . Assuming that ν_θ is absolutely continuous with respect to ν_{data} for all $\theta \in \Theta$, this can be performed by minimizing the Kullback-Leibler divergence

$$D_{\text{KL}}(\nu_\theta || \nu_{\text{data}}) := \int \log\left(\frac{d\nu_\theta}{d\nu_{\text{data}}}\right) d\nu_\theta.$$

In the training of generative adversarial networks, the Kullback-Leibler divergence is often approximated by choosing a set of discriminators $D : \mathbb{R}^{d_{\text{data}}} \rightarrow [0, 1]$ that try to distinguish between samples from the true data distribution ν_{data} and the distribution ν_θ , see [GPAM⁺14]. This leads to the objective function

$$f(\theta) := \max_D \int \log(D(x)) \nu_\theta(dx).$$

Let $\Theta \subset \mathbb{R}^d$ be an open set and assume that, for all $\theta \in \Theta$, ν_θ is a probability measure on $\mathbb{R}^{d_{\text{data}}}$ with density function p_θ . Then, the so-called Fisher information metric is given by

$$g_\theta(e_i, e_j) := - \int \frac{d^2 \log(p_\theta(y))}{dx_i dx_j} p_\theta(y) dy,$$

for $\theta \in \Theta$, where $e_1, \dots, e_d$ denotes the standard basis of the tangent space $T_\theta \Theta \simeq \mathbb{R}^d$. If $F_\theta := (g_\theta(e_i, e_j))_{i,j=1,\dots,d}$ is positive definite for all $\theta \in \Theta$ this defines a Riemannian metric on Θ . One can write the Riemannian gradient of f as $\text{grad } f(\theta) = F_\theta^{-1} Df(\theta)$, where $Df(\theta)$ denotes the Euclidean gradient, i.e. the vector of partial derivatives of $f : \Theta \rightarrow \mathbb{R}$ at θ , see e.g. [AMS08, Section 3.6].

For example, consider the set of univariate normal distributions $\Theta = \mathbb{R} \times \mathbb{R}_{>0} \ni \theta = (\mu, \sigma) \mapsto \mathcal{N}(\mu, \sigma)$. Then, the Fisher information metric is given by the matrix

$$(27) \quad F_{(\mu, \sigma)} = \begin{pmatrix} \frac{1}{\sigma^2} & 0 \\ 0 & \frac{2}{\sigma^2} \end{pmatrix},$$

see e.g. [CSS15]. This leads to a Riemannian metric (27) on the upper half plane Θ with constant negative curvature equal to $-\frac{1}{2}$, see also [CSS15]. Therefore, the geometry of Θ with Riemannian metric (27) is similar to the hyperbolic space considered in Section 6.3. In this situation, Theorem 3.1 can be applied, whereas Theorem 5.1 needs verification of Definition 4.3. Regarding the Fisher information metric and its distance function for multivariate normal distributions and other families of probability distributions see [PSC20, MMC23].

Acknowledgements. We would like to thank Matthias Erbar, Jan Nienhaus and Kevin Poljsak for fruitful discussions during the preparation of this work. BG acknowledges support by the Max Planck Society through the Research Group "Stochastic Analysis in the Sciences (SAiS)". NR acknowledges the support by the Royal Society research professorship of Prof. Martin Hairer, RP\R1\191065. This work was co-funded by the European Union (ERC, FluCo, grant agreement No. 101088488). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or of the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

APPENDIX A. NOTATION

We let M be a d -dimensional $\mathcal{C}^\infty$ -Riemannian manifold that is connected and complete. We denote by $T_x M$ the tangent space of M at point $x \in M$, by $\langle \cdot, \cdot \rangle_x$ the scalar product on $T_x M$ that is given by the Riemannian metric and by $\| \cdot \|_x$ the respective norm on $T_x M$. If it is clear from the context we often omit the reference point x in the above notions and briefly write $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$, respectively. Recall that the tangent bundle $TM := \bigcup_{x \in M} T_x M$ is a $2d$ -dimensional $\mathcal{C}^\infty$ -manifold. For $\alpha \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$, we denote by $\mathfrak{X}^\alpha(M)$ the set of $\mathcal{C}^\alpha$ -vector fields on M , i.e. the set of $\mathcal{C}^\alpha$ -functions $V : M \rightarrow TM$ with $V(x) \in T_x M$. For $V \in \mathfrak{X}^0(M)$ and $g : M \rightarrow \mathbb{R} \in \mathcal{C}^1(M)$ we denote by $Vg : M \rightarrow \mathbb{R}$ the function that is given by $Vg(x) = V(x)g$ for all $x \in M$. For $x \in M$, a $\mathcal{C}^\infty$ -manifold N and a differentiable mapping $\varphi : M \rightarrow N$ we denote by $D_x \varphi : T_x M \rightarrow T_{\varphi(x)} N$

the differential of φ at x , i.e. the linear mapping given by $(D_x\varphi v)g = v(g \circ \varphi)$ for all $v \in T_xM$ and $g \in \mathcal{C}^1(N)$. Furthermore, for $v \in T_xM$ and $W \in \mathfrak{X}^1(M)$ we denote by $\nabla_v W \in T_xM$ the covariant derivative of W along v that is induced by the Levi-Civita connection. For $V \in \mathfrak{X}^0(M)$ and $W \in \mathfrak{X}^1(M)$ we define $\nabla_V W \in \mathfrak{X}^0(M)$ via $\nabla_V W(x) = \nabla_{V(x)} W$. Note that, for brevity, we write $\nabla_{V(x)} W$ instead of $\nabla_{V(x)} W(x)$, whenever the base point is clear from the tangent vector in the first argument of the Levi-Civita connection.

For $\beta \in \mathbb{N}$ and a $\mathcal{C}^\beta$ -path $\gamma : [a, b] \rightarrow M$ and $t \in [a, b]$ we denote by $\dot{\gamma}_t \in T_{\gamma_t}M$ the differential of γ at time t , i.e. $\dot{\gamma}_t = (D_t\gamma)(\frac{d}{ds}|_{s=t})$. Moreover, for $\alpha \in \{0, \dots, \beta\}$ we denote by $\mathfrak{X}^\alpha(\gamma)$ the set of all $\mathcal{C}^\alpha$ -vector fields along γ , i.e. $\mathcal{C}^\alpha$ -mappings $V : [a, b] \rightarrow TM$ with $V_t \in T_{\gamma_t}M$ for all $t \in [a, b]$. For $\alpha \in \{1, \dots, \beta\}$, let $\frac{\nabla}{dt} : \mathfrak{X}^\alpha(\gamma) \rightarrow \mathfrak{X}^{\alpha-1}(\gamma)$ be the Levi-Civita connection on M along γ and $\text{Par}_\gamma : T_{\gamma_a}M \rightarrow T_{\gamma_b}M$ be the parallel transport along γ . We denote by $\exp_x : T_xM \rightarrow M$ the exponential map at $x \in M$.

R denotes the curvature of M , i.e. the $(3, 1)$ -tensor field given by

$$R(U, V)W = \nabla_U \nabla_V W - \nabla_V \nabla_U W - \nabla_{[U, V]} W,$$

for $U, V \in \mathfrak{X}^1(M)$ and $W \in \mathfrak{X}^2(M)$, where $[U, V] \in \mathfrak{X}^0(M)$ denotes the Lie bracket given by

$$[U, V]g = U(Vg) - V(Ug) \quad \text{for all } g \in \mathcal{C}^2(M).$$

APPENDIX B. DERIVATIVES OF HIGHER ORDER

In this section, we introduce the Riemannian gradient and Hessian, as well as derivatives of higher order for real-valued functions and vector fields on M . For more information on the Riemannian gradient and Hessian we refer the reader to Chapter 4 in [Lee12] as well as Chapter 3 and Chapter 5 in [AMS08].

Let $g \in \mathcal{C}^\alpha(M)$ for an $\alpha \in \mathbb{N}$. The Riemannian gradient $\text{grad } g \in \mathfrak{X}^{\alpha-1}(M)$ of g is the unique $\mathcal{C}^{\alpha-1}$ -vector field on M that satisfies for all $x \in M$ and $v \in T_xM$ that

$$\langle \text{grad } g(x), v \rangle = vg.$$

If $\alpha \geq 2$ we define the Riemannian Hessian $\text{Hess } g(x) : T_xM \rightarrow T_xM$ of g at x as the linear mapping given by $(\text{Hess } g(x))(v) = \nabla_v \text{grad } g$ for all $v \in T_xM$. Using that ∇ is a metric connection, this defines a symmetric $(2, 0)$ -tensor field $\nabla^2 g : \mathfrak{X}^1(M) \times \mathfrak{X}^1(M) \rightarrow \mathcal{C}^0(M)$ via

$$\nabla^2 g(V, W) := \langle (\text{Hess } g)V, W \rangle = V(Wg) - (\nabla_V W)g.$$

We have $\|\text{Hess } g(x)\| = \|\nabla^2 g(x)\|$, where the left-hand side denotes the operator norm of the linear mapping $\text{Hess } g(x) : T_xM \rightarrow T_xM$ and the right-hand side denotes the operator norm of the bilinear mapping $\nabla^2 g(x) : T_xM \times T_xM \rightarrow \mathbb{R}$.

This representation allows us to generalize the concept of the Riemannian Hessian to derivatives of higher order. For $3 \leq n \leq \alpha$, we define an $(n, 0)$ -tensor field $\nabla^n g : (\mathfrak{X}^{n-1}(M))^n \rightarrow \mathcal{C}^0(M)$ inductively via

$$\begin{aligned} \nabla^n g(V_1, \dots, V_n) &= V_1((\nabla^{n-1} g)(V_2, \dots, V_n)) \\ &\quad - \sum_{i=2}^n (\nabla^{n-1} g)(V_2, \dots, V_{i-1}, \nabla_{V_1} V_i, V_{i+1}, \dots, V_n). \end{aligned}$$

This canonically defines a multilinear mapping $\nabla^n g(x) : (T_xM)^n \rightarrow \mathbb{R}$ via

$$(\nabla^n g(x))(v_1, \dots, v_n) = (\nabla^n g(V_1, \dots, V_n))(x),$$

where $V_1, \dots, V_n \in \mathfrak{X}^{n-1}(M)$ are vector fields with $V_i(x) = v_i$ for all $1 \leq i \leq n$. We denote by $\|\nabla^n g(x)\|$ the respective operator norm.

If M is an open subset of a Euclidean space $\mathbb{R}^N$ one has $\nabla_{E_i} E_j = 0$ for all $i, j = 1, \dots, N$ where $E_i(x) = \frac{d}{dx_i}|_x$. Thus, in that case $\nabla^n g$ is given by $D^n g$ which denotes the tensor given by all Euclidean derivatives of g of order n .

Analogously, one can define the derivatives of vector fields $V \in \mathfrak{X}^\alpha(M)$ for $\alpha \in \mathbb{N}$. For $n \leq \alpha$ we define an $(n, 1)$ -tensor field $\nabla^n V : (\mathfrak{X}^{n-1}(M))^n \rightarrow \mathfrak{X}^0(M)$ inductively via $(\nabla^1 V)(V_1) = \nabla_{V_1} V$ and

$$\begin{aligned} (\nabla^n V)(V_1, \dots, V_n) = & \nabla_{V_1}((\nabla^{n-1} V)(V_2, \dots, V_n)) \\ & - \sum_{i=2}^n (\nabla^{n-1} V)(V_2, \dots, V_{i-1}, \nabla_{V_1} V_i, V_{i+1}, \dots, V_n). \end{aligned}$$

This induces a multilinear mapping $\nabla^n V(x) : (T_x M)^n \rightarrow T_x M$ and we denote by $\|\nabla^n V(x)\|$ its operator norm.

Definition B.1. (i) We denote by $\mathcal{C}_b^0(M)$ the set of continuous and bounded functions $g : M \rightarrow \mathbb{R}$ and associate to g the norm $\|g\|_{\mathcal{C}_b^0(M)} := \sup_{x \in M} |g(x)|$. Analogously, for $\alpha \in \mathbb{N}$ we denote by $\mathcal{C}_b^\alpha(M)$ the set of functions $g \in \mathcal{C}^\alpha(M)$ that satisfy

$$\|g\|_{\mathcal{C}_b^\alpha(M)} := \sup_{x \in M} \left(|g(x)| \vee \|\text{grad } g(x)\| \vee \max_{2 \leq n \leq \alpha} \|\nabla^n g(x)\| \right) < \infty.$$

(ii) We denote by $\mathfrak{X}_b^0(M)$ the set of continuous and bounded vector fields $V \in \mathfrak{X}^0(M)$ and associate to V the norm $\|V\|_{\mathfrak{X}_b^0(M)} := \sup_{x \in M} \|V(x)\|$. Analogously, for $\alpha \in \mathbb{N}$ we denote by $\mathfrak{X}_b^\alpha(M)$ the set of vector fields $V \in \mathfrak{X}^\alpha(M)$ that satisfy

$$\|V\|_{\mathfrak{X}_b^\alpha(M)} := \sup_{x \in M} \left(\|V(x)\| \vee \max_{1 \leq n \leq \alpha} \|\nabla^n V(x)\| \right) < \infty.$$

Next, we show how the consecutive differentiation of g w.r.t. multiple vector fields can be expressed in terms of the multilinear mappings defined above. Let us start with second order derivatives. Let $V_1, V_2 \in \mathfrak{X}^1(M)$ and $g \in \mathcal{C}^2(M)$. Then, by definition,

$$V_1 V_2 g = \nabla^2 g(V_1, V_2) + \langle (\nabla V_2)(V_1), \text{grad } g \rangle,$$

where the right-hand side only depends on $\nabla^2 g$, $\text{grad } g$, V_1, V_2 and ∇V_2 . Thus, $\|V_1 V_2 g\|_{\mathcal{C}_b^0(M)}$ can be bounded by a constant that only depends $\|g\|_{\mathcal{C}_b^2(M)}$, $\|V_1\|_{\mathfrak{X}_b^0(M)}$ and $\|V_2\|_{\mathfrak{X}_b^1(M)}$. Analogously, for $V_1, V_2, V_3 \in \mathfrak{X}^2(M)$ and $g \in \mathcal{C}^3(M)$ we get

$$\begin{aligned} V_1 V_2 V_3 g = & \nabla^3 g(V_1, V_2, V_3) + \nabla^2 g(\nabla_{V_1} V_2, V_3) + \nabla^2 g(V_2, \nabla_{V_1} V_3) + V_1(\nabla_{V_2} V_3)g \\ = & \nabla^3 g(V_1, V_2, V_3) + \nabla^2 g(\nabla_{V_1} V_2, V_3) + \nabla^2 g(V_2, \nabla_{V_1} V_3) + \nabla^2 g(V_1, \nabla_{V_2} V_3) \\ & + \langle \nabla_{V_1} \nabla_{V_2} V_3, \text{grad } g \rangle, \end{aligned}$$

where the right-hand side of the equation above can be bounded by a constant that only depends on $\|g\|_{\mathcal{C}_b^3(M)}$, $\|V_1\|_{\mathfrak{X}_b^0(M)}$, $\|V_2\|_{\mathfrak{X}_b^1(M)}$ and $\|V_3\|_{\mathfrak{X}_b^2(M)}$. Lastly, for $V_1, V_2, V_3, V_4 \in \mathfrak{X}^3(M)$ and $g \in \mathcal{C}^4(M)$ we get

$$\begin{aligned} V_1 V_2 V_3 V_4 g = & \nabla^4 g(V_1, V_2, V_3, V_4) + \nabla^3 g(\nabla_{V_1} V_2, V_3, V_4) + \nabla^3 g(V_2, \nabla_{V_1} V_3, V_4) \\ & + \nabla^3 g(V_2, V_3, \nabla_{V_1} V_4) + V_1 V_2(\nabla_{V_3} V_4)g + V_1(\nabla_{V_2} V_3)V_4 g - V_1(\nabla_{\nabla_{V_2} V_3} V_4)g \\ & + V_1 V_3(\nabla_{V_2} V_4)g - V_1(\nabla_{V_3} \nabla_{V_2} V_4)g, \end{aligned}$$

where, after a straight-forward computation, the right-hand side of the equation above can be bounded by a constant that only depends on $\|g\|_{\mathcal{C}_b^4(M)}$, $\|V_1\|_{\mathfrak{X}_b^0(M)}$, $\|V_2\|_{\mathfrak{X}_b^1(M)}$, $\|V_3\|_{\mathfrak{X}_b^2(M)}$ and $\|V_4\|_{\mathfrak{X}_b^3(M)}$. For higher derivatives there exist similar expressions.

REFERENCES

- [AM12] P.-A. Absil and J. Malick. Projection-like retractions on matrix manifolds. *SIAM J. Optim.*, 22(1):135–158, 2012.
- [AMS08] P.-A. Absil, R. Mahony, and R. Sepulchre. *Optimization algorithms on matrix manifolds*. Princeton University Press, Princeton, NJ, 2008.
- [AP22] S. Ankirchner and S. Perko. Towards diffusion approximations for stochastic gradient descent without replacement. hal-03527878, 2022.
- [ASM13] A. B. Adcock, B. D. Sullivan, and M. W. Mahoney. Tree-like structure in large social and information networks. In *IEEE 13th international conference on data mining*, pages 1–10. IEEE, 2013.
- [Bon13] S. Bonnabel. Stochastic gradient descent on Riemannian manifolds. *IEEE Trans. Automat. Control*, 58(9):2217–2229, 2013.
- [Bou23] N. Boumal. *An introduction to optimization on smooth manifolds*. Cambridge University Press, 2023.
- [CB19] C. Criscitiello and N. Boumal. Efficiently escaping saddle points on manifolds. In *Advances in Neural Information Processing Systems*, volume 32, 2019.
- [CCD17] B. P. Chamberlain, J. R. Clough, and M. P. Deisenroth. Neural embeddings of graphs in hyperbolic space. In *CoRR. MLG Workshop 2017*, 2017.
- [CRW23] H.-H. Chou, H. Rauhut, and R. Ward. Robust implicit regularization via weight normalization. arXiv:2305.05448, 2023.
- [CSS15] S. I. R. Costa, S. A. Santos, and J. E. Strapasson. Fisher information distance: A geometrical reading. *Discrete Appl. Math.*, 197:59–69, 2015.
- [DC92] M. P. Do Carmo. *Riemannian geometry*, volume 6. Springer, 1992.
- [DJM⁺21] A. Durmus, P. Jiménez, É. Moulines, S. Said, and H.-T. Wai. Convergence analysis of Riemannian stochastic approximation schemes. arXiv:2005.13284, 2021.
- [DJMS21] A. Durmus, P. Jiménez, É. Moulines, and S. Said. On Riemannian stochastic approximation schemes with fixed step-size. In *International Conference on Artificial Intelligence and Statistics*, pages 1018–1026. PMLR, 2021.
- [DK00] J. J. Duistermaat and J. A. C. Kolk. *Lie groups*. Springer-Verlag, Berlin, 2000.
- [DK22a] S. Dereich and S. Kassing. Cooling down stochastic differential equations: almost sure convergence. *Stochastic Process. Appl.*, 152:289–311, 2022.
- [DK22b] S. Dereich and S. Kassing. On minimal representations of shallow ReLU networks. *Neural Networks*, 148:121–128, 2022.
- [DPZ14] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*. Cambridge university press, 2014.
- [EAS99] A. Edelman, T. A. Arias, and S. T. Smith. The geometry of algorithms with orthogonality constraints. *SIAM J. Matrix Anal. Appl.*, 20(2):303–353, 1999.
- [Eld13] J. Eldering. *Normally hyperbolic invariant manifolds: the noncompact case*, volume 2. Springer, 2013.
- [Elw82] K. D. Elworthy. *Stochastic differential equations on manifolds*, volume 70. Cambridge University Press, 1982.
- [FDBD21] X. Fontaine, V. De Bortoli, and A. Durmus. Convergence rates and approximation results for SGD and its continuous-time counterpart. In *Conference on Learning Theory*, pages 1965–2058. PMLR, 2021.
- [FGJ20] B. Fehrman, B. Gess, and A. Jentzen. Convergence rates for the stochastic gradient descent method for non-convex objective functions. *J. Mach. Learn. Res.*, 21:5354–5401, 2020.
- [FGL⁺20] Y. Feng, T. Gao, L. Li, J.-G. Liu, and Y. Lu. Uniform-in-time weak error analysis for stochastic gradient descent algorithms via diffusion approximation. *Commun. Math. Sci.*, 18(1):163–188, 2020.
- [FLL18] Y. Feng, L. Li, and J.-G. Liu. Semigroups of stochastic gradient descent and online principal component analysis: properties and diffusion approximations. *Commun. Math. Sci.*, 16(3), 2018.
- [GK23] B. Gess and S. Kassing. Convergence rates for momentum stochastic gradient descent with noise of machine learning type. arXiv:2302.03550, 2023.
- [GKK24] B. Gess, S. Kassing, and V. Konarovskiy. Stochastic modified flows, mean-field limits and dynamics of stochastic gradient descent. *J. Mach. Learn. Res.*, 25(30):1–27, 2024.
- [GM10] L. Gawarecki and V. Mandrekar. *Stochastic differential equations in infinite dimensions: with applications to stochastic partial differential equations*. Springer Science & Business Media, 2010.

- [GPAM⁺14] I. Goodfellow, J. Pouget-Abadie, M. Mirza, B. Xu, D. Warde-Farley, S. Ozair, A. Courville, and Y. Bengio. Generative adversarial nets. In *Advances in Neural Information Processing Systems*, volume 27, 2014.
- [HHT07] U. Helmke, K. Hüper, and J. Trumpf. Newton’s method on Grassmann manifolds. arXiv:0709.2205, 2007.
- [HKKM23] Y.-P. Hsieh, M. R. Karimi, A. Krause, and P. Mertikopoulos. Riemannian stochastic optimization methods avoid strict saddle points. In *Advances in Neural Information Processing Systems*, volume 37, 2023.
- [HLLL19] W. Hu, C. J. Li, L. Li, and J.-G. Liu. On the diffusion approximation of nonconvex stochastic gradient descent. *Ann. Math. Sci. Appl.*, 4(1), 2019.
- [HSK⁺12] G. E. Hinton, N. Srivastava, A. Krizhevsky, I. Sutskever, and R. R. Salakhutdinov. Improving neural networks by preventing co-adaptation of feature detectors. arXiv:1207.0580, 2012.
- [Hsu02] E. P. Hsu. *Stochastic analysis on manifolds*. Number 38. American Mathematical Soc., 2002.
- [IW89] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*, volume 24 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam; Kodansha, Ltd., Tokyo, second edition, 1989.
- [Jør75] E. Jørgensen. The central limit problem for geodesic random walks. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 32(1):1–64, 1975.
- [KHKM22] M. R. Karimi, Y.-P. Hsieh, P. Mertikopoulos, and A. Krause. Riemannian stochastic approximation algorithms. arXiv:2206.06795, 2022.
- [Lee12] J. M. Lee. *Introduction to Smooth Manifolds*. Springer New York, 2012.
- [LFH⁺20] T. Lin, C. Fan, N. Ho, M. Cuturi, and M. Jordan. Projection robust wasserstein distance and Riemannian optimization. In *Advances in Neural Information Processing Systems*, volume 33, 2020.
- [Li94a] X.-M. Li. Properties at infinity of diffusion semigroups and stochastic flows via weak uniform covers. *Potential Anal.*, 3(4):339–357, 1994.
- [Li94b] X.-M. Li. Strong p -completeness of stochastic differential equations and the existence of smooth flows on noncompact manifolds. *Probab. Theory Related Fields*, 100(4):485–511, 1994.
- [LMA21] Z. Li, S. Malladi, and S. Arora. On the validity of modeling SGD with stochastic differential equations (SDEs). In *Advances in Neural Information Processing Systems*, volume 34, 2021.
- [LR15] W. Liu and M. Röckner. *Stochastic partial differential equations: an introduction*. Springer, 2015.
- [LS21] G. Leobacher and A. Steinicke. Existence, uniqueness and regularity of the projection onto differentiable manifolds. *Ann. Global Anal. Geom.*, 60(3):559–587, 2021.
- [LSW19] H. Liu, A. M.-C. So, and W. Wu. Quadratic optimization with orthogonality constraint: explicit Łojasiewicz exponent and linear convergence of retraction-based line search and stochastic variance-reduced gradient methods. *Math. Program.*, 178(1-2):215–262, 2019.
- [LTE17] Q. Li, C. Tai, and W. E. Stochastic modified equations and adaptive stochastic gradient algorithms. In *International Conference on Machine Learning*, pages 2101–2110. PMLR, 2017.
- [LTE19] Q. Li, C. Tai, and W. E. Stochastic modified equations and dynamics of stochastic gradient algorithms I: Mathematical foundations. *J. Mach. Learn. Res.*, 20:Paper No. 40, 47, 2019.
- [LW22] L. Li and Y. Wang. On uniform-in-time diffusion approximation for stochastic gradient descent. arXiv:2207.04922, 2022.
- [LWA21] Z. Li, T. Wang, and S. Arora. What happens after SGD reaches zero loss ? – a mathematical framework. In *International Conference on Learning Representations*, 2021.
- [MHM96] R. E. Mahony, U. Helmke, and J. B. Moore. Gradient algorithms for principal component analysis. *J. Austral. Math. Soc. Ser. B*, 37(4):430–450, 1996.
- [MMC23] H. K. Miyamoto, F. C. C. Meneghetti, and S. I. R. Costa. On closed-form expressions for the fisher-rao distance. *arXiv preprint arXiv:2304.14885*, 2023.
- [MR22] D. Morwani and H. G. Ramaswamy. Inductive bias of gradient descent for weight normalized smooth homogeneous neural nets. In *International Conference on Algorithmic Learning Theory*, pages 827–880. PMLR, 2022.
- [MTB23] C. Moucer, A. Taylor, and F. Bach. A systematic approach to Lyapunov analyses of continuous-time models in convex optimization. *SIAM J. Optim.*, 33(3):1558–1586, 2023.
- [NK17] M. Nickel and D. Kiela. Poincaré embeddings for learning hierarchical representations. In *Advances in Neural Information Processing Systems*, volume 30, 2017.

- [Oja92] E. Oja. Principal components, minor components, and linear neural networks. *Neural Networks*, 5:927–935, 1992.
- [Per23] S. Perko. Unlocking optimal batch size schedules using continuous-time control and perturbation theory. arXiv:2312.01898, 2023.
- [PLB20] T. Poggio, Q. Liao, and A. Banburski. Complexity control by gradient descent in deep networks. *Nature Communications*, 11(1):1027, 2020.
- [PSC20] J. Pinele, J. E. Strapasson, and S. I. R. Costa. The fisher–rao distance between multivariate normal distributions: Special cases, bounds and applications. *Entropy*, 22(4):404, 2020.
- [Rie11] M. Riedle. Cylindrical Wiener processes. In *Séminaire de Probabilités XLIII*, pages 191–214. Springer, 2011.
- [SDSGR18] F. Sala, C. De Sa, A. Gu, and C. Ré. Representation tradeoffs for hyperbolic embeddings. In *International conference on machine learning*, pages 4460–4469. PMLR, 2018.
- [SFF19] Y. Sun, N. Flammarion, and M. Fazel. Escaping from saddle points on Riemannian manifolds. In *Advances in Neural Information Processing Systems*, volume 32, 2019.
- [Sha21] S. M. Shah. Stochastic approximation on Riemannian manifolds. *Appl. Math. Optim.*, 83:1123–1151, 2021.
- [SHSW23] S. Schwarz, M. Herrmann, A. Sturm, and M. Wardetzky. Efficient random walks on Riemannian manifolds. *Found. Comput. Math.*, 2023. To appear.
- [SI23] H. Sakai and H. Iiduka. Convergence of Riemannian stochastic gradient descent on Hadamard manifold. arXiv:2312.07990, 2023.
- [SK16] T. Salimans and D. P. Kingma. Weight normalization: A simple reparameterization to accelerate training of deep neural networks. In *Advances in Neural Information Processing Systems*, volume 29, 2016.
- [SWRH20] Z. Shen, Z. Wang, A. Ribeiro, and H. Hassani. Sinkhorn natural gradient for generative models. In *Advances in Neural Information Processing Systems*, pages 1646–1656, 2020.
- [SZRM18] T. Salimans, H. Zhang, A. Radford, and D. Metaxas. Improving GANs using optimal transport. In *International Conference on Learning Representations*, 2018.
- [TFBJ18] N. Tripuraneni, N. Flammarion, F. Bach, and M. I. Jordan. Averaging stochastic gradient descent on Riemannian manifolds. In *Conference On Learning Theory*, pages 650–687. PMLR, 2018.
- [Udr94] C. Udriște. *Convex functions and optimization methods on Riemannian manifolds*, volume 297 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1994.
- [WL18] B. Wilson and M. Leimeister. Gradient descent in hyperbolic space. arXiv:1805.08207, 2018.
- [Woj24] S. Wojtowytsch. Stochastic Gradient Descent with Noise of Machine Learning Type Part II: Continuous Time Analysis. *J. Nonlinear Sci.*, 34(1):Paper No. 16, 2024.
- [ZLP⁺22] J. Zhao, A. Lucchi, F. N. Proske, A. Orvieto, and H. Kersting. Batch size selection by stochastic optimal control. In *Has it Trained Yet? NeurIPS 2022 Workshop*, 2022.
- [ZRS16] H. Zhang, S. J. Reddi, and S. Sra. Riemannian SVRG: Fast stochastic optimization on Riemannian manifolds. In *Advances in Neural Information Processing Systems*, volume 29, 2016.
- [ZS16] H. Zhang and S. Sra. First-order methods for geodesically convex optimization. In *Conference on Learning Theory*, pages 1617–1638. PMLR, 2016.

BENJAMIN GESS, INSTITUTE OF MATHEMATICS, TU BERLIN, 10623 BERLIN, GERMANY, AND MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, INSELSTRASSE 22, 04103 LEIPZIG, GERMANY

Email address: benjamin.gess@gmail.com

SEBASTIAN KASSING, FACULTY OF MATHEMATICS, UNIVERSITY OF BIELEFELD, UNIVERSITÄTSSTRASSE 25, 33615 BIELEFELD, GERMANY

Email address: skassing@math.uni-bielefeld.de

NIMIT RANA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK YO10 5DD, UK

Email address: nimit.rana@york.ac.uk