

This is a repository copy of *G-complete reducibility and saturation*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/225148/>

Version: Published Version

Article:

Bate, Michael orcid.org/0000-0002-6513-2405, Böhm, Sören, Litterick, Alastair et al. (2 more authors) (2025) *G-complete reducibility and saturation*. Pacific Journal of Mathematics. pp. 1-24. ISSN 0030-8730

<https://doi.org/10.2140/pjm.2025.337.1>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Pacific Journal of Mathematics

G-COMPLETE REDUCIBILITY AND SATURATION

MICHAEL BATE, SÖREN BÖHM, ALASTAIR LITTERICK,
BENJAMIN MARTIN AND GERHARD RÖHRLE

G -COMPLETE REDUCIBILITY AND SATURATION

MICHAEL BATE, SÖREN BÖHM, ALASTAIR LITTERICK,
BENJAMIN MARTIN AND GERHARD RÖHRLE

Let $H \subseteq G$ be connected reductive linear algebraic groups defined over an algebraically closed field of characteristic $p > 0$. In our first main theorem we show that if a closed subgroup K of H is H -completely reducible, then it is also G -completely reducible in the sense of Serre, under some restrictions on p , generalising the known case for $G = \mathrm{GL}(V)$. Our proof uses R. W. Richardson's notion of reductive pairs to reduce to the $\mathrm{GL}(V)$ case. We study Serre's notion of saturation and prove that saturation behaves well with respect to products and regular subgroups. Our second main theorem shows that if K is H -completely reducible, then the saturation of K in G is completely reducible in the saturation of H in G (which is again a connected reductive subgroup of G), under suitable restrictions on p , again generalising the known instance for $G = \mathrm{GL}(V)$. We also study saturation of finite subgroups of Lie type in G . We show that saturation is compatible with standard Frobenius endomorphisms, and we use this to generalise a result due to Nori from 1987 in the case $G = \mathrm{GL}(V)$.

1. Introduction and main results

Let G be a connected reductive linear algebraic group over an algebraically closed field k of characteristic $p > 0$. Let H be a closed subgroup of G . Following Serre [2004], we say that H is G -completely reducible (G -cr for short) provided that whenever H is contained in a parabolic subgroup P of G , it is contained in a Levi subgroup of P . Further, H is G -irreducible (G -ir for short) provided H is not contained in any proper parabolic subgroup of G at all. Clearly, if H is G -irreducible, it is trivially G -completely reducible; for an overview of this concept, see [Bate et al. 2005; Serre 1998; 2004]. Note in the case $G = \mathrm{GL}(V)$ a subgroup H is G -cr exactly when V is a semisimple H -module and it is G -ir precisely when V is an irreducible H -module. The same equivalence applies to $G = \mathrm{SL}(V)$. The notion of G -complete reducibility is a powerful tool for investigating the subgroup structure of G : see [Litterick and Thomas 2018; Litterick et al. 2025].

MSC2020: 14L24, 20G15.

Keywords: G -complete reducibility, saturation, finite groups of Lie type.

Now suppose H is connected and reductive, and let K be a closed subgroup of H . It is natural to ask the following questions:

Question 1.1. *If K is H -completely reducible, must K be G -completely reducible?*

Question 1.2. *If K is G -completely reducible, must K be H -completely reducible?*

The answer to [Question 1.1](#) is no in general. For instance, if H is a non- G -cr subgroup of G and $K = H$ then K is H -ir but K is not G -cr. For a more complicated example showing that the answer to both [Questions 1.1](#) and [1.2](#) is no, see [\[Bate et al. 2005, Example 3.45\]](#) or [\[Bate et al. 2010, Proposition 7.17\]](#). On the other hand, if $p > 2$ and H is either the special orthogonal group $\mathrm{SO}(V)$ or the symplectic group $\mathrm{Sp}(V)$ with its natural embedding in $G := \mathrm{GL}(V)$ then K is G -cr if and only if K is H -cr (see [\[Bate et al. 2005, Example 3.23\]](#)).

In this paper we consider some variations on [Questions 1.1](#) and [1.2](#). To do this we use two key tools: reductive pairs and saturation. Our first main result shows that the answer to [Question 1.1](#) is yes if we impose some extra conditions on p . To state our result we need some notation. We define an invariant $d(G)$ of G as follows. For G simple let $d(G)$ be as follows:

G	$A_n (n \geq 1)$	$B_n (n \geq 3)$	$C_n (n \geq 2)$	$D_n (n \geq 4)$	E_6	E_7	E_8	F_4	G_2
$d(G)$	$n + 1$	$2n + 1$	$2n$	$2n$	27	56	248	26	7

For G reductive, let $d(G) = \max(1, d(G_1), \dots, d(G_r))$, where G_1, \dots, G_r are the simple components of G . For G simple and simply connected and p good for G , $d(G)$ is the minimal possible dimension of a nontrivial irreducible G -module.

Theorem 1.3. *Let $H \subseteq G$ be connected reductive groups and let K be a closed subgroup of H . Suppose $p \geq d(G)$. If K is H -completely reducible, then K is G -completely reducible.*

[Theorem 1.3](#) generalises [\[Bate et al. 2011, Theorem 1.3\]](#), which is the special case when $G = \mathrm{GL}(V)$. We note that [Theorem 1.3](#) is false without the bound on p : e.g., see [\[Bate et al. 2005, Example 3.44\]](#) or [Example 4.5](#).

To prove [Theorem 1.3](#) we use Richardson's theory of reductive pairs (see [Definition 3.3](#)) to reduce to the case $G = \mathrm{GL}(V)$, and then we apply [\[Bate et al. 2011, Theorem 1.3\]](#). We require a careful analysis of when the Lie algebra of G admits a nondegenerate Ad-invariant bilinear form, building on the discussions in [\[Richardson 1967\]](#) and [\[Slodowy 1997\]](#) (see also [\[Bate et al. 2005, Section 3.5\]](#)). We establish the results we need in [Theorem 3.6](#).

Our second main result involves the concept of saturation due to Serre. Let $h(G)$ and $\tilde{h}(G)$ be as defined in [Section 2.5](#) (if G is simple then $h(G)$ is the Coxeter number of G). If $p \geq h(G)$ then for any unipotent $u \in G$ and any $t \in k$ one obtains a unipotent element u^t of G using versions of the matrix exponential and logarithm

maps; see [Section 5](#) for details. If H is a closed subgroup of G then we call H *saturated* if $u^t \in H$ for every unipotent $u \in H$ and every $t \in k$. We denote by H^{sat} the smallest closed saturated subgroup that contains H . Saturation was used by Serre [\[1998; 2004\]](#) in his study of G -complete reducibility (see [Theorem 5.13](#)).

Theorem 1.4. *Let G , H and K be as in [Theorem 1.3](#). Suppose $p \geq d(G)$ and $p \geq \tilde{h}(G)$. If K is H -completely reducible then K^{sat} is H^{sat} -completely reducible. (Note that K is G -cr by [Theorem 1.3](#).)*

For a direct application of saturation to [Questions 1.1](#) and [1.2](#), see [Proposition 5.14](#).

To prove [Theorem 1.4](#) we establish some results on saturation that are of interest in their own right. These include [Remark 5.9](#), [Lemma 5.10](#) and [Corollary 5.11](#).

We also give some versions of our main results for finite groups of Lie type. Recall that a *Steinberg endomorphism* of G is a surjective morphism $\sigma : G \rightarrow G$ such that the corresponding fixed-point subgroup $G_\sigma := \{g \in G \mid \sigma(g) = g\}$ of G is finite. The latter are the *finite groups of Lie type*; see Steinberg [\[1968\]](#) for a detailed discussion. The set of all Steinberg endomorphisms of G is a subset of the set of all isogenies $G \rightarrow G$ (see [\[Steinberg 1968, 7.1\(a\)\]](#)), which encompasses in particular all generalized Frobenius endomorphisms, i.e., endomorphisms of G some power of which are Frobenius endomorphisms corresponding to some \mathbb{F}_q -rational structure on G . If H is a connected reductive σ -stable subgroup H of G then σ is also a Steinberg endomorphism for H with finite fixed-point subgroup $H_\sigma = H \cap G_\sigma$ [\[Steinberg 1968, 7.1\(b\)\]](#).

Corollary 1.5. *Let $H \subseteq G$ be connected reductive groups. Let $\sigma : G \rightarrow G$ be a Steinberg endomorphism that stabilises H . Suppose $p \geq d(G)$. Then the fixed-point subgroup H_σ is G -completely reducible.*

Theorem 1.6. *Suppose G is simple. Let $p \geq \tilde{h}(G)$. Let σ be a standard Frobenius endomorphism of G and let H be a connected reductive, σ -stable, and saturated subgroup of G . Then $(H_\sigma)^{\text{sat}} = H$.*

[Theorem 1.6](#) generalises a theorem of Nori [\[1987, Theorem B\(2\)\]](#); Nori's result is the special case of [Theorem 1.6](#) for $G = \text{GL}_n$ and $\sigma = \sigma_p$ the standard Frobenius endomorphism of G raising the matrix coefficients to the p -th power. In that context [Proposition 6.1](#) is of independent interest, saying that saturation is compatible with standard Frobenius endomorphisms.

[Section 2](#) contains some preliminary material and [Section 3](#) deals with reductive pairs. We prove [Theorem 1.3](#) and [Corollary 1.5](#) in [Section 4](#). Results on saturation, including the proof of [Theorem 1.4](#), are treated in [Section 5](#). In [Section 6](#) we study saturation for finite subgroups of Lie type. There we prove [Theorem 1.6](#), among other results. Finally, [Section 7](#) explores the connection between saturation and the concept of a semisimplification of a subgroup of G from [\[Bate et al. 2020\]](#).

2. Preliminaries

Throughout, we work over an algebraically closed field k of characteristic $p \geq 0$. All affine varieties are considered over k and are identified with their k -points.

A linear algebraic group H over k has identity component H° ; if $H = H^\circ$, then we say that H is *connected*. We denote by $R_u(H)$ the *unipotent radical* of H ; if $R_u(H)$ is trivial, then we say H is *reductive*.

Throughout, G denotes a connected reductive linear algebraic group over k . All subgroups of G that are considered are closed.

2.1. Good and very good primes. Suppose G is simple. Fix a Borel subgroup B of G containing a maximal torus T . Let $\Phi = \Phi(G, T)$ be the root system of G with respect to T , let $\Phi^+ = \Phi(B, T)$ be the set of positive roots of G , and let $\Sigma = \Sigma(G, T)$ be the set of simple roots of the root system Φ of G defined by B . For $\beta \in \Phi^+$ write $\beta = \sum_{\alpha \in \Sigma} c_{\alpha\beta} \alpha$ with $c_{\alpha\beta} \in \mathbb{N}_0$. A prime p is said to be *good* for G if it does not divide $c_{\alpha\beta}$ for any α and β . A prime p is said to be *very good* for G if p is a good prime for G , and in the case that G is of type A_n , then p does not divide $n + 1$. For G reductive p is *good* (*very good*) for G if p is good (*very good*) for every simple component of G .

2.2. Limits and parabolic subgroups. Let $\phi : k^* \rightarrow X$ be a morphism of algebraic varieties. We say $\lim_{a \rightarrow 0} \phi(a)$ exists if there is a morphism $\hat{\phi} : k \rightarrow X$ (necessarily unique) whose restriction to k^* is ϕ ; if the limit exists, we set $\lim_{a \rightarrow 0} \phi(a) = \hat{\phi}(0)$. As a direct consequence of the definition we have the following:

Remark 2.1. If $\phi : k^* \rightarrow X$ and $h : X \rightarrow Y$ are morphisms of varieties and $x := \lim_{a \rightarrow 0} \phi(a)$ exists then $\lim_{a \rightarrow 0} (h \circ \phi)(a)$ exists, and $\lim_{a \rightarrow 0} (h \circ \phi)(a) = h(x)$.

For an algebraic group G we denote by $Y(G)$ the set of cocharacters of G . For $\lambda \in Y(G)$ we define $P_\lambda := \{g \in G \mid \lim_{a \rightarrow 0} \lambda(a)g\lambda(a)^{-1} \text{ exists}\}$.

Lemma 2.2 [Bate et al. 2005, Lemma 2.4]. *Given a parabolic subgroup P of G and any Levi subgroup L of P , there exists a $\lambda \in Y(G)$ such that the following hold:*

- (i) $P = P_\lambda$.
- (ii) $L = L_\lambda := C_G(\lambda(k^*))$.
- (iii) The map $c_\lambda : P_\lambda \rightarrow L_\lambda$ defined by

$$c_\lambda(g) := \lim_{a \rightarrow 0} \lambda(a)g\lambda(a)^{-1}$$

is a surjective homomorphism of algebraic groups. Moreover, L_λ is the set of fixed points of c_λ and $R_u(P_\lambda)$ is the kernel of c_λ .

Conversely, given any $\lambda \in Y(G)$ the subset P_λ defined above is a parabolic subgroup of G , L_λ is a Levi subgroup of P_λ and the map c_λ as defined in (iii) has the described properties.

2.3. *G*-complete reducibility, products and epimorphisms. Let $f : G_1 \rightarrow G_2$ be a homomorphism of algebraic groups. We say that f is *nondegenerate* provided $(\ker f)^\circ$ is a torus; see [Serre 2004, Corollary 4.3]. In particular, f is nondegenerate if f is an isogeny.

Lemma 2.3 [Bate et al. 2005, Lemma 2.12]. *Let G_1 and G_2 be reductive groups.*

- (i) *Let H be a closed subgroup of $G_1 \times G_2$. Let $\pi_i : G_1 \times G_2 \rightarrow G_i$ be the canonical projection for $i = 1, 2$. Then H is $(G_1 \times G_2)$ -cr if and only if $\pi_i(H)$ is G_i -cr for $i = 1, 2$.*
- (ii) *Let $f : G_1 \rightarrow G_2$ be an epimorphism. Let H_1 and H_2 be closed subgroups of G_1 and G_2 , respectively.*
 - (a) *If H_1 is G_1 -cr, then $f(H_1)$ is G_2 -cr.*
 - (b) *If f is nondegenerate, then H_1 is G_1 -cr if and only if $f(H_1)$ is G_2 -cr, and H_2 is G_2 -cr if and only if $f^{-1}(H_2)$ is G_1 -cr.*

2.4. *Complete reducibility versus reductivity.* Any G -completely reducible subgroup H of G is reductive [Serre 2004, Proposition 4.1]. The converse is true in characteristic 0 — in particular, the answer to both Questions 1.1 and 1.2 is yes in this case — but is false in positive characteristic: for instance, a nontrivial finite unipotent subgroup of G is reductive but can never be G -cr [Serre 2004, Proposition 4.1]. The situation is somewhat nicer for connected H : the converse is true if p is sufficiently large. To be precise, we have the following theorem due to Serre.

Theorem 2.4 [Serre 2004, Theorem 4.4]. *Suppose $p \geq a(G)$ and $(H : H^\circ)$ is prime to p . Then H° is reductive if and only if H is G -completely reducible.*

Here the invariant $a(G)$ of G is defined as follows [Serre 2004, Section 5.2]. For G simple, set $a(G) = \text{rk}(G) + 1$, where $\text{rk}(G)$ is the rank of G . For G reductive, let $a(G) = \max(1, a(G_1), \dots, a(G_r))$, where G_1, \dots, G_r are the simple components of G . In the special case $G = \text{GL}(V)$ we have $a(G) = \dim(V)$, and a subgroup H of G is G -cr if and only if V is a semisimple H -module. We recover a basic result of Jantzen [1997, Proposition 3.2]: if $\rho : H \rightarrow \text{GL}(V)$ is a representation of a connected reductive group H and $p \geq \dim(V)$ then ρ is completely reducible. The finite unipotent example above shows that we cannot expect Theorem 2.4 to carry over completely, even with extra restrictions on p . Even when H is connected, H can fail to be G -cr if p is small.

Remark 2.5. Note that if we assume that K is connected then Theorem 1.3 follows immediately from Theorem 2.4, since $d(G) \geq a(G)$. We need a more elaborate proof with a worse bound on p , as we do not wish to place any restrictions on $(K : K^\circ)$; Theorem 1.3 is only of independent interest when $(K : K^\circ)$ is not prime to p . For an application in such an instance, see Corollary 1.5.

2.5. Coxeter numbers. The invariant $h(G)$ denotes the maximum of the *Coxeter numbers* of the simple components of G [Serre 2004, (5.1)], and we define $\tilde{h}(G)$ to be $h(G)$ if $[G, G]$ is simply connected and $h(G) + 1$ otherwise. Recall that if G is simple, we have $h(G) + 1 = \dim(G)/\text{rk}(G)$; the values of $h(G)$ for the various Dynkin types are as follows:

G	$A_n (n \geq 1)$	$B_n, C_n (n \geq 2)$	$D_n (n \geq 4)$	E_6	E_7	E_8	F_4	G_2
$h(G)$	$n + 1$	$2n$	$2n - 2$	12	18	30	12	6

We thus have

$$(2.6) \quad a(G) \leq h(G) \leq d(G)$$

for any G . But note that we can have $\tilde{h}(G) > d(G)$: for instance, take $G = \text{PGL}_n$.

3. G -complete reducibility, separability and reductive pairs

Now we consider the interaction of subgroups of G with the Lie algebra $\text{Lie } G = \mathfrak{g}$ of G . Much of this material is taken from [Bate et al. 2005].

Definition 3.1 [Bate et al. 2005, Definition 3.27]. For a closed subgroup H of G , the Lie subalgebra $\text{Lie}(C_G(H))$ is contained in $\mathfrak{c}_{\mathfrak{g}}(H)$, the fixed-point space of H on \mathfrak{g} in the adjoint action. In the case of equality (that is, if the scheme-theoretic centralizer of H in G is smooth), we say that H is *separable in G* ; else H is *nonseparable in G* . (See [Bate et al. 2005, Remark 3.32] for an explanation of the terminology.)

Of central importance is the following observation.

Example 3.2 [Bate et al. 2005, Example 3.28]. Any closed subgroup H of $G = \text{GL}(V)$ is separable in G .

Definition 3.3. Following Richardson [1967], we call (G, H) a *reductive pair* provided that H is a reductive subgroup of G and under the adjoint action, $\text{Lie } G$ decomposes as a direct sum

$$\text{Lie } G = \text{Lie } H \oplus \mathfrak{m},$$

for some H -submodule \mathfrak{m} .

For a list of examples of reductive pairs we refer to P. Slodowy [1997, I.3]. For further examples, see [Bate et al. 2005, Example 3.33, Remark 3.34].

Our application of reductive pairs to G -complete reducibility goes via the following result.

Proposition 3.4 [Bate et al. 2005, Corollary 3.36]. *Suppose that $(\text{GL}(V), H)$ is a reductive pair and K is a closed subgroup of H . If V is a semisimple K -module, then K is H -completely reducible.*

We look first at the special case of the adjoint representation.

Example 3.5 [Bate et al. 2005, Example 3.37]. Let H be a simple group of adjoint type and let $G = \mathrm{GL}(\mathrm{Lie } H)$. We have a symmetric nondegenerate Ad -invariant bilinear form on $\mathrm{Lie } G \cong \mathrm{End}(\mathrm{Lie } H)$ given by the usual trace form and its restriction to $\mathrm{Lie } H$ is just the Killing form of $\mathrm{Lie } H$. Since H is adjoint and Ad is a closed embedding, $\mathrm{ad} : \mathrm{Lie } H \rightarrow \mathrm{Lie } \mathrm{Ad}(H)$ is surjective. Thus it follows from the arguments in [Bate et al. 2005, Remark 3.34] that if the Killing form of $\mathrm{Lie } H$ is nondegenerate, then (G, H) is a reductive pair.

Suppose first that H is a simple classical group of adjoint type and $p > 2$. The Killing form is nondegenerate for $\mathfrak{sl}(V)$, $\mathfrak{so}(V)$, or $\mathfrak{sp}(V)$ if and only if p does not divide $2 \dim V$, $\dim V - 2$, or $\dim V + 2$, respectively; see [Bourbaki 1975, Example, Chapter VIII, Section 13.12]. In particular, for H adjoint of type A_n , B_n , C_n , or D_n , the Killing form is nondegenerate if $p > 2$ and p does not divide $n + 1$, $2n - 1$, $n + 1$, or $n - 1$, respectively.

Now suppose that H is a simple exceptional group of adjoint type. If p is good for H , then the Killing form of $\mathrm{Lie } H$ is nondegenerate; this was noted by Richardson [1967, Section 5]. Thus if p satisfies the appropriate condition, then $(\mathrm{GL}(\mathrm{Lie } H), H)$ is a reductive pair and Proposition 3.4 applies.

The next result is new, and gives a more general characterisation of reductive pairs.

Theorem 3.6. *For a simply connected simple algebraic group G in characteristic $p \geq 0$, consider the following conditions:*

- (i) $(\mathrm{GL}(V), \rho(G))$ is a reductive pair, where $\rho : G \rightarrow \mathrm{GL}(V)$ is a nontrivial representation of least dimension.
- (ii) $(\mathrm{GL}(V), \rho(G))$ is a reductive pair, for some nontrivial irreducible representation $\rho : G \rightarrow \mathrm{GL}(V)$.
- (iii) p is very good for G .
- (iv) $(\mathrm{GL}(\mathrm{Lie}(G)), \mathrm{Ad}(G))$ is a reductive pair.
- (v) The Killing form on $\mathrm{Lie}(G)$ is nondegenerate.
- (vi) p is very good for G , and if G has classical type, then $p \nmid e(G)$, where $e(G)$ is as follows:

G	A_n	B_n	C_n	D_n
$e(G)$	2	$2n - 1$	$n + 1$	$n - 1$

Then (i) \iff (ii) \iff (iii) \iff (iv) \iff (v) \iff (vi).

For G of exceptional type, all these conditions are equivalent.

Proof. It is clear that (i) implies (ii). If (ii) holds, then every subgroup of $\rho(G)$ is separable, since every subgroup of $\mathrm{GL}(V)$ is separable, by [Example 3.2](#), and this descends through reductive pairs. This means that p is pretty good for $\rho(G)$ — see [\[Herpel 2013, Definition 2.11\]](#) — which is the same as p being very good for $\rho(G)$ since G is simple. Note that p being very good is insensitive to the isogeny type of G , so (ii) implies (iii).

Next, the implication (iii) \implies (i) for G of types B_n , C_n and D_n is [\[Richardson 1967, Lemma 5.1\]](#). For $\mathrm{SL}(V)$, it is well known that the traceless matrices $\mathfrak{sl}(V)$ and the scalar matrices $\mathfrak{c}(\mathfrak{gl}(V))$ are the only proper, nonzero $\mathrm{GL}(V)$ -submodules (and $\mathrm{SL}(V)$ -submodules) of $\mathfrak{gl}(V)$. Now (iii) means that p is coprime to $\dim V$, which implies that these submodules intersect trivially, so that $\mathfrak{sl}(V)$ is complemented by $\mathfrak{c}(\mathfrak{gl}(V))$. So (i) holds in type A_n .

It remains to consider the exceptional cases $(G, V) = (G_2, V_7), (F_4, V_{26}), (E_6, V_{27}), (E_7, V_{56})$ and (E_8, V_{248}) , where V_j is a minimal-dimensional G -module in each case. Now $\mathrm{Lie}(\mathrm{GL}(V)) = V_j \otimes V_j^*$ contains $\mathrm{Lie}(G)$ as a submodule, which is irreducible since (iii) holds. The known weights of low-dimensional irreducible G -modules, given for instance in [\[Lübeck 2001\]](#), now allow one to calculate the weights of $V_j \otimes V_j^*$ and determine its G -composition factors. In each case, it transpires that $\mathrm{Lie}(G)$ is in fact the unique G -composition factor with a particular highest weight. Since $\mathrm{Lie}(G)$ is a submodule and $V_j \otimes V_j^*$ is self-dual, $\mathrm{Lie}(G)$ also occurs as a quotient of this, and the kernel of this quotient map is then a complement to $\mathrm{Lie}(G)$, so that (i) indeed holds. We illustrate the details in the case that G is of type G_2 in [Remark 3.7\(iv\)](#).

This shows that (i), (ii) and (iii) are equivalent. Now it is clear that (iv) \implies (ii). The equivalence of (v) and (vi) follows from [Example 3.5](#). If (v) holds then the Killing form on $\mathrm{Lie}(G)$ is, up to a nonzero scalar, the restriction of the trace form on $\mathrm{Lie}(\mathrm{GL}(\mathrm{Lie}(G)))$, and hence $\mathrm{Lie}(G)$ has an orthogonal complement, so that (iv) holds. Also, (iii) coincides with (vi) when G has exceptional type, which shows that all the conditions are equivalent in this case. \square

Remarks 3.7. (i) For exceptional groups in good characteristic, one can also check, just as in the case of the minimal module, that $\mathrm{Lie}(G)$ is the unique G -composition factor of $\mathrm{Lie}(G) \otimes \mathrm{Lie}(G)^*$ having a particular high weight, so that (being a submodule) it is a direct summand; this gives an alternative direct proof that (iii) \implies (iv) in [Theorem 3.6](#) for exceptional G .

(ii) For type A_n when $p = 2 \nmid n + 1$, so that $\mathrm{Lie}(G)$ is simple and self-dual but the Killing form vanishes, evidence suggests that $(\mathrm{GL}(\mathrm{Lie}(G)), G)$ is nevertheless a reductive pair. For instance, if $n = 2$ or 4 and $G = \mathrm{SL}_{n+1}$, the module $V = \mathrm{Lie}(G)$ appears with multiplicity 2 as a direct summand of $\mathfrak{gl}(V) = V \otimes V^*$; in fact we have $\mathfrak{gl}(V) \cong V \oplus V \oplus W$ for some indecomposable G -module W . With respect to the

trace form on $\mathfrak{gl}(V)$, the image of the embedding $\mathrm{Lie}(G) \rightarrow \mathfrak{gl}(V)$ gives a totally isotropic subspace isomorphic to V , and the natural isomorphism $\mathfrak{gl}(V)/V^\perp \rightarrow V^*$ then shows the existence of a second composition factor isomorphic to V . One can then use the self-duality of V and $\mathfrak{gl}(V)$ to deduce that $\mathrm{Lie}(G)$ and this second composition factor are each a direct summand of $\mathfrak{gl}(V)$.

(iii) For types B_n, C_n, D_n with p odd but dividing $2n - 1, n + 1, n - 1$, respectively, considering such instances in rank up to 6 suggests that $(\mathrm{GL}(\mathrm{Lie}(G)), G)$ is never a reductive pair. In each case, it transpires that $\mathrm{Hom}(\mathrm{Lie}(G), \mathfrak{gl}(\mathrm{Lie}(G)))$ is 1-dimensional. Thus $\mathfrak{gl}(\mathrm{Lie}(G))$ has a unique G -submodule isomorphic to $\mathrm{Lie}(G)$, which turns out to lie in a self-dual indecomposable G -module direct summand of $\mathfrak{gl}(\mathrm{Lie}(G))$ also having $\mathrm{Lie}(G)$ as its head.

(iv) To illustrate the argument in the proof above, when G has type G_2 and $p \neq 2, 3$, the G -module V_7 is irreducible of highest weight λ_2 , and the 49-dimensional module $V_7 \otimes V_7^*$ has high weights $0, \lambda_1, \lambda_2$ and $2\lambda_2$ when $p \neq 7$; or $0, 0, \lambda_1, \lambda_2$ and $2\lambda_2$ when $p = 7$. In either case, we find a unique composition factor of high weight λ_1 , which is $\mathrm{Lie}(G)$.

4. Proofs of Theorem 1.3 and Corollary 1.5

Proof of Theorem 1.3. Let $\pi : G \rightarrow G/Z(G)^\circ$ be the canonical projection. Owing to Lemma 2.3(ii)(b), we can replace G with $G/Z(G)^\circ$, so without loss we can assume that G is semisimple. Let G_1, \dots, G_r be the simple factors of G . Multiplication gives an isogeny from $G_1 \times \dots \times G_r$ to G . Again by Lemma 2.3(ii)(b), we can replace G with $G_1 \times \dots \times G_r$, so we can assume G is the product of its simple factors. By Lemma 2.3(i) it is thus enough to prove the result when G is simple and simply connected. Of course, as well as replacing G with its (pre)image under an isogeny, we also replace H and K with their (pre)images under that isogeny along the way.

First suppose G is of type A . Then $K \subseteq H \subseteq \mathrm{SL}(V) \subseteq \mathrm{GL}(V)$. By [Bate et al. 2011, Theorem 1.3], if K is H -cr then K is $\mathrm{GL}(V)$ -cr, so K is $\mathrm{SL}(V)$ -cr and we're done. Next suppose G is not of type A . Then, as $p \geq d(G)$, p is good for G . It follows from Theorem 3.6 that $(\mathrm{GL}(V), G)$ is a reductive pair, where V is an irreducible G -module of least dimension. Since $p \geq d(G) = \dim V$, V is semisimple for K , thanks to [Bate et al. 2011, Theorem 1.3], and therefore K is G -cr, by Proposition 3.4. \square

The following is an immediate consequence of [Springer and Steinberg 1970, III 1.19(a)] and [Bate et al. 2022, Theorem 1.3].

Lemma 4.1. *Let σ be a Steinberg endomorphism of G . Then G_σ is G -irreducible.*

Proof of Corollary 1.5. By Lemma 4.1, H_σ is H -ir. The result now follows from Theorem 1.3. \square

Note that [Corollary 1.5](#) is false without the bound on p . See [Example 4.5](#) for an instance when H is G -cr but H_σ is not, where $p = 3 < 8 = d(G)$.

Our next result gives a particular set of conditions on H_σ to guarantee that H_σ and H belong to the same parabolic subgroups and the same Levi subgroups of G . Note that, if $\sigma : H \rightarrow H$ is a Steinberg endomorphism of H , then σ stabilises a maximal torus of H [[Steinberg 1968](#), Corollary 10.10]. Also, for S a torus in G , we have $C_G(S) = C_G(s)$ for some $s \in S$; see [[Borel 1991](#), III Proposition 8.18].

Proposition 4.2. *Let $H \subseteq G$ be connected reductive groups. Let $\sigma : G \rightarrow G$ be a Steinberg endomorphism that stabilises H and a maximal torus T of H . Suppose*

- (i) $C_G(T) = C_G(t)$ for some $t \in T_\sigma$, and
- (ii) H_σ meets every T -root subgroup of H nontrivially.

Then H_σ and H are contained in precisely the same parabolic subgroups of G , and the same Levi factors thereof. In particular, H is G -completely reducible if and only if H_σ is G -completely reducible; similarly, H is G -irreducible if and only if H_σ is G -irreducible.

Proof. First assume $H_\sigma \subseteq P$ for some parabolic subgroup P of G . Then t lies in some maximal torus of P . So we can find a $\lambda \in Y(G)$ such that $P = P_\lambda$ and λ centralizes t . But then λ centralizes T , by (i), so $T \subseteq P$. Now we have a maximal torus T of H and a nontrivial part of each T -root group of H inside P , by (ii), so we can conclude that all of H belongs to P . Similarly, if $H_\sigma \subseteq L_\lambda$ for some $\lambda \in Y(G)$, we get $H \subseteq L_\lambda$. The reverse conclusions are obvious, since $H_\sigma \subseteq H$. \square

In the presence of the conditions in [Proposition 4.2](#) we can improve the bound in [Corollary 1.5](#) considerably; the following is immediate from [Theorem 2.4](#) and [Proposition 4.2](#).

Corollary 4.3. *Suppose G , H and σ satisfy the hypotheses of [Proposition 4.2](#). Suppose in addition that $p \geq a(G)$. Then H_σ is G -completely reducible.*

Note that condition (ii) in [Proposition 4.2](#) is automatically satisfied provided σ induces a standard Frobenius endomorphism on H . In that case [Example 4.4](#) below demonstrates that condition (i) above does hold generically. Nevertheless, [Example 4.5](#) shows that [Proposition 4.2](#) is false in general without condition (i) even when part (ii) is fulfilled.

The following example shows that the conditions in [Proposition 4.2](#) do hold generically.

Example 4.4. Let $\sigma_q : \mathrm{GL}(V) \rightarrow \mathrm{GL}(V)$ be a standard Frobenius endomorphism that stabilises the connected reductive subgroup H of $\mathrm{GL}(V)$ and a maximal torus T of H . Pick $l \in \mathbb{N}$ so that firstly all the different T -weights of V are still distinct when restricted to $T_{\sigma_q^l}$ and secondly that there is a $t \in T_{\sigma_q^l}$, such that $C_{\mathrm{GL}(V)}(T) = C_{\mathrm{GL}(V)}(t)$.

Then for every $n \geq l$, both conditions in [Proposition 4.2](#) are satisfied for $\sigma = \sigma_q^n$. Thus there are only finitely many powers of σ_q for which part (i) can fail. The argument here readily generalises to a Steinberg endomorphism of a connected reductive G which induces a generalized Frobenius morphism on H .

In contrast to the setting in [Example 4.4](#), our next example demonstrates that the conclusion of [Proposition 4.2](#) may fail, if condition (i) is not satisfied. Consequently, the conditions in [Theorem 1.3](#) and [Corollary 1.5](#) are needed in general.

Example 4.5. Let $p = 3$, $q = 9$ and $H = \mathrm{SL}_2$. By Steinberg's tensor product theorem, the simple H -module $V = L(1 + q + q^2)$ is isomorphic to $L(1) \otimes L(1)^{[2]} \otimes L(1)^{[4]}$, the superscripts denoting p -power twists. Thus, after identifying H with its image in $G = \mathrm{GL}(V)$, we see that H is G -cr. Let $\sigma = \sigma_q$ be the standard Frobenius on G . Then $H_\sigma = \mathrm{SL}_2(9)$ is H -cr, by [Lemma 4.1](#). Now as an H_σ -module, V is isomorphic to the H -module $L(1) \otimes L(1) \otimes L(1)$ which admits the nonsimple indecomposable Weyl module of highest weight 3 as a constituent. As the latter is not semisimple for H_σ , V is not semisimple as an H_σ -module and so H_σ is not G -cr.

5. Saturation

Let $u \in \mathrm{GL}(V)$ be unipotent of order p . Then there is a nilpotent element $\epsilon \in \mathrm{End}(V)$ with $\epsilon^p = 0$ such that $u = 1 + \epsilon$. For $t \in \mathbb{G}_a$ we define u^t by

$$(5.1) \quad u^t := (1 + \epsilon)^t = 1 + t\epsilon + \binom{t}{2}\epsilon^2 + \cdots + \binom{t}{p-1}\epsilon^{p-1};$$

see [\[Serre 1998\]](#). Then $\{u^t \mid t \in \mathbb{G}_a\}$ is a closed connected subgroup of $\mathrm{GL}(V)$ isomorphic to \mathbb{G}_a .

Following [\[Nori 1987\]](#) and [\[Serre 1998\]](#), a subgroup H of $\mathrm{GL}(V)$ is *saturated* provided H is closed and for any unipotent element u of H of order p and any $t \in \mathbb{G}_a$ also u^t given by (5.1) belongs to H . The *saturated closure* H^{sat} of H is the smallest saturated subgroup of $\mathrm{GL}(V)$ containing H .

We now recall a notion of saturation for arbitrary connected reductive groups which generalises the one just given for $\mathrm{GL}(V)$. Suppose that $p \geq h(G)$. Then every unipotent element of G has order p ; see [\[Testerman 1995\]](#). Let u be a unipotent element of G . Then for $t \in \mathbb{G}_a$ there is a canonical “ t -th power” u^t of u such that the map $t \mapsto u^t$ defines a homomorphism of the additive group \mathbb{G}_a into G . We recall some results from Serre [\[1998; 2004, Section 5\]](#).

Let \mathcal{U} be the subvariety of G consisting of all unipotent elements of G , and let \mathcal{N} be the subvariety of $\mathrm{Lie}(G)$ consisting of all nilpotent elements of $\mathrm{Lie}(G)$. Fix a maximal torus T of G , a Borel subgroup B of G containing T , and let U be the unipotent radical of B . Since $p \geq h(G)$ and because the nilpotency class of $\mathrm{Lie}(U)$ is at most $h(G)$, we can view $\mathrm{Lie}(U)$ as an algebraic group with multiplication

given by the Baker–Campbell–Hausdorff formula (see [Bourbaki 1972, Chapter II, Section 6]).

Let $\Phi = \Phi(G, T)$ be the root system of G with respect to T . For α a root in Φ , let $x_\alpha : \mathbb{G}_a \rightarrow U_\alpha$ be a parametrization of the root subgroup U_α of G . Let $X_\alpha := \frac{d}{ds}(x_\alpha(s))|_{s=0}$ be a canonical generator of $\text{Lie}(U_\alpha)$. Further, by $\text{Aut}(G)$ we denote the group of algebraic automorphisms of G . We begin with the following result due to Serre; for a detailed proof, see [Balaji et al. 2017, Section 6]. Recall that we define $\tilde{h}(G)$ to be $h(G)$ if $[G, G]$ is simply connected and $h(G) + 1$ otherwise.

Theorem 5.2 [Serre 1998, Theorem 3]. *Let $p \geq \tilde{h}(G)$. There is a unique isomorphism of varieties $\log : \mathcal{U} \rightarrow \mathcal{N}$ such that the following hold:*

- (i) $\log(\sigma u) = d\sigma(\log u)$ for any $\sigma \in \text{Aut}(G)$ and any $u \in \mathcal{U}$.
- (ii) *The restriction of \log to U defines an isomorphism of algebraic groups $U \rightarrow \text{Lie}(U)$ whose tangent map is the identity on $\text{Lie}(U)$.*
- (iii) $\log(x_\alpha(t)) = tX_\alpha$ for any $\alpha \in \Phi$ and any $t \in \mathbb{G}_a$.

Let $\exp : \mathcal{N} \rightarrow \mathcal{U}$ be the inverse morphism to \log . We then define

$$(5.3) \quad u^t := \exp(t \log u),$$

for any $u \in \mathcal{U}$ and any $t \in \mathbb{G}_a$.

Definition 5.4 [Nori 1987; Serre 1998]. Let $p \geq \tilde{h}(G)$. A subgroup H of G is *saturated* (in G) provided H is closed and for any unipotent element u of H and any $t \in \mathbb{G}_a$ also u^t belongs to H . For a subgroup H , its *saturated closure* H^{sat} is the smallest saturated subgroup of G containing H .

We give various fairly straightforward consequences of Theorem 5.2. The third is already recorded in [Serre 1998] for centralizers of subgroups of G .

Corollary 5.5. *Let $p \geq \tilde{h}(G)$. Let $\sigma \in \text{Aut}(G)$. Then the following hold:*

- (i) $\sigma(u^t) = \sigma(u)^t$ for any $u \in \mathcal{U}$ and $t \in \mathbb{G}_a$.
- (ii) *If H is a σ -stable subgroup of G , so is H^{sat} .*
- (iii) *For S a subgroup of $\text{Aut}(G)$, $C_G(S)$ is saturated in G .*

Proof. (i) Since \exp is the inverse to \log , Theorem 5.2(i) gives $\sigma(\exp(X)) = \exp(d\sigma(X))$ for all $X \in \mathcal{N}$. Hence for any $u \in \mathcal{U}$ and $t \in \mathbb{G}_a$,

$$\begin{aligned} \sigma(u^t) &= \sigma(\exp(t \log u)) \\ &= \exp(d\sigma(t \log(u))) = \exp(td\sigma(\log u)) = \exp(t \log \sigma(u)) = \sigma(u)^t. \end{aligned}$$

(ii) If H is σ -stable and M is any saturated subgroup of G containing H , then so is $\sigma(M)$: for, if $u \in \sigma(M)$ is unipotent, then $u = \sigma(v)$ for some $v \in M$ unipotent. Then $u^t = \sigma(v)^t = \sigma(v^t) \in \sigma(M)$ also, by (i) and the fact that M is saturated in G .

Hence H^{sat} , the unique smallest saturated subgroup of G containing H , must also be σ -stable.

(iii) This is immediate by (i). \square

As particular instances of [Corollary 5.5\(iii\)](#), we note that centralizers of graph automorphisms of G are saturated, and also Levi subgroups of parabolic subgroups of G are saturated, since they arise as centralizers of tori in G .

One can use [Theorem 5.2](#) directly to show that parabolic subgroups of G are saturated. The following proof instead uses the language of cocharacters, which also allows us to observe that the process of “taking limits along cocharacters” commutes with saturation.

Proposition 5.6. *Let $\lambda \in Y(G)$ and let $P = P_\lambda$. Then for $u \in P$ unipotent and $v := \lim_{a \rightarrow 0} \lambda(a)u\lambda(a)^{-1}$, we have $\lim_{a \rightarrow 0} \lambda(a)u^t\lambda(a)^{-1} = v^t$. In particular, P is saturated.*

Proof. Observe that for any $t \in \mathbb{G}_a$ the map $h_t : \mathcal{U} \rightarrow \mathcal{U}$ given on points by $h_t(u) := u^t = \exp(t \log(u))$ is an isomorphism of varieties, by [Theorem 5.2](#). Furthermore, by [Corollary 5.5\(i\)](#) we have, for any $a \in k^*$,

$$(\lambda(a)u\lambda(a)^{-1})^t = \lambda(a)u^t\lambda(a)^{-1}.$$

The result now follows from [Remark 2.1](#) and elementary limit calculations. \square

The next result is a slight refinement of a theorem due to Serre.

Theorem 5.7 [[Serre 1998](#), Property 2, Theorem 4]. *Let $p \geq \tilde{h}(G)$. Let H be a saturated connected reductive subgroup of G , and suppose $p \geq \tilde{h}(H)$. Then for any $u \in H$ unipotent, the element u^t , with respect to H , coincides with u^t , with respect to G : that is, saturation in H coincides with saturation in G .*

Remark 5.8. It follows from [[Serre 1998](#), Theorem 4] that if H is any connected reductive subgroup of G — not necessarily saturated — then $h(H) \leq h(G)$. It can happen, however, that $\tilde{h}(H) > \tilde{h}(G) = p$. For instance, let p be an odd prime and let $G = \text{SL}_p \times \text{SL}_p$. Let M be a semisimple subgroup of SL_p such that M is not simply connected, and let $H = \text{SL}_p \times M$. Now $\tilde{h}(G) = h(G) = h(\text{SL}_p) = p$ and $h(H) \geq h(\text{SL}_p) = p$; but H is not simply connected, so $\tilde{h}(H) = h(H) + 1 \geq p + 1 > p = \tilde{h}(G)$. Because of this we have added the hypotheses that $p \geq \tilde{h}(G)$ and $p \geq \tilde{h}(H)$ to [Theorem 5.7](#), although they were not stated explicitly in [[Serre 1998](#), Property 2]. The proof there still holds.

Remark 5.9. Suppose $p \geq \tilde{h}(G)$ and let H be a connected reductive subgroup of G normalized by some maximal torus T of G . We claim that H is saturated in G . Since H and HT contain the same unipotent elements, H is saturated if and only if HT is saturated, so we may assume that H contains T . By [Corollary 5.5\(iii\)](#)

and [Theorem 5.7](#) there is no harm in passing to a minimal Levi subgroup of G containing H , so we may assume that H has maximal semisimple rank in G . Since $p \geq \tilde{h}(G)$, p is good for G , and hence H arises as the centralizer $H = C_G(s)^\circ$ in G of some semisimple element s of G , thanks to Deriziotis' criterion; see [\[Humphreys 1995, Section 2.15\]](#). Thus H is saturated by [Corollary 5.5\(iii\)](#).

Note that this result applies to any connected normal subgroup of G , so in particular to the simple factors of G .

Next we show that saturation is compatible with direct products, in the following sense.

Lemma 5.10. *Suppose $G = G_1 \times \cdots \times G_r$, where each G_i is connected and reductive. Suppose $p \geq \tilde{h}(G)$. Then $p \geq \tilde{h}(G_i)$ and G_i is saturated for $1 \leq i \leq r$. Moreover, if $u_i \in G_i$ is unipotent for $1 \leq i \leq r$ then*

$$\log(u_1 \cdots u_r) = \log_1(u_1) + \cdots + \log_r(u_r),$$

where \log denotes the logarithm map for G and \log_i denotes the logarithm map for G_i .

Proof. For each i , any simple factor of G_i is also a simple factor of G and $[G_i, G_i]$ is simply connected if $[G, G]$ is simply connected, so $\tilde{h}(G) \geq \tilde{h}(G_i)$ and the first assertion follows. Each G_i is normal in G and hence is saturated by [Remark 5.9](#). [Theorem 5.7](#) implies that if $u_i \in G_i$ is unipotent then $\log_i(u_i) = \log(u_i)$. If X and X' are commuting nilpotent elements of $\text{Lie}(G)$ then the Baker–Campbell–Hausdorff product of X and X' is just $X + X'$, so $\exp(X + X') = \exp(X)\exp(X')$. The final assertion now follows easily. \square

Corollary 5.11. *Assume the hypotheses of [Lemma 5.10](#), and let $\pi_i : G \rightarrow G_i$ be the canonical projection. If H is a saturated subgroup of G then $\pi_i(H)$ is a saturated subgroup of G_i .*

Proof. This follows immediately from [Lemma 5.10](#). \square

Proposition 5.12 [\[Serre 2004, Proposition 5.2\]](#). *If H is saturated in G , then $(H : H^\circ)$ is prime to p .*

Theorem 5.13 [\[Serre 2004, Theorem 5.3\]](#). *Let $p \geq \tilde{h}(G)$. For a closed subgroup H of G , the following are equivalent:*

- (i) H is G -completely reducible.
- (ii) H^{sat} is G -completely reducible.
- (iii) $(H^{\text{sat}})^\circ$ is reductive.

The equivalence between (i) and (ii) stems from the fact that both parabolic and Levi subgroups of G are saturated. The equivalence between (ii) and (iii) is an

immediate consequence of [Theorem 2.4](#) and [Proposition 5.12](#), since $\tilde{h}(G) \geq a(G)$ by (2.6).

Proposition 5.14. *Let $K \subseteq H$ be closed subgroups of G with H connected reductive and saturated in G . Suppose $p \geq \tilde{h}(G)$ and $p \geq \tilde{h}(H)$. Then K is H -completely reducible if and only if K is G -completely reducible.*

Proof. By [Theorem 5.13](#), K is H -cr if and only if $(K^{\text{sat}})^{\circ}$ is reductive, where we saturate in H . But saturation in H is the same as saturation in G thanks to [Theorem 5.7](#), so $(K^{\text{sat}})^{\circ}$ is reductive if and only if K is G -cr, again by [Theorem 5.13](#). \square

Note that both implications in the equivalence in [Proposition 5.14](#) may fail if $p < h(G)$; e.g., see [\[Bate et al. 2005, Example 3.45\]](#) and [\[Bate et al. 2010, Proposition 7.17\]](#).

For ease of reference, we recall a connectedness result for H^{sat} from [\[Bate et al. 2011, Corollary 4.2\]](#).

Remark 5.15. Let $p \geq \tilde{h}(G)$. If H is a closed connected subgroup of G , then so is H^{sat} . For, consider the subgroup M of G generated by H and the closed connected subgroups $\{u^t \mid t \in \mathbb{G}_a\} \cong \mathbb{G}_a$ of G for each unipotent element $u \in G$. Then M is connected. By definition, $M \subseteq H^{\text{sat}}$. If $M \neq H^{\text{sat}}$, then by repeating this process with M (possibly several times), we eventually generate all of H^{sat} by H and closed connected subgroups of G isomorphic to \mathbb{G}_a .

Here is a further consequence of [Theorem 5.13](#).

Corollary 5.16. *Let $p \geq \tilde{h}(G)$. Let $K \subseteq H$ be closed subgroups of G with H connected reductive, and suppose that $p \geq \tilde{h}(H)$. Then the following are equivalent:*

- (i) K is H^{sat} -completely reducible.
- (ii) K^{sat} is H^{sat} -completely reducible.
- (iii) $(K^{\text{sat}})^{\circ}$ is reductive.
- (iv) K^{sat} is G -completely reducible.
- (v) K is G -completely reducible.

Proof. Owing to [Remark 5.15](#), H^{sat} is connected. Further, since $\tilde{h}(G) \geq a(G)$, it follows from [Theorems 2.4](#) and [5.13](#) that H^{sat} is reductive.

The equivalence of (i) through (iii) follows from [Theorem 5.13](#) applied to $K \subseteq H^{\text{sat}}$ and [Theorem 5.7](#) and the equivalence of (iii) through (v) is just [Theorem 5.13](#). \square

Proof of Theorem 1.4. Thanks to [Remark 5.15](#), H^{sat} is connected. Since $d(G) \geq h(G) \geq a(G)$ by (2.6), it follows from [Theorems 2.4](#) and [5.13](#) that H^{sat} is reductive.

If K is H -cr, then K is G -cr, by [Theorem 1.3](#), and therefore K^{sat} is H^{sat} -cr, by [Corollary 5.16](#). \square

The following example illustrates that in general connected reductive subgroups are not saturated.

Example 5.17. With the explicit notion of saturation from (5.1) within $\mathrm{GL}(V)$ it is easy to check that the image H of the adjoint representation of SL_p in $G := \mathrm{GL}(\mathrm{Lie}(\mathrm{SL}_p))$ is not saturated in characteristic p ; see [Serre 1998, p. 18]. Evidently, H is contained in the maximal parabolic subgroup P of G that stabilises the H -submodule $\mathfrak{z}(\mathrm{Lie}(\mathrm{SL}_p))$. One checks that its saturation H^{sat} in G includes all of H but also part of the unipotent radical $R_u(P)$ of P . For instance, when $p = 2$ then the adjoint representation of $H := \mathrm{SL}_2$ with respect to a suitable basis is given by

$$\mathrm{Ad} \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = \begin{pmatrix} a^2 & b^2 & 0 \\ c^2 & d^2 & 0 \\ ac & bd & 1 \end{pmatrix}.$$

If

$$u = \mathrm{Ad} \left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \right),$$

then

$$\log u = \begin{pmatrix} 0 & b^2 & 0 \\ 0 & 0 & 0 \\ 0 & b & 0 \end{pmatrix},$$

so

$$u^t = \exp(t \log u) = \begin{pmatrix} 1 & tb^2 & 0 \\ 0 & 1 & 0 \\ 0 & tb & 1 \end{pmatrix} \quad \text{for any } t \in \mathbb{G}_a.$$

We see that if $b \neq 0, 1$ then

$$u^{b^2} \mathrm{Ad} \left(\begin{pmatrix} 1 & b^2 \\ 0 & 1 \end{pmatrix} \right) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & b^2 + b^3 & 1 \end{pmatrix}$$

is a nontrivial element of $H^{\mathrm{sat}} \cap R_u(P)$, where P is the parabolic subgroup of matrices of shape

$$\begin{pmatrix} * & * & 0 \\ * & * & 0 \\ * & * & * \end{pmatrix}.$$

Since the abelian unipotent radical $R_u(P)$ is an irreducible SL_p -module (of highest weight $\lambda_1 + \lambda_{p-1}$, or $2\lambda_1$ when $p = 2$), being a nonzero SL_p -submodule of $R_u(P)$, U is in fact all of $R_u(P)$. So H^{sat} is of the form $H^{\mathrm{sat}} = XR_u(P)$, where X is a subgroup of the Levi subgroup of type SL_{p^2-2} of P . In particular, H^{sat} is not reductive in this case.

We briefly revisit Example 4.5 in the context of saturation.

Example 5.18. With the hypotheses and notation from [Example 4.5](#), a nontrivial unipotent element u from H_σ has order $p = 3$, so we can saturate u in H and in G , according to [\(5.1\)](#) above. Likewise we can saturate nontrivial unipotent elements in H . It turns out that H is not saturated in G . Let $u = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$ be in $\mathrm{SL}_2(9) = H_\sigma$ for a fixed $b \neq 1$. Then one can check that the saturations of u in H and in G do not coincide. Thus, the hypotheses of [Theorem 5.7](#) fail on two accounts, for H is not saturated in G and $p = 3 < h(G) = \dim V = 8$, while $p > h(H) = 2$.

6. Saturation and finite groups of Lie type

In this section we discuss finite subgroups of Lie type in G and their behaviour under saturation. To do this we need to prove the compatibility of the saturation map with standard Frobenius endomorphisms. First recall that if $\sigma_q : G \rightarrow G$ is a standard q -power Frobenius endomorphism of G , then there exists a σ_q -stable maximal torus T and Borel subgroup $B \supseteq T$, and with respect to a chosen parametrization of the root groups as above, we have $\sigma_q(x_\alpha(s)) = x_\alpha(s^q)$ for each $\alpha \in \Phi$ and $s \in \mathbb{G}_a$; see [\[Gorenstein et al. 1998, Theorem 1.15.4\(a\)\]](#).

Proposition 6.1. *Let $p \geq \tilde{h}(G)$. Suppose that $\sigma_q : G \rightarrow G$ is a standard q -power Frobenius endomorphism of G . Then the following hold:*

- (i) $\sigma_q(u^t) = \sigma_q(u)^{t^q}$ for any $u \in \mathcal{U}$ and $t \in \mathbb{G}_a$.
- (ii) If H is a σ_q -stable subgroup of G , then H^{sat} is also σ_q -stable.

Proof. (i) Fix a σ_q -stable Borel subgroup B of G as in the discussion before the statement of the proposition, with unipotent radical U . Since $(gug^{-1})^t = gu^t g^{-1}$ for all $u \in \mathcal{U}$ and $g \in G$ thanks to [Corollary 5.5\(i\)](#), it is enough to show the result for $u \in U$.

There are two ways to define a Frobenius-type map on $\mathrm{Lie}(U)$. Firstly, since the X_α form a basis for $\mathrm{Lie}(U)$ as a k -space, we have the map $F_q : \mathrm{Lie}(U) \rightarrow \mathrm{Lie}(U)$ given by

$$\sum_{\alpha \in \Phi^+} c_\alpha X_\alpha \mapsto \sum_{\alpha \in \Phi^+} c_\alpha^q X_\alpha.$$

For this map, it is clear that $F_q(tX) = t^q F_q(X)$ for every $t \in \mathbb{G}_a$ and $X \in \mathrm{Lie}(U)$. Alternatively, since \exp and \log are mutually inverse group isomorphisms between U and $\mathrm{Lie}(U)$, there is some endomorphism $f_q : \mathrm{Lie}(U) \rightarrow \mathrm{Lie}(U)$ defined by

$$\sigma_q(\exp(X)) = \exp(f_q(X))$$

for all $X \in \mathrm{Lie}(U)$ (or, equivalently, by $\log \sigma_q(u) = f_q(\log u)$ for all $u \in U$). We claim that $f_q = F_q$.

First, note that [Theorem 5.2\(iii\)](#) gives equality straight away for multiples of basis elements: since $\sigma_q(x_\alpha(s)) = x_\alpha(s^q)$ for each $s \in \mathbb{G}_a$ and positive root α , we

have $f_q(sX_\alpha) = s^q X_\alpha = F_q(sX_\alpha)$ for all such s and α . Now recall that if we fix some ordering of the positive roots Φ^+ , then each element $u \in U$ has a unique expression as a product $u = \prod_{\alpha \in \Phi^+} x_\alpha(s_\alpha)$ with the $s_\alpha \in \mathbb{G}_a$ — see [Springer and Steinberg 1970, Chapter I, 1.2(b)] — and hence every $X = \log(u) \in \text{Lie}(U)$ has expression $X = \prod_{\alpha \in \Phi^+} (s_\alpha X_\alpha)$, where this product is calculated using the Baker–Campbell–Hausdorff formula, by Theorem 5.2(ii) and (iii). Thus, to show $f_q = F_q$ it is enough to show that F_q is a group homomorphism.

Let $X = \sum_{\alpha \in \Phi^+} s_\alpha X_\alpha$ and $Y = \sum_{\alpha \in \Phi^+} t_\alpha X_\alpha$ be two elements of $\text{Lie}(U)$. In calculating XY with the Baker–Campbell–Hausdorff formula we get a number of commutators involving the $s_\alpha X_\alpha$ and $t_\beta X_\beta$ for positive roots α and β . Since the Lie bracket is bilinear, we can pull all the coefficients s_α and t_β out to the front of each commutator, and hence write XY as a linear combination of commutators in the X_α . All such commutators of degree greater than 1 can be rewritten in $\text{Lie}(U)$ as a linear combination of the X_α by applying the commutator relations recursively to write any $[X_\beta, X_\gamma]$ in terms of the X_α , and then expanding out and repeating. The coefficients appearing in the commutation relations lie in the finite base field \mathbb{F}_p , and hence are fixed under the q -power map. Thus, for any commutator C in the root elements X_α , we may conclude that $F_q(C) = C$. This is enough to conclude that $F_q(XY) = F_q(X)F_q(Y)$ for any $X, Y \in \text{Lie}(U)$, as claimed.

We can now deduce that for any $t \in \mathbb{G}_a$ and any $X \in \text{Lie}(U)$, we have $f_q(tX) = t^q f_q(X)$, and hence for any $u \in U$ and any $t \in \mathbb{G}_a$,

$$\begin{aligned} \sigma_q(u)^{t^q} &= \exp(t^q \log \sigma_q(u)) = \exp(t^q f_q(\log u)) \\ &= \exp(f_q(t \log u)) = \sigma_q(\exp(t \log u)) = \sigma_q(u^t), \end{aligned}$$

which completes the proof of (i).

(ii) This follows quickly from (i), since if H is σ_q -stable and M is any saturated subgroup of G containing H , then $\sigma_q(M)$ is another saturated subgroup of G containing H : for, if $t \in \mathbb{G}_a$ and $u \in \sigma_q(M)$ is unipotent, then we may find $s \in \mathbb{G}_a$ with $s^q = t$ (since $k = \bar{k}$ is perfect) and $v \in M$ which is unipotent such that $u = \sigma_q(v)$. Then $u^t = \sigma_q(v)^{s^q} = \sigma_q(v^s) \in \sigma_q(M)$ also. Hence H^{sat} , which is the smallest saturated subgroup containing H , must also be σ_q -stable. \square

Combining Corollary 5.5 and Proposition 6.1, we obtain the following.

Corollary 6.2. *Let $p \geq \tilde{h}(G)$. Suppose $\sigma : G \rightarrow G$ is a Steinberg endomorphism of G such that $\sigma = \tau \sigma_q$, where $\tau \in \text{Aut}(G)$ and σ_q is a standard q -power Frobenius endomorphism of G . Then the following hold:*

- (i) $\sigma(u^t) = \sigma(u)^{t^q}$ for any $u \in U$ and $t \in \mathbb{G}_a$.
- (ii) If H is a σ -stable subgroup of G , then H^{sat} is also σ -stable.

Proof. (i) By [Corollary 5.5\(i\)](#) and [Proposition 6.1\(i\)](#), we have for any $u \in \mathcal{U}$, $t \in \mathbb{G}_a$,

$$\sigma(u^t) = \tau(\sigma_q(u^t)) = \tau(\sigma_q(u)^{t^q}) = \tau(\sigma_q(u))^{t^q} = \sigma(u)^{t^q},$$

as desired.

(ii) This part follows from the arguments in the proofs of [Corollary 5.5\(ii\)](#) and [Proposition 6.1\(ii\)](#) along with part (i). \square

We note that in general, a Steinberg endomorphism of a reductive group G need not be of the form given in [Corollary 6.2](#); e.g., see [Example 6.8](#).

We now apply [Theorem 1.4](#) to the case when $K = H_\sigma$ for a Frobenius endomorphism σ of G . The next result is immediate from [Lemma 4.1](#) and [Theorem 1.4](#).

Corollary 6.3. *Let $H \subseteq G$ be connected reductive groups. Suppose $p \geq d(G)$ and $p \geq \tilde{h}(G)$. Let $\sigma : G \rightarrow G$ be a Steinberg endomorphism that stabilises H . Then $(H_\sigma)^{\text{sat}}$ is H^{sat} -completely reducible.*

We can potentially improve the bound on p in the last corollary at the expense of imposing the conditions from [Proposition 4.2](#), as follows.

Corollary 6.4. *Suppose G , H and σ satisfy the hypotheses of [Proposition 4.2](#). Suppose in addition that $p \geq \tilde{h}(G)$. Then $(H_\sigma)^{\text{sat}}$ is H^{sat} -completely reducible.*

Proof. Since $p \geq \tilde{h}(G) \geq a(G)$ by (2.6), H is G -cr, by [Theorem 2.4](#). Thus H_σ is G -cr, by [Proposition 4.2](#). The result now follows from [Corollary 5.16](#). \square

Note that in [Theorem 1.4](#) and [Corollaries 6.3](#) and [6.4](#), H^{sat} is again connected reductive. This follows from the fact that $\tilde{h}(G) \geq a(G)$ and [Theorems 2.4](#), [5.13](#) and [Remark 5.15](#).

[Example 4.4](#) shows that generically the conditions of [Corollary 6.4](#) are fulfilled. Nevertheless, [Example 4.5](#) and [Corollary 5.16](#) show that [Corollary 6.4](#) is false if condition (i) of [Proposition 4.2](#) is not satisfied. In the settings of [Corollaries 6.3](#) and [6.4](#), $((H_\sigma)^{\text{sat}})^\circ$ is reductive.

Assume G is simple for the rest of this section unless specified otherwise. Let $\sigma : G \rightarrow G$ be a Steinberg endomorphism. Then σ is a generalized Frobenius map, i.e., a suitable power of σ is a standard Frobenius map (e.g., see [[Gorenstein et al. 1998](#), Theorem 2.1.11]), and the possibilities for σ are well known [[Steinberg 1968](#), Section 11]: σ is conjugate to either σ_q , $\tau\sigma_q$, $\tau'\sigma_q$ or τ' , where σ_q is a standard Frobenius morphism, τ is an automorphism of algebraic groups coming from a graph automorphism of types A_n , D_n or E_6 , and τ' is a bijective endomorphism coming from a graph automorphism of type B_2 ($p = 2$), F_4 ($p = 2$) or G_2 ($p = 3$). The latter instances only occur in bad characteristic, so are not relevant here. If $\tau = 1$, then we say that G_σ is *untwisted*, else G_σ is *twisted*. Note that, since G is simple, τ and σ_q commute. Note also that $C_G(\tau)$ is again simple (e.g., see [[Gorenstein et al. 1998](#), Theorem 1.15.2(d)]).

Theorem 6.5. *Suppose G is simple. Let $\sigma = \tau\sigma_q$ be a Steinberg endomorphism of G and H a connected semisimple σ -stable subgroup of G . Assume $p \geq \tilde{h}(G)$ and $p \geq \tilde{h}(H)$. Then H^{sat} is also σ -stable, and we have:*

- (i) *If $\tau = 1$, then $(G_\sigma)^{\text{sat}} = G$.*
- (ii) *If $\tau = 1$ and H is saturated in G , then $(H_\sigma)^{\text{sat}} = H$.*
- (iii) *$(G_\sigma)^{\text{sat}} = C_G(\tau)$.*
- (iv) *If H is saturated in G , and both τ and σ_q stabilise H separately, then $(H_\sigma)^{\text{sat}} = C_H(\tau)$.*
- (v) *If H is saturated in G , then $((H_\sigma)^{\text{sat}})_\sigma = H_\sigma$.*

Proof. The fact that H^{sat} is σ -stable follows from [Corollary 6.2](#).

For the rest of the proof, there is no loss in assuming that both G and H are generated by their respective root subgroups relative to some fixed maximal σ -stable tori $T_H \subseteq T_G = T$.

(i) and (ii) If $\tau = 1$, i.e., if $\sigma = \sigma_q$ is standard, then every root subgroup of G meets G_σ nontrivially. (For, each root subgroup U_α of G is σ -stable and the σ -stable maximal torus T acts transitively on U_α . So the result follows from the Lang–Steinberg theorem.) It thus follows from [Theorem 5.2\(iii\)](#) and [\(5.3\)](#) that $(G_\sigma)^{\text{sat}}$ contains each root subgroup of G ; thus (i) follows. The same argument applies for (ii) by considering the simple components of H and the fact that saturation in H coincides with saturation in G , by [Theorem 5.7\(ii\)](#).

(iii) Since τ and σ_q commute, we have $G_\sigma = C_G(\tau)_{\sigma_q}$. Since $C_G(\tau)$ is saturated in G , by [Corollary 5.5\(iii\)](#), the result follows from part (ii).

(iv) Again, since τ and σ_q commute, $H_\sigma = C_H(\tau)_{\sigma_q}$. Now $C_H(\tau)$ is saturated in H , by [Corollary 5.5](#). But since H is saturated in G , saturation in H coincides with saturation in G , by [Theorem 5.7\(ii\)](#), so the result follows from part (ii).

(v) Thanks to [Corollary 6.2](#), $(H_\sigma)^{\text{sat}}$ is σ -stable. Thus, since $H_\sigma \subseteq (H_\sigma)^{\text{sat}}$ and $(H_\sigma)^{\text{sat}} \subseteq H^{\text{sat}} = H$, we have $H_\sigma \subseteq ((H_\sigma)^{\text{sat}})_\sigma \subseteq H_\sigma$, and equality follows. \square

Proof of Theorem 1.6. This follows immediately from [Theorem 6.5\(iii\)](#). \square

Remark 6.6. We note that [Theorem 6.5\(v\)](#) generalises [[Nori 1987](#), Theorem B(1)]: If $G = \text{SL}_n(k)$, $\sigma = \sigma_q$ is a standard Frobenius endomorphism of G , and H is a σ -stable subgroup of G , then it follows directly from [\(5.1\)](#) that H^{sat} is again σ -stable. Thus, in particular, if H is a connected, saturated semisimple σ -stable subgroup of G , then by [Theorem 6.5\(v\)](#) we have $((H_\sigma)^{\text{sat}})_\sigma = H_\sigma$; see [[Nori 1987](#), Theorem B(1)].

We consider some explicit examples for [Theorem 6.5](#).

Example 6.7. Let $G = \mathrm{SL}_n$ and let σ be the Steinberg endomorphism of G given by

$$g \mapsto \sigma_q(n_0({}^t g^{-1})n_0^{-1}),$$

where ${}^t g$ denotes the transpose of the matrix g and n_0 is the antidiagonal permutation matrix of GL_n , normalizing SL_n . Note that $\tau(g) = n_0({}^t g^{-1})n_0^{-1}$ is the graph automorphism of G . Then $\sigma^2 = \sigma_{q^2}$ is a standard Frobenius map of G given by raising coefficients to the q^2 -th power. Note that $G_\sigma = \mathrm{SU}(q)$ is the special unitary subgroup of G . We have $\mathrm{SU}(q) = G_\sigma \subseteq G_{\sigma^2} = \mathrm{SL}(\mathbb{F}_{q^2})$, and since σ_q commutes with τ , we have (assuming $p \geq n$) $(G_\sigma)^{\mathrm{sat}} = C_G(\tau)$, by [Theorem 6.5\(iii\)](#), while $(G_{\sigma^2})^{\mathrm{sat}} = G$, by [Theorem 6.5\(i\)](#).

In the case when G is no longer simple, additional kinds of Steinberg endomorphisms are possible.

Example 6.8. Let H be a semisimple group defined over \mathbb{F}_q and let σ_q be the corresponding standard Frobenius map of H . Let $G = H \times \cdots \times H$ (r factors) and let ΔH be the diagonal copy of H in G . Let π be the r -cycle permuting the r direct copies of H of G cyclically and let $f = (\sigma_q, \mathrm{id}_H, \dots, \mathrm{id}_H) : G \rightarrow G$. Then $\sigma = \pi f$ is a Steinberg endomorphism of G where π and f do not commute. We have $G_\sigma = (\Delta H)_{\sigma_q}$, where by abuse of notation σ_q is a standard Frobenius map on ΔH . (Note that $(\Delta H)_{\sigma_q}$ is isomorphic to $H_{\sigma_q} = H(\mathbb{F}_q)$.) Now suppose $p \geq \tilde{h}(H) = \tilde{h}(G)$. Then ΔH is saturated in G , by [Lemma 5.10](#). Thus $(G_\sigma)^{\mathrm{sat}} = ((\Delta H)_{\sigma_q})^{\mathrm{sat}} = \Delta H$, by [Theorem 6.5\(ii\)](#).

We present an instance where [Theorem 6.5\(i\)](#) can be applied even though G is not simple and σ is a Steinberg endomorphism which is not a generalized Frobenius endomorphism.

Example 6.9. Let $p \geq 2$. Let σ_p and σ_{p^2} be the standard Frobenius maps of SL_2 given by raising coefficients to the p -th and p^2 -th powers, respectively. Let $G = \mathrm{SL}_2 \times \mathrm{SL}_2$. Then the map $\sigma = \sigma_p \times \sigma_{p^2} : G \rightarrow G$ is a Steinberg morphism of G that is not a generalized Frobenius morphism (see the remark following [\[Gorenstein et al. 1998, Theorem 2.1.11\]](#)). We have $G_\sigma = \mathrm{SL}_2(\mathbb{F}_p) \times \mathrm{SL}_2(\mathbb{F}_{p^2})$. The saturation map in G is given by the formula from [\(5.1\)](#). (If $p \geq 5$ then this follows from [Theorem 5.7](#), since the image of the canonical embedding of G into $\mathrm{GL}_4(k)$ is a saturated subgroup of $\mathrm{GL}_4(k)$, but it is easily verified for $p < 5$ also.) By [Lemma 5.10](#), saturating G_σ inside G amounts to saturating each factor of G_σ inside each factor of G . Now, by applying [Theorem 6.5\(i\)](#) to each factor of G , we get $(G_\sigma)^{\mathrm{sat}} = G$.

7. Saturation and semisimplification

Definition 7.1 [\[Bate et al. 2020, Definition 4.1\]](#). Let H be a subgroup of G . We say that a subgroup H' of G is a *semisimplification of H (for G)* if there exists

a parabolic subgroup $P = P_\lambda$ of G and a Levi subgroup $L = L_\lambda$ of P such that $H \subset P_\lambda$ and $H' = c_\lambda(H)$, and H' is G -completely reducible. We say the pair (P, L) yields H' .

The following consequence of [Proposition 5.6](#) shows that passing to a semisimplification of a subgroup of G and saturation are naturally compatible.

Corollary 7.2. *Suppose that $p \geq \tilde{h}(G)$. Let H' be a semisimplification of H yielded by (P, L) . Then a semisimplification of H^{sat} is given by $(H')^{\text{sat}}$ and is yielded also by (P, L) . Moreover, any semisimplification of H^{sat} is G -conjugate to $(H')^{\text{sat}}$.*

Proof. By [Lemma 2.2](#) there exists a $\lambda \in Y(G)$ such that $P = P_\lambda$, $L = L_\lambda$ and $H' = c_\lambda(H)$. According to [Proposition 5.6](#), we have $(H')^{\text{sat}} = (c_\lambda(H))^{\text{sat}} = c_\lambda(H^{\text{sat}})$. Since H' is G -cr, by definition, $(H')^{\text{sat}}$ is G -cr by [Theorem 5.13](#). The final statement follows from [\[Bate et al. 2020, Theorem 4.5\]](#). \square

In [Example 5.17](#), the subgroup H considered is connected, nonsaturated and not G -cr. In our next example, we give a connected, nonsaturated but G -cr subgroup.

Example 7.3. Consider the semisimple SL_2 -module $L(1) \oplus L(p) = L(1) \oplus L(1)^{[p]}$, i.e., SL_2 acting with a Frobenius twist on the second copy of the natural module and without such on the first copy. This defines a diagonal embedding of SL_2 in $M := \text{SL}_2 \times \text{SL}_2 \subseteq G = \text{GL}_4$. The image H of SL_2 in G is G -cr and it is not saturated in G (and also not in the saturated G -cr subgroup M of G). The argument is similar to the one in [Example 5.17](#). Note that M is the saturation of H in G .

We close this section by noting that in general homomorphisms are not compatible with saturation. For instance, take the inclusion of a connected reductive, nonsaturated subgroup H in G . See also [Examples 5.17](#) and [7.3](#).

Acknowledgements

We are grateful to J-P. Serre for some suggestions on an earlier version of the manuscript. The research of this work was supported in part by the DFG (grant #RO 1072/22-1 (project number: 498503969) to G. Röhrle). For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

References

- [Balaji et al. 2017] V. Balaji, P. Deligne, and A. J. Parameswaran, “On complete reducibility in characteristic p ”, *Épjournal Géom. Algébrique* **1** (2017), art. id. 3. [MR](#) [Zbl](#)
- [Bate et al. 2005] M. Bate, B. Martin, and G. Röhrle, “A geometric approach to complete reducibility”, *Invent. Math.* **161**:1 (2005), 177–218. [MR](#) [Zbl](#)
- [Bate et al. 2010] M. Bate, B. Martin, G. Röhrle, and R. Tange, “Complete reducibility and separability”, *Trans. Amer. Math. Soc.* **362**:8 (2010), 4283–4311. [MR](#) [Zbl](#)

- [Bate et al. 2011] M. Bate, S. Herpel, B. Martin, and G. Röhrle, “*G*-complete reducibility and semisimple modules”, *Bull. Lond. Math. Soc.* **43**:6 (2011), 1069–1078. [MR](#) [Zbl](#)
- [Bate et al. 2020] M. Bate, B. Martin, and G. Röhrle, “Semisimplification for subgroups of reductive algebraic groups”, *Forum Math. Sigma* **8** (2020), art. id. e43. [MR](#) [Zbl](#)
- [Bate et al. 2022] M. Bate, B. Martin, and G. Röhrle, “Overgroups of regular unipotent elements in reductive groups”, *Forum Math. Sigma* **10** (2022), art. id. e13. [MR](#) [Zbl](#)
- [Borel 1991] A. Borel, *Linear algebraic groups*, 2nd ed., Grad. Texts Math. **126**, Springer, 1991. [MR](#) [Zbl](#)
- [Bourbaki 1972] N. Bourbaki, *Éléments de mathématique. Fasc. XXXVII. Groupes et algèbres de Lie. Chapitre II: Algèbres de Lie libres. Chapitre III: Groupes de Lie*, Actualités Scientifiques et Industrielles **1349**, Hermann, Paris, 1972. [MR](#) [Zbl](#)
- [Bourbaki 1975] N. Bourbaki, *Éléments de mathématique. Fasc. XXXVIII: Groupes et algèbres de Lie. Chapitre VII: Sous-algèbres de Cartan, éléments réguliers. Chapitre VIII: Algèbres de Lie semi-simples déployées*, Actualités Scientifiques et Industrielles **1364**, Hermann, Paris, 1975. [MR](#) [Zbl](#)
- [Gorenstein et al. 1998] D. Gorenstein, R. Lyons, and R. Solomon, *The classification of the finite simple groups, Number 3, Part I, Chapter A: Almost simple K-groups*, Math. Surv. Monogr. **40.3**, Amer. Math. Soc., Providence, RI, 1998. [MR](#) [Zbl](#)
- [Herpel 2013] S. Herpel, “On the smoothness of centralizers in reductive groups”, *Trans. Amer. Math. Soc.* **365**:7 (2013), 3753–3774. [MR](#) [Zbl](#)
- [Humphreys 1995] J. E. Humphreys, *Conjugacy classes in semisimple algebraic groups*, Math. Surv. Monogr. **43**, Amer. Math. Soc., Providence, RI, 1995. [MR](#) [Zbl](#)
- [Jantzen 1997] J. C. Jantzen, “Low-dimensional representations of reductive groups are semisimple”, pp. 255–266 in *Algebraic groups and Lie groups*, edited by G. Lehrer et al., Austral. Math. Soc. Lect. Ser. **9**, Cambridge Univ. Press, 1997. [MR](#) [Zbl](#)
- [Litterick and Thomas 2018] A. J. Litterick and A. R. Thomas, “Complete reducibility in good characteristic”, *Trans. Amer. Math. Soc.* **370**:8 (2018), 5279–5340. [MR](#) [Zbl](#)
- [Litterick et al. 2025] A. J. Litterick, D. I. Stewart, and A. R. Thomas, “Complete reducibility and subgroups of exceptional algebraic groups”, pp. 191–245 in *Groups St Andrews 2022 in Newcastle*, edited by C. M. Campbell et al., London Math. Soc. Lecture Note Ser. **496**, Cambridge Univ. Press, 2025. [MR](#)
- [Lübeck 2001] F. Lübeck, “Small degree representations of finite Chevalley groups in defining characteristic”, *LMS J. Comput. Math.* **4** (2001), 135–169. [MR](#) [Zbl](#)
- [Nori 1987] M. V. Nori, “On subgroups of $GL_n(\mathbb{F}_p)$ ”, *Invent. Math.* **88**:2 (1987), 257–275. [MR](#) [Zbl](#)
- [Richardson 1967] R. W. Richardson, Jr., “Conjugacy classes in Lie algebras and algebraic groups”, *Ann. of Math. (2)* **86** (1967), 1–15. [MR](#) [Zbl](#)
- [Serre 1998] J-P. Serre, “The notion of complete reducibility in group theory”, Moursund Lectures (University of Oregon), 1998. [arXiv math/0305257v1](#)
- [Serre 2004] J-P. Serre, “Complète réductibilité”, exposé 932, 195–217 in *Séminaire Bourbaki 2003/2004*, Soc. Math. de France, Paris, 2004. [Zbl](#)
- [Slodowy 1997] P. Slodowy, “Two notes on a finiteness problem in the representation theory of finite groups”, pp. 331–348 in *Algebraic groups and Lie groups*, edited by G. Lehrer et al., Austral. Math. Soc. Lect. Ser. **9**, Cambridge Univ. Press, 1997. [MR](#) [Zbl](#)
- [Springer and Steinberg 1970] T. A. Springer and R. Steinberg, “Conjugacy classes”, pp. 167–266 in *Seminar on algebraic groups and related finite groups* (The Institute for Advanced Study, Princeton, 1968/69), Lecture Notes in Math. **131**, Springer, 1970. [MR](#) [Zbl](#)

[Steinberg 1968] R. Steinberg, *Endomorphisms of linear algebraic groups*, Mem. Amer. Math. Soc. **80**, Amer. Math. Soc., Providence, RI, 1968. [MR](#) [Zbl](#)

[Testerman 1995] D. M. Testerman, “ A_1 -type overgroups of elements of order p in semisimple algebraic groups and the associated finite groups”, *J. Algebra* **177**:1 (1995), 34–76. [MR](#) [Zbl](#)

Received October 20, 2024. Revised March 27, 2025.

MICHAEL BATE

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF YORK

YORK

UNITED KINGDOM

michael.bate@york.ac.uk

SÖREN BÖHM

FAKULTÄT FÜR MATHEMATIK

RUHR-UNIVERSITÄT BOCHUM

BOCHUM

GERMANY

soeren.boehm@rub.de

ALASTAIR LITTERICK

SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE

UNIVERSITY OF ESSEX

COLCHESTER

UNITED KINGDOM

a.litterick@essex.ac.uk

BENJAMIN MARTIN

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF ABERDEEN

KING’S COLLEGE

ABERDEEN

UNITED KINGDOM

b.martin@abdn.ac.uk

GERHARD RÖHRLE

FAKULTÄT FÜR MATHEMATIK

RUHR-UNIVERSITÄT BOCHUM

BOCHUM

GERMANY

gerhard.roehrle@ruhr-uni-bochum.de

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2025 is US \$677/year for the electronic version, and \$917/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), PASCAL CNRS Index, [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 337 No. 1 July 2025

<i>G</i> -complete reducibility and saturation	1
MICHAEL BATE, SÖREN BÖHM, ALASTAIR LITTERICK, BENJAMIN MARTIN and GERHARD RÖHRLE	
On gamma factors of generic representations of $U_{2n+1} \times \text{Res}_{E/F} \text{GL}_r$	25
YAO CHENG and CHIAN-JEN WANG	
Atoms and associated spectral properties for positive operators on L^p	87
JEAN-FRANÇOIS DELMAS, KACEM LEFKI and PIERRE-ANDRÉ ZITT	
Regular poles for spinor L -series attached to split Bessel models of $\text{GSp}(4)$	137
MIRKO RÖSNER and RAINER WEISSAUER	