Transmutations from the Covariant Transform on the Heisenberg Group and an Extended Umbral Principle

Vladimir V. Kisil^{1,*}

¹School of Mathematics, University of Leeds, Leeds LS29JT, England (Received May 24, 2023)

We discuss several seemingly assorted objects: the umbral calculus, generalised translations and associated transmutations, symbolic calculus of operators. The common framework for them is representations of the Weyl algebra of the Heisenberg group by ladder operators. Transporting various properties between different implementations we review some classic results and new opportunities.

Dedicated to 100th anniversary of Ivan Aleksandrovich Kipriyanov's birth

То и это, и вон то, вместе с этим тоже —всё равно всегда выходит как одно и то же. Считалочка

I. INTRODUCTION: THE BESSEL OPERATOR

Let us start from the celebrated singular Bessel differential operator $[1, \S1.3]$, $[2, \S2.2.2]$, $[3, \S1.4.1]$ given by:

$$\mathcal{B}_{\nu} = \frac{\mathrm{d}^2}{\mathrm{d}t^2} + \frac{\nu}{t}\frac{\mathrm{d}}{\mathrm{d}t} = t^{-\nu}\frac{\mathrm{d}}{\mathrm{d}t}t^{\nu}\frac{\mathrm{d}}{\mathrm{d}t}.$$
(1)

Due to importance of this operator in theoretical and applied settings there are many different approaches to study \mathcal{B}_{ν} and corresponding boundary value problems. Main tools are various integral transforms [1–3] intertwining \mathcal{B}_{ν} with some other operators, which are more accessible for certain reasons. The special rôle of such integral transforms is often indicated by calling them *transmutations* [4].

For example, the *Poisson operator* is defined for a summable function f(t) by

$$[\mathcal{P}^{\nu}f](x) = \frac{2\Gamma(\frac{\nu+1}{2})}{\sqrt{\pi}\Gamma(\frac{\nu}{2})x^{\nu-1}} \int_{0}^{x} (x^{2} - t^{2})^{\frac{\nu}{2} - 1} f(t) \,\mathrm{d}t.$$
(2)

It transmutes (in other words intertwines) the Bessel operator with the second derivative:

$$\mathcal{P}^{\nu} \circ \mathcal{B}_{\nu} = \frac{\mathrm{d}^2}{\mathrm{d}x^2} \circ \mathcal{P}^{\nu}.$$
(3)

Another example is a Fourier-style decomposition over eigenfunctions of \mathcal{B}_{ν} —the Hankel transform defined by:

$$[\mathcal{H}_{\nu}f](\lambda) = \int_{0}^{\infty} f(t) j_{\frac{\nu-1}{2}}(\sqrt{\lambda}t) t^{\nu} dt, \qquad (4)$$

^{*} E-mail: V.Kisil@leeds.ac.uk; On leave from the Odessa University.

where $j_{\frac{\nu-1}{2}}(\sqrt{\lambda}t)$ is the solution of the boundary value problem:

$$\mathcal{B}_{\nu}j_{\frac{\nu-1}{2}}(\sqrt{\lambda}t) = -\lambda j_{\frac{\nu-1}{2}}(\sqrt{\lambda}t), \quad \text{and} \quad j_{\frac{\nu-1}{2}}(0) = 1, \quad j_{\frac{\nu-1}{2}}'(0) = 0.$$
(5)

The Hankel transform transmutes \mathcal{B}_{ν} with the operator of multiplication:

$$\mathcal{H}_{\nu} \circ \mathcal{B}_{\nu} = (-\lambda I) \circ \mathcal{P}^{\nu}$$

Furthermore, it is fruitful [5, 6] to extend the harmonic analysis analogy with \mathcal{H}_{ν} and introduce the generalised translations T_t^y such that functions $j_{\frac{\nu-1}{2}}(\sqrt{\lambda}t)$ plays the rôle of characters—just like exponents for ordinary translations:

$$T_t^y j_\nu(\lambda t) = j_\nu(\lambda y) j_\nu(\lambda t). \tag{6}$$

There are numerous structural similarities between ordinary and generalised translations [5, 7]. The correspondence between two resembles the *umbral calculus* [8, 9], that is the technique of dealing with combinatorial quantities p_n as "shadows" of power monomials x^n . This intriguing magic may be turn into a solid theory through a systematic usage of certain linear operators, see [8, 9] and §II C below.

In this paper the umbral calculus is unfolded through representations of the Weyl algebra by ladder operators. Covariant transforms intertwine properties of corresponding objects from various representations. It allows us to propagate results and methods from one setting to another. We extend the scope through umbral interpretation of generalised shifts and transmutations. Such connections highlight some gaps and missed opportunities which usually remain in blind spots if the topics are treated in the traditional isolated manner.

The paper *outline* is as follows. We are collecting required preliminary material in Section II. The central place is occupied by representations of Weyl algebra of the Heisenberg group by ladder operators, which are introduced in §II A. Then, we remind basics on the covariant (Berezin) transform and Perelomov's coherent states in §II B. The intertwining properties of the covariant transform are foundations of the *umbral principle* presented in §II C. Convenient complements are the Fourier transform to the momentum picture §II D and the adjoint action of the ladder operators §II E. The final bit is a special case of shift invariant delta operator and the resulting binomial in §II F.

Sufficiently equipped by the previous preparations we interpret several examples through ladder operators in Section III. The combinatorial umbral calculus is our first example in §III A. We extend it to the Delsarte–Levitan's generalised translations and associated transmutations in §III B. These are illustrated by the cases of higher order delta operators—the second derivative in §III B 1 and the Bessel operator \mathcal{B}_{ν} (1) in §III B 2. Finally, we discuss an application of symbolic calculus of operators which stems out from the calculus of pseudo-differential operators (PDO) in §III C.

The final Section IV summarises our finding and proposes an *extended interpretation* of the umbral principle, see illustration (47). We also indicate further developments and opportunities generated by the described connections.

II. PRELIMINARIES

Representation theory, in particular of the Heisenberg group [10], is behind many important calculations in analysis [11–13] and we provide some of its basics in the present section. The group-theoretical foundations of coherent states/wavelets are well-known and widely appreciated [14, 15]. We can widen the applicability of the approach [16] if it is extended to the Banach spaces [17, 18].

A. Abstract representation of the Weyl algebra and the Heisenberg group

Let an infinite-dimensional vector space E have a basis $\{p_n\}$, n = 0, 1, 2, ... Then, one can define the following associated objects:

1. The linear map $\mathcal{Q} : \mathsf{E} \to \mathsf{E}$ such that:

$$\mathcal{Q}p_n = p_{n-1}, \text{ for } n > 0 \quad \text{and} \quad \mathcal{Q}p_0 = 0.$$
 (7)

In a combinatorial context Q is called *delta operator* or simply *delta*.

2. The liner map $\mathcal{P} : \mathsf{E} \to \mathsf{E}$ such that:

$$\mathcal{P}p_n = \iota(n+1)p_{n+1}, \quad \text{for } n = 0, 1, 2, \dots,$$
(8)

for some scalar ι . In many algebraic considerations, e.g. the umbral calculus, it is common to take the simplest value $\iota = 1$. For unitarity in a Hilbert space can employ the imaginary unit $\iota = i$.

3. The linear functional $l_0 \in \mathsf{E}^*$ defined by:

$$\langle l_0, p_n \rangle = \delta_{0n} , \qquad \text{for all } n = 0, 1, 2, \dots$$
 (9)

where δ_{ij} is the Kronecker delta.

The above operators \mathcal{Q} and \mathcal{P} satisfy to the *Heisenberg commutation relation* $[\mathcal{P}, \mathcal{Q}] = -\iota I$. Thus, the set $\{\mathcal{P}, \mathcal{Q}, -\iota I\}$ form a representation of the Weyl Lie algebra \mathfrak{h} of the Heisenberg group \mathbb{H} . In the quantum mechanical framework the action of \mathcal{P} (7) and \mathcal{Q} (8) are known as *ladder operators—creation* and *annihilation*, respectively [19–21]. Their action is visualised as follows:

$$0 \leftarrow \begin{array}{c} & & & \\ & & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & &$$

We can use the standard exponentiation of (7)–(8) to obtain an action of the Heisenberg group. Specifically:

$$e^{yQ}p_n = \sum_{k=0}^{\infty} \frac{y^k Q^k}{k!} p_n = \sum_{k=0}^n \frac{y^k}{k!} p_{n-k};$$
(11)

$$e^{x\mathcal{P}}p_n = \sum_{k=0}^{\infty} \frac{x^k \mathcal{P}^k}{k!} p_n = \sum_{k=0}^{\infty} \binom{n+k}{k} x^k p_{n+k}.$$
 (12)

One may notice a difference between (11) and (12): the former is effectively a finite linear combination of p_n and the latter is infinite. Thus, e^{yQ} makes sense in the linear space E itself, but $e^{x\mathcal{P}}$ needs some additional structure, e.g. a topology to interpret (12) as a convergent series. An undemanding resolution is the "formal power series" framework, that is a collection of all infinite series $f = \sum_{0}^{\infty} a_n p_n$ with $f^{(k)} \to f$ if and only if $a_n^{(k)} \to a_n$ for all n. We will continue to loosely denote some topological extension of E which supports expansions (11)–(12) by the same letter E .

Once actions (11)–(12) are meaningful, they satisfy to the Weyl commutator relation:

$$\mathrm{e}^{y\mathcal{Q}}\,\mathrm{e}^{x\mathcal{P}} = \mathrm{e}^{\iota xy}\,\mathrm{e}^{x\mathcal{P}}\,\mathrm{e}^{y\mathcal{Q}}.$$

Thus, we have a representation of the (polarised) *Heisenberg group* \mathbb{H} of the form:

$$\rho_{\mathsf{E}}(s, x, y) = \mathrm{e}^{-\iota s} \,\mathrm{e}^{-y\mathcal{Q}} \,\mathrm{e}^{-x\mathcal{P}},\tag{13}$$

with the composition law reflecting the multiplication on \mathbb{H} :

$$\rho_{\mathsf{E}}(s, x, y)\rho_{\mathsf{E}}(s', x', y') = \rho_{\mathsf{E}}(s + s' + xy', x + x', y + y'), \tag{14}$$

Example 1. The archetypal model is the linear space of polynomials $\underline{\mathsf{E}}$, say, in a variable t with the monomial basis $\underline{p}_n(t) = \frac{1}{n!}t^n$. Then $\underline{\mathcal{Q}} = \frac{d}{dt}$, $\underline{\mathcal{P}} = \iota t I$ (the operator of multiplication by the variable) and $\langle \underline{l}_0, p \rangle = p(0)$ (the evaluation at 0). An extension of $\underline{\mathsf{E}}$ to a space of power series in t with rapidly decreasing coefficients allows us to write the representation (13) as:

$$\rho_{\iota}(s, x, y)f(t) = e^{-\iota(s+x(t-y))}f(t-y),$$
(15)

which is a cousin of the Schrödinger representation [19, \$1.3].

B. Covariant transform and intertwining properties

There is an extensive literature on the covariant transform (also known as the coherent state transform and many other names) in general [14, Thm. 8.1.3] and for the Heisenberg group in particular [19, 21, 22]. To make a long story short we recall here only the bare minimum of definitions and notations, see provided references for further details.

It is common to consider covariant transform either in inner product spaces or Gelfand triples (ridged Hilbert spaces) [17]. In the present situation it is more convenient to work in a linear space and its dual [18, 23].

Definition 2. [24] Let ρ be a representation of a group G in a vector space V. Take a *fiducial* functional $l \in V^*$ and denote its pairing with $v \in V$ by $\langle l, v \rangle$. The covariant transform is the map:

$$[\mathcal{W}_l v](g) = \langle l, \rho(g) v \rangle, \qquad v \in \mathsf{V}, \quad g \in G, \tag{16}$$

to scalar-valued functions on G. In the case of a Hilbert space V, a fiducial functional is provided by a pairing with a vector $f \in V \sim V^*$ known as a *mother wavelet* or *vacuum state* [14, 15].

The covariant transform \mathcal{W}_l plainly interacts with the left Λ and right *R* regular representations of *G*, which are:

$$\Lambda(g): f(h) \mapsto f(g^{-1}h)$$
 and $R(g): f(h) \mapsto f(hg)$, where $h, g \in G$.

Lemma 3. The covariant transform (16) intertwines the left Λ and right R regular representations of G with the following actions:

$$\Lambda(g) \circ \mathcal{W}_l = \mathcal{W}_l \circ \rho(g) \quad and \quad R(g) \circ \mathcal{W}_l = \mathcal{W}_{\rho^*(g)l} \quad for \ all \ g \in G.$$

$$\tag{17}$$

Here ρ^* is the adjoint representation on V^* , that is $\langle l, \rho(g)v \rangle = \langle \rho^*(g)l, v \rangle$.

For a subgroup $H \subset G$ and its character χ let the fiducial functional l has the *covariance* property:

$$\langle l, \rho(h)v \rangle = \bar{\chi}(h)\langle l, v \rangle, \quad \text{for all } h \in H \quad \text{and all } v \in \mathsf{V}.$$
 (18)

Then, \mathcal{W}_l is a Perelomov-style covariant transform [15], that is only a part of the covariant transform (16) allows to recover all values. To this end, we fix a continuous section $\mathbf{s} : G/H \to G$, which is a right inverse to the natural projection $\mathbf{p} : G \to G/H$.

Definition 4. [16, §5.1] Let $l \in V^*$ intertwine the restriction of ρ to H with a character χ of H, cf. (18). Then, the *induced covariant transform* is:

$$[\mathcal{W}_l v](x) = \langle l, \rho(\mathbf{s}(x))v \rangle, \qquad v \in \mathsf{V}, \quad x \in G/H.$$
(19)

Under our assumptions, the induced covariant transform intertwines ρ with a representation induced from H by the character χ .

In many cases, e.g. for square integrable representations and an admissible mother wavelet $v \in V$, the image space of the covariant transform is a reproducing kernel Hilbert space [14, Thm. 8.1.3]. That means that for any function $l \in W_l V$ we have the integral reproducing formula:

$$v(y) = \int_X v(x) \,\bar{k}_y(x) \,dx,\tag{20}$$

where the reproducing kernel k_y is the twisted convolution with the normalised covariant transform $\mathcal{W}_l(\rho(\mathbf{s}(y)^{-1})l)$ for the mother wavelet l [25].

C. Covariant transforms for the Heisenberg group and the Umbral principle

We can observe that for the linear functional l_0 (9) the identity holds:

$$\langle l_0, \mathrm{e}^{x\mathcal{P}} f \rangle = \mathrm{e}^x \langle l_0, f \rangle,$$

which can be initially verified for $f = p_n$ and then extended to the whole E by linearity. Therefore, there is an induced covariant transform (19) for the subgroup

$$H_y = \{(s, x, 0) \mid s, x \in \mathbb{R}\}$$
 and the respective homogeneous space $\mathbb{H}/H_y \sim \mathbb{R}$. (21)

The representation ρ_h of \mathbb{H} induced from the character $\chi_h(s, x, 0) = e^{\iota s}$ of H_y coincides with the Schrödinger representation (15) [21]:

$$[\rho_h(s, x, y)f](u) = e^{-\iota(s+x(u-y))}f(u-y).$$

Then, the induced covariant transform defined by functional l_0 (9) and the subgroup H_y (21) maps E to a space of functions of one variable:

$$\tilde{f}(u) \coloneqq [\mathcal{W}_0 f](u) = \langle l_0, \mathrm{e}^{u\mathcal{Q}} f \rangle, \quad \text{in particular} \quad \tilde{p}_n(u) = \langle l_0, \mathrm{e}^{u\mathcal{Q}} p_n \rangle = \frac{(\iota u)^n}{n!}, \quad (22)$$

where the last identity follows from (11). The transform \mathcal{W}_0 intertwines the representation ρ_{E} (13) with the representation ρ_{ι} (15), in particular:¹

$$\mathcal{W}_0 \circ \rho_{\mathsf{E}} = \rho_\iota \circ \mathcal{W}_0, \qquad \mathcal{W}_0 \circ \mathcal{Q} = \frac{\mathrm{d}}{\mathrm{d}u} \circ \mathcal{W}_0, \qquad \mathcal{W}_0 \circ \mathcal{P} = (\iota u I) \circ \mathcal{W}_0, \qquad \langle l_0, f \rangle = [\mathcal{W}_0 f](0).$$
(23)

The above relations are the fundamentals of the following principle, cf. §III A below:

Umbral Principle. Any statement on functions of one variable, which is formulated in terms of a linear combination of derivatives, multiplication by monomials and evaluation at 0, corresponds to a statement about elements of E expressed through \mathcal{Q} , \mathcal{P} and l_0 according to the vocabulary (23).

Note that the above principle does not require any additional assumptions about the nature of E or its basis $\{p_n\}$. Once they are fixed, operators \mathcal{Q} (7), \mathcal{P} (8) and the functional l_0 (9) are completely defined and the Umbral principle is fully set.

¹ The archetypal implementation of a covariant transform and these intertwining relations is the Fock-Segal-Bargmann transform, cf. [39]. It intertwines the creation and annihilation operators for a quantum harmonic oscillator with operators of multiplication by the complex variable z and the complex derivative $\frac{d}{dz}$, respectively.

Remark 5. Interestingly, the umbral correspondence \mathcal{W}_0 (23) is defined only through the exponentiation of the delta \mathcal{Q} in (22). Thus, the paired operator \mathcal{P} is missing or at least obscured in many considerations. Therefore, it is tempting to deem the operator \mathcal{P} to be optional or even excessive. However, the importance of \mathcal{P} manifests itself through the possibility to define the entire umbral framework ($\mathsf{E}, \{p_n\}_0^\infty$) through the triple ($\mathsf{E}, \mathcal{P}, p_0$). Indeed, the sequence $\{p_n\}_0^\infty$ is inductively defined, cf. (12) by $p_{n+1} = \frac{1}{\iota(n+1)}\mathcal{P}p_n$, $n = 0, 1, 2, \ldots$ On the other hand, there is no way equally well describe the situation through the delta \mathcal{Q} alone, see the example of different Appell polynomials, which all share the same $\mathcal{Q} = \frac{d}{dt}$, in §III A below. We will meet some more arguments later, e.g. a usage of the operator \mathcal{P} allows an extension of the operator calculus in §III C.

D. The momentum picture and the Fourier transform

In some circumstances it is preferable to have an intertwining property with operators Q and \mathcal{P} swapped in comparison to (23). In the physical language: to use momenta of a particle instead of its coordinates in the configuration space. To this end we replace the functional l_0 (9) by a functional m invariant under the operator e^{uQ}

$$\langle m, \mathrm{e}^{u\mathcal{Q}}f \rangle = \langle m, f \rangle.$$
 (24)

Then, the respective induced covariant transform:

$$[\mathcal{W}_m f](v) = \langle m, \mathrm{e}^{v\mathcal{P}} f \rangle, \text{ intertwines } \mathcal{W}_m \circ \mathcal{Q} = (-\iota vI) \circ \mathcal{W}_m, \text{ and } \mathcal{W}_m \circ \mathcal{P} = \frac{\mathrm{d}}{\mathrm{d}v} \circ \mathcal{W}_m.$$
(25)

The above consideration is purely formal because the functional (24) does not have a non-trivial bounded action on the basis p_n of E . Indeed, $e^{u\mathcal{Q}}p_1 = p_1 + up_0$ and we shall have $\langle m, p_1 \rangle = \langle m, p_1 \rangle + u \langle m, p_0 \rangle$ for all u. Boundedness on p_n implies $\langle m, p_0 \rangle = 0$, which can be extended to $\langle m, p_n \rangle = 0$ for all n by induction.

Yet, such a non-zero functional may exist on a certain subspace of the extended space E. For example, if $\mathcal{Q} = \frac{d}{dt}$ and $e^{u\mathcal{Q}} : f(t) \mapsto f(t+u)$, cf. Ex. 1, a shift-invariant functional is:

$$\langle m, f \rangle = \int_{-\infty}^{\infty} f(t) \, \mathrm{d}t \qquad \text{for } f \in \mathsf{L}_1(\mathbb{R}).$$

Then, the induced covariant transform \mathcal{W}_m (25) effectively becomes the Fourier transform:

$$[\mathcal{W}_m f](v) = \int_{-\infty}^{\infty} e^{-\iota v t} f(t) dt,$$

It intertwines the Schrödinger representation ρ_{ι} (15) with itself through an automorphism of \mathbb{H} [10], [19, §1.3]. On the level of the Weyl algebra representation spanned by operators $\mathcal{Q} = \frac{d}{dt}$, $\mathcal{P} = \iota t I$ from Ex. 1 the automorphism acts as follows:

$$\mathcal{W}_m: \mathcal{Q} \mapsto -\mathcal{P}, \text{ and } \mathcal{W}_m: \mathcal{P} \mapsto \mathcal{Q}, \text{ therefore } [\mathcal{W}_m \mathcal{P}, \mathcal{W}_m \mathcal{Q}] = [\mathcal{P}, \mathcal{Q}] = -\iota I.$$

The above discussed incompatibility of the averaging functional m and the basis $\{p_n\}$ is just another reason why it is preferable to set the umbral framework through operators Q and \mathcal{P} rather than trough a specific basis $\{p_n\}$, cf. Rem. 5.

E. Adjoint action, resolution of the identity and binomial formula

We define adjoint ladder operators $\mathsf{E}^*\to\mathsf{E}^*$ in the usual way:

$$\langle \mathcal{Q}^*l, f \rangle = \langle l, \mathcal{Q}f \rangle, \quad \langle \mathcal{P}^*l, f \rangle = \langle l, \mathcal{P}f \rangle \quad \text{for all } f \in \mathsf{E} \text{ and } l \in \mathsf{E}^*.$$
 (26)

Then, the sequence $\{l_k\}_0^\infty \subset \mathsf{E}^*$ is inductively produced starting from the functional l_0 (9):

$$l_{n+1} = \mathcal{Q}^* l_n, \qquad n = 0, 1, 2, \dots$$
 (27)

Note, that the passage to adjoint operators swaps the creation and annihilation rôles of ladder operators.

The properties (9) and (26) implies bi-orthogonality of sequences $\{p_n\}_0^\infty$ and $\{l_k\}_0^\infty$:

$$\langle l_k, p_n \rangle = \delta_{kn}.\tag{28}$$

Therefore, we have the following resolution of the identity written in the Dirac bra-ket notation:

$$I = \sum_{k=0}^{\infty} |p_k\rangle \langle l_k|, \quad \text{that is} \quad f = \sum_{k=0}^{\infty} \langle l_k, f \rangle p_k, \quad \text{for all } f \in \mathsf{E}.$$
 (29)

The last identity for $f = p_n$ directly follows from (28) and then extends to all elements by linearity.

One can attempt an *unitarisation trick*: define a map $\mathcal{F} : \mathsf{E} \to \mathsf{E}^*$ by the rule $\mathcal{F} : p_n \mapsto l_n$ for all n. Say, if E is a space of polynomials in a single variable, we may look for a measure μ , which implements (28) in the form:

$$\langle \mathcal{F}p_k, p_n \rangle = \int p_k(t) \, p_n(t) \, \mathrm{d}\mu(t) = \delta_{kn}.$$
(30)

That is the classical approach to the *orthogonal polynomials* from ladder operators [20]. Let we have two abstract representations $(\mathsf{E}, \{p_n\}_0^\infty, \mathcal{Q}, \mathcal{P}, l_0)$ and $(\tilde{\mathsf{E}}, \{\tilde{p}_n\}_0^\infty, \tilde{\mathcal{Q}}, \tilde{\mathcal{P}}, \tilde{l}_0)$. Then we may define the correspondence $\mathsf{E} \to \tilde{\mathsf{E}}$ similarly to the resolution of identity (29):

$$\mathcal{V} = \sum_{k=0}^{\infty} |\tilde{p}_k\rangle \langle l_k|, \quad \text{that is} \quad f = \sum_{k=0}^{\infty} \langle l_k, f \rangle \, \tilde{p}_k, \quad \text{for all } f \in \mathsf{E}.$$
(31)

Clearly, (27) implies $\mathcal{V}: p_k \mapsto \tilde{p}_k$ for all $k = 0, 1, 2, \dots$

Similarly we can define generating function between two representations $(\mathsf{E}, \{p_n\}_0^\infty)$ and $(\tilde{\mathsf{E}}, \{\tilde{p}_n\}_0^\infty)$:

$$F(s,t) = \sum_{k=0}^{\infty} k! \,\tilde{p}_k(s) \, p_k(t), \qquad \text{which intertwines} \qquad \mathcal{Q}_t F(s,t) = \tilde{\mathcal{P}}_s F(s,t). \tag{32}$$

In most cases the generating function is taken for $(\tilde{\mathsf{E}}, \{\tilde{p}_n\}_0^\infty)$ being the archetypal model $(\underline{\mathsf{E}}, \{\underline{p}_n\}_0^\infty)$ from Ex. 1, that is:

$$F(s,t) = \sum_{k=0}^{\infty} s^k p_k(t), \qquad \text{such that} \qquad \mathcal{Q}_t F(s,t) = sF(s,t). \tag{33}$$

F. Shift invariance, binomial formula and operator calculus

There is a particular but still important situation of a translation-invariant delta Q.

Proposition 6. Let E be a space of function on the real line, let \mathcal{Q} commute with all translations $T^y: f(t) \mapsto f(t+y), y \in \mathbb{R}$ and $\langle l_0, f \rangle = f(0)$. Then

$$p_n(t+y) = \sum_{k=0}^{n} p_{n-k}(y) \, p_k(t).$$
(34)

Therefore, we can express the translation $T^y = e^{y \frac{d}{dt}}$ as a function of Q:

$$T^{y}f = \sum_{k=0}^{\infty} p_{k}(y) \mathcal{Q}^{k}f. \quad \text{for any } f \in \mathsf{E}.$$
(35)

A proof begins from an application of (29) to $T^y p_n(t) = p_n(t+y)$:

$$p_{n}(t+y) = \sum_{k=0}^{\infty} \langle l_{k}, T^{y} p_{n} \rangle p_{k}(t) = \sum_{k=0}^{\infty} \langle \mathcal{Q}^{*k} l_{0}, T^{y} p_{n} \rangle p_{k}(t) = \sum_{k=0}^{\infty} \langle l_{0}, \mathcal{Q}^{k} T^{y} p_{n} \rangle p_{k}(t)$$
$$= \sum_{k=0}^{\infty} \langle l_{0}, T^{y} \mathcal{Q}^{k} p_{n} \rangle p_{k}(t) = \sum_{k=0}^{n} \langle l_{0}, T^{y} p_{n-k} \rangle p_{k}(t) = \sum_{k=0}^{n} p_{n-k}(y) p_{k}(t),$$

where the series obviously terminates at k = n. Then, (35) for $f = p_n$ is exactly (34), thereafter (35) for a general $f \in \mathsf{E}$ follows from linearity.

In the archetypal case of Ex. 1 with $p_n(t) = \frac{1}{n!}t^n$ identity (34) turns out to be the celebrated binomial formula:

$$\frac{(t+x)^n}{n!} = \sum_{k=0}^n \frac{x^{n-k}}{(n-k)!} \frac{t^k}{k!} \qquad \text{or} \qquad (t+x)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} t^k.$$
(36)

Yet, (34) remains valid for numerous polynomials of binomial types, see below.

Now let us turn to some examples of the above abstract scheme.

III. VARIOUS IMPLEMENTATIONS

We start from the original version of the umbral calculus and then extend the scope by some new illustrations.

A. Finite Operator (Umbral) Calculus

As it often happens, the umbral calculus started from some particular observations in specific circumstances. Thereafter, it took several iterations to separate an abstract core from technical aspects [26] and recognise the rôle of the Heisenberg group [20, 27–33] in the construction. Here we are moving the opposite direction: from the abstract scheme of §II A to its specific implementations.

It is common in combinatorics to take p_n to be a polynomial of degree n. Alternatively, it can be requested that the delta operator Q sends the first order monomial to a constant: Qt = c [34]. Thereafter, some additional assumptions are employed as well, e.g. in the reverse historical order:

- If a delta Q commutes with shifts then $\{p_n\}$ is called a *Sheffer sequence*.
- If $Q = \frac{d}{dt}$ then $\{p_n\}$ is a sequence of *Appell polynomials* (thus, they are special case of Sheffer polynomials)
- If Sheffer polynomials satisfy $\langle l_0, p_n \rangle = p_n(0)$ then $\{p_n\}$ are called sequences of *binomial type*. As we already know, such $\{p_n\}$ shall satisfy to the binomial formula (34), which can be taken as their alternative definition and, clearly, is the source of their name.

Notably, the only Appell polynomials of binomial type are monomials $p_n(t) = \frac{1}{n!} (\iota t)^n$ from Ex. 1. That is the Appell and binomial type polynomials are two different branches springing from the archetypal model and still covered by the same umbrella of Sheffer polynomials.

Example 7. There are numerous sequences of polynomials covered in the literature, cf. [9, 26]:

	$p_n(t)$	Q	\mathcal{P}	$\langle l_0, f \rangle$
Monomials	$\frac{1}{n!}t^n$	$\frac{\mathrm{d}}{\mathrm{d}t}$	tI	f(0)
Lower factorials	$\frac{1}{n!}(t)_n = \frac{1}{n!}t(t-1)(t-2)\cdots(t-n+1)$	f(t+1) - f(t)	(8)	f(0)
Upper factorials	$\frac{1}{n!}t^{(n)} = \frac{1}{n!}t(t+1)(t+2)\cdots(t+n-1)$	f(t) - f(t-1)	(8)	f(0)

with more classical names to follow: the Abel polynomials, the Touchard polynomials, Appell sequences, Hermite polynomials, Bernoulli polynomials, etc. In many cases the simplest description of the operator \mathcal{P} is given by the references to (8). Yet a sort of analytic expressions may be elaborated sometimes as well.

We illustrate the Umbral principle here with one example only. Rewrite the binomial formula (36) using the operator $\frac{d}{dx}$:

$$e^{(y\frac{d}{dx})}\frac{x^n}{n!} = \sum_{k=0}^n \frac{y^k}{k!} \frac{x^{n-k}}{(n-k)!} = \sum_{k=0}^n \frac{y^k}{k!} \left(\frac{d^k}{dx^k} \frac{x^n}{n!}\right).$$

Then an application of the Umbral principle (e.g. the covariant transform) for Appell polynomials p_n with $Q = \frac{d}{dx}$ gives the identity:

$$p_n(t+y) = e^{yQ} p_n(t) = \sum_{k=0}^n \frac{y^k}{k!} Q^k p_n(t) = \sum_{k=0}^n \frac{y^k}{k!} p_{n-k}(t).$$

More illustrations can be found in [9, 26].

B. Delsarte-Levitan's generalised translations

Now we link the umbral principle with *generalised translations* proposed by Delsarte and extensively investigated by Levitan, cf. [7], [2, §2.2], [3, §3.4.3]. We present an adaptation of the original Delsarte's approach, which is well tuned to our theme.

Let L be a linear operator on a space $\mathsf{E}(\mathbb{R})$ of functions in one variable. We also assume that for some neighbourhood Ω of 0 and for any scalar $\lambda \in \Omega$ there is a unique non-zero solution $\phi(\lambda, t)$ of the eigenvalue problem:

$$\begin{cases} L_t \phi(\lambda, t) = \lambda \phi(\lambda, t) & \text{(where } L_t \text{ acts in variable } t); \\ \phi(\lambda, 0) = 1. \end{cases}$$
(37)

Furthermore, let $\phi(\lambda, t)$ be analytic at $\lambda = 0$ and generates functions $\phi_n(t)$ in the power expansion:

$$\phi(\lambda, t) = \sum_{n=0}^{\infty} \lambda^n \phi_n(t) \quad \text{such that (37) implies} \quad L\phi_n = \phi_{n-1} \quad \text{and} \quad \phi_n(0) = \delta_{0n}.$$
(38)

In other words, the collection $(\mathsf{E}(\mathbb{R}), \{\phi_n\}_0^\infty)$ provides the abstract representation of the Heisenberg group from §II A with $\mathcal{Q} = L$ and $\langle l_0, f \rangle = f(0)$. Furthermore, in this setting the decomposition (38) is an implementation of the generating function (33).

Changing the boundary conditions $\phi(\lambda, 0) = 1$ in (38) we get different umbral sequences $\{\phi_n\}_0^\infty$ for the same operator L. This corresponds to various Appell polynomials for the same $\mathcal{Q} = \frac{d}{dt}$ in the classic umbral setting outlined above.

If *L* commutes with all ordinary translations $e^{y\frac{d}{dt}}: f(t) \mapsto f(t+y)$, then $e^{y\frac{d}{dt}} = \sum_{0}^{\infty} \phi_k(y)L_t^k$ by (35). If translation-invariance of *L* is not assumed then the previous formula defines the umbral version for the generalised translation T_t^y :

$$[T_t^y f](t) = \sum_0^\infty \phi_k(y) \, L^k f(t).$$
(39)

By induction the intertwining property (33) of the generating function $\phi(\lambda, t) = \sum_{0}^{\infty} \lambda^{k} \phi_{k}(t)$ (38) implies $L^{k} \phi(\lambda, t) = \lambda^{k} \phi(\lambda, t)$, therefore, cf. (6):

$$[T_t^y \phi](\lambda, t) = \sum_{k=0}^{\infty} \phi_k(y) L^k \phi(\lambda, t) = \sum_{k=0}^{\infty} \phi_k(y) \lambda^k \phi(\lambda, t) = \phi(\lambda, y) \phi(\lambda, t).$$
(40)

That is, the generating function F(s,t) (33) is a character of the generalised translation (39). Now we turn to specific examples of generalised translations.

1. Second order derivative

To begin with, we can drop one of the main assumption of the umbral calculus in combinatorics that p_n is a polynomial of degree n. For example, let us consider even-order monomials $p_n(t) = \frac{(-1)^n t^{2n}}{(2n)!}$, $n = 0, 1, 2, \ldots$ and the linear space E_2 spanned by them. The second derivative $\mathcal{Q} = -\frac{\mathrm{d}^2}{\mathrm{d}t^2}$ has the action $\mathcal{Q}p_n = p_{n-1}$ for n > 1 and $\mathcal{Q}p_0 = 0$. Therefore, to have the commutator $[\mathcal{P}, \mathcal{Q}] = I$ we can define a linear operator \mathcal{P} by the rule

$$\mathcal{P}p_n = (n+1)p_{n+1},$$
 that is $\mathcal{P}t^{2n} = -\frac{1}{2(2n+1)}t^{2n+2}.$

Alternatively, we can employ antiderivatives and get an analytic expression, cf. [32]:

$$[\mathcal{P}f](t) = -\frac{1}{2}t \int_{0}^{t} f(s) \,\mathrm{d}s \qquad \text{for } f \in \mathsf{E}_{2}.$$

$$\tag{41}$$

It is known as a connection between $e^{\mathcal{Q}}$ and the (Gauss–)Weierstrass transform [35]. Therefore, exponentiation of \mathcal{Q} leads to the diffusion semigroup:

$$\mathrm{e}^{u\mathcal{Q}}f(t) = \frac{1}{2\sqrt{\pi u}} \int_{-\infty}^{\infty} f(s) \,\mathrm{e}^{-(t-s)^2/(4u)} \,\mathrm{d}s.$$

Since the functional l_0 is the evaluation $\langle l_0, f \rangle = f(0)$, the covariant transform is:

$$\tilde{f}(u) = \langle l_0, \mathrm{e}^{u\mathcal{Q}} f \rangle = \frac{1}{2\sqrt{\pi u}} \int_{-\infty}^{\infty} f(s) \, \mathrm{e}^{-s^2/(4u)} \, \mathrm{d}s.$$
(42)

This formula also follows from the observation that the intertwining property (23) between $Q = \frac{d^2}{dt^2}$ and $\frac{d}{du}$ requires the integral kernel $k(u, t) = \frac{1}{2\sqrt{\pi u}} e^{-t^2/(4u)}$, which is the fundamental solution of the heat equation $\frac{dk}{du} = \frac{d^2k}{dt^2}$.

We can use the same approach to find the intertwining operator with the momentum representation, cf. §II D, to avoid a challenging exponentiation of the antiderivative operator (41) as required by (25). To intertwine operator $Q = \frac{d^2}{dt^2}$ and vI a kernel k(v,t) shall satisfy to the equation $vk = \frac{d^2k}{dt^2}$ with the initial values k(v,0) = 1 and $k'_t(v,0) = 0$. The same answer appears as the generating function (33):

$$k(v,t) = \cos\left(\sqrt{v}t\right) = \sum_{k=0}^{\infty} \frac{(-v)^k t^{2k}}{(2k)!} \quad \text{and the transform is} \quad \hat{f}(v) = \int_{-\infty}^{\infty} f(t)\cos\left(\sqrt{v}t\right) \mathrm{d}t \,. \tag{43}$$

The operator $\mathcal{Q} = -\frac{\mathrm{d}^2}{\mathrm{d}t^2}$ can be viewed as an "umbra" for a generic Schrödinger operator $\frac{\mathrm{d}^2}{\mathrm{d}t^2} + V(t)I$, see [36] for a related discussion with a usage of the orthogonalisation (30).

2. Bessel operator and umbral calculus

Let $Q = B_{\nu}$ is the singular Bessel differential operator (1). The umbral sequence is formed by even-order monomials with a tailored scaling, cf. (41):

$$p_n(t) = \frac{t^{2n}}{4^n n! \,\Gamma(n + \frac{\nu - 1}{2})} \quad \text{with} \quad \mathcal{P}t^{2n} = \frac{t^{2(n+1)}}{2(2n + \nu + 1)} \quad \text{or} \quad [\mathcal{P}f](t) = \frac{1}{2}s^{1-\nu} \int_0^t s^{\nu} \cdot f(s) \,\mathrm{d}s. \tag{44}$$

The umbral functional is again the evaluation $\langle l_0, f \rangle = f(0)$. Alternatively, an umbral sequence is formed by fractional powers:

$$p_n(t) = \frac{t^{2n+1-\nu}}{4^n \, n! \, \Gamma(n+\frac{1-\nu}{2})} \quad \text{with} \quad \mathcal{P}t^{2n+1-\nu} = \frac{t^{2n+3-\nu}}{2(2n+3-\nu)} \quad \text{or} \quad [\mathcal{P}f](t) = \frac{1}{2} \int_0^t s \cdot f(s) \, \mathrm{d}s.$$

In this case the umbral functional is $\langle l_0, f \rangle = \lim_{t \to 0} (t^{\nu-1} f(t)).$

We are continuing with the first umbral sequence (44). The generating function (33) is also a solution of the eigenvalue problem (37), which is given by the normalised Bessel functions $j_{\frac{\nu-1}{2}}(\sqrt{st})$ (5):

$$j_{\frac{\nu-1}{2}}(\sqrt{st}) = \sum_{k=0}^{\infty} \frac{(-s)^k t^{2k}}{4^k k! \Gamma(k + \frac{1-\nu}{2})}.$$
(45)

A pairing invariant under the action $e^{y\mathcal{B}_{\nu}}$ for all real y is

$$\langle g, f \rangle = \int_{0}^{\infty} g(t) f(t) t^{\nu} dt.$$

The respective induced covariant transform $\left\langle j_{\frac{\nu-1}{2}}(\sqrt{st}), f(t) \right\rangle$ is the Hankel transform (4).

Thereafter, we can define a generalised translation T_t^y (39) for the Bessel operator \mathcal{B}_{ν} . Then, the generating function $j_{\frac{\nu-1}{2}}(\sqrt{st})$ has the character property (40), which takes the form (6). Since the Hankel transform (4) represents elements of E as superpositions of the normalised Bessel functions we can write an integral representation of the generalised translations T_t^y . Finally, the Poisson operator (2) is the intertwining operator (3) between two umbral schemes generated by the Bessel operator and the second derivative considered in the previous paragraph.

C. PDO type calculus of operators

An important tool to study boundary value problems with the Bessel operator \mathcal{B}_{ν} (1) is a *generalised convolution* [1, §1.3], [2, §2.2.3], [3, §3.4.3]. It can be defined through the Hankel transform \mathcal{H}_{ν} (4) treated as a sort of the Fourier transforms:

$$\mathcal{H}_{\nu}(f * g) = \mathcal{H}_{\nu}(f) \cdot \mathcal{H}_{\nu}(g).$$

Equivalently, the operator of convolution $C_f : g \mapsto f * g$ is an integral operator of the generalised translations (6):

$$[\mathcal{C}_f g](t) = \int_0^\infty f(y) \, [T_t^y g](t) \, y^\nu \, \mathrm{d}y = \int_0^\infty f(y) \, \sum_{k=0}^\infty \phi_k(y) \, [\mathcal{B}_\nu^k g](t) \, y^\nu \, \mathrm{d}y.$$

Clearly, this line of thoughts is applicable to any other umbral framework as well. From general principles, a large class of operators commuting with Q is contained in a weak closure of powers Q^k , cf. Prop. 6. Yet, we can use the umbral calculus to get a larger class of operators on E as integrated representations of ρ_{E} (13). That is, for a suitable kernel k(x, y) we define the operator of relative convolution [25, 37]:

$$\rho(k) = \iint k(x, y) e^{y\mathcal{Q}} e^{x\mathcal{P}} dx dy.$$
(46)

Of course, the expression is only meaningful if the operator \mathcal{P} is defined at all, it adds another reason for consideration of \mathcal{P} along with \mathcal{Q} to already presented in Rem. 5. The corresponding calculus of operators is well known [19, 38, 39]. In particular, the composition formula for two operators is

$$\rho(k_1)\rho(k_2) = \rho(k_1 \natural k_2), \quad \text{where} \quad [k_1 \natural k_2](x, y) = \iint k_1(x', y') \, k_2(x - x', y - y') \, \mathrm{e}^{\pi \mathrm{i}(x'y - y'x)} \, \mathrm{d}x' \, \mathrm{d}y',$$

and $k_1 \not\models k_2$ is called the *twisted convolution* for the Heisenberg group. A further facilitation is provided through considering operators of the form $a(\mathcal{P}, \mathcal{Q}) := \rho(\hat{a})$ (46) with a kernel being the Fourier transform of a function a(x, y). If ρ is the Schrödinger representation (15) then the operator $\rho(\hat{a})$ is a *pseudodifferential operator* (PDO) $a(\mathcal{P}, \mathcal{Q})$ with the *symbol* a(x, y) [19, 38, 39].

For other umbral frameworks we obtain different realisations of operators (46). However, all of them enjoy the same set of properties which are inherited from PDO through the Umbral principle. To make this approach successful for a specific class of operators one needs to make a suitable choice of an umbral model. On one hand selected operators \mathcal{P} and \mathcal{Q} need to be accessible themselves, on the other hand \mathcal{P} and \mathcal{Q} shall be sufficiently versatile to represent the desired class of operators as PDO $a(\mathcal{P}, \mathcal{Q})$ with a manageable type of symbols a(x, y). A similar dilemma is elaborated for a different PDO-like calculus of operators in [40, 41].

IV. DISCUSSION AND CLOSING REMARKS

In this paper we present umbral calculus foundations through representations of the Heisenberg group. Although this approach was around for a while [20, 27–33, 42] it largely remains out of the mainstream theory [9, 43]. The discussed viewpoint on the umbral techniques covers some additional areas, e.g. generalised translations [1–3, 5, 7]. Such inclusion rewards the umbral approach with a removal of some unessential limitations, e.g. on the degree of a delta operator. Thus, it is stimulating to consider the umbral calculus not just as a one-way road to replicate some results from the archetypal model of power series to other situations. A more fruitful umbral ideology creates a hub, which facilities all-to-all exchanges between various umbral implementations, schematically depicted as follows:



Here the contravariant transform $\mathcal{M}_A : \underline{\mathsf{E}} \to \mathsf{E}_A$ is the adjoint of the covariant transform $\mathcal{W}_A : \mathsf{E}_A \to \underline{\mathsf{E}}$. Then, the transmutation $\mathcal{T}_{BA} = \mathcal{M}_B \circ \mathcal{W}_A : \mathsf{E}_A \to \mathsf{E}_B$ intertwines respective ladder operator:

$$\mathcal{T}_{BA} \circ \mathcal{Q}_A = \mathcal{Q}_B \circ \mathcal{T}_{BA}$$
 and $\mathcal{T}_{BA} \circ \mathcal{P}_A = \mathcal{P}_B \circ \mathcal{T}_{BA}$

Such construction of transmutations falls within the *composition method* [2, Ch. 5], [3, Ch. 6].

Now we can formulate the extension of the original Umbral Principle.

Umbral Principle (Extended). For any two implementations of the umbral model intertwining operators (transmutations) $\mathcal{M}_B \circ \mathcal{W}_A$ and $\mathcal{M}_A \circ \mathcal{W}_B$ (47) allow us to exchange scopes, problems, ideas, methods, results, etc. between those implementations.

Here are few examples of such cross-fertilisation:

- The generalised translations for the Bessel operators shows that the umbral delta operator shall not be restricted by the condition that it reduces the order of polynomials by one. It will be interesting to see if the Bessel operator (1) may be useful as a delta in some combinatorial and enumeration problems.
- The generalised translations T^y naturally create respective convolutions. Yet, we can transport a technique for calculus of larger classes (not necessarily T^y -invariant) operators from the theory of PDO [19, 38, 39] generated by the archetypal umbral model from Ex. 1.

• It is possible to transport the idea of the Feynman path integral from quantum mechanics to combinatorics using the umbral framework [42]. It may be interesting to see the meaning of path integrals for generalised translations as well as other umbral implementations.

Despite of the exceptional rôle of the Heisenberg group in the analysis [10] an umbral approach may be build on other group representations as well. The next natural candidate to implement a sort of ladder actions (7)–(8) would be the group $SL_2(\mathbb{R})$. Indeed, for a given basis p_n of a linear space E there are many essentially different possibilities to chose sequences of numbers a_n , b_n , c_n such that the extended set $\{\mathcal{Q}, \mathcal{P}, \mathcal{Z}\}$ of linear operators defined by [44, 45]

$$\mathcal{Q}p_n = a_n p_{n-1}, \qquad \mathcal{P}p_n = b_n p_{n+1}, \qquad \mathcal{Z}p_n = c_n p_n$$

$$\tag{48}$$

make a representation of the Lie algebra of $SL_2(\mathbb{R})$:

$$[\mathcal{Q},\mathcal{P}] = \lambda \mathcal{Z}, \qquad [\mathcal{Z},\mathcal{Q}] = \lambda_{-}\mathcal{Q}, \qquad [\mathcal{Z},\mathcal{P}] = \lambda_{+}\mathcal{P}, \qquad \text{for some scalars} \quad \lambda, \ \lambda_{-}, \ \lambda_{+}. \tag{49}$$

A graphical illustration of action (48) is:

$$0 \underbrace{-\begin{array}{c} & & \\$$

where, in comparison to (10), the reflexive arrows for \mathbb{Z} -action are added. Such ladders include much more than just the unitary representations of $SL_2(\mathbb{R})$ by the discrete holomorphic series [44, 45]. The additional representations have various applications, e.g. for quasi-exact solvable quantum systems [46].

Interestingly, the Heisenberg ladder (10) can be a source of some $SL_2(\mathbb{R})$ representations of the form (50) through the quadratic algebra concept [47, §2.2.4]. Indeed, if $[\mathcal{P}, \mathcal{Q}] = -\iota I$ then operators $\mathcal{P}_2 = \mathcal{P}^2$, $\mathcal{Q}_2 = \mathcal{Q}^2$ and $\mathcal{Z}_2 = \mathcal{P}\mathcal{Q}$ satisfies commutators (49) and form a *metaplectic representation* of $SL_2(\mathbb{R})$ [19, 44]. Therefore, the even-numbered elements of the ladder (10) becomes nodes of the action (50) depicted as follows:



A transition in the opposite direction—from the $SL_2(\mathbb{R})$ -action (50) to its embedding in an extended ladder (51)—can be viewed as a sort of factorisation of a second-order operator. This provides a more detailed resolutions, say, for the Bessel operator \mathcal{B}_{ν} (1).

Overall, the umbral approach in the context of $SL_2(\mathbb{R})$ may be an exciting topic with an extensive exchange of ideas between various fields and deserves a further study.

 I. A. Kipriyanov. Singular Boundary Value Problems [Сингулярные Эллиптические Краевые Задачи]. Fizmatlit "Nauka", Moscow, 1997. ISBN 5-02-014799-0. In Russian. 1, 12, 13

S. Sitnik and E. Shishkina. Method of Transmutations for Differential Equations with Bessel Operators. Fizmatlit, Moscow, 2019. (in Russian). 1, 9, 12, 13

^[3] E. Shishkina and S. Sitnik. Transmutations, Singular and Fractional Differential Equations with Applications to Mathematical Physics. Mathematics in Science and Engineering. Elsevier/Academic Press, London, 2020. ISBN 978-0-12-819781-3. 1, 9, 12, 13

- [4] V. V. Kravchenko and S. M. Sitnik. Some recent developments in the transmutation operator approach. In V. V. Kravchenko and S. M. Sitnik, editors, *Transmutation Operators and Applications*, pages 3–9. Springer International Publishing, Cham, 2020. ISBN 978-3-030-35914-0. URL https://doi.org/10.1007/978-3-030-35914-0_1.1
- [5] B. M. Levitan. The application of generalized displacement operators to linear differential equations of the second order. Uspehi Matem. Nauk (N.S.), 4(1(29)):3–112, 1949. ISSN 0042-1316. 2, 13
- [6] B. M. Levitan. Expansion in Fourier series and integrals with Bessel functions. Uspehi Matem. Nauk (N.S.), 6(2(42)):102-143, 1951. ISSN 0042-1316.
- [7] В. М. Levitan. Теория операторов обобщенного сдвига. Izdat. "Nauka", Moscow, 1973. 2, 9, 13
- [8] G.-C. Rota. Finite Operator Calculus. Academic Press, Inc., New York, 1975. 2
- [9] J. P. S. Kung, G.-C. Rota, and C. H. Yan. Combinatorics: the Rota way. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2009. ISBN 978-0-521-73794-4. 2, 9, 13
- [10] R. Howe. On the role of the Heisenberg group in harmonic analysis. Bull. Amer. Math. Soc. (N.S.), 3 (2):821–843, 1980. ISSN 0002-9904. 2, 6, 14
- [11] V. V. Kisil. Möbius transformations and monogenic functional calculus. Electron. Res. Announc. Amer. Math. Soc., 2(1):26–33, 1996. ISSN 1079-6762. On-line. 2
- [12] V. V. Kisil. Spectrum as the support of functional calculus. In Functional analysis and its applications, volume 197 of North-Holland Math. Stud., pages 133–141, Amsterdam, 2004. Elsevier. arXiv:math.FA/0208249.
- [13] V. V. Kisil. Operator covariant transform and local principle. J. Phys. A: Math. Theor., 45:244022, 2012. doi:10.1088/1751-8113/45/24/244022. arXiv:1201.1749. On-line. 2
- [14] S. T. Ali, J.-P. Antoine, and J.-P. Gazeau. Coherent States, Wavelets and Their Generalizations. Graduate Texts in Contemporary Physics. Springer-Verlag, New York, 2000. ISBN 0-387-98908-0. 2, 4, 5
- [15] A. Perelomov. Generalized Coherent States and Their Applications. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1986. ISBN 3-540-15912-6. 2, 4
- [16] V. V. Kisil. Erlangen programme at large: An overview. In S. Rogosin and A. Koroleva, editors, Advances in Applied Analysis, chapter 1, pages 1–94. Birkhäuser Verlag, Basel, 2012. arXiv:1106.1686. 2, 4
- [17] H. G. Feichtinger and K. Gröchenig. A unified approach to atomic decompositions via integrable group representations. In *Function spaces and applications (Lund, 1986)*, volume 1302 of *Lecture Notes in Math.*, pages 52–73. Springer, Berlin, 1988. doi:10.1007/BFb0078863. URL https://doi.org/10.1007/BFb0078863. 2, 4
- [18] V. V. Kisil. Wavelets in Banach spaces. Acta Appl. Math., 59(1):79–109, 1999. ISSN 0167-8019. arXiv:math/9807141, On-line. 2, 4
- G. B. Folland. Harmonic Analysis in Phase Space, volume 122 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1989. ISBN 0-691-08527-7; 0-691-08528-5. 3, 4, 6, 12, 13, 14
- [20] P. Feinsilver. Lie algebras and recurrence relations. I. Acta Appl. Math., 13(3):291-333, 1988. ISSN 0167-8019. doi:10.1007/BF00046967. URL https://doi.org/10.1007/BF00046967. 7, 8, 13
- [21] V. V. Kisil. Hypercomplex representations of the Heisenberg group and mechanics. Internat. J. Theoret. Phys., 51(3):964-984, 2012. ISSN 0020-7748. doi:10.1007/s10773-011-0970-0. URL http://dx.doi.org/10.1007/s10773-011-0970-0. arXiv:1005.5057. Zbl1247.81232. 3, 4, 5
- [22] A. A. Al Ameer and V. V. Kisil. Tuning co- and contra-variant transforms: the Heisenberg group illustration. SIGMA, 18(065):21 p., 2022. arXiv:2105.13811. 4
- [23] V. V. Kisil. The real and complex techniques in harmonic analysis from the point of view of covariant transform. *Eurasian Math. J.*, 5:95–121, 2014. arXiv:1209.5072. On-line. 4
- [24] V. V. Kisil. Wavelets beyond admissibility. In M. Ruzhansky and J. Wirth, editors, Progress in Analysis and its Applications, pages 219–225. World Sci. Publ., Hackensack, NJ, 2010. URL http://dx.doi.org/10.1142/9789814313179_0029. arXiv:0911.4701. Zbl1269.30052. 4
- [25] V. V. Kisil. Calculus of operators: Covariant transform and relative convolutions. Banach J. Math. Anal., 8(2):156-184, 2014. URL http://www.emis.de/journals/BJMA/tex_v8_n2_a15.pdf. arXiv:1304.2792, on-line. 5, 12
- [26] J. P. Kung, editor. Gian-Carlo Rota on Combinatorics: Introductory Papers and Commentaries, volume 1 of Contemporary Mathematicians. Birkhäuser Verlag, Boston, 1995. 8, 9, 16

- [27] J. Cigler. Some remarks on Rota's umbral calculus. Nederl. Akad. Wetensch. Proc. Ser. A, 81:27–42, 1978. 8, 13
- [28] V. V. Kisil. Umbral calculus and cancellative semigroup algebras. Z. Anal. Anwendungen, 19(2): 315–338, 2000. ISSN 0232-2064. arXiv:funct-an/9704001. Zbl0959.43004.
- [29] V. V. Kisil. Tokens: an algebraic construction common in combinatorics, analysis, and physics. In Ukrainian Mathematics Congress-2001 (Ukrainian), pages 146–155. Natsīonal. Akad. Nauk Ukraïni Inst. Mat., Kiev, 2002. arXiv:math.FA/0201012.
- [30] A. K. Kwaśniewski. Main theorems of extended finite operator calculus. Integral Transforms Spec. Funct., 14(6):499–516, 2003. ISSN 1065-2469. doi:10.1080/10652460290029743. URL https://doi.org/10.1080/10652460290029743.
- [31] D. Levi, P. Tempesta, and P. Winternitz. Umbral calculus, difference equations and the discrete Schrödinger equation. J. Math. Phys., 45(11):4077–4105, 2004. ISSN 0022-2488. doi:10.1063/1.1780612. URL https://doi.org/10.1063/1.1780612.
- [32] G. Dattoli, D. Levi, and P. Winternitz. Heisenberg algebra, umbral calculus and orthogonal polynomials. J. Math. Phys., 49(5):053509, 19, 2008. ISSN 0022-2488. doi:10.1063/1.2909731. URL https://doi.org/10.1063/1.2909731. 10
- [33] N. Faustino and G. Ren. (Discrete) Almansi type decompositions: An umbral calculus framework based on osp(1|2) symmetries. Math. Methods Appl. Sci., 34(16):1961-1979, 2011. ISSN 0170-4214. doi:10.1002/mma.1498. URL https://doi.org/10.1002/mma.1498. 8, 13
- [34] R. Mullin and G.-C. Rota. On the foundation of combinatorial theory (III): Theory of binomial enumeration. In B.Harris, editor, *Graph Theory and Its Applications*, pages 167–213. Academic Press, Inc., New York, 1970. Reprinted in [26, pp. 118–147]. 8
- [35] G. G. Bilodeau. The Weierstrass transform and Hermite polynomials. Duke Math. J., 29:293-308, 1962. ISSN 0012-7094. URL http://projecteuclid.org/euclid.dmj/1077470134. 11
- [36] I. M. Gel' fand and B. M. Levitan. On the determination of a differential equation from its spectral function. *Izvestiya Akad. Nauk SSSR. Ser. Mat.*, 15:309–360, 1951. ISSN 0373-2436. 11
- [37] V. V. Kisil. Relative convolutions. I. Properties and applications. Adv. Math., 147(1):35–73, 1999.
 ISSN 0001-8708. arXiv:funct-an/9410001, On-line. Zbl933.43004. 12
- [38] R. Howe. Quantum mechanics and partial differential equations. J. Funct. Anal., 38(2):188–254, 1980. ISSN 0022-1236. 12, 13
- [39] V. V. Kisil. Cross-Toeplitz operators on the Fock-Segal-Bargmann space and two-sided convolutions on the Heisenberg group. Ann. Funct. Anal., 14(38), 2023. to appear. arXiv:2108.13710. doi: 10.1007/s43034-022-00249-7. 5, 12, 13
- [40] J. Delgado, M. Ruzhansky, and N. Tokmagambetov. Schatten classes, nuclearity and nonharmonic analysis on compact manifolds with boundary. J. Math. Pures Appl. (9), 107(6):758-783, 2017. ISSN 0021-7824. doi:10.1016/j.matpur.2016.10.005. URL https://doi.org/10.1016/j.matpur.2016.10.005. 13
- [41] D. Cardona, V. Kumar, M. Ruzhansky, and N. Tokmagambetov. Global functional calculus, lower/upper bounds and evolution equations on manifolds with boundary, 2021. 13
- [42] V. V. Kisil. Polynomial sequences of binomial type and path integrals. Ann. Comb., 6(1):45–56, 2002.
 ISSN 0218-0006. arXiv:math/9808040 Zbl1009.05013. 13, 14
- [43] R. P. Stanley. *Enumerative Combinatorics. Vol. 1.* Cambridge University Press, Cambridge, 1997.
 ISBN 0-521-55309-1. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original. 13
- [44] R. Howe and E.-C. Tan. Nonabelian Harmonic Analysis. Applications of SL(2, R). Springer-Verlag, New York, 1992. ISBN 0-387-97768-6. 14
- [45] V. V. Kisil. Erlangen Programme at Large 3.2: Ladder operators in hypercomplex mechanics. Acta Polytechnica, 51(4):44–53, 2011. on-line. arXiv:1103.1120. 14
- [46] A. V. Turbiner. Quasi-exactly-solvable problems and sl(2) algebra. Comm. Math. Phys., 118(3):467-474, 1988. ISSN 0010-3616. URL http://projecteuclid.org/euclid.cmp/1104162094. 14
- [47] J.-P. Gazeau. Coherent States in Quantum Physics. Wiley-VCH Verlag, 2009. ISBN 9783527407095. doi:10.1002/9783527628285. 14