



UNIVERSITY OF LEEDS

This is a repository copy of *A Frobenius splitting and cohomology vanishing for the cotangent bundles of the flag varieties of GL_n* .

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/226375/>

Version: Accepted Version

Article:

Tange, R. orcid.org/0000-0003-0867-1573 (Accepted: 2025) A Frobenius splitting and cohomology vanishing for the cotangent bundles of the flag varieties of GL_n . Journal of Algebra. ISSN 0021-8693 (In Press)

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

A FROBENIUS SPLITTING AND COHOMOLOGY VANISHING FOR THE COTANGENT BUNDLES OF THE FLAG VARIETIES OF GL_n

RUDOLF TANGE

ABSTRACT. Let k be an algebraically closed field of characteristic $p > 0$, let $G = GL_n$ be the general linear group over k , let P be a parabolic subgroup of G , and let \mathfrak{u}_P be the Lie algebra of its unipotent radical. We show that the Kumar-Lauritzen-Thomsen splitting of the cotangent bundle $G \times^P \mathfrak{u}_P$ of G/P has top degree $(p-1)\dim(G/P)$. The component of that degree is therefore given by the $(p-1)$ -th power of a function f . We give a formula for f and deduce that it vanishes on the exceptional locus of the resolution $G \times^P \mathfrak{u}_P \rightarrow \overline{O}$ where \overline{O} is the closure of the Richardson orbit of P . As a consequence we obtain that the higher cohomology groups of a line bundle on $G \times^P \mathfrak{u}_P$ associated to a dominant weight are zero. The splitting of $G \times^P \mathfrak{u}_P$ given by f^{p-1} can be seen as a generalisation of the Mehta-Vander Kallen splitting of $G \times^B \mathfrak{u}$.

INTRODUCTION

Let G be a reductive group over an algebraically closed field k of positive characteristic p . For a parabolic P containing the positive Borel and P -module M , we denote by $H^i(G/P, M)$ the i -th cohomology group of the sheaf $\mathcal{L}_{G/P}(M)$ on G/P associated to M . It is an open problem whether we have for all parabolic subgroups P and all dominant characters λ of P that

$$H^i(G/P, S(\mathfrak{u}_P^*) \otimes k_{-\lambda}) = 0 \quad \text{for all } i > 0, \quad (*)$$

where the most important case is $\lambda = 0$, see e.g. [2, Introduction to Ch 5]. In characteristic 0 this is an easy consequence of the Grauert-Riemenschneider Theorem, see [4, Thm 2.2]. In characteristic p (*) is known for $P = B$, for arbitrary P and “ P -regular” dominant λ , see [12], and for P corresponding to sets of pairwise orthogonal short simple roots and $\lambda = 0$, see [16].¹

It is easy to write a formula for the Euler character

$$\sum_{i \geq 0} (-1)^i \operatorname{ch} H^i(G/P, S(\mathfrak{u}_P^*) \otimes k_{-\lambda}),$$

see [10, Sect 8.14-8.16] and [3, Prop 2.1], so if (*) holds we get a formula for $\operatorname{ch} H^0(G/P, S(\mathfrak{u}^*) \otimes k_{-\lambda})$.

For computing cohomology of Frobenius kernels of G , (*), or a special case of it, is often used, see [9, II.12.12-15], [12, Thm 8], [1, Sect 7] and [13, Sect 7].

2020 *Mathematics Subject Classification*. 14F17, 14M15, 14L30.

Key words and phrases. cohomology, cotangent bundle, flag variety, Frobenius splitting.

¹The case $P = B$ and $\lambda = 0$ is [9, Lem II.12.12].

When $\mathcal{L}_{G/P}(\lambda) = \mathcal{L}_{G/P}(k_{-\lambda})$ is ample, i.e. λ “ P -regular” dominant, one gets (*) from the fact that $G \times^P \mathfrak{u}_P$ is Frobenius split. One can also use Frobenius splittings to prove (*) for $\lambda = 0$ via a characteristic p -version of the Grauert-Riemenschneider Theorem [14, Thm 1.2], since the canonical bundle of $G \times^P \mathfrak{u}_P$ is trivial. But then the map from $G \times^P \mathfrak{u}_P$ to the Richardson orbit closure has to be birational and the splitting has to be a $(p-1)$ -th power of a section σ of the anti-canonical bundle which vanishes on the exceptional locus. This is the approach we will follow.

When I asked Thomsen about the case $G = \mathrm{GL}_n$, he told me he expected that the pushforward to $G \times^P \mathfrak{u}_P$ of the splitting of $G \times^B \mathfrak{u}_P$ induced by the “MVdK-splitting” of $G \times^B \mathfrak{u}$ from [15] is the homogeneous component of degree $(p-1)\dim(G/P)$ of the “KLT-splitting”, see Section 1.2, from [12]. Although we can not prove this conjecture, we can show that the above component is in fact the top degree component and therefore a $(p-1)$ -th power. From this we can then deduce that this homogeneous splitting vanishes on the exceptional locus of the resolution $\varphi : [g, X] \mapsto gXg^{-1} : G \times^P \mathfrak{u}_P \rightarrow \overline{\mathcal{O}}$, where $\overline{\mathcal{O}}$ is the closure of the Richardson orbit corresponding to P , see Theorem 1 in Section 2. Finally, we then deduce that (*) holds in type A , see Theorem 2. In fact we can formulate this as a result for arbitrary reductive groups.

The main idea of the proof is as follows. The “KLT-splitting” from [12] is the $(p-1)$ -th power of the pullback along φ of the function which maps an $n \times n$ matrix X to

$$\prod_{i=1}^{n-1} \det((I_n + X)_{\leq i, \leq i}), \quad (1)$$

where $Y_{\leq i, \leq i}$ denotes the submatrix of Y given by the first i rows and columns, see [2, Example 5.1.15].² Unlike in the case $P = B$, the degree of the i -th factor may be less than i . In Lemma’s 1(ii) and 3 we determine the degree of the i -th factor and from that it follows that the product (1) has degree $\dim(G/P)$.

1. PRELIMINARIES

1.1. Notation. Let k be an algebraically closed field of characteristic $p > 0$ and let G be a reductive group over k . We fix a Borel subgroup $B \leq G$ and maximal torus $T \leq B$. We denote by R the set of roots of T in the Lie algebra $\mathfrak{g} = \mathrm{Lie}(G)$ of G , and we denote the unipotent radical of B by U . We call the roots of T in $\mathfrak{u} = \mathrm{Lie}(U)$ positive and we denote the corresponding set of simple roots by S . For a subset I of S we denote the root system spanned by I by R_I . Furthermore, we denote the corresponding parabolic subgroup containing B and its Levi subgroup containing T by P_I and L_I . Denote the character group of an algebraic group H by $X(H)$. For $I \subseteq S$ we identify $X(P_I)$ and $X(L_I)$ with $\{\lambda \in X(T) \mid \langle \lambda, \alpha^\vee \rangle = 0 \text{ for all } \alpha \in I\}$.

For P a parabolic of G and M a P -module we write $\mathcal{L}(M)$ for the G -linearised sheaf on G/P associated to M . For $\lambda \in X(P) \leq X(T)$ we put $\mathcal{L}(\lambda) = \mathcal{L}(k_{-\lambda})$, it is the sheaf of sections of the line bundle $G \times^P k_{-\lambda}$ on G/P . We use the same

²Apart from the degree computation, the arguments there work for any parabolic.

symbol $\mathcal{L}(\lambda)$ to denote the sheaf of sections of the pullback of this line bundle to $G \times^P V$ for any P -variety V . We also write $H^i(G/P, M)$ for

$$H^i(G/P, \mathcal{L}(M)) \simeq R^i \mathrm{ind}_P^G(M),$$

see [9, I.5.12]. We have that

$$H^i(G \times^P \mathfrak{u}_P, \mathcal{L}(\lambda)) = H^i(G/P, k[\mathfrak{u}_P] \otimes k_{-\lambda}),$$

see [2, Lem 5.2.2].

If $p = \mathrm{char} k$ is good for G , then we have $\mathfrak{g}/\mathfrak{p} \simeq \mathfrak{u}_P^*$ as P -modules and $G \times^P \mathfrak{u}_P$ is the cotangent bundle $T^\vee(G/P)$ of G/P , see [2, 5.1.8-11].

1.2. Frobenius splittings. By [2, Lem 5.1.1] the canonical bundle of $G \times^P \mathfrak{u}_P$ is trivial, so we can choose a nowhere zero global section: a volume form. It is easy to see that such a section is unique up to a scalar multiple, see [2, 5.1.2]. This means that we can think of Frobenius splittings (up to a scalar multiple) of $G \times^P \mathfrak{u}_P$ as certain regular functions on $G \times^P \mathfrak{u}_P$.

In [12, Thm 1] it was proved that, when p is good for G , the cotangent bundle $T^\vee(G/P)$ of G/P is Frobenius split, see also [2, Thm 5.1.3]. We will refer to the B -canonical splitting $\psi_P(f_- \otimes f_+)$ as the “KLT-splitting” of $T^\vee(G/P)$, where ψ_P, f_-, f_+ are as defined in [2, Ch 5]. Actually this is only a splitting up to a scalar multiple, but in the case $G = \mathrm{GL}_n$ we assume that the chosen volume form on $T^\vee(G/P)$ is such that the pullback along φ of the function given by (1), φ defined as in the introduction, defines a splitting. That formula is all we need to know about the KLT-splitting in this paper.

The standard grading of $k[\mathfrak{u}_P] = S(\mathfrak{u}_P^*)$ gives a grading on $k[G \times^P \mathfrak{u}_P]$, and in [2, 5.1.14] it is explained that the homogeneous component of degree $(p-1)\dim(G/P)$ of a splitting σ of $G \times^P \mathfrak{u}_P$ is again a splitting of $G \times^P \mathfrak{u}_P$. This component is B -canonical if σ is B -canonical.

1.3. A result on cohomology vanishing. The following result may be well-known, but for lack of reference we give a proof.

Proposition 1. *Assume p is good for G , let P be a parabolic of G , let $\lambda \in X(P)$ be dominant, let Q be the parabolic of G containing P such that $\lambda \in X(Q)$ and $\mathcal{L}_{G/Q}(\lambda)$ is ample, and let L be the Levi subgroup of Q containing T . If $H^i(L/L \cap P, S(\mathfrak{l}/\mathfrak{l} \cap \mathfrak{p})) = 0$ for all $i > 0$, then $H^i(G/P, S(\mathfrak{g}/\mathfrak{p}) \otimes k_{-\lambda}) = 0$ for all $i > 0$.*

Proof. By [12, Cor 3 to Thm 4] or [2, Thm 5.3] $G \times^P \mathfrak{u}_P$ is Frobenius split, so by [2, Lemma 1.2.7(i)] it is enough to show the vanishing for $m\lambda$, $m \gg 0$ (in fact we only need it for $p^m\lambda$ and some $m \geq 0$).

Some of the arguments below are adaptations of arguments from the proof of [2, Lem 5.2.7].

Each $S^j(\mathfrak{g}/\mathfrak{p})$ has a filtration with sections $S^r(\mathfrak{q}/\mathfrak{p}) \otimes S^s(\mathfrak{g}/\mathfrak{q})$, $r + s = j$, so it is enough to show that $R^i \mathrm{ind}_P^G(S(\mathfrak{q}/\mathfrak{p}) \otimes S(\mathfrak{g}/\mathfrak{q}) \otimes k_{-\lambda}) = 0$ for all $i > 0$. We have $P = (L \cap P)U_Q$ and $\mathfrak{q}/\mathfrak{p} \simeq \mathfrak{l}/\mathfrak{l} \cap \mathfrak{p}$. Note that U_Q acts trivially on $\mathfrak{q}/\mathfrak{p}$. Combining [9, I.6.11] and our assumption with a standard spectral sequence

argument, we have

$$R^i \text{ind}_P^G(S(\mathfrak{q}/\mathfrak{p}) \otimes S(\mathfrak{g}/\mathfrak{q}) \otimes k_{-\lambda}) \simeq R^i \text{ind}_Q^G(\text{ind}_P^Q S(\mathfrak{q}/\mathfrak{p}) \otimes S(\mathfrak{g}/\mathfrak{q}) \otimes k_{-\lambda}). \quad (2)$$

We have $\mathfrak{p} = \mathfrak{l} \cap \mathfrak{p} \oplus \mathfrak{u}_Q$, $\mathfrak{u}_P = \mathfrak{u}_{L \cap P} \oplus \mathfrak{u}_Q$, $(\mathfrak{g}/\mathfrak{p})^* \simeq \mathfrak{u}_P$, $(\mathfrak{g}/\mathfrak{q})^* \simeq \mathfrak{u}_Q$, and $(\mathfrak{q}/\mathfrak{p})^* \simeq \mathfrak{u}_{L \cap P}$. By the arguments of [10, p94] there exists an affine Q -variety V_0 such that $k[V_0] \simeq k[Q \times^P \mathfrak{u}_{L \cap P}] = \text{ind}_P^Q S(\mathfrak{q}/\mathfrak{p})$, Q -equivariantly (U_Q acting trivially). Put $V = V_0 \times \mathfrak{u}_Q$. Then $k[V] = \text{ind}_P^Q S(\mathfrak{q}/\mathfrak{p}) \otimes S(\mathfrak{g}/\mathfrak{q})$. Now the morphism $G \times^Q V \rightarrow G/Q$ is affine, so by [7, Ex III.8.2] the RHS of (2) is isomorphic to

$$H^i(G \times^Q V, \mathcal{L}(\lambda)). \quad (3)$$

By [6, 5.1.12] $\mathcal{L}(\lambda)$ is ample on $G \times^Q V$, since $G \times^Q V \rightarrow G/Q$ is affine. The morphism $V_0 \rightarrow \overline{Q \cdot \mathfrak{u}_{L \cap P}}$ is finite, see [10, p94], so the same is true for the morphisms $V \rightarrow \overline{Q \cdot \mathfrak{u}_{L \cap P}} \times \mathfrak{u}$ and $G \times^Q V \rightarrow G \times^Q (\overline{Q \cdot \mathfrak{u}_{L \cap P}} \times \mathfrak{u}_Q)$. So composing the latter with the projective morphism $G \times^Q (\overline{Q \cdot \mathfrak{u}_{L \cap P}} \times \mathfrak{u}_Q) \rightarrow \mathfrak{g}$, given by the embedding of $\overline{Q \cdot \mathfrak{u}_{L \cap P}} \times \mathfrak{u}_Q$ in \mathfrak{g} , we obtain a proper morphism $G \times^Q V \rightarrow \mathfrak{g}$. Now [7, III.5.3] tells us that (3) is 0 if we replace λ by $m\lambda$, $m \gg 0$. \square

2. THE MAIN RESULTS

Throughout this section, except in Theorem 2 and its proof, $G = \text{GL}_n = \text{GL}(k^n)$ and T is the subgroup of diagonal matrices in G . As simple roots we choose the usual characters $\varepsilon_i - \varepsilon_{i+1}$, $1 \leq i \leq n-1$, where we used additive notation for characters, and ε_i is the i -th coordinate function on T . Then B consists of the upper triangular matrices in G . As is well-known, the conjugacy classes of parabolic subgroups of G are labelled by the compositions of n , see e.g. [8, 3.2]. By ν we denote a composition of n and $P = P_\nu \supseteq B$ is the standard parabolic whose block sizes are given in order by ν . If A_ν is the set $\{\nu_1, \nu_1 + \nu_2, \dots, \sum_{j=1}^{s-1} \nu_j\}$, s the length of ν , then $P_\nu = P_{I_\nu}$, the parabolic associated to the set of simple roots $I_\nu = \{\varepsilon_i - \varepsilon_{i+1} \mid i \in \{1, \dots, n-1\} \setminus A_\nu\}$. We denote by λ the transposed partition of the weakly descending sorted version of ν . It is well-known that the Richardson orbit of P_ν is \mathcal{O}_λ , the nilpotent orbit whose Jordan block sizes are given by λ , see e.g. [8, Thm 3.3(a)].

It is well-known that the map $\varphi : [g, X] \mapsto gXg^{-1} : G \times^P \mathfrak{u}_P \rightarrow \overline{\mathcal{O}_\lambda}$ is birational. Indeed the group centraliser G_X of any $X \in \mathfrak{g}$ is the set of invertible elements in the Lie algebra centraliser \mathfrak{g}_X , so is connected. Now see [10, 4.9 and 8.8 Remark]. It is also well-known that $\overline{\mathcal{O}_\lambda}$ is normal, see e.g. [5] or [15, Sect 4.7].

For $i \in \{1, \dots, n-1\}$ we denote by $d_{\lambda,i}$ the number of nonzero positions on the $(n-i)$ -th upper codiagonal of \mathfrak{u}_P . So for $\nu = (2, 1, 2)$ we have $d_{\lambda,1}, d_{\lambda,2}, d_{\lambda,3}, d_{\lambda,4} = 1, 2, 3, 2$, see the figure of \mathfrak{u}_p below.

$$\begin{bmatrix} 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Since each diagonal $j \times j$ block of P takes away $j - i$ nonzero positions from the i -th upper codiagonal, we have, if j occurs m_j times in ν ,

$$d_{\lambda, n-i} = n - i - \sum_{j>i} (j - i)m_j = n - i - \sum_{j>i} \lambda_j = - \sum_{j=i+1}^n (\lambda_j - 1) = \sum_{j=1}^i (\lambda_j - 1).$$

Therefore, $d_{\lambda, i} = i - \sum_{j>n-i} \lambda_j = \sum_{j=1}^{n-i} (\lambda_j - 1)$. So indeed the $d_{\lambda, i}$ only depend on λ , moreover, they determine λ .

For a square matrix X we denote by $X_{\leq i, \leq i}$ the submatrix of X given by the first i rows and columns. For an $i \times i$ matrix Y we denote by $s_j(Y)$ the trace of the j -th exterior power of Y , i.e. the sum of the diagonal $j \times j$ minors of Y . As is well-known, $\det(aI_i - Y) = a^i + \sum_{j=1}^i (-1)^j a^{i-j} s_j(Y)$, where I_i is the $i \times i$ identity matrix. So the largest j with $s_j(Y) \neq 0$ is the number of nonzero eigenvalues of Y , counted with (algebraic) multiplicity. This number also equals the rank of Y^l for l sufficiently big. We will call it the *stable rank* of Y .

Lemma 1. *Let $X \in \mathcal{O}_\lambda$.*

- (i) *Any i -dimensional subspace W of $V = k^n$ contains an X -invariant subspace U of dimension $\geq \sum_{j>n-i} \lambda_j$.*
- (ii) *$X_{\leq i, \leq i}$ has stable rank $\leq d_{\lambda, i}$.*

Proof. (i). We show this by induction on n . It is trivial when $i \leq n - r$, r the length of λ , in particular when $n = 0$. Assume $i > n - r$. Then W has nonzero intersection with $\mathrm{Ker}(X)$ for dimension reasons. Pick v nonzero in that intersection. First note that the transformation \bar{X} induced on V/kv by X has partition μ which is obtained from λ by subtracting 1 from one part of λ and then sorting the result in weakly descending order. Indeed if we decompose V as a direct sum of X -Jordan blocks and we pick a X -Jordan block of minimal size with the property that v has nonzero component in it, then we can replace that X -Jordan block by an X -Jordan block of the same size which contains v . Now we apply the induction hypothesis to V/kv and W/kv , noting that $(n - 1) - (i - 1) = n - i$, to obtain an \bar{X} -invariant subspace U/kv of W/kv of dimension $\geq \sum_{j>n-i} \mu_j \geq \sum_{j>n-i} \lambda_j - 1$. Now U is the X -invariant subspace we want.

(ii). The linear map $(X_{\leq i, \leq i})^i$ coincides with X^i on any X -invariant subspace U of $k^i \leq k^n$ and therefore kills it. Choosing U as in (i), it induces a linear map $k^i/U \rightarrow k^i$ and therefore has rank $\leq i - \sum_{j>n-i} \lambda_j = d_{\lambda, i}$. \square

Lemma 3 below follows from Lemma 1(ii) and the existence of the KLT-splitting, but we prefer to give a direct proof.

Lemma 2. *For any $h \in \{1, \dots, i - 1\}$ there exists a regular nilpotent $i \times i$ matrix X such that $X_{\leq h, \leq h}$ is invertible.*

Proof. Let (e_1, \dots, e_i) be the standard basis of k^i . Then the regular nilpotent matrix X given by $X(e_j) = e_{j-1}$ for $2 \leq j \leq i$, $X(e_2) = e_1 + e_{h+1}$ and $X(e_1 + e_{h+1}) = 0$ has the desired property. \square

Remark 2.1. Of course it follows from Lemma 2 that there exists a regular $i \times i$ matrix X such that $X_{\leq h, \leq h}$ is invertible for all $h \in \{1, \dots, i-1\}$, but we won't need this.

Lemma 3. *There exists $X \in \mathcal{O}_\lambda$ such that $X_{\leq i, \leq i}$ has stable rank $d_{\lambda, i}$.*

Proof. First choose any $Y \in \mathfrak{g}$ nilpotent with partition λ and decompose k^n into Y -Jordan blocks with sizes $\lambda_1, \lambda_2, \dots, \lambda_r$, where r is the length of λ . It suffices to find an ordered basis \mathcal{B} of k^n such that the upper left $i \times i$ -block Z of the matrix of Y relative to this basis has stable rank $d_{\lambda, i}$.

Determine $s \leq r$ maximal with $\sum_{j=1}^s (\lambda_j - 1) \leq i$ and put $h = i - \sum_{j=1}^s (\lambda_j - 1)$. Using Lemma 2 choose for each $j \leq s$ a basis of the j -th block such that the upper left $(\lambda_j - 1) \times (\lambda_j - 1)$ block of the matrix of Y relative to this basis is invertible, if $s < r$ and $h > 0$ choose a basis of the $(s+1)$ -th block such that the upper left $h \times h$ block of the matrix of Y relative to this basis is invertible, and for the remaining blocks choose any basis.

We now form \mathcal{B} as follows. First consider the case $i \leq n - r$. For each $j \leq s$ we pick the first $\lambda_j - 1$ basis vectors from the j -th block, if $s < r$ we append the first h basis vectors from the $(s+1)$ -th block, and finally we append all remaining $n - i$ basis vectors. Now Z is in block diagonal form with invertible diagonal block of sizes $\lambda_1 - 1, \dots, \lambda_s - 1, h$, where h has to be omitted if $h = 0$. Now consider the case $i > n - r$. For each $j \leq n - i$ we pick the first $\lambda_j - 1$ basis vectors from the j -th block, then we append the basis vectors from the next $r - (n - i)$ blocks, and finally we append all remaining $n - i$ basis vectors. Now Z is in block diagonal form with diagonal block sizes $\lambda_1 - 1, \dots, \lambda_{n-i} - 1, \lambda_{n-i+1}, \dots, \lambda_r$ where the first $n - i$ blocks are invertible, and the others nilpotent. In both cases we obtain that Z has stable rank $d_{\lambda, i}$ (when $i \leq n - r$ we have $d_{\lambda, i} = i$). \square

Below we will denote a function $X \mapsto E(X)$ on a closed subvariety of \mathfrak{g} just by the expression $E(X)$.

Theorem 1. *The degree $(p-1) \dim(G/P)$ component of the KLT splitting of $G \times^P \mathfrak{u}_P$ is the top degree component and equals the $(p-1)$ -th power of the pullback of $\prod_{i=1}^{n-1} s_{d_{\lambda, i}}(X_{\leq i, \leq i}) \in k[\overline{\mathcal{O}_\lambda}]$ along the resolution $\varphi : G \times^P \mathfrak{u}_P \rightarrow \overline{\mathcal{O}_\lambda}$. This pullback vanishes on the exceptional locus of φ .*

Proof. The KLT splitting is the pullback along φ of the function given by (1). Furthermore, we have $\det(I_i + Y) = \sum_{j=0}^i s_j(Y)$ for any $i \times i$ matrix Y , and, of course, $s_j(X_{\leq i, \leq i}) \neq 0$ on $\mathcal{O}_\lambda \iff s_j(X_{\leq i, \leq i}) \neq 0$ on $\overline{\mathcal{O}_\lambda}$. So by Lemma's 1(ii) and 3 the top degree component of the i -th factor in (1) is $s_{d_{\lambda, i}}(X_{\leq i, \leq i})$. So the KLT-splitting has top degree $p-1$ times $\sum_{i=1}^{n-1} d_{\lambda, i} = \dim \mathfrak{u}_P = \dim(G/P)$, and the top degree component is the $(p-1)$ -th power of the pullback along φ of the function given by the stated formula.

To prove the second assertion, put $f_{\lambda, i}(X) = s_{d_{\lambda, i}}(X_{\leq i, \leq i})$ and $f_\lambda = \prod_{i=1}^{n-1} f_{\lambda, i}$. The exceptional locus is $\varphi^{-1}(\overline{\mathcal{O}_\lambda} \setminus \mathcal{O}_\lambda)$, so it suffices to show that f_λ vanishes on any $\mathcal{O}_\mu \subseteq \overline{\mathcal{O}_\lambda} \setminus \mathcal{O}_\lambda$. We have $\dim(\mathfrak{u}_Q) = \frac{1}{2} \dim(\mathcal{O}_\mu) < \frac{1}{2} \dim(\mathcal{O}_\lambda) = \dim(\mathfrak{u}_P)$, where Q is a standard parabolic whose Richardson orbit is \mathcal{O}_μ , see [10, 4.9].

So for some i we must have $d_{\mu,i} < d_{\lambda,i}$ which means that $f_{\lambda,i}$ and therefore f_λ vanishes on \mathcal{O}_μ . \square

Theorem 2. *Let G be any reductive group for which p is good, let $\lambda \in X(T)$ be dominant, put $I = \{\alpha \in S \mid \langle \lambda, \alpha^\vee \rangle = 0\}$. Then $H^i(T^\vee(G/P_J), \mathcal{L}(\lambda)) = 0$ for all $J \subseteq I$ such that R_J contains all irreducible components of R_I not of type A .*

Proof. By Proposition 1 we may assume that $\lambda = 0$ and that all irreducible components of R have type A . Since we are dealing with cotangent bundles we may assume that G is semisimple and simply connected. By the Künneth formula [11, Prop 9.2.4] we may then assume $G = \mathrm{SL}_n$ and finally we may assume $G = \mathrm{GL}_n$. Now the result follows from Theorem 1 and [14, Thm 1.2], bearing in mind that the canonical bundle of $T^\vee(G/P)$ is trivial, see [2, Lem 5.1.1], and that $R^i\varphi_*(\mathcal{O}_{T^\vee(G/P)})$ is the sheaf associated with the cohomology group $H^i(T^\vee(G/P), \mathcal{O}_{T^\vee(G/P)})$, since φ is affine. \square

We remind the reader that a proper birational morphism $\psi : X \rightarrow Y$ is called a *rational resolution* if $\psi_*\mathcal{O}_X = \mathcal{O}_Y$ and the higher direct images of \mathcal{O}_X and ω_X are 0, see [2, Def 3.4.1]. We assume again that $G = \mathrm{GL}_n$.

Corollary. *The resolution $\varphi : G \times^P \mathfrak{u}_P \rightarrow \overline{\mathcal{O}_\lambda}$ is rational.*

Proof. This follows from a standard argument, see e.g. [9, Lem 14.5], and Theorem 2. \square

Remarks 2.2. 1. If $P = B$, then $d_{(n),i} = i$ for all i , so the splitting from Theorem 1 equals the $(p-1)$ -th power of the pullback of $\prod_{i=1}^{n-1} \det(X_{\leq i, \leq i})$ along $\varphi : G \times^B \mathfrak{u} \rightarrow \mathcal{N}$. This is the MVdK splitting of $G \times^B \mathfrak{u}$, see [15].

2. Thomsen mentioned to me another proof of Lemma 1(ii): One can easily deduce it from the following result which can be proved by induction on n . For $X \in \mathfrak{u}_P$ let $y_{ij} = \delta_{ij} + x_{ij}$ be the (i, j) -th entry of $I_n + X$. Then any monomial $y_{i_1 j_1} y_{i_2 j_2} \cdots y_{i_s j_s}$ with the i_l all distinct and the j_l all distinct has degree $\leq d_{\lambda,s}$ in the x_{ij} .

3. In [15, Sect 4.9] there is also a proof of the above corollary for certain parabolics, but that relies on the existence of a principal effective divisor D which is a subdivisor of (σ) (σ^{p-1} is the MVdK splitting) and contains the exceptional locus. This is claimed in [15, Prop 4.5], but the proof of that result is incomplete and it seems rather unlikely that such a divisor exists for the set of parabolics in question. The proof of the above corollary sketched in [2, Exercise 5.3.E(b)] is also problematic: after pushing the splitting of Exercise 5.1.E.6 forward from $G \times^B \mathfrak{u}_P$ to $G \times^P \mathfrak{u}_P$ it's no longer clear that the splitting is a $(p-1)$ -th power, so one can't apply [2, Thm 1.3.14].

Conjecture (Thomsen). *The pushforward to $G \times^P \mathfrak{u}_P$ of the splitting of $G \times^B \mathfrak{u}_P$ induced by the MVdK splitting is the top degree component of the KLT splitting.*

ACKNOWLEDGEMENT

I would like to thank Jesper Funch Thomsen for helpful discussions.

REFERENCES

- [1] C. Bendel, D. Nakano, B. Parshall, C. Pillen, *Cohomology for quantum groups via the geometry of the nullcone*, Mem. Amer. Math. Soc. **229** (2014), no. 1077.
- [2] M. Brion, S. Kumar, *Frobenius splitting methods in geometry and representation theory*, Progress in Mathematics, 231, Birkhäuser Boston, Inc., Boston, MA, 2005.
- [3] A. Broer, *Line bundles on the cotangent bundle of the flag variety*, Invent. Math. **113** (1993), no. 1, 1-20.
- [4] A. Broer, *Normality of some nilpotent varieties and cohomology of line bundles on the cotangent bundle of the flag variety*, in “Lie Theory and Geometry”, Prog. Math. **123** (1994), Birkhäuser, Boston, 1-19.
- [5] S. Donkin, *The normality of closures of conjugacy classes of matrices*, Invent. Math. **101** (1990), no. 3, 717-736.
- [6] A. Grothendieck, *Eléments de géométrie algébrique II, Étude globale élémentaire de quelques classes de morphismes*, Inst. Hautes Études Sci. Publ. Math. No. 8, 1961.
- [7] R. Hartshorne, *Algebraic Geometry*, GTM **52**, Springer-Verlag, New York-Heidelberg, 1977.
- [8] W. H. Hesselink, *Polarizations in the classical groups*, Math. Z. **160** (1978), no. 3, 217-234.
- [9] J. C. Jantzen, *Representations of algebraic groups*, Second edition, American Mathematical Society, Providence, RI, 2003.
- [10] J. C. Jantzen, *Nilpotent orbits in representation theory*, “Lie theory”, Progr. Math. **228**, Birkhäuser Boston, 2004, 1-211.
- [11] G. R. Kempf, *Algebraic Varieties*, London Math. Soc. Lecture Note Ser. 172, Cambridge University Press, Cambridge, 1993.
- [12] S. Kumar, N. Lauritzen, J. F. Thomsen, *Frobenius splitting of cotangent bundles of flag varieties*, Invent. Math. **136** (1999), no.3, 603-621.
- [13] Z. Lin, D. Nakano, *Realizing Rings of Regular Functions via the Cohomology of Quantum Groups*, Preprint.
- [14] V. B. Mehta, W. van der Kallen, *On a Grauert-Riemenschneider vanishing theorem for Frobenius split varieties in characteristic p* , Invent. Math. **108** (1992), no. 1, 11-13.
- [15] V. B. Mehta, W. van der Kallen, *A simultaneous Frobenius splitting for closures of conjugacy classes of nilpotent matrices*, Compositio Math. **84** (1992), no. 2, 211-221.
- [16] J. F. Thomsen, *Normality of certain nilpotent varieties in positive characteristic*, J. Algebra **227** (2000), no. 2, 595-613.

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LS2 9JT, LEEDS, UK

Email address: R.H.Tange@leeds.ac.uk